

A comment on the OPERA result and CPT

Benjamin Koch

*Pontificia Universidad Católica de Chile.
Av. Vicuña Mackenna 4860. Macul.
Santiago de Chile.**

(Dated: February 25, 2024)

We consider the possibility that the superluminal neutrino propagation reported by the OPERA collaboration originates from a violation of CPT. On this basis we compare our actual knowledge concerning the CPT theorem to the nuclear reaction chain between CNGS and OPERA.

A recent experimental result from the OPERA collaboration [1] is that neutrinos of energies around $\langle E_\nu \rangle = 17$ GeV seem to travel at velocities of $v_\nu \approx 1.00002c$. If this observation is confirmed it is surprising in various ways: First, it implies a violation of conventional Lorentz variance at the order of $\epsilon_\Lambda = (v_\nu - c)/c = (2.48 \pm 0.58) \times 10^{-5}$. Second, if the violation would originate from quantum gravitational effects one would expect it to be suppressed by the Planck mass $m_\nu/M_{Pl} \sim 10^{-28}$. In this sense the reported effect is many orders of magnitude larger than predicted by quantum gravity of approaches. Third, if one tries to accommodate this effect in an extended theory of relativity that by construction allows for superluminal propagation, the observed velocity difference appears highly fine tuned to a value much less than one but different from zero. Thus, from this perspective the value of ϵ_Λ is much smaller than expected. Therefore, the value of ϵ_Λ is unnatural for both, quantum gravity and extended Lorentz approaches. Estimating the mass scale corresponding to this effect by $\epsilon_\Lambda \sim m_\nu/M_X$ one finds that the scale M_X should actually lie in the MeV range. The violation of CP in this energy regime is well studied within the standard model [2], therefore it is natural to consider also the possibility of a violation of CPT [3]. Since it has been shown that a violation of CPT implies a violation of Lorentz invariance [4] this fact could be relevant for an understanding of the origin and value of ϵ_Λ .

The idea of this comment is to study possible CPT [3] violations as source of LI violations. Please note that a violation of CPT implies a violation of Lorentz invariance [4] while the inverse statement is not necessarily true. In order to obtain an estimate for the value of CPT violation one has to consider the total nuclear reaction chain between CNGS and OPERA.

This reaction chain $p \rightarrow (\text{mesons} + \dots) \rightarrow (\nu + \dots)$ consists basically of three stages of propagation and two intermediate stages of conversion. It is hard to imagine that the conversions (decays) produce a Lorentz violating effect that results in an advance of $\delta l = 18$ meters. Also the velocity of the proton is known and measured to high precision leaving only the stages of meson and neutrino propagation as possible sources of Lorentz violation. Estimating the amount of Lorentz violation due to neutral Kaons [5, 6] one gets $\epsilon_\Lambda^{K^0} \approx \text{Re}(\delta_{CPT}^{K^0})$ with $0.2 \times 10^{-4} \leq \text{Re}(\delta_{CPT}^{K^0}) \leq 4.8 \times 10^{-4}$, which would be in agreement with the OPERA value. However, such an estimate is too naive. For example one knows that the relevant mesonic multiplicity consists of 99 percent pions and only about one percent neutral Kaons [7]. Although it is sufficient to have only a fraction of the events with superluminal velocities [8], it is clear that this fraction should exceed one percent.

There are also good experimental bounds on the amount CPT violation in pions [9]. However, the limit on CPT violation in pions measured at ~ 0.3 GeV [10] is $\Delta\tau_\pi = \epsilon_\pi\tau_\pi \approx 5 \cdot 10^{-4}\tau_\pi$. By using the corresponding Lorentz γ factor, the corresponding limit on the advance within the lifetime of a pion is $\delta l \leq \Delta\tau_\pi c\gamma \approx 0.5$ m, which is a factor of thirtysix smaller, than the advance of $\delta l = 18$ m needed to explain the OPERA data. Of course one has to notice that this comparison involves different energy scales (0.3 GeV and 17 GeV). Considering a cinematic explanation of this discrepancy already a linear energy dependence δ_{CPT}^π would be sufficient to get an agreement between $\epsilon_\pi|_{CPT}$ and $\epsilon_\Lambda|_{OPERA}$.

The remaining possibility is the propagation of the neutrino. It is known that the CPT violating terms in a renormalizable Lorentz violating standard model extension [11, 12] can not generate superluminal neutrino velocities. This implies that apparently the only realization of this idea in the neutrino sector will imply non-renormalizable terms. Unfortunately there is basically no model independent data on CPT violation in the neutrino sector. Therefore, the general comment has to end at this point.

As continuation a special toy model for CPT violation in the neutrino sector will be mentioned. For other models and explanations the reader is referred to [13–20]. A possible toy model for this idea can be constructed in the spirit of [21] where a mass splitting between neutrino mass m and anti-neutrino mass \tilde{m} is the source of CPT violation. The

*Electronic address: bkoch@fis.puc.cl

hamiltonian of a fermionic lagrangian is

$$: H_0 := \int \frac{d^3p}{(2\pi)^3} \sum_s ((\mathbf{p}^2 + m^2) a_{\mathbf{p}}^{s\dagger} a_{\mathbf{p}}^s + (\mathbf{p}^2 + \tilde{m}^2) b_{\mathbf{p}}^{s\dagger} b_{\mathbf{p}}^s) \quad (1)$$

Since the CPT operator converts the operators $a_{\mathbf{p}}^s$ and $b_{\mathbf{p}}^s$ into each other, any difference between m^2 and \tilde{m}^2 immediately generates a violation of CPT. This idea has already shown its utility in the context of explaining neutrino oscillations and the LSND results [22, 23]. As a slight modification one can consider that from a certain energy scale M_X on a new interaction with width Γ introduces a momentum dependent splitting of the type

$$\tilde{m}^2 - m^2 = 2\epsilon \cdot M_X^2 (\text{Erf}((\mathbf{p} - M_X)/\Gamma) - 1) \quad . \quad (2)$$

Due to the error function the splitting is just generated above the momentum scale M_X , while the CPT is conserved below M_X .

The violation parameter ϵ has mass dimension zero and it is assumed that $\epsilon \ll 1$. The dispersion relation of the neutrino would then be

$$E^2 = \mathbf{p}^2 + \epsilon (\text{Erf}((\mathbf{p} - M_X)/\Gamma) - 1) + m^2 \quad . \quad (3)$$

Thus, the group velocity for neutrinos with mass m is

$$\frac{v}{c} = \partial_p E = 1 + \epsilon \frac{M_X^2}{\sqrt{\pi}\Gamma|\mathbf{p}|} \exp(-(|\mathbf{p}| - M_X)^2/\Gamma^2) + \mathcal{O}\left(\epsilon \frac{m^2}{\mathbf{p}^2}, \epsilon^2, \frac{m^4}{\mathbf{p}^4}\right) \quad (4)$$

where an expansion for large momentum and small CPT violating parameter was used. One observes that around M_X , the enhancement is $\Delta v/c \sim \epsilon p/\Gamma$, whereas for larger values of p it is $\sim \epsilon$. Thus, the relation (4) has the effect that around the scale M_X a large enhance of group velocity occurs, whereas before and after, this effect is zero or negligible as it is shown in figure (1). Such a cinematic behavior could explain the absence of significant superluminal




FIG. 1: Example for (4) with $m \ll M_X = 17$ GeV, $\Gamma = 1$ GeV, and $\epsilon = 2.5 \cdot 10^{-5}$

velocities of neutrinos from the supernova 1987a. This energy dependence would also be essentially different from the power law deviations and respective bounds studied in [24, 25]. As explained in [22, 26] CPT violating mass differences could also have effects on special types of neutrino oscillation experiments [27]. Thus, any model like (2) should be confronted with all available neutrino data [28–30], which goes beyond the scope of this comment.

Although the relation (2) seems to work potentially well, the point of this paper is not to put forward a specific model. The idea is to emphasize the possible connection between a tiny CPT violation ϵ and the small deviation from c reported by OPERA. It is found that for a constant CPT violation in pions ϵ_π is not sufficient to be the source of the observed superluminal propagation. This restriction, does however not apply if one considers a ϵ_π that increases linearly with the energy. Additionally, also the propagation of the neutrino has the potential of describing such a sizable observational effect. Model independent tests of CPT violation in neutrinos are however missing. For both, the neutrino and the pion sector one would have to invoke some kind of energy dependent CPT violation in order to evade contradiction between different experiments.

Many thanks to M.A. Diaz, J.S. Diaz, and J. Gamboa for valuable hints. The work of B. K. was supported by CONICYT project PBCTNRO PSD-73.

-
- [1] OPERA, [arXiv:1109.4897 [hep-ex]].
 - [2] L. Wolfenstein (1989). CP violation. NorthHolland Publishing. ISBN: 0444-88081X; J. H. Christenson, J. W. Cronin, V. L. Fitch, R. Turlay Phys. Rev. Lett. **13**, 138 (1964); A. Hocker and Z. Ligeti, Ann. Rev. Nucl. Part. Sci. **56**, 501 (2006) [arXiv:hep-ph/0605217].
 - [3] R. F. Streater and A. S. Wightman, Princeton University Press, ISBN-10: 0691070628.
 - [4] O. W. Greenberg, Phys. Rev. Lett. **89**, 231602 (2002). [hep-ph/0201258].
 - [5] A. Angelopoulos *et al.* [CPLEAR Collaboration], Phys. Rept. **374**, 165-270 (2003).
 - [6] C. Amsler *et al.* [PDG], Phys. Lett.B **667**,1 (2006); F. Ambrosino *et al.* [KLOE Collaboration], JHEP **12**, 011 (2008); A. Di Domenico *et al.* [KLOE Collaboration], J. Phys. Conf. Ser. **171**, 012008 (2009).
 - [7] M. Bleicher, E. Zabrodin, C. Speies, S. A. Bass, C. Ernst, S. Soff, L. Bravina, M. Belkacem *et al.*, J. Phys. G **G25**, 1859-1896 (1999). [hep-ph/9909407].
 - [8] W. Winter, [arXiv:1110.0424 [hep-ph]].
 - [9] K. Nakamura *et al.* (Particle Data Group), J. Phys. G **37**, 075021 (2010).
 - [10] A. J. Greenberg, D. S. Ayres, A. M. Cormack, R. W. Kenney, D. O. Caldwell, V. B. Elings, W. P. Hesse, R. J. Morrison, Phys. Rev. Lett. **23**, 1267 (1969).
 - [11] V. A. Kostelecky, Phys. Rev. Lett. **80**, 1818 (1998). [hep-ph/9809572]; V. A. Kostelecky, R. Lehnert, Phys. Rev. **D63**, 065008 (2001). [hep-th/0012060].
 - [12] J. S. Diaz, V. A. Kostelecky, Phys. Lett. **B700**, 25-28 (2011). [arXiv:1012.5985 [hep-ph]]; J. S. Diaz, A. Kostelecky, [arXiv:1108.1799 [hep-ph]]; J. S. Diaz, [arXiv:1109.4620 [hep-ph]]; J. S. Diaz, private communication.
 - [13] J. Alfaro, Phys. Rev. Lett. **94**, 221302 (2005). [hep-th/0412295].
 - [14] H. Pas, S. Pakvasa and T. J. Weiler, Phys. Rev. D **72**, 095017 (2005) [arXiv:hep-ph/0504096].
 - [15] J. Dent, H. Pas, S. Pakvasa and T. J. Weiler, arXiv:0710.2524 [hep-ph].
 - [16] F. R. Klinkhamer, [arXiv:1109.5671 [hep-ph]].
 - [17] P. Wang, H. Wu and H. Yang, arXiv:1109.6930 [hep-ph].
 - [18] P. Wang, H. Wu, H. Yang, [arXiv:1110.0449 [hep-ph]].
 - [19] J. Alfaro, [arXiv:1110.3540 [hep-ph]].
 - [20] F. Giacosa and P. K. S. Lottini, arXiv:1110.3642 [hep-ph].
 - [21] G. Barenboim, L. Borisso, J. D. Lykken, A. Y. Smirnov, JHEP **0210**, 001 (2002). [hep-ph/0108199].
 - [22] G. Barenboim, N. E. Mavromatos, JHEP **0501**, 034 (2005). [hep-ph/0404014].
 - [23] N. E. Mavromatos, arXiv:1110.3729 [hep-ph].
 - [24] G. F. Giudice, S. Sibiryakov and A. Strumia, arXiv:1109.5682 [hep-ph].
 - [25] J. Alexandre, J. Ellis, N. E. Mavromatos, [arXiv:1109.6296 [hep-ph]].
 - [26] H. Murayama and T. Yanagida, Phys. Lett. B **520**, 263 (2001) [arXiv:hep-ph/0010178].
 - [27] A. Aguilar *et al.* [LSND Collaboration], Phys. Rev. **D64**, 112007 (2001). [hep-ex/0104049].
 - [28] D. Fargion, D. D'Armiento, [arXiv:1109.5368 [astro-ph.HE]].
 - [29] X. -J. Bi, P. -F. Yin, Z. -H. Yu, Q. Yuan, [arXiv:1109.6667 [hep-ph]].
 - [30] N. D. H. Dass, [arXiv:1110.0351 [hep-ph]].