

Gauge-Higgs unification models in six dimensions with S^2/Z_2 extra space and GUT gauge symmetry

Cheng-Wei Chiang,^{1,2,*} Takaaki Nomura,^{1,†} and Joe Sato^{3,‡}

¹*Department of Physics and Center for Mathematics and Theoretical Physics,
National Central University, Chungli, Taiwan 320, R.O.C.*

²*Institute of Physics, Academia Sinica, Taipei, Taiwan 115, R.O.C.*

³*Department of Physics, Saitama University,
Shimo-Okubo, Sakura-ku, Saitama 355-8570, Japan*

(Dated: February 18, 2022)

Abstract

In this article, we review gauge-Higgs unification models based on gauge theories defined on six-dimensional spacetime with S^2/Z_2 topology in the extra spatial dimensions. On the extra S^2/Z_2 space, non-trivial boundary conditions are imposed. This review considers two scenarios for constructing a four-dimensional theory from the six-dimensional model. One scheme utilizes the $SO(12)$ gauge symmetry with a special symmetry condition imposed on the gauge field, whereas the other employs the E_6 gauge symmetry without requiring the additional symmetry condition. Both models lead to a Standard Model-like gauge theory with the $SU(3) \times SU(2)_L \times U(1)_Y (\times U(1)^2)$ symmetry and SM fermions in four dimensions. The Higgs sector of the model is also analyzed. The electroweak symmetry breaking can be realized, and the weak gauge boson and Higgs boson masses are obtained.

*e-mail: chengwei@ncu.edu.tw

†e-mail: nomura@ncu.edu.tw

‡e-mail: joe@phy.saitama-u.ac.jp

I. INTRODUCTION

The Higgs sector of the Standard Model (SM) plays an essential role in the spontaneous symmetry breaking (SSB) from the $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge group down to $SU(3)_C \times U(1)_{EM}$, thereby giving masses to the SM elementary particles. However, the SM does not address the most fundamental nature of the Higgs sector, such as the mass and self-coupling constant of the Higgs boson. Therefore, the Higgs sector is not only the last territory in the SM to be discovered, but will also provide key clues to new physics at higher energy scales.

Gauge-Higgs unification is one of many attractive approaches to physics beyond the SM in this regard [1–3] (for recent approaches, see Refs. [5–21]). In this approach, the Higgs particles originate from the extra-dimensional components of the gauge field defined on spacetime with the number of dimensions greater than four. In other words, the Higgs sector is embraced into the gauge interactions in the higher-dimensional model, and many fundamental properties of Higgs boson are dictated by the gauge interactions.

In our recent studies, we have shown interesting properties of one type of gauge-Higgs unification models based on grand unified gauge theories defined on six-dimensional (6D) spacetime, with the extra-dimensional space having the topological structure of two-sphere orbifold S^2/Z_2 [22, 23].

In the usual coset space dimensional reduction (CSDR) approach [1, 24–27], one imposes on the gauge fields the symmetry condition which identifies the gauge transformation as the isometry transformation of S^2 due to its coset space structure $S^2 = SU(2)/U(1)$. The dimensional reduction is explicitly carried out by applying the solution of the symmetry condition. A background gauge field is introduced as part of the solution [1]. Such a background gauge field is also necessary for obtaining chiral fermions in four dimensional (4D) spacetime, even without the symmetry condition. After the dimensional reduction, no Kaluza-Klein (KK) mode appears because of the imposed symmetry condition. The symmetry condition also restricts the gauge symmetry and the scalar contents originated from the extra gauge field components in the 4D spacetime. Moreover, a suitable potential for the scalar sector can be obtained to induce SSB at tree level.

In this article, we consider two scenarios for constructing the 4D theory from a 6D model: one utilizing the symmetry condition for the gauge field, whereas the other without it. In the first scenario, however, we do not impose the condition on the fermions as used in other

CSDR models. We then have massive KK modes for fermions but not the gauge and scalar fields in 4D. We can thus obtain a dark matter candidate under assumed KK parity. In the case without the symmetry condition, we find that the background gauge field restricts the gauge symmetry and massless particle contents in 4D. Also, there are KK modes for each field, with the mass spectrum determined according to the model. Generally, massless modes do not appear in the KK mass spectrum because of the positive curvature of the S^2 space [28]. With the help of the background gauge field, however, we obtain massless KK modes for the gauge bosons and fermions.

Generally, the gauge symmetry of a grand unified theory (GUT) tends to remain in 4D in these dimensional reduction approaches [25, 29–33]. Also, it is difficult to obtain the appropriate Higgs potential to break the GUT gauge symmetry because of the gauge group structure. A GUT gauge symmetry can be broken to the SM gauge symmetry by the non-trivial boundary conditions (for cases with orbifold extra space, see, for example, [5–9, 12, 13, 17–19, 34, 35]). Therefore, to solve the above-mentioned problems, we impose on the fields of the 6D model a set of non-trivial boundary conditions on the S^2/Z_2 space. Therefore, the gauge symmetry, scalar contents and massless fermions are determined by these boundary conditions and the background gauge field. We find that in both scenarios, with or without the symmetry condition for the gauge field, the electroweak symmetry breaking (EWSB) can be realized, and the Higgs boson mass is predicted by analyzing the Higgs potential in the respective model.

This paper is organized as follows. In Sec. II, we review the two scenarios for constructing the 4D theory from gauge theories on 6D spacetime whose extra space has the S^2/Z_2 topology with a set of non-trivial boundary conditions. In Sec. III, we show the models based on $SO(12)$ and E_6 gauge symmetries, with the former imposed by the symmetry condition on the gauge field and the latter without. We summarize our results in Sec. IV.

II. THE 6D GAUGE-HIGGS UNIFICATION MODEL CONSTRUCTION SCHEME WITH EXTRA-SPACE S^2/Z_2

In this section, we recapitulate the scheme for constructing a 4D theory from a gauge theory on 6D spacetime which has extra-space as two-sphere orbifold S^2/Z_2 . We consider two cases; one case uses the symmetry condition and the other case does not use symmetry

condition. We apply non-trivial boundary condition for both cases.

A. A Gauge theory on 6D spacetime with S^2/Z_2 extra-space

1. The 6D spacetime with S^2/Z_2 extra-space

We begin with properties of a 6D spacetime M^6 . The spacetime M^6 is assumed to be a direct product of the 4D Minkowski spacetime M^4 and two-sphere orbifold S^2/Z_2 such that $M^6 = M^4 \times S^2/Z_2$. The two-sphere S^2 is a unique two-dimensional coset space, and can be written as $S^2 = \text{SU}(2)_I/\text{U}(1)_I$, where $\text{U}(1)_I$ is the subgroup of $\text{SU}(2)_I$. This coset space structure of S^2 requires that S^2 has the isometry group $\text{SU}(2)_I$, and that the group $\text{U}(1)_I$ is embedded into the group $\text{SO}(2)$ which is a subgroup of the Lorentz group $\text{SO}(1,5)$. We denote the coordinate of M^6 by $X^M = (x^\mu, y^\theta = \theta, y^\phi = \phi)$, where x^μ and $\{\theta, \phi\}$ are M^4 coordinates and S^2 spherical coordinates, respectively. The spacetime index M runs over $\mu \in \{0, 1, 2, 3\}$ and $\alpha \in \{\theta, \phi\}$. The orbifold S^2/Z_2 is defined by the identification of (θ, ϕ) and $(\pi - \theta, -\phi)$ [39]. The two fixed points are $(\pi/2, 0)$ and $(\pi/2, \pi)$. The metric of M^6 , denoted by g_{MN} , can be written as

$$g_{MN} = \begin{pmatrix} \eta_{\mu\nu} & 0 \\ 0 & -g_{\alpha\beta} \end{pmatrix}, \quad (1)$$

where $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$ and $g_{\alpha\beta} = \text{diag}(1, \sin^{-2}\theta)$ are metric of M^4 and S^2 respectively. Notice that we omit the radius R of S^2 in this discussion. We define the vielbein e_A^M that connects the metric of M^6 and that of the tangent space of M^6 , denoted by h_{AB} , as $g_{MN} = e_M^A e_N^B h_{AB}$. Here $A = (\mu, a)$, where $a \in \{4, 5\}$, is the index for the coordinates of tangent space of M^6 . The explicit form of the vielbeins are summarized as

$$e_\theta^1 = 1, \quad e_\phi^2 = \sin \theta, \quad e_\phi^1 = e_\theta^2 = 0. \quad (2)$$

Also the non-zero components of spin connection are

$$R_\phi^{12} = -R_\phi^{21} = -\cos \theta. \quad (3)$$

2. A Lagrangian on 6D spacetime with S^2/Z_2 extra-space

We then show the general structure of a gauge theory on M^6 . We introduce a gauge field $A_M(x, y) = (A_\mu(x, y), A_\alpha(x, y))$, which belongs to the adjoint representation of the gauge

group G , and fermions $\psi(x, y)$, which lies in a representation F of G . The action of this theory is given by

$$S = \int dx^4 \sin \theta d\theta d\phi \left(\bar{\Psi} i \Gamma^\mu D_\mu \Psi + \bar{\Psi} i \Gamma^a e_a^\alpha D_\alpha \Psi - \frac{1}{4g^2} g^{MN} g^{KL} \text{Tr}[F_{MK} F_{NL}] \right), \quad (4)$$

where $F_{MN} = \partial_M A_N(X) - \partial_N A_M(X) - [A_M(X), A_N(X)]$ is the field strength, D_M is the covariant derivative including spin connection, and Γ_A represents the 6-dimensional Clifford algebra. Here D_M and Γ_A can be written explicitly as,

$$D_\mu = \partial_\mu - A_\mu, \quad D_\theta = \partial_\theta - A_\theta, \quad D_\phi = \partial_\phi - i \frac{\Sigma_3}{2} \cos \theta - A_\phi, \quad (5)$$

$$\Gamma_\mu = \gamma_\mu \otimes \mathbf{I}_2, \quad \Gamma_4 = \gamma_5 \otimes \sigma_1, \quad \Gamma_5 = \gamma_5 \otimes \sigma_2, \quad (6)$$

where $\{\gamma_\mu, \gamma_5\}$ are the 4-dimensional Dirac matrices, $\sigma_i (i = 1, 2, 3)$ are Pauli matrices, \mathbf{I}_d is $d \times d$ identity, and Σ_3 is defined as $\Sigma_3 = \mathbf{I}_4 \otimes \sigma_3$. The covariant derivative D_ϕ has the spin connection term $i \frac{\Sigma_3}{2} \cos \theta$ which is needed for space with non-zero curvature like S^2 and applied for only fermions. In 6D spacetime, we can define chirality of fermions and the projection operators are defined as

$$\Gamma_\pm = \frac{1 \pm \Gamma_7}{2} \quad (7)$$

where $\Gamma_7 \equiv \gamma_5 \otimes \sigma_3$ is the chiral operator. We can write chiral fermion on 6D spacetime such as

$$\Psi_\pm = \Gamma_\pm \Psi, \quad \Gamma_7 \Psi_\pm = \pm \Psi_\pm \quad (8)$$

where ψ is a Dirac fermion on 6D spacetime. The 6D chiral fermion can be also written in terms of 4D chiral fermions $\psi_{L(R)}$ as

$$\Psi_+ = \begin{pmatrix} \psi_R \\ \psi_L \end{pmatrix}, \quad (9)$$

$$\Psi_- = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix}. \quad (10)$$

3. Non-trivial boundary conditions on the two-sphere orbifold

On the two-sphere orbifold, one can consider parity operations $P : (\theta, \phi) \rightarrow (\pi - \theta, -\phi)$ and azimuthal translation $T_\phi : (\theta, \phi) \rightarrow (\theta, \phi + 2\pi)$. Here we note that the periodicity

$\phi \rightarrow \phi + 2\pi$ is not associated with the orbifolding. We can impose the following two types of boundary conditions on both gauge and fermion fields under the two operations:

$$A_\mu(x, \pi - \theta, -\phi) = P_1 A_\mu(x, \theta, \phi) P_1, \quad A_\mu(x, \pi - \theta, 2\pi - \phi) = P_2 A_\mu(x, \theta, \phi) P_2, \quad (11)$$

$$A_\alpha(x, \pi - \theta, -\phi) = -P_1 A_{\theta, \phi}(x, \theta, \phi) P_1, \quad A_\alpha(x, \pi - \theta, 2\pi - \phi) = -P_2 A_\alpha(x, \theta, \phi) P_2, \quad (12)$$

$$\Psi(x, \pi - \theta, -\phi) = \pm \gamma_5 P_1 \Psi(x, \theta, \phi), \quad \Psi(x, \pi - \theta, 2\pi - \phi) = \pm \gamma_5 P_2 \Psi(x, \theta, \phi), \quad (13)$$

or

$$A_\mu(x, \pi - \theta, -\phi) = P_1 A_\mu(x, \theta, \phi) P_1, \quad A_\mu(x, \theta, \phi + 2\pi) = P_2 A_\mu(x, \theta, \phi) P_2, \quad (14)$$

$$A_\alpha(x, \pi - \theta, -\phi) = -P_1 A_{\theta, \phi}(x, \theta, \phi) P_1, \quad A_\alpha(x, \theta, \phi + 2\pi) = P_2 A_\alpha(x, \theta, \phi) P_2, \quad (15)$$

$$\Psi(x, \pi - \theta, -\phi) = \pm \gamma_5 P_1 \Psi(x, \theta, \phi), \quad \Psi(x, \theta, \phi + 2\pi) = \pm P_2 \Psi(x, \theta, \phi), \quad (16)$$

where former conditions are associated with P operation and combination of P and T_ϕ operations, and latter conditions are associated with P and T_ϕ operation itself. These boundary conditions are determined by requiring the invariance of the six-dimensional action under the transformation $(\theta, \phi) \rightarrow (\pi - \theta, -\phi)$ and $\phi \rightarrow \phi + 2\pi$.

The projection matrices $P_{1,2}$ act on the gauge group representation space and have eigenvalues ± 1 . They assign different parities for different representation components. For fermion boundary conditions, the sign in front of γ_5 can be either $+$ or $-$ since the fermions always appear in bilinear forms in the action. The 4-dimensional action is then restricted by these parity assignments.

B. The dimensional reduction scheme with symmetry condition

Here we review the dimensional reduction scheme which apply symmetry condition of gauge field [22].

1. The symmetry condition

We impose on the gauge field $A_M(X)$ the symmetry which connects $SU(2)_I$ isometry transformation on S^2 and the gauge transformation on the fields in order to carry out dimensional reduction, and the non-trivial boundary conditions of S^2/Z_2 to restrict 4D theory. The symmetry requires that the $SU(2)_I$ coordinate transformation should be compensated

by a gauge transformation [1, 24]. The symmetry further leads to the following set of the symmetry condition on the fields:

$$\xi_i^\beta \partial_\beta A_\mu = \partial_\alpha W_i + [W_i, A_\mu], \quad (17)$$

$$\xi_i^\beta \partial_\beta A_\alpha + \partial_\alpha \xi_i^\beta A_\beta = \partial_\alpha W_i + [W_i, A_\alpha], \quad (18)$$

where ξ_i^α is the Killing vectors generating $SU(2)_I$ symmetry and W_i are some fields which generate an infinitesimal gauge transformation of G . Here index $i = 1, 2, 3$ corresponds to that of $SU(2)$ generators. The explicit forms of ξ_i^α s for S^2 are:

$$\begin{aligned} \xi_1^\theta &= \sin \phi, & \xi_1^\phi &= \cot \theta \cos \phi, \\ \xi_2^\theta &= -\cos \phi, & \xi_2^\phi &= \cot \theta \sin \phi, \\ \xi_3^\theta &= 0, & \xi_3^\phi &= -1. \end{aligned} \quad (19)$$

The LHSs of Eq (17,18) are infinitesimal isometry $SU(2)_I$ transformation and the RHSs of those are infinitesimal gauge transformation.

2. The dimensional reduction and a Lagrangian in 4D spacetime

The dimensional reduction of gauge sector is explicitly carried out by applying the solutions of the symmetry condition Eq (17,18). These solutions are given by Manton [1] as

$$A_\mu = A_\mu(x), \quad A_\theta = -\Phi_1(x), \quad A_\phi = \Phi_2(x) \sin \theta - \Phi_3 \cos \theta, \quad (20)$$

$$W_1 = -\Phi_3 \frac{\cos \phi}{\sin \theta}, \quad W_2 = -\Phi_3 \frac{\sin \phi}{\sin \theta}, \quad W_3 = 0, \quad (21)$$

and satisfy the following constraints:

$$[\Phi_3, A_\mu] = 0, \quad (22)$$

$$[-i\Phi_3, \Phi_i(x)] = i\epsilon_{3ij}\Phi_j(x), \quad (23)$$

where $\Phi_1(x)$ and $\Phi_2(x)$ are scalar fields, and $-i\Phi_3$ are chosen as generator of $U(1)_I$. Note that the Φ_3 term for A_ϕ corresponds to the background gauge field [36]. Substituting the solutions Eq (20) into $A_M(X)$ in action Eq (4), we can easily integrate coordinates θ and ϕ

in the gauge sector. We then obtain a four dimensional action as

$$S_{4D}^{(gauge)} = \int d^4x \left(-\frac{1}{4g^2} Tr[F_{\mu\nu}F^{\mu\nu}(x)] \right. \\ \left. -\frac{1}{2g^2} Tr[D'_\mu\Phi_1(x)D'^\mu\Phi_1(x) + D'_\mu\Phi_2(x)D'^\mu\Phi_2(x)] \right. \\ \left. -\frac{1}{2g^2} Tr[(\Phi_3 + [\Phi_1(x), \Phi_2(x)])(\Phi_3 + [\Phi_1(x), \Phi_2(x)])] \right), \quad (24)$$

where $D'_\mu\Phi = \partial_\mu - [A_\mu, \Phi]$. The fermion sector of 4D action is obtained by expanding fermions in normal modes of S^2/Z_2 and then integrating S^2/Z_2 coordinate in 6D action. Thus, the fermions have massive KK modes which would be a candidate of dark matter. Generally, the KK modes do not have massless mode because of the positive curvature of S^2 [28]. We, however, can show that the fermion components satisfying the following condition have massless mode:

$$-i\Phi_3\psi = \frac{\Sigma_3}{2}\psi. \quad (25)$$

Square mass of the KK modes are eigenvalues of square of extra-dimensional Dirac-operator $-i\hat{D}$. In the S^2 case, $-i\hat{D}$ is written as

$$-i\hat{D} = -ie^{\alpha a}\Gamma_a D_\alpha \\ = -i\left[\Sigma_1\left(\partial_\theta + \frac{\cot\theta}{2}\right) + \Sigma_2\left(\frac{1}{\sin\theta}\partial_\phi + \Phi_3\cot\theta\right)\right], \quad (26)$$

where $\Sigma_i = \mathbf{I}_4 \times \sigma_i$. Square of $-i\hat{D}$ can be explicitly calculated:

$$(-i\hat{D})^2 = -\left[\frac{1}{\sin\theta}\partial_\theta(\sin\theta\partial_\theta) + \frac{1}{\sin^2\theta}\partial_\phi^2 + i(2(-i\Phi_3) - \Sigma_3)\frac{\cos\theta}{\sin^2\theta}\partial_\phi \right. \\ \left. -\frac{1}{4} - \frac{1}{4\sin^2\theta} + \Sigma_3(-i\Phi_3)\frac{1}{\sin^2\theta} - (-i\Phi_3)^2\cot^2\theta\right]. \quad (27)$$

We then act this operator on a fermion $\psi(X)$ which satisfy Eq. (25), and obtain the relation

$$(-i\hat{D})^2\psi = -\left[\frac{1}{\sin\theta}\partial_\theta(\sin\theta\partial_\theta) + \frac{1}{\sin^2\theta}\partial_\phi^2\right]\psi. \quad (28)$$

The eigenvalues of the RHS operator are less than or equal to zero. Thus the fermion components satisfying Eq. (25) have massless mode, while other components only have massive KK mode. Note that the massless mode ψ_0 should be independent of S^2 coordinates θ and ϕ :

$$\psi_0 = \psi(x). \quad (29)$$

The existence of massless fermion may indicate the meaning of the symmetry condition; though the energy density of the gauge sector in the appearance of the background fields is higher than that of no background fields, since we have massless fermions, it may consist a ground state as a total in the presence of fermions. We also note that we could impose symmetry condition on fermions [25, 37]. In that case, we obtain the massless condition Eq. (25) from symmetry condition of fermion, and the solution of symmetry condition is independent from S^2 coordinate: $\psi = \psi(x)$ with no massive KK mode. Therefore, we can apply the same discussion for this case as our case if we only focus on the massless mode in our scheme.

3. A gauge symmetry and particle contents in 4D spacetime

The symmetry conditions and the non-trivial boundary conditions substantially constrain the four dimensional gauge group and its representations for the particle contents. The gauge symmetry and particle contents in 4D spacetime must satisfy the constraints Eq (22),(23),(25) and be consistent with the boundary conditions Eq (14)-(16). We show the prescriptions to identify 4D gauge symmetry and particle contents below.

First, we show the prescriptions to identify gauge symmetry and field components which satisfy the constraints Eq (22),(23),(25). The gauge group H that satisfy the constraint Eq (22) is identified as

$$H = C_G(U(1)_I) \quad (30)$$

where $C_G(U(1)_I)$ denotes the centralizer of $U(1)_I$ in G [24]. Note that this implies $G \supset H = H' \times U(1)_I$, where H' is some subgroup of G .

Second, the scalar field components which satisfy the constraints Eq. (23) are specified by the following prescription. Suppose that the adjoint representations of $SU(2)_I$ and G are decomposed according to the embeddings $SU(2)_I \supset U(1)_I$ and $G \supset H' \times U(1)_I$ as

$$3(\text{adj } SU(2)) = (0(\text{adj } U(1)_R)) + (2) + (-2), \quad (31)$$

$$\text{adj } G = (\text{adj } H)(0) + 1(0(\text{adj } U(1)_R)) + \sum_g h_g(r_g), \quad (32)$$

where h_g s denote representation of H' , and r_g s denote $U(1)_I$ charges. The scalar components satisfying the constraints belong to h_g s whose corresponding r_g s in the decomposition Eq. (32) are ± 2 .

Third, the fermion components which satisfy the constraints Eq. (25) are determined as follows [37]. Let the group $U(1)_I$ be embedded into the Lorentz group $SO(2)$ in such a way that the vector representation 2 of $SO(2)$ is decomposed according to $SO(2) \supset U(1)_I$ as

$$2 = (2) + (-2). \quad (33)$$

This embedding specifies a decomposition of the weyl spinor representation $\sigma_6=4$ of $SO(1,5)$ according to $SO(1,5) \supset SU(2) \times SU(2) \times U(1)_I$ as

$$\sigma_6 = (2, 1)(1) + (1, 2)(-1), \quad (34)$$

where $SU(2) \times SU(2)$ representations (2,1) and (1,2) correspond to left-handed and right-handed spinors, respectively. We note that this decomposition corresponds to Eq. (9)(or Eq. (10)). We then decompose F according to $G \supset H' \times U(1)_I$ as

$$F = \sum_f h_f(r_f). \quad (35)$$

Now the fermion components satisfying the constraints are identified as $h_f s$ whose corresponding $r_f s$ in the decomposition Eq. (35) are (1) for left-handed fermions and (-1) for right-handed fermions.

Finally, we show which gauge symmetry and field components remain in 4D spacetime by surveying the consistency between the boundary conditions Eq. (14)-(16), the solutions Eq. (20), and fermion massless mode Eq. (29). We then apply Eq (20) and Eq. (29) to Eq. (14)-(16), and obtain the parity conditions

$$A_\mu(x) = P_{1(2)} A_\mu(x) P_{1(2)}, \quad (36)$$

$$-\Phi_1(x) = -P_1(-\Phi_1(x))P_1, \quad (37)$$

$$-\Phi_1(x) = P_2(-\Phi_1(x))P_2, \quad (38)$$

$$\Phi_2(x) + \Phi_3 \cos \theta = -P_1 \Phi_2(x) P_1 + P_1 \Phi_3 P_1 \cos \theta, \quad (39)$$

$$\Phi_2(x) - \Phi_3 \cos \theta = P_2 \Phi_2(x) P_2 - P_2 \Phi_3 P_2 \cos \theta, \quad (40)$$

$$\Psi(x) = \gamma^5 P_1 \Psi(x), \quad (41)$$

$$\Psi(x) = P_2 \Psi(x). \quad (42)$$

We find that gauge fields, scalar fields and massless fermions in 4D spacetime should be even for $P_1 A_\mu P_1$ and $P_2 A_\mu P_2$; $-P_1 \Phi_{1,2} P_1$ and $P_2 \Phi_{1,2} P_2$; $\gamma_5 P_1 \psi$ and $P_2 \psi$, respectively. Φ_3 always

remains since it is proportional to an $U(1)_I$ generator and commutes with $P(P')$. Therefore the particle contents are identified as the components which satisfy both the constraints Eq (22),(23),(25) and the parity conditions Eq (36)-(42). The gauge symmetry remained in 4D spacetime can also be identified by observing which components of the gauge fields remain.

C. The dimensional reduction scheme without symmetry condition

Here we review the dimensional reduction scheme which does not apply symmetry condition of gauge field [23].

1. Background gauge field and gauge group reduction

For the case without symmetry condition, we consider the background gauge field $A_\phi^B \equiv \tilde{A}_\phi^B \sin \theta$ that corresponds to a Dirac monopole [36]

$$\tilde{A}_\phi^B = -Q \frac{\cos \theta \mp 1}{\sin \theta} , \quad (- : 0 \leq \theta < \frac{\pi}{2} , \quad + : \frac{\pi}{2} \leq \theta \leq \pi) \quad (43)$$

where Q is proportional to the generator of a $U(1)$ subgroup of the original gauge group. This background gauge field A_ϕ^B is corresponding to $\Phi_3 \cos \theta \subset A_\phi$ in Eq. (20).

The background gauge field is chosen to belong to $U(1)_I$ group which is a subgroup of original gauge group G such as

$$G \supset G_{\text{sub}} \otimes U(1)_I, \quad (44)$$

where G_{sub} is subgroup of G . We then find that there is no massless mode for gauge field components which have non-zero $U(1)_I$ charge. In fact, these components acquire masses due to the background field from the term proportional to $F_{\mu\phi} F_\phi^\mu$

$$\begin{aligned} & Tr \left[-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2R^2 \sin^2 \theta} F_{\mu\phi} F_\phi^\mu \right] \\ & \rightarrow Tr \left[-\frac{1}{4} (\partial_\mu A_\nu - \partial_\nu A_\mu) (\partial^\mu A^\nu - \partial^\nu A^\mu) - \frac{1}{2R^2 \sin^2 \theta} [A_\mu, A_\phi^B] [A^\mu, A_\phi^B] \right] . \end{aligned} \quad (45)$$

For the components of A_μ with nonzero $U(1)_I$ charge, we have

$$A_\mu^i Q_i + A_{i\mu} Q^i \in A_\mu , \quad (46)$$

where Q_i ($Q^i = Q_i^\dagger$) are generators corresponding to distinct components in Eq. (97) that have nonzero $U(1)_I$ charges, and $A_{i\mu}$ ($A_\mu^i = A_{i\mu}^\dagger$) are the corresponding components of A_μ . We then find the term

$$\begin{aligned} \frac{1}{\sin^2 \theta} \text{Tr}[[A_\mu, A_\phi^B][A^\mu, A_\phi^B]] &= \frac{(\cos \theta \mp 1)^2}{\sin^2 \theta} \text{Tr}[[A_\mu^i Q_i + A_{i\mu} Q^i, Q][A^{i\mu} Q_i + A_i^\mu Q^i, Q]] \\ &= -2|q|^2 \frac{(\cos \theta \mp 1)^2}{\sin^2 \theta} A^{i\mu} A_{i\mu} , \end{aligned} \quad (47)$$

where q is the Q charge of the relevant component. Use of the facts that A_ϕ^B belongs to $U(1)_I$ and that $\text{Tr}[Q_i Q^i] = 2$ has been made in the above equation. A mass is thus associated with the lowest modes of those components of A_μ with nonzero $U(1)_I$ charges:

$$\begin{aligned} \int d\Omega \text{Tr} \left[-\frac{1}{4} (\partial_\mu A_\nu - \partial_\nu A_\mu) (\partial^\mu A^\nu - \partial^\nu A^\mu) - \frac{1}{2R^2 \sin^2 \theta} [A_\mu, A_B][A^\mu, A_B] \right] \Big|_{\text{lowest}} \\ \rightarrow -\frac{1}{2} [\partial_\mu A_{i\nu}(x) - \partial_\nu A_{i\mu}(x)] [\partial^\mu A^{i\nu}(x) - \partial^\nu A^{i\mu}(x)] + m_B^2 A_\mu^i(x) A^{i\mu}(x) , \end{aligned} \quad (48)$$

where the subscript ‘lowest’ means that only the lowest KK modes are kept. Here the lowest KK modes of A_μ correspond to the term $A_\mu(x)/\sqrt{4\pi}$ in the KK expansion. In summary, any representation of A_μ carrying a nonzero $U(1)_I$ charge acquires a mass m_B from the background field contribution after one integrates over the extra spatial coordinates. More explicitly,

$$m_B^2 = \frac{|q|^2}{4\pi R^2} \int d\Omega \frac{(\cos \theta \mp 1)^2}{\sin^2 \theta} \simeq 0.39 \frac{|q|^2}{R^2} \quad (49)$$

for the zero mode. Therefore the gauge group G is reduced to $G_{\text{sub}} \otimes U(1)_I$ by the existence of the background gauge field. We note that this condition is same as the case with the symmetry condition.

2. Scalar field contents in 4D spacetime

The scalar contents in 4D spacetime are obtained from the extra-dimensional components of the gauge field $\{A_\theta, A_\phi\}$ after integrating out the extra spatial coordinates. The kinetic term and potential term of $\{A_\theta, A_\phi\}$ are obtained from the gauge sector containing these

components

$$\begin{aligned}
S_{\text{scalar}} &= \int dx^4 d\Omega \left(\frac{1}{2g^2} \text{Tr}[F_{\mu\theta} F_{\theta}^{\mu}] + \frac{1}{2g^2 \sin^2 \theta} \text{Tr}[F_{\mu\phi} F_{\phi}^{\mu}] \right. \\
&\quad \left. - \frac{1}{2g^2 R^2 \sin^2 \theta} \text{Tr}[F_{\theta\phi} F_{\theta\phi}] \right) \\
&\rightarrow \int dx^4 d\Omega \left(\frac{1}{2g^2} \text{Tr}[(\partial_{\mu} A_{\theta} - i[A_{\mu}, A_{\theta}])^2] \right. \\
&\quad + \frac{1}{2g^2} \text{Tr}[(\partial_{\mu} A_{\theta} - i[A_{\mu}, \tilde{A}_{\phi}])^2] \\
&\quad - \frac{1}{2g^2 R^2} \text{Tr} \left[\left(\frac{1}{\sin \theta} \partial_{\theta} (\sin \theta \tilde{A}_{\phi} + \sin \theta \tilde{A}_{\phi}^B) \right. \right. \\
&\quad \left. \left. - \frac{1}{\sin \theta} \partial_{\phi} A_{\theta} - i[A_{\theta}, \tilde{A}_{\phi} + \tilde{A}_{\phi}^B] \right)^2 \right] \Bigg], \tag{50}
\end{aligned}$$

where we have taken $A_{\phi} = \tilde{A}_{\phi} \sin \theta + \tilde{A}_{\phi}^B \cos \theta$. In the second step indicated by the arrow in Eq. (50), we have omitted terms which do not involve A_{θ} and \tilde{A}_{ϕ} from the right-hand side of the first equality. It is known that one generally cannot obtain massless modes for physical scalar components in 4D spacetime [15, 40]. One can see this by noting that the eigenfunction of the operator $\frac{1}{\sin \theta} \partial_{\theta} \sin \theta$ with zero eigenvalue is not normalizable [15]. In other words, these fields have only KK modes. However, an interesting feature is that it is possible to obtain a negative squared mass when taking into account the interactions between the background gauge field \tilde{A}_{ϕ}^B and $\{A_{\theta}, \tilde{A}_{\phi}\}$. This happens when the component carries a nonzero $U(1)_I$ charge, as the background gauge field belongs to $U(1)_I$. In this case, the $(\ell = 1, m = 1)$ modes of these real scalar components are found to have a negative squared mass in 4D spacetime. They can be identified as the Higgs fields once they are shown to belong to the correct representation under the SM gauge group. Here the numbers (ℓ, m) are the angular momentum quantum number on S^2/Z_2 , and each KK mode is characterized by these numbers. One can show that the $(\ell = 1, m = 0)$ mode has a positive squared mass and is not considered as the Higgs field. A discussion of the KK masses with general (ℓ, m) will be given in Section III B 5.

3. Chiral fermions in 4D spacetime

We introduce fermions as the Weyl spinor fields of the 6D Lorentz group $SO(1,5)$. They can be written in terms of the $SO(1,3)$ Weyl spinors as Eq. (9) and Eq. (10). In general, fermions on the two-sphere do not have massless KK modes because of the positive curvature

of the two-sphere. The massless modes can be obtained by incorporating the background gauge field (43) though, for it can cancel the contribution from the positive curvature. In this case, the condition for obtaining a massless fermion mode is

$$Q\Psi = \pm \frac{1}{2}\Psi , \quad (51)$$

where Q comes from the background gauge field and is proportional to the $U(1)_I$ generator [36, 39, 40]. We observe that the upper [lower] component on the RHS of Eq. (9) [(10)] has a massless mode for the $+$ ($-$) sign on the RHS of Eq. (51).

4. The Higgs potential

The Lagrangian for the Higgs sector is derived from the gauge sector that contains extra-dimensional components of the gauge field $\{A_\theta, \tilde{A}_\phi\}$, as given in Eq. (50), by considering the lowest KK modes of them. The kinetic term and potential term are, respectively,

$$L_K = \frac{1}{2g^2} \int d\Omega \left(Tr[(\partial_\mu A_\theta - i[A_\mu, A_\theta])^2] + Tr[(\partial_\mu A_\theta - i[A_\mu, \tilde{A}_\phi])^2] \right) \Big|_{\text{lowest}} , \quad (52)$$

$$V = \frac{1}{2g^2 R^2} \int d\Omega Tr \left[\left(\frac{1}{\sin \theta} \partial_\theta (\sin \theta \tilde{A}_\phi + \sin \theta \tilde{A}_\phi^B) - \frac{1}{\sin \theta} \partial_\phi A_\theta - i[A_\theta, \tilde{A}_\phi + \tilde{A}_\phi^B] \right)^2 \right] \Big|_{\text{lowest}} . \quad (53)$$

In our model, scalar components other than the Higgs field have vanishing VEV because only the Higgs field has a negative mass-squared term, coming from the interaction with the background gauge field at tree level. Therefore, only the Higgs field contributes to the spontaneous symmetry breaking. Consider the $(1, 1)$ mode of the $\{(1, 2)(3, -3, 3) + \text{h.c.}\}$ representation in Eq. (98) as argued in the previous section. The gauge fields are given by the following KK expansions

$$A_\theta = -\frac{1}{\sqrt{2}} [\Phi_1(x) \partial_\theta Y_{11}^-(\theta, \phi) + \Phi_2(x) \frac{1}{\sin \theta} \partial_\phi Y_{11}^-(\theta, \phi)] + \dots , \quad (54)$$

$$\tilde{A}_\phi = \frac{1}{\sqrt{2}} [\Phi_2(x) \partial_\theta Y_{11}^-(\theta, \phi) - \Phi_1(x) \frac{1}{\sin \theta} \partial_\phi Y_{11}^-(\theta, \phi)] + \dots , \quad (55)$$

where \dots represents higher KK mode terms [39]. The function $Y_{11}^- = -1/\sqrt{2}[Y_{11} + Y_{1-1}]$ is odd under $(\theta, \phi) \rightarrow (\pi/2 - \theta, -\phi)$. We will discuss their higher KK modes and masses in

the existence of the background gauge field in Section III B 5. With Eqs. (54) and (55), the kinetic term becomes

$$L_K(x) = \frac{1}{2g^2} \left(Tr[D_\mu \Phi_1(x) D^\mu \Phi_1(x)] + Tr[D_\mu \Phi_2(x) D^\mu \Phi_2(x)] \right), \quad (56)$$

where $D_\mu \Phi_{1,2} = \partial_\mu \Phi_{1,2} - i[A_\mu, \Phi_{1,2}]$ is the covariant derivative acting on $\Phi_{1,2}$. The potential term, on the other hand, is

$$\begin{aligned} V = \frac{1}{2g^2 R^2} \int d\Omega Tr \bigg[& \left(-\sqrt{2} Y_{11}^- \Phi_2(x) + Q \right. \\ & + \frac{i}{2} [\Phi_1(x), \Phi_2(x)] \{ \partial_\theta Y_{11}^- \partial_\theta Y_{11}^- + \frac{1}{\sin^2 \theta} \partial_\phi Y_{11}^- \partial_\phi Y_{11}^- \} \\ & \left. + \frac{i}{\sqrt{2}} [\Phi_1(x), \tilde{A}_\phi^B] \partial_\theta Y_{11}^- + \frac{i}{\sqrt{2}} [\Phi_2(x), \tilde{A}_\phi^B] \frac{1}{\sin \theta} \partial_\phi Y_{11}^- \right)^2 \bigg], \end{aligned} \quad (57)$$

where $\partial_\theta(\sin \theta \tilde{A}_\phi^B) = Q \cos \theta$ from Eq. (43) is used. Expanding the square in the trace, we get

$$\begin{aligned} V = \frac{1}{2g^2 R^2} \int d\Omega Tr \bigg[& 2(Y_{11}^+)^2 \Phi_2^2(x) + Q^2 \\ & - \frac{1}{4} [\Phi_1(x), \Phi_2(x)]^2 \left(\partial_\theta Y_{11}^- \partial_\theta Y_{11}^- + \frac{1}{\sin^2 \theta} \partial_\phi Y_{11}^- \partial_\phi Y_{11}^- \right)^2 \\ & - \frac{1}{2} [\Phi_1(x), \tilde{A}_\phi^B]^2 (\partial_\theta Y_{11}^-)^2 - \frac{1}{2} [\Phi_2(x), \tilde{A}_\phi^B]^2 \left(\frac{1}{\sin \theta} \partial_\phi Y_{11}^- \right)^2 \\ & - 2i \Phi_2(x) [\Phi_1(x), \tilde{A}_\phi^B] Y_{11}^- \partial_\theta Y_{11}^- \\ & - [\Phi_1(x), \tilde{A}_\phi^B] [\Phi_2(x), \tilde{A}_\phi^B] \partial_\theta Y_{11}^- \frac{1}{\sin \theta} \partial_\phi Y_{11}^- \\ & + iQ [\Phi_1(x), \Phi_2(x)] \left(\partial_\theta Y_{11}^- \partial_\theta Y_{11}^- + \frac{1}{\sin^2 \theta} \partial_\phi Y_{11}^- \partial_\phi Y_{11}^- \right) \bigg], \end{aligned} \quad (58)$$

where terms that vanish after the $d\Omega$ integration are directly omitted. In the end, the potential is simplified to

$$\begin{aligned} V = \frac{1}{2g^2 R^2} Tr \bigg[& 2\Phi_2^2(x) + 4\pi Q^2 - \frac{3}{10\pi} [\Phi_1(x), \Phi_2(x)]^2 + \frac{5i}{2} Q [\Phi_1(x), \Phi_2(x)] \\ & + \mu_1 [Q, \Phi_1(x)]^2 + \mu_2 [Q, \Phi_2(x)]^2 \bigg], \end{aligned} \quad (59)$$

where use of $\tilde{A}_\phi^B = -Q(\cos \theta \mp 1)/\sin \theta$ has been made and $\mu_1 = 1 - \frac{3}{2} \ln 2$ and $\mu_2 = \frac{3}{4}(1 - 2 \ln 2)$.

We now take the following linear combination of Φ_1 and Φ_2 to form a complex Higgs doublet,

$$\Phi(x) = \frac{1}{\sqrt{2}}(\Phi_1(x) + i\Phi_2(x)) , \quad (60)$$

$$\Phi(x)^\dagger = \frac{1}{\sqrt{2}}(\Phi_1(x) - i\Phi_2(x)) . \quad (61)$$

It is straightforward to see that

$$[\Phi_1(x), \Phi_2(x)] = i[\Phi(x), \Phi^\dagger(x)] . \quad (62)$$

The kinetic term and the Higgs potential now become

$$L_K = \frac{1}{g^2} \text{Tr}[D_\mu \Phi^\dagger(x) D^\mu \Phi(x)] , \quad (63)$$

$$V = \frac{1}{2g^2 R^2} \text{Tr} \left[2\Phi_2^2(x) + 4\pi Q^2 + \frac{3}{10\pi} [\Phi(x), \Phi^\dagger(x)]^2 - \frac{5}{2} Q[\Phi(x), \Phi^\dagger(x)] \right. \\ \left. + \mu_1 [Q, \Phi_1(x)]^2 + \mu_2 [Q, \Phi_2(x)]^2 \right] . \quad (64)$$

III. THE MODELS BASED ON OUR SCHEMES

In this section, we show concrete models based on the scheme introduced previous section. We introduce the model based on $\text{SO}(12)$ gauge symmetry for the scheme with symmetry condition and introduce the model based on E_6 gauge symmetry for the scheme without symmetry condition [22, 23].

A. The $\text{SO}(12)$ model with symmetry condtion

Here we show a model based on a gauge group $G=\text{SO}(12)$ and a representation $F=32$ of $\text{SO}(12)$ for fermions, under the scheme with symmetry condition [22]. The choice of $G=\text{SO}(12)$ and $F=32$ is motivated by the study based on CSDR which leads to an $\text{SO}(10) \times \text{U}(1)$ gauge theory with one generation of fermion in 4D spacetime [29] (for $\text{SO}(12)$ GUT see also [38]).

1. A gauge symmetry and particle contents

First, we show the particle contents in 4D spacetime without parities Eq. (14)-(16). We assume that $U(1)_I$ is embedded into $SO(12)$ such as

$$SO(12) \supset SO(10) \times U(1)_I. \quad (65)$$

Thus we identify $SO(10) \times U(1)_I$ as the gauge group which satisfy the constraints Eq (22), using Eq. (30). We identify the scalar components which satisfy Eq. (23) by decomposing adjoint representation of $SO(12)$:

$$SO(12) \supset SO(10) \times U(1)_I : 66 = 45(0) + 1(0) + 10(2) + 10(-2). \quad (66)$$

According to the prescription below Eq. (30) in sec. II, the scalar components $10(2)+10(-2)$ remains in 4D spacetime. We also identify the fermion components which satisfy Eq. (25) by decomposing 32 representations of $SO(12)$ as

$$SO(12) \supset SO(10) \times U(1)_I : 32 = 16(1) + \overline{16}(-1). \quad (67)$$

According to the prescription below Eq. (32) in sec. II, we have the fermion components as $16(1)$ for a left-handed fermion and $\overline{16}(-1)$ for a right-handed fermion, respectively, in 4D spacetime.

Next, we specify the parity assignment of $P(P')$ in order to identify the gauge symmetry and particle contents that actually remain in 4D spacetime. We choose a parity assignment so as to break gauge symmetry as $SO(12) \supset SO(10) \times U(1)_I \supset SU(5) \times U(1)_X \times U(1)_I \supset SU(3) \times SU(2)_L \times U(1)_Y \times U(1)_X \times U(1)_I$, and to maintain Higgs-doublet in 4D spacetime. The parity assignment is written in 32 dimensional spinor basis of $SO(12)$ such as

$$\begin{aligned} SO(12) \supset SU(3) \times SU(2)_L \times U(1)_Y \times U(1)_X \times U(1)_I \\ 32 = (3, 2)^{(+ -)}(1, -1, 1) + (\bar{3}, 2)^{(+ -)}(-1, 1, -1) \\ + (3, 1)^{(- -)}(4, 1, -1) + (\bar{3}, 1)^{(- -)}(-4, -1, 1) \\ + (3, 1)^{(- +)}(-2, -3, -1) + (\bar{3}, 1)^{(- +)}(2, 3, 1) \\ + (1, 2)^{(+ +)}(3, -3, -1) + (1, 2)^{(+ +)}(-3, 3, 1) \\ + (1, 1)^{(- -)}(6, -1, 1) + (1, 1)^{(- -)}(-6, 1, -1) \\ + (1, 1)^{(- +)}(0, -5, 1) + (1, 1)^{(- +)}(0, 5, -1), \end{aligned} \quad (68)$$

where e.g. $(+, -)$ means that the parities (P, P') of the associated components are (even, odd). We find the gauge symmetry in 4D spacetime by surveying parity assignment for the gauge field. The parity assignments of the gauge field under $A_\mu \rightarrow PA_\mu P(P'A_\mu P')$ are:

$$\begin{aligned}
66 = & (8, 1)^{++}(0, 0, 0) + (1, 3)^{++}(0, 0, 0) + (1, 1)^{++}(0, 0, 0) \\
& + (1, 1)^{++}(0, 0, 0) + (1, 1)^{++}(0, 0, 0) \\
& + [(3, 2)^{(-+)}(-5, 0, 0) + (\bar{3}, 2)^{(-+)}(5, 0, 0) \\
& + (3, 2)^{(-)}(1, 4, 0) + (\bar{3}, 2)^{(-)}(-1, -4, 0) \\
& + (3, 1)^{(+)}(4, -4, 0) + (\bar{3}, 1)^{(+)}(-4, 4, 0) \\
& + \underline{(3, 1)^{(+)}(-2, 2, 2)} + \underline{(\bar{3}, 1)^{(+)}(2, -2, -2)} \\
& + \underline{(3, 1)^{++}(-2, 2, -2)} + \underline{(\bar{3}, 1)^{++}(2, -2, 2)} \\
& + \underline{(1, 2)^{(-)}(3, 2, 2)} + \underline{(1, 2)^{(-)}(-3, -2, -2)} \\
& + \underline{(1, 2)^{(-+)}(3, 2, -2)} + \underline{(1, 2)^{(-+)}(-3, -2, 2)} \\
& + (1, 1)^{(+)}(6, 4, 0) + (1, 1)^{(+)}(-6, -4, 0)].
\end{aligned} \tag{69}$$

The components with an underline are originated from $10(2)$ and $10(-2)$ of $SO(10) \times U(1)_I$, which do not satisfy constraints Eq. (22), and hence these components do not remain in 4D spacetime. Thus we have the gauge field with $(+, +)$ parity components without an underline in 4D spacetime, and the gauge symmetry is $SU(3) \times SU(2)_L \times U(1)_Y \times U(1)_X \times U(1)_I$.

The scalar particle contents in 4D spacetime are determined by the parity assignment,

under $\Phi_{1,2} \rightarrow -P\Phi_{1,2}P$ and $P'\Phi_{1,2}P'$:

$$\begin{aligned}
66 = & (8, 1)^{(-+)}(0, 0, 0) + (1, 3)^{(-+)}(0, 0, 0) + (1, 1)^{(-+)}(0, 0, 0) \\
& + (1, 1)^{(-+)}(0, 0, 0) + (1, 1)^{(-+)}(0, 0, 0) \\
& + [(3, 2)^{(++)}(-5, 0, 0) + (\bar{3}, 2)^{(++)}(5, 0, 0) \\
& + (3, 2)^{(+)}(1, 4, 0) + (\bar{3}, 2)^{(+)}(-1, -4, 0) \\
& + (3, 1)^{(-)}(4, -4, 0) + (\bar{3}, 1)^{(-)}(-4, 4, 0) \\
& + \underline{(3, 1)^{(-)}(-2, 2, 2) + (\bar{3}, 1)^{(-)}(2, -2, -2)} \\
& + \underline{(3, 1)^{(-)}(-2, 2, -2) + (\bar{3}, 1)^{(-)}(2, -2, 2)} \\
& + \underline{(1, 2)^{(+)}(3, 2, 2) + (1, 2)^{(+)}(-3, -2, -2)} \\
& + \underline{(1, 2)^{++}(3, 2, -2) + (1, 2)^{++}(-3, -2, 2)} \\
& + (1, 1)^{(-)}(6, 4, 0) + (1, 1)^{(-)}(-6, -4, 0)].
\end{aligned} \tag{70}$$

Note that the relative sign for the parity assignment of P is different from Eq. (69), and that the only underlined parts satisfy the constraints Eq. (23). Thus the scalar components in 4D spacetime are $(1, 2)(3, 2, -2)$ and $(1, 2)(-3, -2, 2)$.

We find massless fermion contents in 4D spacetime, by surveying the parity assignment for each components of fermion fields. We introduce two types of left-handed Weyl fermions that belong to 32 representation of $SO(12)$, which have parity assignment $\psi^{(P')} \rightarrow \gamma_5 P \psi^{(P')} (P' \psi^{(P')})$ and $\psi^{(-P')} \rightarrow \gamma_5 P \psi^{(-P')} (-P' \psi^{(-P')})$ respectively. They have the parity

assignment as

$$\begin{aligned}
32_L^{(P')} = & \underline{(3, 2)^{(--)}(1, -1, 1)_L} + \underline{(\bar{3}, 2)^{(--)}(-1, 1, -1)_L} \\
& + \underline{(\bar{3}, 1)^{(+)}(-4, -1, 1)_L} + \underline{(3, 1)^{(+)}(4, 1, -1)_L} \\
& + \underline{(\bar{3}, 1)^{(++)}(2, 3, 1)_L} + \underline{(3, 1)^{(++)}(-2, -3, -1)_L} \\
& + \underline{(1, 2)^{(-)}(-3, 3, 1)_L} + \underline{(1, 2)^{(-)}(3, -3, -1)_L} \\
& + \underline{(1, 1)^{(+)}(6, -1, 1)_L} + \underline{(1, 1)^{(+)}(-6, 1, -1)_L} \\
& + \underline{(1, 1)^{(++)}(0, -5, 1)_L} + \underline{(1, 1)^{(++)}(0, 5, -1)_L}, \tag{71}
\end{aligned}$$

$$\begin{aligned}
32_R^{(P')} = & \underline{(3, 2)^{(+)}(1, -1, 1)_R} + \underline{(\bar{3}, 2)^{(+)}(-1, 1, -1)_R} \\
& + \underline{(\bar{3}, 1)^{(--)}(-4, -1, 1)_R} + \underline{(3, 1)^{(--)}(4, 1, -1)_R} \\
& + \underline{(\bar{3}, 1)^{(-)}(2, 3, 1)_R} + \underline{(3, 1)^{(-)}(-2, -3, -1)_R} \\
& + \underline{(1, 2)^{(++)}(-3, 3, 1)_R} + \underline{(1, 2)^{(++)}(3, -3, -1)_R} \\
& + \underline{(1, 1)^{(--)}(6, -1, 1)_R} + \underline{(1, 1)^{(--)}(-6, 1, -1)_R} \\
& + \underline{(1, 1)^{(-)}(0, -5, 1)_R} + \underline{(1, 1)^{(-)}(0, 5, -1)_R}, \tag{72}
\end{aligned}$$

and

$$\begin{aligned}
32_L^{(-P')} = & \underline{(3, 2)^{(-)}(1, -1, 1)_L} + \underline{(\bar{3}, 2)^{(-)}(-1, 1, -1)_L} \\
& + \underline{(\bar{3}, 1)^{(++)}(-4, -1, 1)_L} + \underline{(3, 1)^{(++)}(4, 1, -1)_L} \\
& + \underline{(\bar{3}, 1)^{(+)}(2, 3, 1)_L} + \underline{(3, 1)^{(+)}(-2, -3, -1)_L} \\
& + \underline{(1, 2)^{(--)}(-3, 3, 1)_L} + \underline{(1, 2)^{(--)}(3, -3, -1)_L} \\
& + \underline{(1, 1)^{(++)}(6, -1, 1)_L} + \underline{(1, 1)^{(++)}(-6, 1, -1)_L} \\
& + \underline{(1, 1)^{(+)}(0, -5, 1)_L} + \underline{(1, 1)^{(+)}(0, 5, -1)_L}, \tag{73}
\end{aligned}$$

$$\begin{aligned}
32_R^{(-P')} = & \underline{(3, 2)^{(++)}(1, -1, 1)_R} + \underline{(\bar{3}, 2)^{(++)}(-1, 1, -1)_R} \\
& + \underline{(\bar{3}, 1)^{(-)}(-4, -1, 1)_R} + \underline{(3, 1)^{(-)}(4, 1, -1)_R} \\
& + \underline{(\bar{3}, 1)^{(+)}(2, 3, 1)_R} + \underline{(3, 1)^{(+)}(-2, -3, -1)_R} \\
& + \underline{(1, 2)^{(+)}(-3, 3, 1)_R} + \underline{(1, 2)^{(+)}(3, -3, -1)_R} \\
& + \underline{(1, 1)^{(-)}(6, -1, 1)_R} + \underline{(1, 1)^{(-)}(-6, 1, -1)_R} \\
& + \underline{(1, 1)^{(--)}(0, -5, 1)_R} + \underline{(1, 1)^{(--)}(0, 5, -1)_R}, \tag{74}
\end{aligned}$$

where L(R) means left-handedness(right-handedness) of fermions in 4D spacetime, and the underlined parts correspond to the components which satisfy constraints Eq. (25).

Note the relative sign for parity assignment of P between left-handed fermion and right-handed fermion, and that of P' between $32^{(P')}$ and $32^{(-P')}$. The difference between $32^{(P')}$ and $32^{(-P')}$ is allowed because of the bilinear form of the fermion sector. We thus find that the massless fermion components in 4D spacetime are one generation of SM-fermions with right-handed neutrino: $\{(3,2)(1,-1,1)_L, (3,1)(4,1,-1)_R, (3,1)(-2,-3,-1)_R, (1,2)(-3,3,1)_L, (1,1)(-6,1,-1)_R, (1,1)(0,5,-1)_R\}$.

2. The Higgs sector of the model

We analyze the Higgs-sector of our model. The Higgs-sector L_{Higgs} is the last two terms of Eq. (24):

$$L_{\text{Higgs}} = -\frac{1}{2g^2} \text{Tr}[D'_\mu \Phi_1(x) D'^\mu \Phi_1(x) + D'_\mu \Phi_2(x) D'^\mu \Phi_2(x)] - \frac{1}{2g^2} \text{Tr}[(\Phi_3 + [\Phi_1(x), \Phi_2(x)])(\Phi_3 + [\Phi_1(x), \Phi_2(x)])], \quad (75)$$

where the first term of LHS is the kinetic term of Higgs and the second term gives the Higgs potential. We then rewrite the Higgs-sector in terms of genuine Higgs field in order to analyze it.

We first note that the Φ_i s are written as

$$\Phi_i = i\phi_i = i\phi_i^a Q_a, \quad (76)$$

where Q_a s are generators of gauge group $\text{SO}(12)$, since Φ_i s are originated from gauge fields $A_\alpha = iA_\alpha^a Q_a$; for the gauge group generator we assume the normalization $\text{Tr}(Q_a Q_b) = -2\delta_{ab}$. Note that we assumed the $-i\Phi_3$ as the generator of $\text{U}(1)_I$ embedded in $\text{SO}(12)$,

$$-i\Phi_3 = Q_I. \quad (77)$$

We change the notation of the scalar fields according to Eq. (31) such that,

$$\phi_+ = \frac{1}{2}(\phi_1 + i\phi_2), \quad \phi_- = \frac{1}{2}(\phi_1 - i\phi_2), \quad (78)$$

in order to express solutions of the constraints Eq. (23) clearly. The constraints Eq. (23) is then rewritten as

$$[Q_I, \phi_+] = \phi_+, \quad [Q_I, \phi_-] = -\phi_-. \quad (79)$$

The kinetic term L_{KE} and potential $V(\phi)$ term are rewritten in terms of ϕ_+ and ϕ_- :

$$L_{KE} = -\frac{1}{g^2} \text{Tr}[D'_\mu \phi_+(x) D'^\mu \phi_-(x)], \quad (80)$$

$$V = -\frac{1}{2g^2} \text{Tr}[Q_I^2 - 4Q_I[\phi_+, \phi_-] + 4[\phi_+, \phi_-][\phi_+, \phi_-]], \quad (81)$$

where covariant derivative D'_μ is $D'_\mu \phi_\pm = \partial_\mu \phi_\pm - [A_\mu, \phi_\pm]$.

Next, we change the notation of SO(12) generators Q_a according to decomposition Eq (69) such that

$$\begin{aligned} Q_G = \{ & Q_i, Q_\alpha, Q_Y, Q, Q_I, Q_{ax(-500)}, Q^{ax(500)} \\ & Q_{ax(140)}, Q^{ax(-1-40)}, Q_{a(4-40)}, Q^{a(-440)} \\ & Q_{a(-22-2)}, Q^{a(2-22)}, Q_{a(-222)}, Q^{a(2-2-2)} \\ & Q_{x(322)}, Q^{x(-3-2-2)}, Q_{x(32-2)}, Q^{x(-3-22)} \\ & Q(640), Q(-6-40)\}, \end{aligned} \quad (82)$$

where the order of generators corresponds to Eq (69), index $i = 1 - 8$ corresponds to SU(3) adjoint rep, index $\alpha = 1 - 3$ corresponds to SU(2) adjoint rep, index $a = 1 - 3$ corresponds to SU(3)-triplet, and index $x = 1, 2$ corresponds to SU(2)-doublet. We write ϕ_\pm in terms of the genuine Higgs field ϕ_x which belongs to $(1,2)(3,2,-2)$, such that

$$\phi_+ = \phi_x Q^{x(-3-22)} \quad (83)$$

$$\phi_- = \phi^x Q_{x(32-2)}, \quad (84)$$

where $\phi^x = (\phi_x)^\dagger$. We also write gauge field $A_\mu(x)$ in terms of Q s in Eq. (108) as

$$A_\mu(x) = i(A_\mu^i Q_i + A_\mu^\alpha Q_\alpha + B_\mu Q_Y + C_\mu Q + E_\mu Q_I). \quad (85)$$

We then need commutation relations of $Q^{x(-3-22)}$, $Q_{x(32-2)}$, Q_α , Q_Y , Q and Q_I in order to analyze the Higgs sector; we summarized them in Table II.

Finally, we obtain the Higgs sector with genuine Higgs field by substituting Eq. (139)-(114) into Eq. (80, 81) and rescaling the fields $\phi \rightarrow g/\sqrt{2}\phi$ and $A_\mu \rightarrow gA_\mu$, and the couplings $\sqrt{2}g = g_2$ and $\sqrt{6/5}g = g_Y$,

$$L_{Higgs} = |D_\mu \phi_x|^2 - V(\phi), \quad (86)$$

$[Q^{x(-3-22)}, Q_{y(32-2)}] = -\sqrt{\frac{3}{10}} \delta_y^x Q_Y + -\sqrt{\frac{1}{5}} \delta_y^x Q + \delta_y^x Q_I + \frac{1}{\sqrt{2}} (\sigma_\alpha^*)^x_y Q_\alpha$	
$[Q_\alpha, Q_x] = -\frac{1}{\sqrt{2}} (\sigma_\alpha)_x^y Q_y$	$[Q_\alpha, Q^x] = \frac{1}{\sqrt{2}} (\sigma_\alpha^*)^x_y Q^y$
$[Q_x, Q_y] = 0$	$[Q_Y, Q^x] = -\sqrt{\frac{3}{10}} Q^x$
$[Q, Q^x] = -\sqrt{\frac{1}{5}} Q^x$	$[Q_I, Q^x] = Q^x$

TABLE I: commutation relations of $Q^{x(-3-22)}$, $Q_{x(32-2)}$, Q_α , Q_Y , Q and Q_I

where the covariant derivative $D_\mu \phi_x$ and potential $V(\phi)$ are

$$D_\mu \phi_x = \partial_\mu \phi_x + ig_2 \frac{1}{2} (\sigma_\alpha)_x^y A_{\alpha\mu} \phi_y + ig_Y \frac{1}{2} B_\mu \phi_x + i\sqrt{\frac{1}{5}} g C_\mu \phi_x - ig E_\mu \phi_x, \quad (87)$$

$$V = -\frac{2}{R^2} \phi^x \phi_x + \frac{3g^2}{2} (\phi^x \phi_x)^2, \quad (88)$$

respectively. Notice that we explicitly write radius R of S^2 in the Higgs potential, and that we omitted the constant term in the Higgs potential. We note that the $SU(2)_L \times U(1)_Y$ parts of the Higgs sector has the same form as the SM Higgs sector. Therefore we obtain the electroweak symmetry breaking $SU(2)_L \times U(1)_Y \rightarrow U(1)_{EM}$. The Higgs field ϕ^x acquires vaccume expectation value(VEV) as

$$\langle \phi \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}, \quad (89)$$

$$v = \sqrt{\frac{4}{3}} \frac{1}{gR}, \quad (90)$$

and W boson mass m_W and Higgs mass m_H are given in terms of radius R

$$m_W = g_2 \frac{v}{2} = \sqrt{\frac{2}{3}} \frac{1}{R}, \quad (91)$$

$$m_H = \sqrt{3} g v = \sqrt{4} \frac{1}{R}. \quad (92)$$

The ratio between m_W and m_H is predicted

$$\frac{m_H}{m_W} = \sqrt{6}. \quad (93)$$

We thus find $m_H \sim 196 \text{ GeV}$ in this model. We also find the Weinberg angle is given by

$$\begin{aligned} \sin^2 \theta_W &= \frac{g_Y^2}{g_2^2 + g_Y^2} \\ &= \frac{6/5}{2 + 6/5} \\ &= \frac{3}{8} \end{aligned} \quad (94)$$

which is same as $SU(5)$ GUT case.

B. The E_6 model without symmetry condition

Here we show a model based on a gauge group $G=E_6$ and a representation 27 of E_6 for fermions, under the scheme without symmetry condition [23].

1. Gauge group reduction

We consider the following gauge group reduction

$$\begin{aligned}
E_6 &\supset SO(10) \times U(1)_I \\
&\supset SU(5) \times U(1)_X \times U(1)_I \\
&\supset SU(3) \times SU(2) \times U(1)_Y \times U(1)_X \times U(1)_I .
\end{aligned} \tag{95}$$

The background gauge field in Eq. (43) is chosen to belong to the $U(1)_I$ group. This choice is needed in order to obtain chiral SM fermions in 4D spacetime to be discussed later. There are two other symmetry reduction schemes. One can prove that the results in those two schemes are effectively the same as the one considered here once we require the correct $U(1)$ combinations for the hypercharge and the background field.

We then impose the parity assignments with respect to the fixed points, Eqs. (11)-(16). The parity assignments for the fundamental representation of E_6 is chosen to be

$$\begin{aligned}
\mathbf{27} = & (1, 2)(-3, -2, -2)^{(+,+)} + (1, 2)(3, 2, -2)^{(-,-)} + (1, 2)(-3, 3, 1)^{(+,-)} \\
& + (1, 1)(6, -1, 1)^{(+,+)} + (1, 1)(0, -5, 1)^{(-,-)} + (1, 1)(0, 0, 4)^{(-,+)} \\
& + (3, 2)(1, -1, 1)^{(-,+)} + (3, 1)(-2, 2, -2)^{(+,-)} + (\bar{3}, 1)(-4, -1, 1)^{(+,+)} \\
& + (\bar{3}, 1)(2, 3, 1)^{(+,+)} + (\bar{3}, 1)(2, -2, 2)^{(-,+)} ,
\end{aligned} \tag{96}$$

where, for example, $(+, -)$ means that the parities under P_1 and P_2 are (even, odd). By the requirement of consistency, we find that the components of A_μ in the adjoint representation

have the parities under $A_\mu \rightarrow P_1 A_\mu P_1$ ($P_2 A_\mu P_2$) as follows:

$$\begin{aligned}
\mathbf{78}|_{A_\mu} = & \underline{(8, 1)(0, 0, 0)^{(+, +)} + (1, 3)(0, 0, 0)^{(+, +)}} \\
& + \underline{(1, 1)(0, 0, 0)^{(+, +)} + (1, 1)(0, 0, 0)^{(+, +)} + (1, 1)(0, 0, 0)^{(+, +)}} \\
& + (3, 2)(-5, 0, 0)^{(-, +)} + (\bar{3}, 2)(5, 0, 0)^{(-, +)} \\
& + (3, 2)(1, 4, 0)^{(+, -)} + (\bar{3}, 2)(-1, -4, 0)^{(+, -)} \\
& + (3, 1)(4, -4, 0)^{(-, -)} + (\bar{3}, 1)(-4, 4, 0)^{(-, -)} \\
& + (1, 1)(-6, -4, 0)^{(-, -)} + (1, 1)(6, 4, 0)^{(-, -)} \\
& + (3, 2)(1, -1, -3)^{(+, +)} + (\bar{3}, 2)(-1, 1, 3)^{(+, +)} \\
& + (3, 1)(4, 1, 3)^{(-, +)} + (\bar{3}, 1)(-4, -1, -3)^{(-, +)} \\
& + (3, 1)(-2, -3, 3)^{(+, -)} + (\bar{3}, 1)(2, 3, -3)^{(+, -)} \\
& + (1, 2)(-3, 3, -3)^{(-, -)} + (1, 2)(3, -3, 3)^{(-, -)} \\
& + (1, 1)(-6, 1, 3)^{(-, +)} + (1, 1)(6, -1, -3)^{(-, +)} \\
& + (1, 1)(0, -5, -3)^{(+, -)} + (1, 1)(0, 5, 3)^{(+, -)}, \tag{97}
\end{aligned}$$

where the underlined components correspond to the adjoint representations of $SU(3) \times SU(2) \times U(1)_Y \times U(1)_X \times U(1)_I$, respectively. We note that the components with parity $(+, +)$ can have massless zero modes in 4D spacetime. Such components include the adjoint representations of $SU(3) \times SU(2) \times U(1)^3$, $(3, 2)(1, -1, -3)$ and its conjugate. The latter components seem problematic since they do not appear in the low-energy spectrum due to non-zero $U(1)_I$ charge.

2. Scalar field contents in 4D spacetime

With the parity assignments with respect to the fixed points, Eqs. (12) and (15), we have for the A_θ and A_ϕ fields

$$\begin{aligned}
\mathbf{78}|_{A_{\theta,\phi}} = & (8, 1)(0, 0, 0)^{(-,-)} + (1, 3)(0, 0, 0)^{(-,-)} \\
& + (1, 1)(0, 0, 0)^{(-,-)} + (1, 1)(0, 0, 0)^{(-,-)} + (1, 1)(0, 0, 0)^{(-,-)} \\
& + (3, 2)(-5, 0, 0)^{(+,-)} + (\bar{3}, 2)(5, 0, 0)^{(+,-)} \\
& + (3, 2)(1, 4, 0)^{(-,+)} + (\bar{3}, 2)(-1, -4, 0)^{(-,+)} \\
& + (3, 1)(4, -4, 0)^{(+,+)} + (\bar{3}, 1)(-4, 4, 0)^{(+,+)} \\
& + (1, 1)(-6, -4, 0)^{(+,+)} + (1, 1)(6, 4, 0)^{(+,+)} \\
& + (3, 2)(1, -1, -3)^{(-,-)} + (\bar{3}, 2)(-1, 1, 3)^{(-,-)} \\
& + (3, 1)(4, 1, 3)^{(+,-)} + (\bar{3}, 1)(-4, -1, -3)^{(+,-)} \\
& + (3, 1)(-2, -3, 3)^{(-,+)} + (\bar{3}, 1)(2, 3, -3)^{(-,+)} \\
& + (1, 2)(-3, 3, -3)^{(+,+)} + (1, 2)(3, -3, 3)^{(+,+)} \\
& + (1, 1)(-6, 1, 3)^{(+,-)} + (1, 1)(6, -1, -3)^{(+,-)} \\
& + (1, 1)(0, -5, -3)^{(-,+)} + (1, 1)(0, 5, 3)^{(-,+)} .
\end{aligned} \tag{98}$$

Components with $(+, -)$ or $(-, +)$ parity do not have KK modes since they are odd under $\phi \rightarrow \phi + 2\pi$ and the KK modes of gauge field are specified by integer angular momentum quantum numbers ℓ and m on the two-sphere. We then concentrate on the components which have either $(+, +)$ or $(-, -)$ parity and nonzero $U(1)_I$ charges as the candidate for the Higgs field. These include $\{(1, 2)(3, -3, 3) + \text{h.c.}\}$ and $\{(3, 2)(1, -1, -3) + \text{h.c.}\}$ with parities $(+, +)$ and $(-, -)$, respectively. The representations $(1, 2)(-3, 3, -3)$ and $(1, 2)(3, -3, 3)$ have the correct quantum numbers for the SM Higgs doublet. Therefore, we identify the $(1, 1)$ mode of these components as the SM Higgs fields in 4D spacetime.

3. Chiral fermion contents in 4D spacetime

In our model, we choose the fermions as the Weyl fermions Ψ_- belonging to the **27** representation of E_6 . The **27** representation is decomposed as in Eq. (96) under the group reduction, Eq. (95). In this decomposition, we find that our choice of the background gauge

field of $U(1)_I$ is suitable for obtaining massless fermions since all such components have $U(1)_I$ charge 1. In the fundemantal representation, the $U(1)_I$ generator is

$$Q_I = \frac{1}{6} \text{diag}(-2, -2, -2, -2, 1, 1, 1, 1, 4, 1, 1, 1, 1, 1, 1, -2, -2, -2, 1, 1, 1, 1, 1, 1, -2, -2, -2) ,$$

according to the decomposition Eq. (96). By identifying $Q = 3Q_I$, we readily obtain the condition

$$Q\Psi_- = \frac{1}{2}\Psi_- . \quad (99)$$

Therefore, the chiral fermions ψ_L in 4D spacetime have zero modes.

Next, we consider the parity assignments for the fermions with respect to the fixed points of S^2/Z_2 . The boundary conditions are given by Eqs. (13) and (16). It turns out that four **27** fermion copies with different boundary conditions are needed in order to obtain an entire generation of massless SM fermions. They are denoted by $\Psi^{(1,2,3,4)}$ with the following parity assignments

$$\Psi_{\pm}^{(i)}(x, \pi - \theta, -\phi) = \xi \gamma_5 P_1 \Psi_{\pm}^{(i)}(x, \theta, \phi) , \quad (100)$$

$$\Psi_{\pm}^{(i)}(x, \pi - \theta, 2\pi - \phi) = \eta \gamma_5 P_2 \Psi_{\pm}^{(i)}(x, \theta, \phi) , \quad (101)$$

where γ_5 is the chirality operator, and $(\xi, \eta) = (+, +), (-, -), (+, -)$ and $(-, +)$ for $i =$

1, 2, 3, 4, respectively. From these fermions we find that $\psi_{1,2,3,4}$ have the parity assignments

$$\begin{aligned}
\mathbf{27}_{\psi_L^{(1)}} = & (1, 2)(-3, -2, -2)^{(-,-)} + (1, 2)(3, 2, -2)^{(+,+)} + (1, 2)(-3, 3, 1)^{(-,+)} \\
& + (1, 1)(6, -1, 1)^{(-,-)} + \underline{(1, 1)(0, -5, 1)^{(+,+)}} + (1, 1)(0, 0, 4)^{(+,-)} \\
& + (3, 2)(1, -1, 1)^{(+,-)} + (3, 1)(-2, 2, -2)^{(-,+)} + (\bar{3}, 1)(-4, -1, 1)^{(-,-)} \\
& + (\bar{3}, 1)(2, 3, 1)^{(-,-)} + (\bar{3}, 1)(2, -2, 2)^{(+,-)} \tag{102}
\end{aligned}$$

$$\begin{aligned}
\mathbf{27}_{\psi_L^{(2)}} = & (1, 2)(-3, -2, -2)^{(+,+)} + (1, 2)(3, 2, -2)^{(-,-)} + (1, 2)(-3, 3, 1)^{(+,-)} \\
& + \underline{(1, 1)(6, -1, 1)^{(+,+)}} + (1, 1)(0, -5, 1)^{(-,-)} + (1, 1)(0, 0, 4)^{(-,+)} \\
& + (3, 2)(1, -1, 1)^{(-,+)} + (3, 1)(-2, 2, -2)^{(+,-)} + \underline{(\bar{3}, 1)(-4, -1, 1)^{(+,+)}} \\
& + \underline{(\bar{3}, 1)(2, 3, 1)^{(+,+)}} + (\bar{3}, 1)(2, -2, 2)^{(-,+)} \tag{103}
\end{aligned}$$

$$\begin{aligned}
\mathbf{27}_{\psi_L^{(3)}} = & (1, 2)(-3, -2, -2)^{(-,+)} + (1, 2)(3, 2, -2)^{(+,-)} + (1, 2)(-3, 3, 1)^{(-,-)} \\
& + (1, 1)(6, -1, 1)^{(-,+)} + (1, 1)(0, -5, 1)^{(+,-)} + (1, 1)(0, 0, 4)^{(+,+)} \\
& + \underline{(3, 2)(1, -1, 1)^{(+,+)}} + (3, 1)(-2, 2, -2)^{(-,-)} + (\bar{3}, 1)(-4, -1, 1)^{(-,+)} \\
& + (\bar{3}, 1)(2, 3, 1)^{(-,+)} + (\bar{3}, 1)(2, -2, 2)^{(+,+)} \tag{104}
\end{aligned}$$

$$\begin{aligned}
\mathbf{27}_{\psi_L^{(4)}} = & (1, 2)(-3, -2, -2)^{(+,-)} + (1, 2)(3, 2, -2)^{(-,+)} + \underline{(1, 2)(-3, 3, 1)^{(+,+)}} \\
& + (1, 1)(6, -1, 1)^{(+,-)} + (1, 1)(0, -5, 1)^{(-,+)} + (1, 1)(0, 0, 4)^{(-,-)} \\
& + (3, 2)(1, -1, 1)^{(-,-)} + (3, 1)(-2, 2, -2)^{(+,+)} + (\bar{3}, 1)(-4, -1, 1)^{(+,-)} \\
& + (\bar{3}, 1)(2, 3, 1)^{(+,-)} + (\bar{3}, 1)(2, -2, 2)^{(-,-)} , \tag{105}
\end{aligned}$$

where the underlined components have even parities and $U(1)_I$ charge 1. One can readily identify one generation of SM fermions, including a right-handed neutrino, as the zero modes of these components.

A long-standing problem in the gauge-Higgs unification framework is the Yukawa couplings of the Higgs boson to the matter fields. Here we discuss about the Yukawa couplings in our model. As mentioned before, the SM Higgs is the $(\ell = 1, |m| = 1)$ KK mode of the extra-spatial component of the gauge field, the Yukawa term at tree level has the following form

$$L_{\text{Yukawa}} \supset \bar{\psi}_L^{00} \Phi^{11} \psi_R^{\ell 1} + \bar{\psi}_L^{\ell 1} \Phi^{11} \psi_R^{00} + \text{h.c.} , \tag{106}$$

where $\psi^{\ell m}$ s are the fermionic KK modes with the $(\ell = 0, m = 0)$ modes appearing as the chiral fermions and Φ^{11} denotes the SM Higgs field. We here identify the left-handed fermionic zero modes as $SU(2)$ doublets and the right-handed fermionic zero modes as $SU(2)$

singlets, as in the SM. Therefore, the $(\ell, |m| = 1)$ modes and the $(\ell = 0, |m| = 0)$ modes mix after spontaneous symmetry breaking. One needs to diagonalize the mass terms to obtain physical eigenstates. The Yukawa couplings in our model are thus more complicated than other gauge-Higgs unification models. However, the difficulty of obtaining the realistic fermion mass spectrum remains since the Yukawa couplings all arise from gauge interactions. One way to get realistic Yukawa couplings is to introduce SM fermions localized at an orbifold fixed point and make use of nonlocal interactions with Wilson lines [4]. Another possible solution is to consider fermions in 6D spacetime belonging to a higher dimensional representation of the original E_6 gauge group, rendering more than one generation of SM fermions. In that case, mixing among generations will be obtained from gauge interactions and is given by Clebsch-Gordan coefficients. We expect that realistic Yukawa couplings could be obtained using a combination of these methods. A detailed analysis of this issue is beyond the scope of the paper and left for a future work.

4. Higgs potential of the model

Here we analyze the Higgs potential for the E_6 model. To further simplify the Higgs potential, we need to find out the algebra of the gauge group generators. Note that the E_6 generators are chosen according to the decomposition of the adjoint representation given in Eq. (97)

$$\begin{aligned}
& \{Q_i, Q_\alpha, Q_Y, Q_X, Q_I, \\
& Q_{ax(-5,0,0)}, Q^{ax(5,0,0)}, Q_{ax(1,4,0)}, Q^{ax(-1,-4,0)}, \\
& Q_{a(4,-4,0)}, Q^{a(-4,4,0)}, Q_{(-6,-4,0)}, Q_{(6,4,0)}, \\
& Q_{ax(1,-1,-3)}, Q^{ax(-1,1,3)}, Q_{a(4,1,3)}, Q^{a(-4,-1,-3)}, \\
& Q_{a(-2,-3,3)}, Q^{a(2,3,-3)}, Q_{x(3,-3,3)}, Q^{x(-3,3,-3)}, \\
& Q_{(-6,1,3)}, Q_{(6,-1,-3)}, Q_{(0,-5,-3)}, Q_{(0,5,3)}\} ,
\end{aligned} \tag{107}$$

$$\begin{aligned}
& [Q_{x(3,-3,3)}, Q^{y(-3,3,-3)}] = \frac{1}{2}\delta_x^y Q_I - \frac{1}{2}\sqrt{\frac{3}{5}}\delta_x^y Q_X + \frac{1}{\sqrt{10}}\delta_x^y Q_Y + \frac{1}{\sqrt{6}}(\sigma_\alpha)_x^y Q_\alpha \\
& [Q_\alpha, Q_{x(3,-3,3)}] = \frac{1}{\sqrt{6}}(\sigma_\alpha)_x^y Q_{y(3,-3,3)} \quad [Q_\alpha, Q^{x(-3,3,-3)}] = -\frac{1}{\sqrt{6}}(\sigma_\alpha^*)_x^y Q^{y(-3,3,-3)} \\
& [Q_{x(3,-3,3)}, Q_{y(3,-3,3)}] = 0 \quad [Q_I, Q_{x(3,-3,3)}] = \frac{1}{2}Q_{x(3,-3,3)} \\
& [Q_X, Q_{x(3,-3,3)}] = -\frac{1}{2}\sqrt{\frac{3}{5}}Q_{x(3,-3,3)} \quad [Q_Y, Q_{x(3,-3,3)}] = \frac{1}{\sqrt{10}}Q_{x(3,-3,3)}
\end{aligned}$$

TABLE II: Commutation relations of Q_α , $Q_{X,Y,Z}$, $Q_{x(3,-3,3)}$ and $Q^{x(-3,3,-3)}$, where σ_i are the Pauli matrices.

where the generators are listed in the corresponding order of the terms in Eq. (97) and the indices

$$i = 1, \dots, 8 : \text{SU}(3) \text{ adj rep index} \Rightarrow Q_i : \text{SU}(3) \text{ generators} , \quad (108)$$

$$\alpha = 1, 2, 3 : \text{SU}(2) \text{ adj rep index} \Rightarrow Q_\alpha : \text{SU}(2) \text{ generators} , \quad (109)$$

$$Q_{X,Y,Z} : U(1)_{X,Y,Z} \text{ generators} , \quad (110)$$

$$x = 1, 2 : \text{SU}(2) \text{ doublet index} , \quad (111)$$

$$a = 1, 2, 3 : \text{SU}(3) \text{ color index} . \quad (112)$$

Here we take the standard normalization for generators, $\text{Tr}[QQ^\dagger] = 2$. The Higgs fields are in the representations of $(1, 2)(3, -3, 3)$ and $(1, 2)(-3, 3, -3)$. We write

$$\Phi(x) = \phi^x Q_{x(3,-3,3)} \quad (\Phi^\dagger(x) = \phi_x Q^{x(-3,3,-3)}) . \quad (113)$$

Likewise, the gauge field $A_\mu(x)$ in terms of the Q 's in Eq. (108) is

$$A_\mu(x) = A_\mu^i Q_i + A_\mu^\alpha Q_\alpha + B_\mu Q_Y + C_\mu Q_X + E_\mu Q_I. \quad (114)$$

The commutation relations between the generators Q_α , $Q_{X,Y,Z}$, $Q_{x(3,-3,3)}$ and $Q^{x(-3,3,-3)}$ are summarized in Table. II.

Finally, we obtain the Lagrangian associated with the Higgs field by applying Eqs. (113, 114) to Eqs. (63, 64) and carrying out the trace. Furthermore, to obtain the canonical form of kinetic terms, the Higgs field, the gauge field, and the gauge coupling need to be rescaled

in the following way:

$$\phi \rightarrow \frac{g}{\sqrt{2}}\phi \quad (115)$$

$$A_\mu \rightarrow \frac{g}{R}A_\mu \quad (116)$$

$$\frac{g}{\sqrt{6\pi R^2}} = g_2 , \quad (117)$$

where g_2 denotes the SU(2) gauge coupling. The Higgs sector is then given by

$$\mathcal{L}_{\text{Higgs}} = |D_\mu \phi|^2 - V(\phi) \quad (118)$$

where

$$D_\mu \phi = \left[\partial_\mu + ig_2 \frac{\sigma_\alpha}{2} A_{\alpha\mu} + ig \frac{1}{\sqrt{40\pi R^2}} B_\mu - ig \frac{1}{2} \sqrt{\frac{3}{20\pi R^2}} C_\mu + ig \frac{1}{2\sqrt{4\pi R^2}} E_\mu \right] \phi , \quad (119)$$

$$V = -\frac{\chi}{8R^2} \phi^\dagger \phi + \frac{3g^2}{40\pi R^2} (\phi^\dagger \phi)^2 , \quad (120)$$

where $\chi = 7 + 9\mu_1 + 9\mu_2$. We have omitted the constant term in the Higgs potential. Comparing the potential derived above with the standard form $\mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2$ in the SM, we see that the model has a tree-level μ^2 term that is negative and proportional to R^{-2} . Moreover, the quartic coupling $\lambda = 3g^2/(40\pi R^2)$ is related to the 6D gauge coupling g and grants perturbative calculations because it is about 0.16, using the value of R to be extracted in the next section. Therefore, the order parameter in this model is controlled by a single parameter R , the compactification scale.

In fact, the $(1, 1)$ mode of the $\{(3, 2)(1, -1, -3) + \text{h.c.}\}$ representation also has a negative squared mass term because it has the same Q_I charge as the $\{(1, 2)(3, -3, 3) + \text{h.c.}\}$ representation. Therefore, it would induce not only electroweak symmetry breaking but also color symmetry breaking. This undesirable feature can be cured by adding brane terms

$$\frac{\alpha}{R^2 \sin^2 \theta} (F_{\theta\phi}^a F^{a\theta\phi})^2 \delta\left(\theta - \frac{\pi}{2}\right) [\delta(\phi) + \delta(\phi - \pi)] , \quad (121)$$

where a denotes the group index of the $\{(3, 2)(1, -1, -3) + \text{h.c.}\}$ representation. These brane terms preserve the Z'_2 symmetry which corresponds to the symmetry under the transformation $(\phi \rightarrow \phi + \pi)$. With an appropriate choice of the dimensionless constant α , the squared mass of the $(1, 1)$ can be lifted to become positive and sufficiently large.

Due to a negative mass term, the Higgs potential in Eq. (120) can induce the spontaneous symmetry breakdown: $SU(2) \times U(1)_Y \rightarrow U(1)_{EM}$ in the SM. The Higgs field acquires a vacuum expectation value (VEV)

$$\langle \phi \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix} \text{ with } v = \sqrt{\frac{5\pi\chi}{3}} \frac{1}{g} \simeq \frac{4.6}{g}. \quad (122)$$

One immediately finds that the W boson mass

$$m_W = \frac{g_2}{2} v = \frac{1}{6} \sqrt{\frac{5\chi}{2}} \frac{1}{R} \simeq \frac{0.53}{R}, \quad (123)$$

from which the compactification scale $R^{-1} \simeq 152$ GeV is inferred. Moreover, the Higgs boson mass at the tree level is

$$m_H = \sqrt{\frac{3}{20\pi}} \frac{gv}{R} = 3\sqrt{\frac{2}{5}} m_W = \frac{\sqrt{\chi}}{2} \frac{1}{R}, \quad (124)$$

which is about 152 GeV, numerically very close to the compactification scale. Since the hypercharge of the Higgs field is $1/2$, the $U(1)_Y$ gauge coupling is derived from Eq. (119) as

$$g_Y = \frac{g}{\sqrt{10\pi R^2}}. \quad (125)$$

The Weinberg angle is thus given by

$$\sin^2 \theta_W = \frac{g_Y^2}{g_2^2 + g_Y^2} = \frac{3}{8}, \quad (126)$$

and the Z boson mass

$$m_Z = \frac{m_W}{\cos \theta_W} = m_W \sqrt{\frac{8}{5}}, \quad (127)$$

both at the tree level. These relations are the same as the $SU(5)$ GUT at the unification scale. This is not surprising because this part only depends on the group structure.

5. *KK mode spectrum of each field*

Since we did not impose symmetry condition, we have KK mode for each field in this model. Here we show KK mass spectrum under the existence of background field for our E_6 model. The masses are basically controlled by the compactification radius R of the two-sphere. They receive two kinds of contributions: one arising from the angular momentum in the S^2 space, and the other coming from the interactions with the background field.

The KK masses for fermions have been given in Refs. [36, 39, 40]. We give them in terms of our notation here:

$$M_{\ell m}^{KK}(\psi_L) = \frac{1}{R} \sqrt{\ell(\ell+1) - \frac{4q^2-1}{4}}, \quad (128)$$

where q is proportional to the $U(1)_I$ charge of fermion and determined by the action of $Q = 3Q_I$ on fermions as $Q\Psi = q\Psi = 3q_I\Psi$. Note that the mass does not depend on the quantum number m . The lightest KK mass, corresponding to $\ell = 1$ and $q_I = 1/6$, is about 214 GeV at the tree level. The range of ℓ is

$$\frac{2q \pm 1}{2} \leq \ell \quad (+ : \text{ for } \psi_{R(L)} \text{ in } \Psi_{+(-)}, \quad - : \text{ for } \psi_{L(R)} \text{ in } \Psi_{-(+)}). \quad (129)$$

We thus can have zero mode for $Q\Psi = \pm \frac{1}{2}\Psi$, where this condition is given in Eq. (51).

For the 4D gauge field A_μ , its kinetic term and KK mass term are obtained from the terms

$$L = \int d\Omega Tr \left[-\frac{1}{4} F_{\mu\nu}^2 + \frac{1}{2R^2} F_{\mu\theta} F_\theta^\mu + \frac{1}{2R^2 \sin^2 \theta} F_{\mu\phi} F_\phi^\mu \right]. \quad (130)$$

Taking terms quadratic in A_μ , we get

$$L_{\text{quad}} = \int d\Omega Tr \left[-\frac{1}{4} (\partial_\mu A_\nu - \partial_\nu A_\mu) (\partial^\mu A^\nu - \partial^\nu A^\mu) + \frac{1}{2R^2} \partial_\theta A_\mu \partial_\theta A^\mu \right. \\ \left. + \frac{1}{2R^2 \sin^2 \theta} \partial_\phi A_\mu \partial_\phi A^\mu - \frac{1}{2R^2} [A_\mu, \tilde{A}_\phi^B] [A^\mu, \tilde{A}_\phi^B] \right], \quad (131)$$

where \tilde{A}_ϕ^B is the background gauge field given in Eq. (43). The KK expansion of A_μ is

$$A_\mu = \sum_{\ell m} A_\mu^{\ell m}(x) Y_{\ell m}^\pm(\theta, \phi) \quad (132)$$

where $Y_{\ell m}^\pm(\theta, \phi)$ are the linear combinations of spherical harmonics satisfying the boundary condition $Y_{\ell m}^\pm(\pi - \theta, -\phi) = \pm Y_{\ell m}^\pm(\theta, \phi)$. Their explicit forms are [39]

$$Y_{\ell m}^+(\theta, \phi) \equiv \frac{(i)^{\ell+m}}{\sqrt{2}} [Y_{\ell m}(\theta, \phi) + (-1)^\ell Y_{\ell -m}(\theta, \phi)] \quad \text{for } m \neq 0 \quad (133)$$

$$Y_{\ell m}^-(\theta, \phi) \equiv \frac{(i)^{\ell+m+1}}{\sqrt{2}} [Y_{\ell m}(\theta, \phi) - (-1)^\ell Y_{\ell -m}(\theta, \phi)] \quad \text{for } m \neq 0 \quad (134)$$

$$Y_{\ell 0}^{+(-)}(\theta) \equiv \begin{cases} Y_{\ell 0}(\theta) & \text{for } m = 0 \text{ and } \ell = \text{even (odd)} \\ 0 & \text{for } m = 0 \text{ and } \ell = \text{odd (even)}. \end{cases} \quad (135)$$

Note that we do not have KK mode functions that are odd under $\phi \rightarrow \phi + 2\pi$ since the KK modes are specified by the integer angular momentum quantum numbers ℓ and m of gauge

field A_M on the two-sphere. Thus, the components of A_μ and $A_{\theta,\phi}$ with $(+, -)$ or $(-, +)$ parities do not have corresponding KK modes. Applying the KK expansion and integrating about $d\Omega$, we obtain the kinetic and KK mass terms for the KK modes of A_μ

$$\begin{aligned}
L_M = & -\frac{1}{2} [\partial_\mu A_\nu^{\ell m}(x) - \partial_\nu A_\mu^{\ell m}(x)] [\partial^\mu A^{\ell m \nu}(x) - \partial^\nu A^{\ell m \mu}(x)] \\
& + \frac{\ell(\ell+1)}{R^2} A_\mu^{\ell m}(x) A^{\ell m \mu}(x) \\
& + \frac{9q_I^2}{R^2} \left[\int d\Omega \frac{(\cos \theta \pm 1)^2}{\sin^2 \theta} (Y_{\ell m}^\mp)^2 \right] A_\mu^{\ell m}(x) A^{\ell m \mu}(x) ,
\end{aligned} \tag{136}$$

where we have used $Tr[Q_i Q^i] = 2$ and $[A_\mu(x), Q_I] = q_I(A_\mu^i(x)Q_i - A_{i\mu}(x)Q^i)$. Therefore, the KK masses of A_μ are

$$M_{\ell m}^{KK}(A_\mu) = \frac{1}{R} \sqrt{\ell(\ell+1) + (m_{\ell m}^B)^2} , \tag{137}$$

$$(m_{\ell m}^B)^2 = 9q_I^2 \int d\Omega \frac{(\cos \theta \pm 1)^2}{\sin^2 \theta} (Y_{\ell m}^\mp)^2 , \tag{138}$$

where $m_{\ell m}^B$ corresponds to the contribution from the background gauge field. Note that Eq. (137) agrees with Eq. (49) when $\ell = 0$. Also, since the SM gauge bosons have $q_I = 0$, their KK masses are simply $\sqrt{\ell(\ell+1)}/R$ at the tree level.

The kinetic and KK mass terms of A_θ and A_ϕ are obtained from the terms in the higher dimensional gauge sector

$$\begin{aligned}
L = & \frac{1}{2g^2} \int d\Omega \left\{ \left(Tr[(\partial_\mu A_\theta - i[A_\mu, A_\theta])^2] + Tr[(\partial_\mu A_\theta - i[A_\mu, \tilde{A}_\phi])^2] \right) \right. \\
& - \frac{1}{R^2} Tr \left[\left(\frac{1}{\sin \theta} \partial_\theta (\sin \theta \tilde{A}_\phi + \sin \theta \tilde{A}_\phi^B) \right. \right. \\
& \left. \left. - \frac{1}{\sin \theta} \partial_\phi A_\theta - i[A_\theta, \tilde{A}_\phi + \tilde{A}_\phi^B] \right)^2 \right] \left. \right\} .
\end{aligned} \tag{139}$$

The first line on the right-hand side of Eq. (139) corresponds to the kinetic terms, and the second line corresponds to the potential term. Applying the background gauge field Eq. (43), the potential becomes

$$L_V = -\frac{1}{2g^2 R^2} \int d\Omega Tr \left[\left(\frac{1}{\sin \theta} \partial_\theta (\sin \theta \tilde{A}_\phi) + Q - \frac{1}{\sin \theta} \partial_\phi A_\theta - i[A_\theta, \tilde{A}_\phi + \tilde{A}_\phi^B] \right)^2 \right] \tag{140}$$

For A_θ and A_ϕ we use the following KK expansions to obtain the KK mass terms,

$$A_\theta(x, \theta, \phi) = \sum_{\ell m (\neq 0)} \frac{-1}{\sqrt{\ell(\ell+1)}} [\Phi_1^{\ell m}(x) \partial_\theta Y_{\ell m}^\pm(\theta, \phi) + \Phi_2^{\ell m}(x) \frac{1}{\sin \theta} \partial_\phi Y_{\ell m}^\pm(\theta, \phi)] , \quad (141)$$

$$A_\phi(x, \theta, \phi) = \sum_{\ell m (\neq 0)} \frac{1}{\sqrt{\ell(\ell+1)}} [\Phi_2^{\ell m}(x) \partial_\theta Y_{\ell m}^\pm(\theta, \phi) - \Phi_1^{\ell m}(x) \frac{1}{\sin \theta} \partial_\phi Y_{\ell m}^\pm(\theta, \phi)] , \quad (142)$$

where the factor of $1/\sqrt{\ell(\ell+1)}$ is needed for normalization. These particular forms are convenient in giving diagonalized KK mass terms [39]. Applying the KK expansions Eq. (141) and Eq. (142), we obtain the kinetic term

$$L_K = \frac{1}{2g^2} \sum_{\ell m (\neq 0)} \text{Tr} \left[\partial_\mu \Phi_1^{\ell m}(x) \partial^\mu \Phi_1^{\ell m}(x) + \partial_\mu \Phi_2^{\ell m}(x) \partial^\mu \Phi_2^{\ell m}(x) \right] \quad (143)$$

where only terms quadratic in $\partial_\mu \Phi$ are retained. The potential term

$$\begin{aligned} L_V = & -\frac{1}{2g^2 R^2} \sum_{\ell m (\neq 0)} \int d\Omega \text{Tr} \left[\left(\frac{\Phi_2^{\ell m}}{\sqrt{\ell(\ell+1)}} \frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta Y_{\ell m}^\pm) + Q \right. \right. \\ & + \frac{\Phi_2^{\ell m}}{\sqrt{\ell(\ell+1)}} \frac{1}{\sin^2 \theta} \partial_\phi^2 Y_{\ell m}^\pm \\ & - \frac{i}{\ell(\ell+1)} \left[-\Phi_1^{\ell m} \partial_\theta Y_{\ell m}^\pm - \Phi_2^{\ell m} \frac{1}{\sin \theta} \partial_\phi Y_{\ell m}^\pm, \right. \\ & \left. \left. \Phi_2^{\ell m} \partial_\theta Y_{\ell m}^\pm - \Phi_1^{\ell m} \frac{1}{\sin \theta} \partial_\phi Y_{\ell m}^\pm + \sqrt{\ell(\ell+1)} A_\phi^B \right] \right)^2 \Big] , \quad (144) \end{aligned}$$

where only terms diagonal in (ℓ, m) are consider. Using the relation $\frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta Y_{\ell m}) + \frac{1}{\sin^2 \theta} \partial_\phi^2 Y_{\ell m} = -\ell(\ell+1) Y_{\ell m}$, the potential term is simplified as

$$\begin{aligned} L_V = & -\frac{1}{2g^2 R^2} \sum_{\ell m (\neq 0)} \int d\Omega \text{Tr} \left[\left(-\sqrt{\ell(\ell+1)} \Phi_2^{\ell m} Y_{\ell m}^\pm + Q \right. \right. \\ & + \frac{i}{\ell(\ell+1)} [\Phi_1^{\ell m}, \Phi_2^{\ell m}] (\partial_\theta Y_{\ell m}^\pm \partial_\theta Y_{\ell m}^\pm + \frac{1}{\sin^2 \theta} \partial_\phi Y_{\ell m}^\pm \partial_\phi Y_{\ell m}^\pm) \\ & \left. \left. + \frac{i}{\sqrt{\ell(\ell+1)}} [\Phi_1^{\ell m}, \tilde{A}_\phi^B] \partial_\theta Y_{\ell m}^\pm + \frac{i}{\sqrt{\ell(\ell+1)}} [\Phi_2^{\ell m}, \tilde{A}_\phi^B] \frac{\partial_\phi Y_{\ell m}^\pm}{\sin \theta} \right)^2 \right] . \quad (145) \end{aligned}$$

To obtain the mass term, we focus on terms quadratic in $\Phi_{1,2}$:

$$\begin{aligned}
L_M = & -\frac{1}{2g^2 R^2} \int d\Omega \text{Tr} \left[\ell(\ell+1) (\Phi_2^{\ell m})^2 (Y_{\ell m}^\pm)^2 \right. \\
& + \frac{2iQ}{\ell(\ell+1)} [\Phi_1^{\ell m}, \Phi_2^{\ell m}] \left(\partial_\theta Y_{\ell m}^\pm \partial_\theta Y_{\ell m}^\pm + \frac{1}{\sin^2 \theta} \partial_\phi Y_{\ell m}^\pm \partial_\phi Y_{\ell m}^\pm \right) \\
& + 2i\tilde{A}_\phi^B [\Phi_1^{\ell m}, \Phi_2^{\ell m}] Y_{\ell m}^\pm \partial_\theta Y_{\ell m}^\pm - \frac{1}{\ell(\ell+1)} [\Phi_1^{\ell m}, \tilde{A}_\phi^B]^2 (\partial_\theta Y_{\ell m}^\pm)^2 \\
& \left. - \frac{1}{\ell(\ell+1)} [\Phi_2^{\ell m}, \tilde{A}_\phi^B]^2 \frac{(\partial_\phi Y_{\ell m}^\pm)^2}{\sin^2 \theta} \right].
\end{aligned} \tag{146}$$

Note that we have dropped the term proportional to $[\Phi_1, \tilde{A}_\phi^B][\Phi_2, \tilde{A}_\phi^B]$ because this term vanishes after turning the field into the linear combinations of Φ and Φ^\dagger , Eqs. (60) and (60):

$$\begin{aligned}
\text{Tr}[[\Phi_1, \tilde{A}_\phi^B][\Phi_1, \tilde{A}_\phi^B]] & \rightarrow \text{Tr}[(\Phi + \Phi^\dagger), Q][(\Phi - \Phi^\dagger), Q] \\
& \propto \text{Tr}[(\Phi - \Phi^\dagger)(\Phi + \Phi^\dagger)] \\
& \propto \text{Tr}[\Phi\Phi^\dagger] - \text{Tr}[\Phi^\dagger\Phi] = 0
\end{aligned} \tag{147}$$

Integrating the second term of Eq. (146) by part, we obtain

$$\begin{aligned}
L_M = & -\frac{1}{2g^2 R^2} \left(\ell(\ell+1) \text{Tr}[(\Phi_2^{\ell m})^2] + 2i \text{Tr}[Q[\Phi_1^{\ell m}, \Phi_2^{\ell m}]] \right. \\
& - 2i \text{Tr}[Q[\Phi_1^{\ell m}, \Phi_2^{\ell m}]] \int d\Omega \frac{\cos \theta \mp 1}{\sin \theta} Y_{\ell m}^\pm \partial_\theta Y_{\ell m}^\pm \\
& - \frac{1}{\ell(\ell+1)} [\Phi_1^{\ell m}, Q]^2 \int d\Omega \frac{(\cos \theta \mp 1)^2}{\sin^2 \theta} (\partial_\theta Y_{\ell m}^\pm)^2 \\
& \left. - \frac{1}{\ell(\ell+1)} [\Phi_2^{\ell m}, Q]^2 \int d\Omega \frac{(\cos \theta \mp 1)}{\sin^2 \theta} \frac{(\partial_\phi Y_{\ell m}^\pm)^2}{\sin^2 \theta} \right).
\end{aligned} \tag{148}$$

Therefore, the KK masses depend on the $U(1)_I$ charges of the scalar fields.

For components with zero $U(1)_I$ charge, we write $\Phi_{1(2)}(x)$ as $\phi_{1(2)}(x)Q$ where Q is the corresponding generator of E_6 in Eq. (97) with zero $U(1)_I$ charge. Taking the trace, we have the following kinetic and KK mass terms instead:

$$L = \sum_{\ell m (\neq 0)} \left(\partial_\mu \phi_1^{\ell m}(x) \partial^\mu \phi_1^{\ell m}(x) + \partial_\mu \phi_2^{\ell m}(x) \partial^\mu \phi_2^{\ell m}(x) + \ell(\ell+1) \phi_2^{\ell m}(x) \phi_2^{\ell m}(x) \right) \tag{149}$$

where we have made the substitution $\phi_i \rightarrow g\phi_i$. Note that ϕ_1 is considered as a massless Nambu-Goldstone (NG) boson in this case.

For components with nonzero $U(1)_I$ charge, we use Eq. (60) and (61) and write $\Phi(x)$ as $\phi^i(x)Q_i$ where Q_i is the corresponding generator of E_6 in Eq. (97) with nonzero $U(1)_I$ charge. The commutator between Q and Φ is

$$[Q, \Phi] = 3[Q_I, Q_i]\phi^i = 3q_I\phi^i, \quad (150)$$

where we have used $Q = 3Q_I$ and that q_I is a constant determined by the $U(1)_I$ charge of the corresponding component. Finally, the Lagrangian becomes

$$\begin{aligned} L = \sum_{\ell m (\neq 0)} \Big\{ & \partial_\mu \phi_{\ell m}^\dagger \partial^\mu \phi_{\ell m} \\ & - \frac{1}{4R^2} \left[2\ell(\ell+1)\phi_{\ell m}^\dagger \phi_{\ell m} - 12q_I\phi_{\ell m}^\dagger \phi_{\ell m} \right. \\ & + 12q_I\phi_{\ell m}^\dagger \phi_{\ell m} \int d\Omega \frac{\cos\theta \mp 1}{\sin\theta} Y_{\ell m}^\pm \partial_\theta Y_{\ell m}^\pm \\ & \left. + \frac{18q_I^2}{\ell(\ell+1)} \phi_{\ell m}^\dagger \phi_{\ell m} \int d\Omega \frac{(\cos\theta \mp 1)^2}{\sin^2\theta} \left((\partial_\theta Y_{\ell m}^\pm)^2 + \frac{(\partial_\phi Y_{\ell m}^\pm)^2}{\sin^2\theta} \right) \right] \Big\}. \end{aligned} \quad (151)$$

where the subscript i is omitted for simplicity. The KK masses of the complex scalar field ϕ are then

$$\begin{aligned} M_{\ell m}^{KK}(\phi) &= \frac{1}{R} \sqrt{\frac{\ell(\ell+1)}{2} + (m_{\ell m}^B)^2}, \\ (m_{\ell m}^B)^2 &= -3q_I + 3q_I \int d\Omega \frac{\cos\theta \mp 1}{\sin\theta} Y_{\ell m}^\pm \partial_\theta Y_{\ell m}^\pm \\ &\quad + \frac{9q_I^2}{2\ell(\ell+1)} \int d\Omega \frac{(\cos\theta \mp 1)^2}{\sin^2\theta} (\partial_\theta Y_{\ell m}^\pm)^2 \\ &\quad + \frac{9q_I^2}{2\ell(\ell+1)} \int d\Omega \frac{(\cos\theta \mp 1)^2}{\sin^2\theta} \frac{(\partial_\phi Y_{\ell m}^\pm)^2}{\sin^2\theta}. \end{aligned} \quad (152)$$

The squared KK mass $(M_{\ell m}^{KK})^2$ is always positive except for the lowest mode ($\ell = 1, m = 1$). In fact, the squared KK mass of the $(1, 1)$ mode agrees with the coefficient of quadratic term in the Higgs potential (120).

IV. SUMMARY AND DISCUSSIONS

We have reviewed a gauge theory defined on 6D spacetime with the S^2/Z_2 topology on the extra space. Two scenarios are considered to construct a 4D theory from the 6D model.

One scenario based on the $SO(12)$ gauge group requires a symmetry condition for the gauge field. The other involves the E_6 gauge group, but does not need the symmetry condition. Non-trivial boundary conditions on the extra space are imposed in both scenarios.

We explicitly give the prescriptions to identify the gauge field and the scalar field remaining in 4D spacetime after the dimensional reduction. We show that the $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_X \times U(1)_I$ gauge symmetry remains in 4D spacetime, and that the SM Higgs doublet with a suitable potential for electroweak symmetry breaking can be derived from the gauge sector in both models. The Higgs boson mass is also predicted in such models. Massless fermion modes are also successfully obtained as the SM fermions by introducing appropriate field contents in 6D spacetime, with suitable parity assignments on the S^2/Z_2 extra dimension, and incorporating the background gauge field. We also discuss about the massive KK modes of fermions for the scenario with the symmetry condition and the KK modes of all fields for the one without the symmetry condition. The lightest fermionic KK mode can serve as a dark matter candidate. In general, they may give rise to rich phenomena in collider experiments and implications in cosmological studies.

To make our models more realistic, there are several challenges such as eliminating the extra $U(1)$ symmetries and constructing the realistic Yukawa couplings, which are the same as other gauge-Higgs unification models. We, however, can get Kaluza-Klein modes in our models. This suggests that we obtain the dark matter candidate in our model. Thus it is very important to study these models further such as dark matter physics and collider physics.

Acknowledgement

This research was supported in part by the Grant-in-Aid for the Ministry of Education, Culture, Sports, Science, and Technology, Government of Japan, No. 20540251 (J.S.), and the National Science Council of R.O.C. under Grant No. NSC-100-2628-M-008-003-MY4.

-
- [1] N.S. Manton, Nucl. Phys. B 158 (1979) 141.
 - [2] D.B. Fairlie, Phys. Lett. B 82 (1979) 97.
 - [3] D.B. Fairlie, J. Phys. G 5 (1979) L55.

- [4] C. Csaki, C. Grojean and H. Murayama, Phys. Rev. D **67**, 085012 (2003) [arXiv:hep-ph/0210133].
- [5] L.J. Hall, Y. Nomura, and D.R. Smith, Nucl. Phys. B 639 (2002) 307.
- [6] G. Burdman and Y. Nomura, Nucl. Phys. B 656 (2003) 3.
- [7] I. Gogoladze, Y. Mimura, and S. Nandi, Phys. Lett. B 562 (2003) 307.
- [8] C.A. Scrucca, M. Serone, L. Silvestrini, and A. Wulzer, JHEP 02 (2004) 049.
- [9] N. Haba, Y. Hosotani, Y. Kawamura, and T. Yamashita, Phys. Rev. D 70 (2004) 015010.
- [10] C. Biggio and M. Quiros, Nucl. Phys. B 703 (2004) 199.
- [11] K. Hasegawa, C. S. Lim and N. Maru, Phys. Lett. B 604 (2004) 133.
- [12] N. Haba, S. Matsumoto, N. Okada, and T. Yamashita, JHEP 02 (2006) 073.
- [13] Y. Hosotani, S. Noda, Y. Sakamura, and S. Shimasaki, Phys. Rev. D 73 (2006) 096006.
- [14] M. Sakamoto and K. Takenaga, Phys. Rev. D 75 (2007) 045015.
- [15] C. S. Lim, N. Maru and K. Hasegawa, J.Phys.Soc.Jap.77 (2008) 074101.
- [16] Y. Hosotani and Y. Sakamura, Prog. Theor. Phys. 118 (2007) 935.
- [17] Y. Sakamura, Phys. Rev. D 76 (2007) 065002.
- [18] A.D. Medina, N.R. Shah, and C.E.M. Wagner, Phys. Rev. D 76 (2007) 095010.
- [19] C.S. Lim and N. Maru, Phys.Lett.B653 (2007) 320-324.
- [20] Y. Adachi, C.S. Lim, and N. Maru. Phys. Rev. D 76 (2007) 075009.
- [21] I. Gogoladze, N. Okada, and Q. Shafi, Phys. Lett. B 659 (2008) 316.
- [22] T. Nomura and J. Sato, Nucl. Phys. B **811**, 109 (2009) [arXiv:0810.0898 [hep-ph]].
- [23] C. W. Chiang T. Nomura, Nucl. Phys. B **842**, 362 (2011) [arXiv:1006.4446 [hep-ph]].
- [24] P. Forgacs and N.S. Manton, Commun. Math. Phys. 72 (1980) 15.
- [25] D. Kapetanakis and G. Zoupanos, Phys. Rept. 219 (1992) 1.
- [26] A. Chatzistavrakidis, P. Manousselis, N. Prezas, and G. Zoupanos, Phys. Lett. B 656 (2007) 152.
- [27] George Douzas, Theodoros Grammatikopoulos and George Zoupanos, arXiv:0808.3236[hep-th]
- [28] A.A. Abrikosov, arXiv:hep-th/0212134.
- [29] G. Chapline and R. Slansky, Nucl. Phys. B 209 (1982) 461.
- [30] K. Farakos, D. Kapetanakis, G. Koutsoumbas and G. Zoupanos, Phys. Lett. B211 (1988) 322.
- [31] D. Kapetanakis and G. Zoupanos, Phys. Lett. B249 (1990) 66.
- [32] B.E. Hanlon and G.C. Joshi, Phys. Rev. D48 (1993) 2204.

- [33] T. Jittoh, M. Koike, T. Nomura, J. Sato and T. Shimomura, arXiv:0803.0641 [hep-ph].
- [34] Y. Kawamura, Prog. Theor. Phys. 105 (2001) 999.
- [35] Y. Kawamura, Prog. Theor. Phys. 105 (2001) 691.
- [36] S. Randjbar-Daemi and E. Percacci, Phys. Lett. B117 (1982) 41.
- [37] N.S. Manton, Nucl. Phys. B 193 (1981) 502.
- [38] S. Rajpoot and Sithikong Phys. Rev. D23 (1981) 1649-1656.
- [39] N. Maru, T. Nomura, J. Sato and M. Yamanaka Nucl. Phys. B **830**, 414 (2010)
[arXiv:0810.0898 [hep-ph]]
- [40] H. Dohi and K. Oda, arXiv.1004.3722 (2010)
- [41] G. von Gersdorff, N. Irges and M. Quiros Phys. Lett. B **551**, 351 (2003) [ArXiv:hep-ph/0210134]