

Superluminality and a Curious Phenomenon in the Relativistic Quantum Hamilton-Jacobi Equation

Marco Matone

Dipartimento di Fisica “G. Galilei” and Istituto Nazionale di Fisica Nucleare
Università di Padova, Via Marzolo, 8 – 35131 Padova, Italy
e-mail: matone@pd.infn.it Phone: +39 049 827 7142, Fax: +39 049 827 7102

Dedicated to Shawn Lane

Abstract

A basic problem in the relativistic quantum Hamilton-Jacobi theory is to understand whether it may admit superluminal solutions. Here we consider the averaging of the speed on a period of the oscillating term which is similar to Dirac’s averaging of the oscillating part of the free electron’s speed. Such an averaging solves the problem of getting the $\hbar \rightarrow 0$ limit of the speed of the free particle, and leads to solutions that, depending on the integration constants, may be superluminal.

A basic issue related to the relativistic quantum Hamilton-Jacobi equation is to investigate its superluminal solutions. This is a subtle question involving the quantum potential and the problem of defining the classical limit in the case of a free particle. The main idea here is to average on the oscillating part, a procedure which is reminiscent of Dirac's averaging of the oscillating part of the free electron's speed $\frac{i}{2}c\hbar\alpha_1^0 e^{-2iHt/\hbar} H^{-1}$ [1]. This suggests a new view on the quantum Hamilton-Jacobi equations.

The quantum version of the non-relativistic and relativistic Hamilton-Jacobi equations [2] has been derived by first principles in [3, 4, 5], strictly related to classical-quantum and Legendre dualities [6, 7, 8, 9]. In [3] it has been shown that energy quantization follows as a theorem without using any axiomatic interpretation of the wave function.

A basic outcome of the formulation proposed in [3, 4, 5], is that the quantum potential, that plays the role of intrinsic energy, is never trivial. This is true even in the case of the free particle classically at rest.

Consider the Klein-Gordon equation in the stationary case

$$(-\hbar^2 c^2 \Delta + m^2 c^4 - E^2)\psi = 0 . \quad (1)$$

The relativistic stationary quantum Hamilton-Jacobi equation follows by setting $\psi = Re^{\frac{i}{\hbar}S_0}$

$$(\nabla S_0)^2 + m^2 c^2 - \frac{E^2}{c^2} - \hbar^2 \frac{\Delta R}{R} = 0 , \quad (2)$$

where S_0 and R satisfy the continuity equation

$$\nabla \cdot (R^2 \nabla S_0) = 0 . \quad (3)$$

In terms of the quantum potential

$$Q = -\frac{\hbar^2}{2m} \frac{\Delta R}{R} ,$$

and of the conjugate momentum

$$p = \nabla S_0 ,$$

Eq.(2) reads

$$E^2 = p^2 c^2 + m^2 c^4 + 2mQc^2 . \quad (4)$$

For our purpose it is sufficient to consider the 1 + 1 dimensional case. The continuity equation gives $R = 1/\sqrt{S_0'}$, so that

$$Q = \frac{\hbar^2}{4m} \{S_0, q\} ,$$

where $\{f, q\} = \frac{f'''}{f'} - \frac{3}{2} \left(\frac{f''}{f'}\right)^2$ is the Schwarzian derivative of f . Therefore (2) and (3) reduce to the single equation

$$(\partial_q S_0)^2 + m^2 c^2 - \frac{E^2}{c^2} + \frac{\hbar^2}{2} \{S_0, q\} = 0 . \quad (5)$$

Eq.(5) is formally equivalent to the quantum Hamilton-Jacobi equation which follows from the stationary Schrödinger equation, namely

$$\frac{1}{2m} (\partial_q S_0)^2 + V - E + \frac{\hbar^2}{4m} \{S_0, q\} = 0 , \quad (6)$$

where $V(q)$ is the potential. It follows that the general solution of (5) has the same form of the one of (6), that is [3, 4, 5]

$$e^{\frac{2i}{\hbar}\mathcal{S}_0\{\delta\}} = e^{i\alpha\frac{w+i\bar{\ell}}{w-i\ell}}, \quad (7)$$

where $w = \psi^D/\psi \in \mathbb{R}$ with ψ and ψ^D two real linearly independent solutions of the Klein-Gordon equation (1) in the 1+1 dimensional case. Furthermore, we have $\delta = \{\alpha, \ell\}$, with $\alpha \in \mathbb{R}$ and $\ell = \ell_1 + i\ell_2$ integration constants. The crucial point here is to note that $\ell_1 \neq 0$ even when $E = mc^2$, equivalent to having $\mathcal{S}_0 \neq cnst$, which is a necessary condition to define $\{\mathcal{S}_0, q\}$. This implies a non-trivial \mathcal{S}_0 , even for a particle classically at rest.

Following Floyd [10], time parametrization is defined by Jacobi's theorem

$$t = \frac{\partial\mathcal{S}_0}{\partial E}, \quad (8)$$

that, since $p = \partial_q\mathcal{S}_0$, implies $v = \partial E/\partial p$, which is the group velocity. It should be stressed that (8) is different from the time parametrization which follows by the conventional Bohmian mechanics, where

$$v = \frac{p}{m} \neq \left(\frac{\partial^2\mathcal{S}_0}{\partial q\partial E}\right)^{-1}.$$

Furthermore, the formulation of the quantum Hamilton-Jacobi equation derived in [3, 4] solves the problem, first observed by Einstein, of considering the classical limit in the case of bound states, such as in the case of the harmonic oscillator.

Set $L = k\ell$, $L_1 = \Re L$, $L_2 = \Im L$ where

$$k = \frac{1}{\hbar c}\sqrt{E^2 - m^2c^4}.$$

Note that L may depend on the particle quantum numbers. Even if, as shown in [3, 4, 5], L may depend on the energy and fundamental constants as well, here we consider the case of L independent of E . Two linearly independent solutions of (1) are $\sin(kq)$ and $\cos(kq)$, so that by (7) we have that in the case of \mathcal{S}_0 , solution of (5), the mean speed is

$$v = \frac{q}{t} = c\frac{\sqrt{E^2 - m^2c^4}}{E}H_E(L_1, L_2; q), \quad (9)$$

with

$$H_E(L_1, L_2; q) = \frac{|\sin(kq) - iL\cos(kq)|^2}{L_1}. \quad (10)$$

We see that for $L = 1$

$$H_E(1, 0; q) = 1,$$

so that one recovers the classical relativistic relation

$$v = c\frac{\sqrt{E^2 - m^2c^4}}{E}.$$

The situation changes considerably if $L_1 \neq 1$ and/or L has a non-trivial imaginary part, $L_2 \neq 0$, even small. In this case, due to the oscillating terms, the limit $\hbar \rightarrow 0$ is not

well-defined. Such a problem also arises in the non-relativistic case. Actually, the above expressions can be obtained directly from the ones of free non-relativistic particle replacing E by $\frac{E^2}{2mc^2} - \frac{mc^2}{2}$. In [3, 4, 5] it has been shown that taking the $\hbar \rightarrow 0$ limit requires introducing fundamental constants, in particular the Planck length. This is of particular interest since the analysis concerned the non-relativistic formulation of the quantum Hamilton-Jacobi equation and may be seen as a signal that at the level of foundations of quantum mechanics, special relativity should be taken into account. A related issue is that in spite of the fact that the Heisenberg uncertainty principle is fundamental, there is no a rigorous formulation of an analog relativistic principle leading to the uncertainty relations in the $c \rightarrow \infty$ limit. Its heuristic formulation follows by the observation that whether c is the maximal speed, then the relativistic version of the Heisenberg uncertainty relations should have an upper bound on the uncertainty of the momentum, namely $\Delta p \sim mc$. In particular [11]

$$\Delta q \sim \frac{\hbar}{mc} .$$

We now show that there is a natural way of getting the $\hbar \rightarrow 0$ limit which interestingly leads to solutions which also admit superluminality. First, one may easily check that there are values of q , L_1 and L_2 for which

$$H_E(L_1, L_2; q) > 1 , \tag{11}$$

which is a basic hint that superluminal solutions exist. Furthermore, we note that there is a natural way to get rid of the oscillatory terms which opens a new view on the quantum Hamilton-Jacobi theory. In particular, averaging on the short period leads to an expression which is independent of \hbar , so providing an alternative to the undefined $\hbar \rightarrow 0$ limit. Let us restrict to the case $|L| = 1$, that is set

$$L_1 = \cos \theta , \quad L_2 = \sin \theta , \tag{12}$$

so that Eq.(9) reduces to

$$v = \frac{q}{t} = c \frac{\sqrt{E^2 - m^2 c^4}}{E} \frac{1 + \sin \theta \sin(2kq)}{\cos \theta} . \tag{13}$$

Note that the term $\sin(2kq)$ is very rapidly oscillating (e.g., in the case of [12], the neutrino Compton wavelength λ_ν is approximately 10^{11} times k^{-1}). This justifies averaging v on the interval $[q, q + \frac{\pi}{k}]$, so that Eq.(13) becomes

$$\langle v \rangle = \frac{k}{\pi} \int_q^{q+\frac{\pi}{k}} v(q') dq' = c \frac{\sqrt{E^2 - m^2 c^4}}{E} \frac{1}{\cos \theta} . \tag{14}$$

This is the analog of Dirac's averaging of the oscillating part of the free electron's speed [1]. By (14) we see that, depending on the values of θ , E and m , one may have $\langle v \rangle > c$.

Another interesting consequence of (14) is that, since it corresponds to (13) at $q = j\pi/2k$, it may indicate a discrete space-time. In this respect, it would be very interesting to formulate the relativistic quantum Hamilton-Jacobi equation on a space-time lattice and check if it may lead to an analogue of Eq.(14).

Acknowledgements. It is a pleasure to heartily thank A.E. Faraggi for the collaboration over the years.

References

- [1] P. A. M. Dirac, *The Principles of Quantum Mechanics*, Oxford University Press.
- [2] P. R. Holland, *The Quantum Theory of Motion: an Account of the de Broglie-Bohm Causal Interpretation of Quantum Mechanics*, Cambridge University Press, 1993; *Phys. Rept.* **224**, 95 (1993).
- [3] A. E. Faraggi and M. Matone, *Phys. Lett. B* **450**, 34 (1999); *Phys. Lett. B* **437**, 369 (1998); *Phys. Lett. A* **249**, 180 (1998); *Phys. Lett. B* **445**, 77 (1998); *Phys. Lett. B* **445**, 357 (1999); *Int. J. Mod. Phys. A* **15**, 1869 (2000); arXiv:0912.1225, Contribution to the book on Quantum Trajectories, Edited by Pratim Chattaraj, Taylor&Francis/CRC press, 2010.
- [4] G. Bertoldi, A. E. Faraggi and M. Matone, *Class. Quant. Grav.* **17**, 3965 (2000).
- [5] M. Matone, *Found. Phys. Lett.* **15**, 311 (2002); hep-th/0212260; hep-th/0502134.
- [6] A. E. Faraggi and M. Matone, *Phys. Rev. Lett.* **78**, 163 (1997).
- [7] M. Matone, *Phys. Lett. B* **357**, 342 (1995).
- [8] M. Matone, *Phys. Rev. D* **53**, 7354 (1996).
- [9] M. Matone, *Nucl. Phys. B* **656**, (2003) 78.
- [10] E. R. Floyd, *Phys. Rev. D* **25**, 1547 (1982); *Phys. Rev. D* **26**, 1339 (1982); *Phys. Rev. D* **29**, 1842 (1984); *Phys. Rev. D* **34**, 3246 (1986); *Int. J. Theor. Phys.* **27**, 273 (1988); *Phys. Lett. A* **214**, 259 (1996); *Found. Phys. Lett.* **9**, 489 (1996); *Found. Phys. Lett.* **13**, 235 (2000); *Int. J. Mod. Phys. A* **14**, 1111 (1999); *Int. J. Mod. Phys. A* **15**, 1363 (2000); quant-ph/0302128; quant-ph/0307090.
- [11] L. D. Landau and E. M. Lifshitz, *Théorie Quantique Relativiste, I*, Éditions de Moscou, 1972.
- [12] T. Adam *et al.*, arXiv:1109.4897.