

OPERA Collaboration have observed phase
speed of neutrino wave function, while
advanced time displacement is mainly due to
finite life time of pion

Shi-Yuan Li

School of Physics, Shandong University, Jinan, 250100, PRC

October 11, 2018

Abstract

We show that the OPERA Collaboration have measured the phase velocity of the neutrino wave function, based on the analysis of the experimental method. On the other hand, the significant advanced time displacement δt reported by OPERA is mainly due to the finite life time of the pions (w.r.t. its flying time in the 1 km long tunnel) which decay to the muon neutrinos.

1. The OPERA Collab. [1] announced to have measured the ‘time of flight’ T (TOF_ν in [1]) of the neutrino from CERN to Gran Sasso. Dividing the distance between these two places L (from BCT to OPERA [1]) by T , one can get a ‘velocity’ whose value turned out to be larger than the speed of light in vacuum, c , i.e., $L/T > c$. Such a result, at first glance, could be a challenge to Einstein’s Special Relativity, though one is aware that c is one of the basic constant in modern physics whose physical meaning is much more deeper than light’s ‘speed’ whose physical meaning is quite ambiguous. Only in the case of strict classical mechanics, when the position (element of some mathematical structure on which differential w.r.t. the parameter ‘time’ can be defined) is considered as the basic observable of a particle, and as a single value function of time, the velocity is a well-defined quantity. According to Nöther’s theorem, a basic and well defined space-time related observable is

the generator of some kind of space-time symmetry operation. In quantum theory, the movement of a particle is described by state vector, and generally position is not a good quantum number. So in position representation we encounter a distribution described by wave function of the particle (leaving out the more complexity of particle creation and annihilation). Hence velocity is too complex and case-dependent, generally can not be taken as basic observable related to some kind of space-time symmetry, but as a quantity defined by others, depending on the concrete cases. E.g. (suppose Lorentz invariance kept), for a free particle in energy-momentum eigenstate $|E \mathbf{P} \rangle$, with $E^2 = P^2 + m^2$, its velocity can be defined by \mathbf{P}/E , but is not easy to relate to some unambiguous distance over time interval as in classical mechanics, even with the knowledge of the wave function. Another intuitive example is the electron bound in hydrogen atom, whose 3-momentum is not a ‘good quantum number’, i.e., not commuted with the Hamiltonian hence not the movement integral. In this case the above definition of velocity fails; and because the electron is ‘off-shell’, if we define some ratio between momentum and energy, or space and time, it is very possible to get some ‘velocity’ which is large than c . One may call the electron in such case as tachyon if one likes, but we know that the basic physical principles like causality of the hydrogen system, e.g., in process of transition between different energy levels, always keep. Hence we should pay attention that some ‘velocity’ defined by ratio between some measured space and time intervals is not always straightforward pointing to the basic principle about the property of space-time. On aware of the above, we can well understand that even the terminology ‘speed of light in vacuum’ is purely a traditional name. Since we should ask what kind of ‘light’: a photon? a light pulse? etc. For each case the definition of velocity varies. We know that c is further used, e.g., to scale the momentum, mass, energy of a free particle so that $E^2 = P^2 + M^2$, etc., etc. However, if the wave function of a particle can be deduced from some experiments, one of course can discuss some ‘displacements’ in space as well as time based on the wave function. Their ratio could be interesting provided a careful investigation on what such kind of measurement the experiment in fact makes, and what/whether basic physical principles can be further deduced from it.

In the following we will recall that the phase speed of the wave function can very naturally be larger than c (section 2). With the effort to analyze the key experiment method, i.e., fitting the wave forms (average time distributions in the time interval of the extraction [1]) of the proton as well as the neutrino event to get the time displacement w.r.t. ‘flying in the speed of c ’,

we show that the OPERA Collab. have measured the phase of the particle wave function and its speed hence is possibly larger than c (section 3). Since neutrino mass is very small, whether phase velocity or group velocity, if properly measured, both should not vary from c significantly, no matter larger or smaller than c . However, due to the finite life time of the pions (w.r.t. its flying time in the 1 km long tunnel) which decay to the muon neutrinos, the wave form for neutrino is deformed from that of the proton. Such an effect is estimated to lead to an advanced time displacement of the order of 100ns, which coincide with the measured δt reported in [1] (section 4).

2. The basic things measured by OPERA Collab. are two time distributions in two space-time positions. At position x_0 (CERN) and time t_0 , one measures the proton to deduce the wave function of the neutrino (for the problems see section 4) and then at t_1 and position x_1 (Gran Sasso) measures the wave function by the OPERA detector (of course we can only deduce the wave function by measuring its modulus square). To make the discussions simple, we do not discuss the neutrino oscillation. The good time correspondence of the neutrino events and the proton extractions as presented by the Collab. [1] indicates that the neutrino is well propagating and could be modeled as a plane wave for the simplest consideration. We will employ wave packet to describe the neutrino in Section 3 and show that the measured result L/T is possibly understood as the phase speed of the propagating wave function since their key measurement is to calibrate the pace/phase for the particle in these two space-time positions. L/T larger than c is the straightforward result from that the physical velocity of neutrino is smaller than c and well obeys the energy-momentum relation of a massive particle. Suppose a likely experiment is done for a much more massive ‘dark matter’ particle, such a phase speed may be significantly larger than c . However, this does not say such measurement is meaningless. Detailed analysis on such measurement may help to improve the experiments on neutrino oscillation or even absolute value of neutrino mass.

The wave function is

$$e^{-i\omega t + i\kappa \cdot \mathbf{x}}, \quad (1)$$

and we find that

$$\frac{\omega}{|\kappa|} = \frac{|\Delta \mathbf{x}|}{\Delta t} = \frac{|\mathbf{x}_1 - \mathbf{x}_0|}{t_1 - t_0} \quad (2)$$

This is the well-known phase velocity of the wave of eq. (1). Here one encounters the similar problem that Louis de Broglie encountered almost

100 years ago [2]. When ω and κ correspond to energy and momentum of a particle, according to the Einstein relation, we will find that this phase velocity is never smaller than c but always larger, except for light it equals to c . And the wise solution of employing more reasonable group velocity by de Broglie is the key corner stone for his matter wave hypothesis which is one of the most important turning points from Bohr's old quantum theory to the modern quantum mechanics.

3. A plane wave as above extends infinitely. If nothing can mark its phase, our discussions in Section 2 are just for an 'ideal experiment'. By the above we only show that for a plane wave, the interval between two space-time positions with the same phase could be space like. For the mechanical wave like that on a long shaking rope, we can see the oscillation (say, up and down) of the points of the rope with different phases, so that we can observe the propagation. It turns out that the time structure of the proton PDF (probability density function [1], but in fact number density as to be clarified) and that of the OPERA neutrino events can mark the phase to be measurable, which we will make clear in the following based on the analysis of the experiment. It is our key point in this section to make clear what the experiment measures with a simple model. Now it is more of reality to describe the propagation of neutrino by wave packet,

$$\Psi(x, t) = \int dk \Psi(k) e^{-i\omega t + ikx}, \quad (3)$$

where $\omega = \omega(k)$ is defined by the on-shell Einstein relation for four momentum (In this section the space related variables like k , x are not specially denoted to show they are SO(3) vectors but indicated). Here we do not discuss the details of $\Psi(k)$, only this wave packet is assumed well localized: For space, even the whole neutrino source system of CERN at Preveessin and the detectors in laboratory at Gran Sasso can be taken as point comparing to the distance between these two places. So here we need not discuss how well the position localized which is in fact well investigated by OPERA Collab. For time, we have no problem to say that this particle is localized much smaller than $10.5\mu s$, based on the PDF time structure. Same as [1], we employ the time when the proton going through the BCT to mark the time of the wave packet (e.g., the central peak of the wave packet of Eq. (3)). Taking into account the reaction time of the BCT or by carefully analyzing the PDF, one can extract a basic time length l_T (say, around 5 ns according to Fig.4 OF [1]). Here a proper $\Psi(k)$ need to lead that $\Psi(x, t + \Delta t)$ vanishing when

$|\Delta t| > l_T$. Since the wave packet could expand en route of propagation, so this requirement is only for the time and place at CERN, where the protons employed to produce neutrino by its reaction with the target is carefully measured, one of the most important instrument for the this measurement is the BCT [1].

This is the key point here. We would like to remind the reader that if he/she understood that we assign the distribution of the PDF or such kinds as the time distribution of the wave (packet) function of each proton as well as that of the neutrino, he/she completely misunderstood. Particles produced in CERN in the processes described in Section 2 of [1], of which proton is the most especially typical, can all be considered as that their wave packet size is as small as possible (in time and space) so that the inner structure of the wave packet function is not possible seen by all detectors. The PDF time structures exactly is the numbers of particles varying with time. However, this varying value can mark the PHASE of the wave packet of the particle.

Since in no way to assume any difference among all the protons, we can employ the same form of wave packet to describe all the protons as well as neutrinos. Only that, each of the particles (proton/neutrino) corresponding to each ‘point’ (l_T is the unit) t' at the time axis of PDF distribution has a time displacement w.r.t. the beginning of the extraction:

$$e^{-iHt'} \Psi(x, t) = \int dk \Psi(k) e^{-i\omega(t+t') + ikx}, \quad (4)$$

with $0 < t' < 10.5\mu s$. This phase in other word can be called the pace of the protons/neutrinos, i.e., to mark their order of ‘step’ in time of reactionreaction and reaction of each extraction. (Actually this only marks the neutrino’s mother particles, mainly pions and kaons, the problems caused by the infinite life time during the decay is discussed in details in section 4.) As the experiment has put together and makes statistics for all the particles in different ‘extractions’, they are to project on a same t' region by plus/minus n time of 50 ms (the time interval between extractions) for their real time t . Here we emphasize again that the wave packet is so well located that any ‘interference’ between different wave packets (particles) are neglected, as can be assumed by the experiment (or effects as beam interaction has been corrected by experiment). We emphasize this is to point out that those kind of effects are of NO relation with our discussions. And we also point out that though in experiment each event and wave form has been stamped by the real time, but what to be used in analysis is the averaged results obtained by

large number of samples of extractions and events projected together on the interval of the extraction. We in fact can not say which proton (neutrino) has which phase but just employ the different numbers of particles (events) in this duration to represent the fact that each wave packet of the particle has certain definite phase, marked by t' , relative to the, e.g, the beginning time of the extraction.

The wave packet will expand during propagating, but the experiment show they still keep some peak structure so that can be detected in Gran Sasso at some certain time. This exact value of time but is only used to extract the t' to get the event time structure for comparing with the source time structure (PDF), then to calibrate the phase parameter corresponding to t' . The method is, as described by the OPERA Collab., to employ the maximum likelihood method to coincide/tune the same pace for these two distributions. This is just to determine which phase group of proton PDF the particles/events detected by OPERA belong to. Then it is possible to find the same phase positions (t_0, x_0) and (t_1, x_1) when the phase/pace is calibrated to coincide. The space-time interval ratio hence can be now understood as the velocity of the phase of the wave packet marked by t' moves from (t_0, x_0) to (t_1, x_1) (i.e., from CERN to Gran Sasso). $x_1 - x_0 = L$ is the length of the baseline presented from [1], while $t_1 - t_0 = T$ is also possible deduced from the measured quantities in [1]. In other words, the experiment is comparing the step order of the proton and reaction events, so that to find two space-time position with the same phase. Here we would like to emphasize again that the PDF and the coinciding neutrino signal distribution is not the form of wave function but only show the fact that the wave packet's phase marked by t' properly propagates from (x_0, t_0) to (x_1, t_1) . Their interval could be space like as discussed in the following.

The movement of the wave packet from CERN to Gran Sasso,

$$e^{-iH(t_1-t_0)}\Psi(x_0, t_0, t') \tag{5}$$

in practice is too complex to be employed to calculate the phase velocity and information (or 'particle') velocity based on the real form of $\Psi(k)$, even we can get it. Here we only employ the (maybe too) simplified qualitative expressions to discuss: Suppose that the effective integral region of k (where $\Psi(k)$ significantly different from zero), Δk , is much smaller than k , i.e.,

$$\Delta k \ll k, \text{ which also renders } \Delta\omega \ll \omega. \tag{6}$$

This may be valid in high energies. Now we can rewrite the wave function $\Psi(x, t, t')$ as

$$\Delta k \Psi(k) e^{-i\omega(t+t')+ikx} e^{-i\Delta\omega t + i\Delta k x}. \quad (7)$$

$\Delta\omega t'$ is higher order infinitesimal and neglected from Eq. 7, comparing to a long propagation time, since t' is fixed as a small value between 0 to $10.5\mu s$. So here t' only appear in the ‘carrier’ wave. From this quite simple expression we see the phase propagates with a velocity ω/k , and the same phase space-time position can be marked by t' . While the wave packet propagates at the group velocity $\Delta\omega/\Delta k$. The above simplification is just a reproduction of the classical discussion from Brillouin’ book, chapter 1 [3]. In the most simple case the phase velocity is large than c while the group velocity is smaller, just to refer Section 2.

The same narrow k distribution assumption (6) for large k approximately leads to

$$|\Psi(x, t + t')|^2 \cong \delta(v_g(t + t') - x), \quad (8)$$

with $v_g = \Delta\omega/\Delta k$ the group velocity in Eq. (7). This is just the distribution function of a classical particle and this interpretes why in most high energy experiments, one can treat the space-time movement of a single particle as classical with velocity P/E , even with the fact it is almost in energy momentum eigenstate rather than position eigenstate. Such a result also confirms what we have emphasized above, that in our treatment, we need not to assume anything new from the conventional treatment about the proton beam, as well as the pions, kaons, and neutrinos in the production tunnel [1]. However, one should be aware that since in quantum mechanics, a single microscopic particle is not able to be marked and measured (e.g, the values of the position) many times without dramatically changing its state. For a static particle flow, this delta function description is of no difference from the plane wave description: no signal of movement like classical one particle displacement in space-time can be observed.

Now comes to the description of the time structure of the proton beam, which is also assumed that of the neutrino. This is an inherent property of the proton beam system. Here we assumed that many times average (average particle number $\langle N(t') \rangle$) can cancel the fluctuations from environment, i.e., $\langle N(t') \rangle$ is just the PDF [1]. So the particle system can be described as direct product of the the single particle wave functions, with the interactions

among them canceled.

$$\Phi(x, t + t') = \prod_{j=1}^{\langle N(t') \rangle} \left(\int dk_j \Psi(k_j) \right) e^{-i \sum_{j=1}^{\langle N(t') \rangle} [\omega_j(t+t') + ik_j x]}, \quad (9)$$

which renders

$$|\Phi(x, t + t')|^2 \cong \delta^{\langle N(t') \rangle} (v_g(t + t') - x). \quad (10)$$

Eq. (10) shows that the BCT signal is proportional to the number of the protons $N(t')$ as they pass through the BCT at position x and time $t+t'$. When we make the measurement for the times structure of the number of particles to mark the same-phase space-time positions as the OPERA experiment, with the static particle flow rather than one particle movement, we are in touch with a space-time distribution $B(x, t+t')$, which is function of the phase, i.e.,

$$B = B \left(\sum_{j=1}^{\langle N(t') \rangle} \omega_j(t + t') - k_j x \right) \quad (11)$$

The experiments measured two space time positions with the same time structure, as one sees the different points of a rope with the same pace of oscillation.

The real information signal, is not the averaged number of $\langle N(t') \rangle$, but a real time change of this $\langle N(t') \rangle$, which propagates with the velocity v_g . Supposing a distance large enough so that $x/v_g \gg 50\mu s$, then at some time we change from the static flow from $\langle N_1(t') \rangle$, to $\langle N_2(t') \rangle$. and keep the new flow for a time τ , with $50 \ll \tau \ll x/v_g$, then change back to $\langle N_1(t') \rangle$. These two places can also communicate with light signal, and then comparing the propagation time for light and neutrino to get the velocity. In fact this is just the analysis based on supernova 1987A observation [4].

Such a simple model shows what the measurement the OPERA Collab. have got. Of course, further and deeper analysis, e.g., considering the propagation as pulse and employ more practical formulations, especially combined with oscillation, could help us to improve the measurement of the neutrino sector.

The above discussion in fact is not restricted to the neutrino sector: It is a very general discussion based on quantum mechanics principles. However, other particle is not possible interact so weak so that can not travel such a

long distance. On the other hand, it is not very easy to discuss the causality from the formulae here, since the causal or Feynman propagator is composed of positive and negative frequency parts, or say retarded as well as advanced parts to keep causality. Here we only employ positive frequency part. So a more practical formulation to describe the wave packet of a relativistic particle in the quantum field theory framework still need exploration. And a precise observable in quantum field theory to define the pace of the particle is also in need.

4. Since the neutrino mass is very small, whether the ‘velocity’ measured by OPERA is the phase one or group one, it should be very near the value of c , not as significantly different as reported in [1]. Once the physical meaning of the velocity measured by OPERA is clear, the question on what systematics which is missed by the OPERA data analysis cause the advanced time displacement $\delta t \sim 60ns$ need to be clarified. Such an investigation is very important for this kind of measurements is reported to be projected by other Collab., such as MINOS. Here we show that the systematics comes from the finite life of the pions which decay to the muon neutrinos, ν_μ 's.

As is introduced by [1], the ν_μ is produced in a 1 km long tunnel, from the decay of the production of the proton-target collisions, mainly pion and kaon. In hadron-hadron collisions, pion production is dominant. However, for higher energy region, the rate of kaon increases. For a neutrino with certain energy, the energy of its mother pion or kaon is not quite different in average, but the $\gamma = E/m$ for kaon is much smaller than that of pion. This leads to that the effective (Lorentz retarded) life time of the kaon is much smaller than that of the pion and two results are deduced: 1) This effectively increase the contribution of kaon to ν_μ ; 2) the kaon can be taken as instantly decayed so that the ν_μ from kaon can be considered as having the same wave form as the proton.

Let's give the numerical estimation for the γ 's and hence the life times of various particles which decay to ν_μ with energy 17 GeV as example, comparing with their flying time in the 1km tunnel, $T_f \sim 0.3 \times 10^{-5}$. For the case of pion, dominantly decaying to $\mu\nu_\mu$, we choose the case that the momentum of ν_μ in the same direction of that of the pion. We find that γ must larger than several hundreds and its life time T_π is larger than $10^{-5}s$, which is comparable with (and larger than) T_f , hence the neutrinos from pions have a deformed distribution from that of the proton. For the Kaon sector, with the same example case and same method, we can get the conclusion $T_K \sim 10^{-7}s \ll T_f$ and hence the above conclusions. But the K_s should be treated separately.

Since it mainly decays into two pions instantly (comparing to T_f), so its case is same as that of the pion.

Other particles can also contribute, of them muon is very abundant. For the case of muon, it decays into ν_μ through a 3-body channel. Take the case that ν_μ energy is half of the mass of muon and momentum is parallel to the muon momentum, one can get the γ is of the order 10^2 . So T_μ is of the order $10^{-4}s \gg T_f$. So for simplicity and a coarse estimation we can ignore the contribution from the muon. If taken into account, the muon number can be taken as constant during T_f so will not lead to deformation of the wave form.

So the time of fly in the tunnel T_f is the very key point leads to the systematics for the distribution. Mainly the case of pion to ν_μ has to be considered. Since the neutrino production is proportional to the number of the pions, one can propose a exponentially decreasing distribution for the neutrino production time in the tunnel. Besides directly to calculate the wave form deformation caused by the exponential form of the time distribution of the neutrinos from the pion, the effect can be estimated by the displacement of the average production time of these neutrinos from the flat ones in the flying interval, with the latter $T_f/2$. Only considering the effect of pion, it leads to the value of the order of 100 ns, which is coincide with the OPERA result. The result is calculated from the following equations:

$$\frac{\int_0^1 dt t e^{-t/T_\pi}}{\int_0^1 dt e^{-t/T_\pi}}. \quad (12)$$

Here for simplicity we take T_f as unit. We also emphasize that this effect is not sensitive to the wave form of the proton, whether the old long time one and the new short time (pulse) one.

However, here we must point out that such an effect is significant but missed by the Collab. CRUCIALLY depends on the special property of this kind of measurement and even more CRUCIALLY depends on the understanding for the physical meaning of this experiment: First, as clarified in section 3, this experiment compares phase/pase (‘wave form’) to measure the phase velocity. Second, the recorded neutrino events are quite rare, statistically the number of neutrino events recorded by OPERA per proton extraction is much much smaller than one. In such a special case, the decay process in the tunnel for each extraction is not possible to be tracked by the neutrino events (requiring number of neutrino events per extraction ≥ 2), so

statistically the phase displacement in the whole tunnel flying will deform the recorded neutrino event wave form. Needless to say, if one insists to understand this experiment as to directly measure the flying of each neutrino recorded by OPERA, in the same way for a classical one, the decay effect is very small because $1 - \beta_\pi \sim 10^{-4}$ and pion only fly not more than 1km, which HAS been taken into account in [1].

The MINOS [5] experiment is much less affected by the decay effect discussed above since its energy is much lower hence a much smaller γ . All the mother particles can be taken as to decay instantly.

5. After the release of the OPERA data, there have been more than 100 papers discussing this result [6]. This has stimulated the study/review of the Lorentz invariance violation in various ways, which could be a good window for the future new physics. On the other hand, there are several papers devoted to study the systematics of the experiment or suggesting the velocity could be unphysical one. However, such kind of investigation should include several key points: first to clarify what the observable is measured by analyzing the basic experimental method; second to see whether the measured value agree or not with the suggested, if not, where is the systematic. This paper is trying to do in this way. This is a thing that one can not escape from for exploring Lorentz invariance violation, not relating to any conservative attitude of believing Einstein, but something learned from Einstein's attitude to experiments, no matter sooner or later his theories are found to be broken through.

The author thanks Prof. Dr. WANG Meng for explaining some aspects of the OPERA experiments. This work is partially supported by NSFC(10935012), SFSD (JQ201101) and SDU (2010JQ006).

References

- [1] T. Adam *et al.* [OPERA Collaboration], [arXiv:1109.4897 [hep-ex]].
- [2] Louis de Broglie, *Recherches sur la théorie des quanta*, PhD Thesis (Paris), 1924.

- [3] L. Brillouin, *Wave propagation and group velocity*, Academic Press, 1960.
- [4] M. J. Longo, Phys. Rev. D 36 (1987) 3276, Phys. Rev. Lett. 60 (1988) 173; L. Stodolsky, Phys. Lett. B **201** (1988) 353. The references for the observations on the neutrino as well as light signals from Supernova 1987A are to be found in these three papers.
- [5] P. Adamson *et al.* [MiNOS Collaboration], Phys. Rev. D 76 (2007) 072005.
- [6] Most of the papers are to be found from
<http://www.slac.stanford.edu/spires/find/hep?c=ARXIV:1109.4897>

This figure "1.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/1110.0302v3>

This figure "2.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/1110.0302v3>