

OPERA neutrinos and deformed special relativity

Giovanni Amelino-Camelia^a, Laurent Freidel^b, Jerzy Kowalski-Glikman^c, Lee Smolin^b

^aDipartimento di Fisica, Università “La Sapienza” and Sez. Roma1 INFN, P.le A. Moro 2, 00185 Roma, Italy

^bPerimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo, Ontario N2J 2Y5, Canada

^cInstitute for Theoretical Physics, University of Wrocław, Pl. Maxa Born 9, 50-204 Wrocław, Poland

In a recent study, Cohen and Glashow argue that superluminal neutrinos of the type recently reported by OPERA should be affected by anomalous Cherenkov-like processes. This causes them to lose much of their energy before reaching the OPERA detectors. Related concerns were reported also by Gonzalez-Mestres, Bi et al, and Cowsik et al, who argued that pions cannot decay to superluminal neutrinos over part of the energy range studied by OPERA. We observe here that these arguments are set within a framework in which Lorentz symmetry is broken, by the presence of a preferred frame. We further show that these anomalous processes are forbidden if Lorentz symmetry is instead “deformed”, preserving the relativity of inertial frames. These deformations add non-linear terms to energy momentum relations, conservation laws and Lorentz transformations in a way that is consistent with the relativity of inertial observers.

The OPERA collaboration recently reported [1] evidence of superluminal behavior for μ neutrinos with energies of a few tens of GeVs: $v - 1 \simeq 2.4 \cdot 10^{-5}$, with a significance of six standard deviations (we use units such that the speed of light is $c = 1$).

As usual in science, when a particularly striking experimental result is first reported, the most likely hypothesis is that some unknown bias or source of uncertainty affects these OPERA data. However, the result would be of such potential importance that it behoves us to at least investigate what is the second most likely hypothesis that is consistent with all the relevant data, including OPERA, and gives a reasonable path for physics to go ahead. In the event that the OPERA data is right, we must, by the development of such a hypothesis, come to understand special relativity as an approximation to deeper physics. This is of no small importance because a particular hypothesis that can explain all the relevant data can serve to predict other experiments. These predictions, if confirmed, would strengthen the case that there really are departures from special relativity.

However, a recent letter by Cohen and Glashow [2] appears to indicate that there cannot be such a second hypothesis, since it claims that the OPERA results are self-contradictory. Cohen and Glashow argue that neutrinos in OPERA’s CNGS beam, if superluminal as described in Ref. [1], would lose much of their energy via Cherenkov-like processes on their way from CERN to Gran Sasso. They then could not be detected with energy in excess of 12GeV , contrary to what is also reported [1] by OPERA. And a similar message is contained in studies by Gonzalez-Mestres [3], by Bi et al [4], and by Cowsik et al [5], which argued that the reported superluminality of μ neutrinos would even prevent, over part of the energy range studied by OPERA, the pion-decay processes at CERN, which are partly responsible for the flux of neutrinos reaching Gran Sasso. If these arguments are correct, then it seems unavoidable that the OPERA results are mistaken.

But, as we show here, there is a loophole¹ in the arguments reported in Refs. [2–5], due to an implicit assumption they make about the fate of Lorentz invariance.

As already established in Ref. [8], the possibility that these neutrinos are tachyons, consistently with special relativity, must be excluded, if we take into account other available data on the possible dependence of the neutrino maximal speed with energy. Any attempt to interpret the OPERA anomaly as actual evidence of neutrinos with superluminal behavior must therefore accommodate departures from Lorentz symmetry.

Before we begin such an analysis it is crucial to recall that Lorentz invariance can be superseded in two ways. It can be “broken” [9–11] in the sense that there is a preferred frame of reference. Or it can be “deformed” [12–14], so that the principle of relativity of inertial frames is preserved, but the action of Lorentz transformations on physical states is deformed. In such a deformation of special relativity the energy-momentum relations, conservation laws, and Poincaré transformations are all modified by non-linear terms, which are mutually consistent so that the relativity of inertial frames is preserved.

However, as we show below, the Cherenkov-like processes considered in Ref. [2] and the features of pion decay discussed in Refs. [3–5], necessarily assume that Lorentz invariance is broken. Hence, their claims that the OPERA results contradict known physics are only valid in a framework in which Lorentz invariance is broken by the presence of a preferred frame.

Furthermore, as we also show below, if one instead interprets the OPERA results within a framework in which the relativity of inertial frames is unbroken, but the laws of special relativistic kinematics are modified, then the objections of Refs. [2–5] are not valid. Superluminal neutrinos may be produced by pion decay over the whole range of OPERA energies and, once produced, are stable against loss of energy through Cherenkov-like processes. These results, combined with the ones previously

¹ Cohen and Glashow mention the possibility of a different loophole in which electrons and photons share with neutrinos “unconventional dispersion relations... such that, in the energy domain of the OPERA experiment, these particles travel with a common velocity” [2]. This would indeed render the OPERA data no longer self-contradictory; however, our understanding is that this is in contradiction with established upper bounds on the speed of multi-GeV electrons[6] and of multi-GeV photons[7].

reported in Refs. [2–5], imply that any attempt to interpret the OPERA anomaly in terms of superluminal neutrinos should favor interpretations in which Lorentz symmetry is deformed but not broken.

We now will contrast the interpretation of the OPERA experiment in two scenarios, in which the relativity of inertial frames is respectively, broken and deformed. These two frameworks have in common that the energy-momentum relation is modified by non-linear terms governed by a characteristic mass scale M_* . We start with a general form of dispersion relation

$$E^2 f^2(E, p; \mu^2) - p^2 g^2(E, p; \mu^2) = m^2 \quad (1)$$

with

$$f(E, p; \mu^2) = 1 + \phi(E, p; \mu^2), \quad g(E, p; \mu^2) = 1 + \gamma(E, p; \mu^2)$$

where ϕ, γ are functions, much smaller than 1 for in the relevant regime, codifying departures from special relativity, m is the particle rest mass and μ^2 is a deformation scale that, *crucially for what follows, can be different for different particles*.

Assuming (will return to this point in the closing remarks) that the observer adopts spacetime coordinates conjugate to the momentum-space variables E, p one easily finds that as a consequence of (1) the speed of a particle depends on energy as follows

$$v(E) = \frac{dE}{dp} = \frac{\bar{p}}{E} - \frac{d}{dE} (\bar{p} \phi(E, \bar{p}; \mu^2)) + \frac{\bar{p}^2}{E^2} \frac{d}{dE} (\bar{p} \gamma(E, \bar{p}; \mu^2)) \quad (2)$$

where $\bar{p} \equiv \sqrt{E^2 - m^2}$ and we worked at leading order in ϕ, γ .

In the framework with a preferred frame these equations hold only in that special frame. In this case it is legitimate to hypothesize that energy momentum conservation remains linear in all frames of reference. However the energy-momentum relations (1) alter their form [12] in other than the preferred frame. This, as we will show below, is the context in which Cohen and Glashow, Gonzalez-Mestres, Bi et al, and Cowsik et al, derive their conclusions.

However, if we want the relativity of inertial frames to be preserved then we want (2) and (1) to hold in all inertial frames. This is possible [12–14] if the action of the generators of Lorentz boosts \mathcal{N}_j is modified in such a way that (without modifying the standard algebra of Lorentz generators)

$$[\mathcal{N}_j^{(\mu)}, E^2 f^2(E, p; \mu^2) - p^2 g^2(E, p; \mu^2)] = 0 \quad (3)$$

The next step is to realize that if the action of the boosts is modified, then the energy-momentum conservation laws must modified as well, so that they stay covariant [12, 14, 15].

The key differences between the broken-Lorentz and the deformed-Lorentz frameworks are two: in the broken case the conservation laws remain the standard linear laws in every frame but the modified energy-momentum relation (1) hold only in the preferred frame. However in the deformed Lorentz framework the modified energy-momentum relations hold in all inertial frames and the conservation laws become non-linear as well.

The arguments of Refs. [2–5] adopt modified dispersion relations but relied on unmodified energy-momentum conservation, so they are situated in the framework of broken Lorentz invariance.

One consequence of the presence of a preferred frame is that processes with only one incoming particle can be allowed or forbidden, depending on whether the energy of the incoming particle is above or below a certain threshold value. This cannot happen in relativistic frameworks, even in cases in which the Lorentz transformations have been deformed [16, 17].

This is illustrated in the example of the process

$$\nu_\mu \rightarrow \nu_\mu + e^+ + e^- \quad (4)$$

which is the main focus of Ref. [2]. Cohen and Glashow find that this process is forbidden at low energies, but above a certain threshold energy E_{thresh} , in the preferred frame, the superluminality of the neutrino becomes sufficient to render this process allowed.

To see that this requires a preferred frame note that the threshold in question is not Lorentz invariant as is the case of thresholds found in special relativistic kinematics. Consider the case in which the preferred frame observer Alice sees neutrinos with energy just over the threshold, $E_{Alice} = E_{thresh} + \Delta E$. According to Alice, this neutrino can undergo $\nu_\mu \rightarrow \nu_\mu + e^+ e^-$.

Now consider an observer, Bob, moving with respect to Alice in the same direction as the neutrino, who sees the neutrino to have a lower energy than Alice. Such observers exist for which, $E_{Bob} < E_{thresh}$. If Bob were to use the same dispersion relation, and hence the same formula for a threshold, he would conclude that the neutrino does not emit an electron-positron pair. There is clearly a contradiction here. There are two ways to resolve it. Either there is a preferred frame in which case only that preferred observer, Alice, correctly computes the threshold energy. Or, if we insist that the energy-momentum relation is observer independent, and that both observers agree on the physics, there cannot be such an observer-dependent threshold.

As a consequence of this general argument the framework of deformed Lorentz symmetry is immune from such anomalous thresholds.

This suffices to conclude in full generality that the concerns for neutrino superluminality reported in Refs. [2–5] do not apply to the case of deformed Lorentz symmetry. We now illustrate this general result with an explicit calculation. For this purpose we consider a particularly simple toy model with only neutrinos and electrons/positrons, such that neutrinos are affected by a deformation of the form

$$E^2 = p^2 + m_\nu^2 + \frac{2E^2 p^2}{\mu^2}, \quad (5)$$

whereas electrons are ordinarily special relativistic.

From Eq. (5) it follows that the speed of ultrarelativistic neutrinos (2) is given by

$$v = 1 - \frac{m_\nu^2}{2p^2} + \frac{3p^2}{\mu^2}, \quad (6)$$

and evidently this gives $v > 1$ whenever $p > \sqrt{m_\nu \mu}/6^{1/4}$.

We stress that we consider this scenario only to illustrate the argument that the concerns reported in Refs. [2–5] are automatically evaded by any theory with deformed Lorentz invariance. It is however intriguing that for $\mu^2 = m M_{Planck}$, with $m^2 = \Delta m^2$, one of the neutrino mass differences, this simple scenario produces estimates compatible with the OPERA data [18, 19] (however it does not easily fit the SN1987a data).

The dispersion relation (5) is evidently not invariant under ordinary, undeformed Lorentz boosts. But it is invariant,

$$\left[\mathcal{N}_j^{(\mu)}, E^2 - p^2 - \frac{2E^2 p^2}{\mu^2} \right] = 0,$$

under the action of deformed boosts with the following generators $\mathcal{N}_j^{(\mu)}$

$$\delta_j E \equiv [\mathcal{N}_j^{(\mu)}, E] = p_j + \frac{p^2 p_j}{\mu^2} + \frac{2E^2 p_j}{\mu^2}, \quad (7)$$

$$\delta_j p_k \equiv [\mathcal{N}_j^{(\mu)}, p_k] = \left(E - \frac{p^2 E}{\mu^2} \right) \delta_{jk}. \quad (8)$$

The final step is to introduce conservation laws that are covariant under the action of these boosts on neutrino momenta, and the action of standard undeformed boosts on electron/positron momenta. Since we choose as illustrative case the process $\nu_\mu \rightarrow \nu_\mu + e^+ + e^-$ analyzed by Cohen and Glashow, we explicitly show a suitable conservation law for that process:

$$\vec{p} = \vec{p}' + \vec{k}_- + \vec{k}_+ \quad (9)$$

$$E = E' + \Omega_+ + \Omega_- + \alpha \left(\frac{E p^2}{\mu^2} - \frac{E' p'^2}{\mu^2} \right). \quad (10)$$

where \vec{p}, E (resp \vec{p}', E') are for the incoming (resp outgoing) neutrino whereas \vec{k}_\pm, Ω_\pm are for the outgoing electron-positron. The parameter α has been introduced so that $\alpha = 1$ for the deformed symmetry framework and $\alpha = 0$ for the case of broken Lorentz symmetry.

It is easy to verify, using (7), (8), (9), (10) for the neutrino momenta (and undeformed boosts for electron/positron momenta), that our conservation laws, for $\alpha = 1$, are covariant.

In order to see that the process $\nu_\mu \rightarrow \nu_\mu + e^+ + e^-$ is allowed by broken Lorentz symmetry but is forbidden by deformed Lorentz symmetry, it is convenient to consider the total momentum of the outgoing electron-positron pair: $\Omega_T \equiv \Omega_+ + \Omega_-$ and $\vec{k}_T = \vec{k}_+ + \vec{k}_-$. Since in our illustrative toy model electrons/positrons are ordinarily special relativistic also \vec{k}_T, Ω_T is on a special-relativistic shell: $M_T^2 = \Omega_T^2 - \vec{k}_T^2$. And while the rest energy of the pair is of course context dependent, for any given electron-positron pair it will be an invariant and it will be strictly positive: $M_T^2 \geq (2m_e)^2 > 0$.

In terms of \vec{k}_T, Ω_T and M_T one can use the conservation laws (9),(10) to obtain the following

$$p^2 = p'^2 + k_T^2 + 2p' k_T \cos \theta_* \quad (11)$$

$$E = E' + \Omega_T + \alpha \left(\frac{E p^2}{\mu^2} - \frac{E' p'^2}{\mu^2} \right), \quad (12)$$

which also involve the opening angle θ between the outgoing-neutrino spatial momentum \vec{p}' and the total spatial momentum of the outgoing electron-positron pair.

Next we combine these two equations and use dispersion relations to arrive (also assuming for simplicity that the particles involved are ultrarelativistic) at

$$\cos\theta = \frac{2E'\Omega_T + M_T^2 + 2(\alpha - 1)\frac{(E' + \Omega_T)^4 - E'^4}{\mu^2} - 2\alpha\Omega_T\frac{E'^3}{\mu^2}}{2E'\Omega_T - M_T^2\frac{E'}{\Omega_T} - m_\nu^2\frac{\Omega_T}{E'} - 2\Omega_T\frac{E'^3}{\mu^2}}. \quad (13)$$

As it is inevitable in light of the general argument we discussed earlier in this Letter, the process is indeed forbidden in the deformed-Lorentz-symmetry case: for $\alpha = 1$ one formally finds that the process requires $\cos\theta > 1$, which is not possible for physical opening angles. Of course, the same holds in undeformed special relativity: also for $\mu \rightarrow \infty$ (the limit where also the neutrinos of our toy model become ordinarily special relativistic) one would need $\cos\theta > 1$. But the process is possible, as already observed by Cohen and Glashow, when Lorentz symmetry is broken: for finite μ and $\alpha = 0$, there are combinations of E' and Ω_T in the physical phase space compatible with physical opening angles, such that $\cos\theta \leq 1$.

The derivations proceed in analogous way for verifying that also the issues for pion decay raised by by Gonzalez-Mestres [3], Bi et al [4], and Cowsik et al [5] do not apply to the deformed-Lorentz-symmetry case.

In conclusion, Refs. [2–5] claim that the OPERA result cannot be interpreted in term of superluminal propagation. In this paper we observed that their conclusions require a breaking of lorentz invariance by the introduction of a preferred frame. In addition we showed that any description of neutrino superluminality based on a deformation of special relativity that preserves the relativity of inertial frames would automatically evade their concerns. The key to this evasion is an interplay between the deformations of the dispersion relations, lorentz transformations and conservation laws.

In order to illustrate our argument we made use of a simple, unrealistic model of particle dependent deformed lorentz symmetry. We have not addressed the question of whether there is a physically realistic model with deformed Lorentz transformations that can explain the data on neutrino speeds².

To explain the challenges faced by the construction of such a model we point out that there are again two possible classes of models. The first class involves non-linear realizations of the Lorentz transformations obtainable by nonlinear changes of variables in momentum space (see, *e.g.*, Refs. [14]–[17]). In this approach the momentum space picture is straightforward, but it is not clear how the processes are to be interpreted in spacetime. The toy model we here used to illustrate our observation belongs to this class. It exploits, however, the previously unexplored feature of having deformations of the lorentz transformations that are particle dependent. In particular only the neutrino dispersion relations and transformation laws need be deformed in relation to the OPERA anomaly. The associated spacetime picture should formally allow for neutrinos to live in a slightly different spacetime than the other particles. Interpreting v in eqs. (2) and (6) as a velocity and insisting also on the relativity of inertial frames one has to face the apparent non-locality of distant events [23–26]. This may be tolerable in the neutrino sector, since very little has been established experimentally about the spacetime localization of neutrinos.

The second approach and class of models goes under the name of *relative locality framework* [21, 22]. This new principle is realized in a class of models in which the deformations arise from modifications of the geometry of momentum space such as curvature. Also in these models different particles would appear to live in different spacetimes, because x and t are conjugate to coordinates on a curved momentum space. In this case there is also a large class of curved momentum space diffeomorphisms transformations, and it turns out that apparent non-localities affecting processes distant from an observer (characteristic of scenarios with particles “living in different spacetimes”) are gauge artifacts. Gauge invariant quantities corresponding to physical arrival times can be computed and reproduce the velocities (2) and (6) [27, 28]. In such a relative locality model, one could hope to provide a comprehensive explanation of the OPERA anomaly without running into the problems raised by Refs. [2–5].

This remains a task for the future. What we have shown here is that deformed Lorentz symmetry provides a large class of counter examples to the issues raised in [2–5]. If the OPERA results are confirmed, theorists will face a formidable challenge (see Refs. [8, 18, 29–34] for some of the attempts to deal with this challenge.) These challenges include developing quantum field theories consistent with relative locality; efforts to do so are underway. What we have observed in this paper is that the challenge of understanding the OPERA results-if they are confirmed-is more likely to succeed within the framework of deformed rather than broken Lorentz invariance.

ACKNOWLEDGEMENTS

We are grateful to many members of the Perimeter Institute community for their insights, comments and encouragement and also thank Andrew Cohen for correspondence. GAC and JKG thank Perimeter Institute for hospitality during their visits in September 2011. The work of JKG was supported in part by grants 182/N-QGG/2008/0 and 2011/01/B/ST2/03354. Research

² Neutrinos in the context of deformed special relativity have been studied in [20].

at Perimeter Institute for Theoretical Physics is supported in part by the Government of Canada through NSERC and by the Province of Ontario through MRI.

-
- [1] T. Adam *et al*, arXiv:1109.4897
 - [2] A.G. Cohen and S.L. Glashow, arXiv:1109.6562, Phys. Rev. Lett. **107** (2011) 181803
 - [3] L. Gonzalez-Mestres, arXiv:1109.6630
 - [4] X.-J. Bi, P.-F. Yin, Z.-H. Yu and Q. Yuan, arXiv:1109.6667, Phys. Rev. Lett. **107** (2011) 241802
 - [5] R. Cowsik, S. Nussinov and U. Sarkar, arXiv:1110.0241, Phys. Rev. Lett. **107** (2011) 251801.
 - [6] Z.G.T. Guiragossian, G.B. Rothbart, M.R. Yearian, R. Gearhart and J.J. Murray, Phys. Rev. Lett. **34** (1975) 335.
 - [7] A.A. Abdo *et al*, Nature **462** (2009) 331.
 - [8] G. Amelino-Camelia, G. Gubitosi, N. Loret, F. Mercati, G. Rosati, P. Lipari, arXiv:1109.5172, Int. J. Mod. Phys. **D20** (2011) 2623.
 - [9] D. Colladay, V. A. Kostelecky, hep-ph/9703464, Phys. Rev. **D55** (1997) 6760
 - [10] S.R. Coleman and S.L. Glashow, hep-ph/9703240, Phys.Lett. **B405** (1997) 249
 - [11] G. Amelino-Camelia, J. Ellis, N.E. Mavromatos, D.V. Nanopoulos and S. Sarkar, astro-ph/9712103, Nature **393** (1998) 763
 - [12] G. Amelino-Camelia, hep-th/0012238, Phys. Lett. **B510** (2001) 255; gr-qc/0012051, Int. J. Mod. Phys. **D11** (2002) 35
 - [13] J. Kowalski-Glikman, hep-th/0102098, Phys. Lett. **A286** (2001) 391.
 - [14] J. Magueijo, L. Smolin, hep-th/0112090, Phys. Rev. Lett. **88** (2002) 190403; gr-qc/0207085, Phys. Rev. **D67** (2003) 044017.
 - [15] S. Judes and M. Visser, Phys. Rev. **D68** (2003) 045001
 - [16] G. Amelino-Camelia, gr-qc/0212002, New J.Phys. **6** (2004) 188
 - [17] D. Heyman, F. Hinteleitner, and S. Major, gr-qc/0312089, Phys. Rev. **D69** (2004) 105016
 - [18] J. Magueijo, arXiv:1109.6055.
 - [19] L. Smolin, *Note on the OPERA neutrino speed data and relative locality*, unpublished, Sept 25, 2011.
 - [20] M. Blasone, J. Magueijo and P. Pires-Pacheco, Europhys. Lett. **70** (2005) 600; Braz. J. Phys. **35** (2005) 447-454.
 - [21] G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman and L. Smolin, arXiv:1101.0931 Phys. Rev. **D84** (2011) 084010.
 - [22] G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman and L. Smolin, arXiv:1106.0313, Gen. Relativ. Gravit. **43** (2011) 2547.
 - [23] R. Schutzhold and W. G. Unruh, gr-qc/0308049, JETP Lett. **78** (2003) 431.
 - [24] S. Hossenfelder, arXiv:1004.0418, Phys. Rev. Lett. **104** (2010) 140402
 - [25] G. Amelino-Camelia, M. Matassa, F. Mercati and G. Rosati, arXiv:1006.2126, Phys. Rev. Lett. **106** (2011) 071301
 - [26] L. Smolin, arXiv:1007.0718.
 - [27] L. Freidel and L. Smolin, arXiv:1103.5626.
 - [28] G. Amelino-Camelia, M. Arzano, J. Kowalski-Glikman, G. Rosati and G. Trevisan, arXiv:1107.1724, Class. Quant. Grav. (2012, in press)
 - [29] G. Cacciapaglia, A. Deandrea and L. Panizzi, arXiv:1109.4980, JHEP **11** (2011) 137
 - [30] G. F. Giudice, S. Sibiryakov and A. Strumia, arXiv:1109.5682
 - [31] J. Alexandre, J. Ellis and N.E. Mavromatos, arXiv:1109.6296
 - [32] C. Pfeifer and M.N.R. Wohlfarth, arXiv:1109.6005
 - [33] A. Drago, I. Masina, G. Pagliara and R. Tripiccone, arXiv:1109.5917
 - [34] G. Dvali and A. Vikman arXiv:1109.5685