Chiral phase transition at finite temperature and conformal dynamics in large N_f QCD

Kohtaroh Miura[a], Maria Paola Lombardo[a,b], Elisabetta Pallante[c]

[a] *INFN Laboratori Nazionali di Frascati, I-00044, Frascati (RM), Italy*
[b] *Humboldt-Universität zu Berlin, Institut für Physik, D-12489 Berlin, Germany*
[c] *Centre for Theoretical Physics, University of Groningen, 9747 AG, Netherlands*

arXiv:1110.3152v2 [hep-lat] 9 Mar 2012

Abstract

We investigate the chiral phase transition at finite temperature (T) in colour SU($N_c = 3$) Quantum Chromodynamics (QCD) with a variable number of fermions N_f in the fundamental representation by using lattice QCD. For $N_f = 6$ we study the approach to asymptotic scaling by considering lattices with several temporal extensions N_t. We then extract the dimensionless ratio $T_c/\Lambda_{\rm L}$ ($\Lambda_{\rm L}$ = Lattice Lambda-parameter) for $N_f = 6$ and $N_f = 8$, the latter relying on our earlier results. Further, we collect the (pseudo) critical couplings $\beta_{\rm L}^{\rm c}$ for the chiral phase transition at $N_f = 0$ (quenched), and $N_f = 4$ at a fixed $N_t = 6$. The results are consistent with enhanced fermionic screening at larger N_f. The ratio $T_c/\Lambda_{\rm L}$ depends very mildly on N_f in the $N_f = 0 - 4$ region, starts increasing at $N_f = 6$, and becomes significantly larger at $N_f = 8$, close to the edge of the conformal window. We discuss interpretations of these results as well as their possible interrelation with preconformal dynamics in the light of a functional renormalization group analysis.

Keywords: Lattice Gauge Theory, Conformal Symmetry, Chiral Symmetry, Finite Temperature

INTRODUCTION Conformal invariance is anticipated to emerge in asymptotically free non-Abelian gauge theories when the number of flavours exceeds a critical value $N_f = {N_f}^c$. The approach to conformality for $N_f \lesssim {N_f}^c$, close to the edge of the conformal window, is in principle associated with a walking behaviour of the running coupling, which has been advocated as a basis for strongly interacting mechanisms of electroweak symmetry breaking [1].

Recent lattice studies [2] focused on the computation of ${N_f}^c$ and the analysis of the conformal window itself, either with fundamental fermions [3, 4, 5, 6, 7, 8, 9, 10] or other representations [11, 12, 13, 14]. Among the many interesting results with fundamental fermions, we single out the observation that QCD with three colours and eight flavours is still in the hadronic phase [4, 5], while $N_f = 12$ seems to be close to the critical number of flavours, with some groups favouring conformality [3, 4, 6, 7], and others chiral symmetry breaking [9].

In comparison, much less effort has been devoted to the analysis of the phenomenologically relevant subcritical region [15, 16, 17, 18]. Here, we would like to learn where and how QCD at the edge of the conformal window displays walking, and its associated manifestations like separation of scales and approximate scale invariance. This Letter is one step in this direction.

Building on the above mentioned results, we decided to concentrate ourselves on $N_f \leq 8$, so to be safely in the hadronic region, but not too far from the edge of the conformal window. A recent study [15] noted an enhancement of the zero temperature ratio $\langle\bar{\psi}\psi\rangle/F^3$, where F is the pseudoscalar decay constant. This suggests that $N_f = 6$ might indeed be the onset of new strong dynamics.

In this letter we study the thermal transition of QCD with N_f = 0,4,6, and combine our findings with those of our early work for $N_f = 8$ [5]. We confirm the expected enhanced screening when the number of flavours increases, and discuss the interrelation of our results with a possible emergence of a new, preconformal dynamics.

As a general remark, we note that using the thermal transition as a tool for investigating preconformal dynamics was largely inspired by a renormalization group analysis [19], as we will review below. Further reasons of interest for finite temperature studies of large N_f include a connection between the quark-gluon plasma phase and the cold conformal region, which might lend support to analyses of quark-gluon plasma based on the AdS/CFT correspondence.

BRIEF OVERVIEW OF ANALYTIC RESULTS A second zero of the two-loop beta-function of a non-Abelian gauge theory implies, at least perturbatively, the appearance of an infrared fixed point (IRFP) and the restoration of conformal symmetry [20, 21]. In colour SU(3) gauge theory with N_f massless fermions in the fundamental representation, the second zero appears for $N_f \gtrsim 8.05$, before the loss of asymptotic freedom (LAF) at $N_f^{\rm LAF} = 16.5$.

One expects that conformality should emerge when the renormalized coupling at the would be IRFP is not strong enough to break chiral symmetry. This condition provides the lower bound N_f^c of a so called conformal window in N_f. Analytic studies based on the Schwinger-Dyson equation with rainbow resummations [22, 23, 24] or the functional renormalization group method [19] suggest $N_f^c \sim 12$. An all-order pertur-

bative beta-function [25] inspired by the NSVZ beta-function of SQCD [26] has been conjectured, leading to a bound $N_f^c > 8.25$; N_f^c has also been estimated for different fermion representations [27]. In addition to the lower bound of the conformal window, walking dynamics in the preconformal region is another interesting subject in relation to proposed scenarios of walking technicolor. Instanton studies at large N_f [28] claimed a qualitative change of behaviour at $N_f = 6$.

All the above phenomena happen well into the strong coupling regime, rendering a perturbative prediction unreliable and a non-perturbative analysis mandatory. The genuinely non-perturbative lattice formulation of gauge theories is thus a natural candidate for this study.

Recently, the functional renormalization group (FRG) method has been applied to finite T QCD with varying number of flavours, and the critical temperature for the chiral phase transition was obtained as a function of N_f [19]. In this $T - N_f$ phase diagram, the onset of the conformal window has been estimated by locating the vanishing critical temperature.

Most interestingly, the critical exponents associated with the behaviour of the beta function at the IRFP manifest themselves also in the shape of the thermal critical line in the vicinity of the critical number of flavours N_f^c. In more detail, the line is almost linear with N_f for small N_f, and displays a singular behaviour when approaching N_f^c. As emphasised by the authors in Ref. [19], the result clearly elucidates the universality of the critical behaviour at zero and non-zero temperature in the vicinity of N_f^c. It thus seems a promising direction to extend the knowledge of finite T lattice QCD to the larger N_f region, by using the FRG results as analytic guidance.

In this work we investigate the thermal chiral phase transition for $N_f = 0, 4, 6, 8$ colour SU($N_c = 3$) QCD by using lattice QCD Monte Carlo simulations with staggered fermions. $N_f = 6$ is expected to be in the important regime as suggested by the results in Refs. [3, 28]. This work includes the first study of $N_f = 6$ staggered fermions at finite T, and it provides an important ingredient to a broader project that studies the emergence of the conformal window in the $T - N_f$ phase diagram. In addition to $N_f = 6$, we compute the (pseudo) critical coupling for $N_f = 0$ (quenched) and $N_f = 4$ at $N_t = 6$, and use the results from Ref. [5] for $N_f = 8$.

In short, this work explores a largely uncharted territory: the chiral transition of strong interactions at high temperature, and large number of light flavours. Our goal is to observe, and understand possible qualitative differences with the very well known behaviour of QCD thermodynamics. To this end, we have for the first time collected and analysed results for different number of flavours, and the same lattice action, so to be able to meaningfully compare them. We ask the question: are we still finding just small differences among theories with different number of flavours, or are we going to observe some significant trend? As we will see, our exploratory, and in many respects qualitative analysis, will indicate that $N_f = 6$, and even more $N_f = 8$, are serious candidates for a different chiral dynamics. Obviously, our observations call for detailed quantitative studies which are already underway.

SETUP Simulations have been performed by utilising the publicly available MILC code [29]. The setup explained below is the same as the one used for $N_f = 8$ in Ref. [5]. We use an improved version of the staggered action, the Asqtad action, with a one-loop Symanzik [30, 31] and tadpole [34] improved gauge action,

$$S = -\frac{N_f}{4}\mathrm{Tr}\log M[am, U, u_0] + \sum_{i=p,r,pg}\beta_i(g_{\rm L}^2)\mathrm{Re}[1 - U_{C_i}]\,, \quad (1)$$

where $g_{\rm L}$ is the lattice bare coupling, and β_i are defined as

$$(\beta_p, \beta_r, \beta_{pg}) = \left(\frac{10}{g_{\rm L}^2}, -\frac{\beta_p(1 - 0.4805\alpha_s)}{20u_0^2}, -\frac{\beta_p}{u_0^2}0.03325\alpha_s\right) \quad (2)$$

$$\alpha_s = -4\log\frac{u_0}{3.0684}\,, \quad u_0 = \langle U_{C_p}\rangle^{1/4}\,. \quad (3)$$

The plaquette coupling $\beta_p = 10/g_{\rm L}^2 \equiv \beta_{\rm L}$ is a simulation input. The $M[am, U, u_0]$ in Eq. (1) denotes the matrix for a single flavour Asqtad fermion with bare lattice mass am, and U_{C_i} represents the trace of the ordered product of link variables along C_i, for the 1×1 plaquettes ($i = p$), the 1×2 and 2×1 rectangles ($i = r$), and the $1\times1\times1$ parallelograms ($i = pg$), respectively - all divided by the number of colours. The tadpole factor u_0 is determined by performing zero temperature simulations on the 12^4 lattice, and used as an input for finite temperature simulations.

To generate configurations with mass degenerate dynamical flavours, we have used the rational hybrid Monte Carlo algorithm (RHMC) [32], which allows to simulate an arbitrary number of flavours through varying the number of pseudo-fermions. Simulations for $N_f = 6$ have been performed by using two pseudo-fermions, and subsets of trajectories for the chiral condensates and Polyakov loop have been compared with those obtained by using three pseudo-fermions with the same Monte Carlo time step $d\tau$ and total time length τ of a single trajectory. We have observed very good agreement between the two cases for both evolution and thermalization. We have monitored the Metropolis acceptance and reject ratio, and adjusted $\tau = 0.2 - 0.24$ and $d\tau = 0.008 - 0.018$ to realize the best performance. For each parameter set, we have collected a number of trajectories ranging from a one thousand to five thousand - the latter closer to the critical region.

The focus of this letter is the analysis of the chiral transition. The fundamental observable is then the order parameter for chiral symmetry, the chiral condensate:

$$a^3\langle\bar{\psi}\psi\rangle = \frac{N_f}{4N_s^3N_t}\left\langle\mathrm{Tr}[\mathrm{M}^{-1}]\right\rangle\,, \quad (4)$$

where N_s (N_t) represents the number of lattice sites in the spatial (temporal) direction and $U_{4,\mathbf{tx}}$ is the temporal link variable. We have also measured connected and disconnected chiral susceptibilities,

$$a^2\chi_{\rm conn} = -\frac{N_f}{4N_s^3N_t}\langle\mathrm{Tr}\left[(MM)^{-1}\right]\rangle\,,$$
$$a^2\chi_{\rm disc} = \frac{N_f^2}{16N_s^3N_t}\left[\langle\mathrm{Tr}\left[M^{-1}\right]^2\rangle - \langle\mathrm{Tr}\left[M^{-1}\right]\rangle^2\right]\,, \quad (5)$$

and we have considered the logarithmic derivative,

$$R_\pi \equiv \chi_\sigma/\chi_\pi \,, \tag{6}$$

where,

$$\chi_\sigma \equiv \chi = \frac{\partial\langle\bar\psi\psi\rangle}{\partial m} = \chi_{\rm conn} + \chi_{\rm disc} \,, \tag{7}$$

and

$$\chi_\pi = \frac{\langle\bar\psi\psi\rangle}{m} \,. \tag{8}$$

As discussed in previous work [5, 33], R_π is a probe of chiral symmetry which is particularly useful for numerical investigations. To further characterize the critical region, we also measured the Polyakov loop,

$$L = \frac{1}{N_c N_s^3}\sum_{\mathbf{x}} {\rm Re}\Big\langle {\rm tr}_c \prod_{t=1}^{N_t} U_{4,t\mathbf{x}}\Big\rangle \,, \tag{9}$$

and tr_c denotes the trace in colour space.

The temperature T is related to the inverse of the lattice temporal extension,

$$T \equiv \frac{1}{a(\beta_{\rm L})\cdot N_t} \,. \tag{10}$$

We measure $\langle\bar\psi\psi\rangle$, the chiral susceptibilities and L at various temperatures. The output of this measurement is the (pseudo) critical coupling $\beta_{\rm L}^{\rm c}$ for the chiral phase transition for a given value of N_t. We underscore that all the measurements of the pseudo-critical couplings which will be used in our discussion are only based on fermionic observables.

RESULTS All results have been obtained for a fermion bare lattice mass $am = 0.02$. In Figs. 1 and 2, the expectation values of the chiral condensate $a^3\langle\bar\psi\psi\rangle$, and the Polyakov loop L are displayed as a function of $\beta_{\rm L}$ for several N_t, respectively.It is found that different N_t give a different behaviour of $a^3\langle\bar\psi\psi\rangle$ and L. In particular, this indicates that their rapid crossover with increasing $\beta_{\rm L}$ is not to be attributed to a bulk transition. The asymptotic scaling analysis below will confirm that it corresponds instead to a thermal chiral phase transition (or crossover) in the continuum limit. [1]

For $N_t = 4$, it is possible to extract $\beta_{\rm L}^{\rm c} = 4.65(25)$ from the peak position of the first derivative $-a^3 d\langle\bar\psi\psi\rangle/d\beta_{\rm L}$. We note that at this level of accuracy we cannot disentangle the peak position for the derivatives of the chiral condensate and the Polyakov loop $dL/d\beta_{\rm L}$.

For $N_t = 6$, we find a small jump between $\beta_{\rm L} = 5.0$ and 5.05, where the Polyakov loop also shows a significant enhancement.

For $N_t = 8$, the Polyakov loop L shows a clear signal as indicated in Fig. 3. In particular, we observe a drastic increase of the Polyakov loop L around $5.2 < \beta_{\rm L} < 5.3$. The histogram of the chiral condensate around the rapid increase of the Polyakov loop is shown in Fig. 4. At $\beta_{\rm L} = 5.2$, the histogram exhibits a broadening that suggests the increase of fluctuations around the pseudo-critical point.

For $N_t = 12$ the results are particularly smooth. Note that in this case the aspect ratio is only two, and larger volumes would be required to reach a comparable clarity in the signal, although the onset for the Polyakov loop at $\beta = 5.50(5)$ is still appreciable.

The results for R_π are collected in Fig.5. R_π has a clear jump for $\beta_c = (4.65(5), 5.05(5), 5.20(5))$ for $N_t = (4, 6, 8)$. For $N_t = 12$, the behaviour is again rather smooth. We have then searched for an inflection point in R_π by fitting to a hyperbolic tangent in several intervals, varying the extrema of the integrations. The results are reasonably stable in the errors. We will quote as a central value the average of the fit results within different intervals, and for the error, we will use the conservative estimate of half the difference between the largest and smallest result, obtaining $\beta_c = 5.46(14)$. All values of the (pseudo) critical lattice coupling $\beta_{\rm L}^{\rm c}$ are summarized in Table 1.

These results can be analyzed and interpreted in terms of the two-loop asymptotic scaling law. Let us consider the two-loop lattice beta function,

$$\beta(g) = -(b_0 g^3 + b_1 g^5) \,, \tag{11}$$

$$b_0 = \frac{1}{(4\pi)^2}\left(\frac{11C_2[G]}{3} - \frac{4T[F]N_f}{3}\right), \tag{12}$$

$$b_1 = \frac{1}{(4\pi)^4}\left(\frac{34(C_2[G])^2}{3} - \left(\frac{20C_2[G]}{3} + 4C_2[F]\right)T[F]N_f\right), \tag{13}$$

for fundamental fermions in $SU(N_c)$ ($C_2[G]$, $C_2[F]$, $T[F]$) = (N_c, $(N_c^2-1)/(2N_c)$, $1/2$). From Eq. (11) we obtain the well known two-loop asymptotic scaling law,

$$\Lambda_{\rm L}\, a(\beta_{\rm L}) = \left(\frac{2N_c b_0}{\beta_{\rm L}}\right)^{-b_1/(2b_0^2)} \exp\left[\frac{-\beta_{\rm L}}{4N_c b_0}\right]. \tag{14}$$

Here $\Lambda_{\rm L}$ is the so called lattice Lambda-parameter, and $\beta_L = 2N_c/g^2$.

The above relation is valid in the massless limit. In the following, we will use it to analyze results obtained at finite mass. This assumes that the shift of the (pseudo) critical coupling induced by a non-zero mass is smaller than other errors. This assumption should ultimately be tested by performing simulations with different masses and extrapolating to the chiral limit.

The equation (10) can be written as

$$\frac{1}{N_t} = \frac{T_c}{\Lambda_{\rm L}} \times \left(\Lambda_{\rm L}\, a(\beta_{\rm L}^{\rm c})\right). \tag{15}$$

[1] A note on the mass dependence is in order. According to the Pisarski-Wilczek scenario, the most likely possibility for $N_f \geq 3$ is a first order chiral transition in the chiral limit. Standard arguments indicate that first order phase transitions are robust against explicit breaking: when introducing a bare quark mass, then, we expect a first order phase transition which will eventually end in a genuine singularity at some critical point (T_c, m_c). By further increasing the bare mass, the transition will turn into a crossover. In the unexpected situation of a second order transition in the chiral limit, any nonzero quark mass will immediately produce a crossover. Even in the case of a first order transition, though, a finite lattice will turn it into a crossover. All in all, as in any lattice study, we are never dealing directly with a genuine criticality. Rather, we are locating a pseudo-critical point, and only by considering several masses and several volumes, we can assess with confidence the nature of the phase transition in the infinite volume and in the chiral limit. Discriminating among these different behaviours is however beyond the scope of this study.

The left-hand side is a given number, and we have obtained the corresponding $\beta_{\rm L}^{\rm c}$ by lattice simulations. Hence, the quantity $T_c/\Lambda_{\rm L}$ on the right-hand side of Eq. (15) can be extracted, and must be unique as long as the asymptotic scaling law Eq. (14) is verified for a given $\beta_{\rm L}^{\rm c}$.

If we use the lattice bare coupling to carry out the above program, we notice appreciable scaling violations. Indeed, Eq. (14) holds true up to non-universal scaling-violating terms. One possibility is then to follow earlier work [35] and to parametrize scaling violations as

$$\Lambda_{\rm L}\, a(\beta_{\rm L}) \equiv [\Lambda_{\rm L} a(\beta_{\rm L})]_{\rm 2loops}\Big(1 + h[\Lambda_{\rm L} a(\beta_{\rm L})]^2_{\rm 2loops}\Big)\,. \qquad (16)$$

where $[\Lambda_{\rm L} a(\beta_{\rm L})]_{\rm 2loops}$ is defined by Eq. (14).

Alternatively, we can trade the bare lattice coupling $g_{\rm L}$ for the boosted coupling introduced in our previous work [5] $g = \sqrt{2N_c/10}\cdot g_{\rm L}$. We will show below that this prescription leads to a rather accurate two-loop scaling, equivalent to the enhancement of the scaling behaviour obtained by considering Eq. (14). Our prescription is similar in spirit to the Parisi-Lepage-Mackenzie [34] boosted coupling. Our boosted coupling, however, cannot be derived in (tadpole improved) perturbation theory. In our future work, we plan to perform zero temperature measurements which will allow an independent estimate of the lattice spacing, and a complete discussions of scaling, and asymptotic scaling. Hopefully, this will shed light on the reasons why our simple approach works so well. At this stage, admittedly, it remains a heuristic, ad-hoc prescription which effectively incorporates the scaling violations in the range we have explored. This warning issued, we will continue the discussion by using our boosted coupling.

In Fig. 6, we show the $N_t^{-1} - \Lambda_{\rm L} a(\beta_{\rm L}^{\rm c})$ plot. The slope of the line connecting the origin and the data points corresponds to $T_c/\Lambda_{\rm L}$. The $N_t = 6,\ 8$, and 12 points have a common slope to a very good approximation, while the $N_t = 4$ result falls on a smaller slope.

The latter is interpreted as a scaling violation effect due to the use of a too small N_t. The existence of a common $T_c/\Lambda_{\rm L}$ for $N_t \geq 6$ indicates that the data are consistent with the two-loop asymptotic scaling Eq. (14), confirms the thermal nature of the transition and that $N_f = 6$ is outside the conformal window, as expected from a previous $N_f = 8$ study [5]. A linear fit provides $T_c/\Lambda_{\rm L} = 1.02(12)\times 10^3$, which can be interpreted as the value in the continuum limit for $N_f = 6$ QCD. Note that this compares very well with the result obtained by using $\beta_{\rm L}^{\rm c}$ obtained by $N_t = 8$ simulations. In the following, we can then use it as a representative result for six flavours.

In order to have a more complete overview, we have performed simulations for the theory with $N_f = 0$ (quenched) and $N_f = 4$, only at $N_t = 6$. These theories are of course very well investigated, however we have not found in the literature results for the same action as ours. Table 1 shows a summary of our results for the (pseudo) critical coupling $\beta_{\rm L}^{\rm c}$ of the chiral phase transition at finite temperature for $N_f = 0,\ 4,\ 6$, and 8 - the latter from Ref. [5].

DISCUSSION In Fig. 7, we display the (pseudo) critical values of the lattice coupling $g_c = \sqrt{2N_c/\beta_{\rm L}^{\rm c}}$ from Table 1 in




Figure 1: The chiral condensate $a^3\langle\bar{\psi}\psi\rangle$ for $N_f = 6$ and $am = 0.02$ in lattice units, as a function of $\beta_{\rm L}$, for $N_t = 4,\ 6,\ 8$, and 12. Error-bars are smaller than symbols.




Figure 2: The Polyakov loop L for $N_f = 6$ and $am = 0.02$ in lattice units, as a function of $\beta_{\rm L}$, for $N_t = 4,\ 6,\ 8$, and 12. Error-bars are smaller than symbols.

the Miransky-Yamawaki phase diagram.

Consider the $N_t = 6$ results: it is expected that an increasing number of flavours favors chiral symmetry restoration. Indeed, we find that, on a fixed lattice, the (pseudo) critical coupling increases with N_f in agreement with early studies and naive reasoning. The precise dependence of the (pseudo) critical coupling on N_f at fixed N_t is not known. It is, however, amusing to note that the results seem to be smoothly connected by an almost straight line: the brown line in the plot is a linear fit to the data. Comparing the trend for $N_f = 6$ to the one for $N_f = 8$, for varying N_t one can infer a decreasing in magnitude (and small) step scaling function, hence a walking behaviour. Further study is needed at larger N_f, and by using the same action used for $N_f = 0 - 8$, to confirm or disprove it.

Next, we study the N_f dependence of the ratio $T_c/\Lambda_{\rm L}$ and related quantities. We recall that the simulations for $N_f = 4$ and $N_f = 0$ have been performed by using only $N_t = 6$. Hence, in these two cases, the results will hold true barring strong scaling violations at $N_t = 6$. We note that in a previous lattice study with improved staggered fermions [36], asymptotic scaling was

Table 1: Summary of the (pseudo) critical lattice couplings β_L^c for the theories with $N_f = 0,\ 4,\ 6,\ 8$, $am = 0.02$ and varying $N_t = 4,\ 6,\ 8,\ 12$. All results are obtained using the same lattice action.

$N_f \backslash N_t$	4	6	8	12
0	-	7.88 ± 0.05	-	-
4	-	5.89 ± 0.03	-	
6	4.65 ± 0.05	5.05 ± 0.05	5.2 ± 0.05	5.45 ± 0.15
8	-	4.1125 ± 0.0125	-	4.34 ± 0.04




Figure 3: Zoom-in of the chiral condensate $a^3\langle\bar{\psi}\psi\rangle$ and the Polyakov loop L shown in Figs. 1 and 2 in the critical region at $N_t = 8$, with spatial volume 24^3.




Figure 4: Distribution of the chiral condensate $a^3\langle\bar{\psi}\psi\rangle$ for $N_f = 6$, $am = 0.02$ and spatial volume 24^3, in the vicinity of the chiral phase transition at $N_t = 8$.

indeed observed using a boosted coupling for $N_t \geq 6$ for $0 \leq N_f \leq 4$.

Ideally, we would like to convert our results to $T_c/\Lambda_{\overline{\rm MS}}$. Unfortunately, to our knowledge, the conversion from Λ_L to $\Lambda_{\overline{\rm MS}}$ for a generic number of flavours is only available for Wilson fermions [37].

Here we consider a simplified procedure, aiming at capturing at least the basic features induced by setting a UV scale. For this purpose, we introduce a reference coupling $\beta_L^{\rm ref}$ and an associated reference energy scale $\Lambda_{\rm ref}$. Then Eq. (14) is generalized as

$$\Lambda_{\rm ref}(\beta_L^{\rm ref})\, a(\beta_L) = \left(\frac{b_1}{b_0^2}\,\frac{\beta_L + 2N_c b_1/b_0}{\beta_L^{\rm ref} + 2N_c b_1/b_0}\right)^{b_1/(2b_0^2)} \exp\left[-\frac{\beta_L - \beta_L^{\rm ref}}{4N_c b_0}\right]. \qquad (17)$$

At leading order of perturbation theory $b_1 \to 0$, Λ_L and $\Lambda_{\rm ref}$ are related via

$$\frac{\Lambda_{\rm ref}}{\Lambda_L} = \exp\left[\frac{\beta_L^{\rm ref}}{4N_c b_0}\right]. \qquad (18)$$

This equation would be analogous of the ratio $\Lambda_L/\Lambda_{\rm MS}$ derived in [37] for Wilson fermions up to a further linear dependence on N_f in the numerator of the exponent. In a nutshell, the difference originates from the fact that we are fixing a bare reference coupling $\beta_L^{\rm ref}$, which will be specified later. Notice that by construction $\Lambda_{\rm ref}$ reproduces the lattice Lambda-parameter Λ_L in the limit

$$\Lambda_{\rm ref}(\beta_L^{\rm ref} \to 0) = \Lambda_L\Big(1 + O(1/\beta_L^c)\Big). \qquad (19)$$

In summary, when trading Λ_L for $\Lambda_{\rm ref}$, we are moving towards a more UV scale.

Let us consider first T_c/Λ_L. The values of T_c/Λ_L are summarized in Table 2, and plotted in Fig. 8. The ratio does not show a significant N_f dependence in the region $0 \leq N_f \leq 4$, it starts increasing at $N_f = 6$, and undergoes a rapid rise around $N_f = 8$. The chiral phase transition would happen when T becomes comparable to a typical energy scale $M_\chi = C\Lambda_L$. The nearly constant nature of T_c/Λ_L in the region $N_f \leq 4$ indicates that the role of such energy scale is not significantly changed by the variation of N_f (see [38] for a detailed discussion of this point.) In turn, the increase of T_c/Λ_L in the region $N_f \geq 6$ might well imply that the chiral dynamics becomes different from the one for $N_f \leq 4$. Indeed, a recent lattice study [15] indicates that $N_f = 6$ is close to the threshold for preconformal dynamics.

We now consider $T_c/\Lambda_{\rm ref}$. The N_f dependence of the ratio $R(N_f) \equiv (T_c/\Lambda_{\rm ref})(N_f)$ is shown for several $\beta_L^{\rm ref}$ in Fig. 9, where the vertical axis is normalized by $R(0) = (T_c/\Lambda_{\rm ref})(N_f = 0)$ for each $\beta_L^{\rm ref}$. $T_c/\Lambda_{\rm ref}$ is now a decreasing function of N_f for a larger $\beta_L^{\rm ref}$, *i.e.* for a more UV reference scale $\Lambda_{\rm ref}$. This result is consistent with the FRG study [19], where the decreasing $T_c(N_f)$ has been obtained by using the τ-lepton mass m_τ as a common UV reference scale with a common coupling $\alpha_s(m_\tau)$.

The $\Lambda_{\rm ref}$ scale associated with a $\beta_L^{\rm ref} \gg \beta_*$ where β_* is eval-




Figure 5: The ratio of scalar and pseudo-scalar contributions to the susceptibility, defined in Eq. (6) as a function of β_L.




Figure 6: The thermal scaling behaviour of the (pseudo) critical lattice coupling β_L^c. Data points for $\Lambda_L\ a(\beta_L^c)$ at a given $1/N_t$ are obtained by using β_L^c from Table 1 as input for extracting $\Lambda_L\ a(\beta_L^c)$ in the two-loop expression Eq. (14). The dashed line is a linear fit with zero intercept to the data with $N_t > 4$.

uated at the infra-red fixed point should provide a UV scale well-separated from the IR dynamics. If we assume the lower bound of the conformal window to be $N_f^c \simeq 12$, the two-loop beta-function leads to $\beta_* = -2N_c b_1/b_0 \simeq 0.63$. Indeed Fig. 9 shows that the decreasing nature of $(T_c/\Lambda_{\rm ref})(N_f)$ is still weak at $\beta_L^{\rm ref} = 1.0$. In the limit $\beta_L^{\rm ref} \to 0$, $T_c/\Lambda_{\rm ref}$ reproduces Fig. 8, and the resultant increasing feature should be attributed to the vanishing of Λ_L due to infra-red dynamics. We also notice that $\beta_L^{\rm ref}$ must always be smaller than β at the UV cutoff, $\beta_{\rm UV} = \beta_L^c(N_f)$. As shown in Table 1, the lowest value of the (pseudo) critical coupling is given by $\beta_L^c(N_f = 8, N_t = 6) = 4.1125 \pm 0.0125$, hence we constrain our analyses to $\beta_L^{\rm ref} \leq 4.0$. In summary, Figs. 8 and 9 together show the effects of shifting the reference scales from the IR to the UV.

With the use of a UV reference scale, we should observe the predicted critical behavior [19]

$$T_c(N_f) = K|N_f - N_f^c|^{-1/\theta}\ . \qquad (20)$$

By choosing the critical exponent θ in the range predicted by




Figure 7: (Pseudo) critical values of the lattice coupling $g_c = \sqrt{2N_c/\beta_L^c}$ for theories with $N_f = 0,\ 4,\ 6,\ 8$ and for several values of N_t in the Miransky-Yamawaki phase diagram. The dashed (brown) line is a linear fit to the $N_t = 6$ results.

FRG: $1.1 < 1/|\theta| < 2.5$, our data are consistent with the values $N_f^c = 9(1)$ for $\beta_L^{\rm ref} = 4.0$ and $N_f^c = 11(2)$ for $\beta_L^{\rm ref} = 2$. We plan to extend and refine this analysis in the future, and here we only notice a reasonable qualitative behaviour.

Table 2: T_c/Λ_L for several N_f. Results are obtained by using the same lattice action. For $N_f = 6$, we have used the $N_t = 8$ result as a representative value. The values for $N_f = 8$ are extracted from Ref. [5].

N_f	T_c/Λ_L
0	600 ± 34
4	620 ± 28
6	1000 ± 92
8	2098 ± 191

SUMMARY We have investigated the chiral phase transition and its asymptotic scaling for $N_f = 6$ colour SU(3) QCD by using lattice QCD Monte Carlo simulations with improved staggered fermions. This study provides an important ingredient to a broader project that studies the emergence of a conformal window in the $T - N_f$ phase diagram. We have determined the (pseudo) critical lattice coupling β_L^c for several lattice temporal extensions N_t. We have extracted the dimensionless ratio T_c/Λ_L (Λ_L =Lattice Lambda-parameter) for the theory with $N_f = 6$ using two-loop asymptotic scaling. The analogous result for $N_f = 8$ has been extracted from Ref. [5]. T_c/Λ_L for $N_f = 0$ and $N_f = 4$ has been measured at fixed $N_t = 6$, barring asymptotic scaling violations. Then we have discussed the N_f dependence of the ratios T_c/Λ_L and $T_c/\Lambda_{\rm ref}$, where $\Lambda_{\rm ref}$ is a UV reference energy scale, related to Λ_L as in Eq. (18).

We have observed that T_c/Λ_L shows an increase in the region $N_f = 6$–8, while it is approximately constant in the region $N_f \leq 4$. We have discussed this qualitative change for $N_f \geq 6$ and a possible relation with a preconformal phase. We repeat that all results have been obtained by working at one value of the quark mass and this is a potential weakness of our calculations.




Figure 8: The ratio $T_c/\Lambda_{\rm L}$, for $N_f = 0$, 4, 6 and 8 and lattice bare mass $am = 0.02$.




Figure 9: The N_f dependence of $R(N_f)/R(0)$ for several finite fixed $\beta_{\rm L}^{\rm ref}$. Here, $R(N_f) \equiv (T_c/\Lambda_{\rm ref})(N_f)$. The limit $\beta_{\rm L}^{\rm ref} \to 0$ reproduces the results shown in Fig. 8 up to a renormalization factor and and up to a corrections $O(1/\beta_{\rm L}^{\rm c})$.

The ratio $T_c/\Lambda_{\rm ref}$ is a decreasing function of N_f. This behaviour is consistent with the result obtained in the functional renormalization group analysis [19], where a common UV reference scale was used to study the chiral phase boundary in the $T - N_f$ phase diagram.

Next steps of the current project involve a scale setting at zero temperature by measuring a common UV observable. It would also be desirable to have the relation between $\Lambda_{\rm L}$ and $\Lambda_{\overline{\rm MS}}$ for our action.

This, together with a more extended set of flavour numbers, will allow a quantitative analysis of the critical behaviour. We expect the resultant $T_c - N_f$ phase diagram to play an essential role in the study of the conformal window.

ACKNOWLEDGMENTS We thank Holger Gies and Jens Braun for fruitful discussions and most useful suggestions. We have enjoyed discussing these topics with Koichi Yamawaki, Masafumi Kurachi, Hiroshi Ohki, Michael Müller-Preussker, Marc Wagner, Biagio Lucini, Volodya Miransky, Albert Deuzeman and Tiago Nunes da Silva. Kohtaroh Miura thanks Michael Müller-Preussker and the theory group in the Humboldt University for their hospitality. Kohtaroh Miura is partially supported by EU I3HP2-WP22. This work was in part based on the MILC Collaboration's public lattice gauge theory code. See http://www.physics.indiana.edu/~sg/milc.html for details. The numerical calculations were carried out on the IBM-SP6 at CINECA, Italian-Grid-Infrastructures in Italy, and the Hitachi SR-16000 at YITP, Kyoto University in Japan.

References

[1] For a recent review, see F. Sannino, Acta Phys. Polon. B **40** (2009) 3533.
[2] For recent reviews, see L. Del Debbio, PoS **LATTICE2010** (2010) 004; E. Pallante, PoS **LATTICE2009** (2009) 015.
[3] T. Appelquist, G. T. Fleming, M. F. Lin, E. T. Neil, D. A. Schaich, Phys. Rev. **D84**, 054501 (2011).
[4] T. Appelquist, G. T. Fleming and E. T. Neil, Phys. Rev. D **79** (2009) 076010;
Phys. Rev. Lett. **100** (2008) 171607 [Erratum-ibid. **102** (2009) 149902].
[5] A. Deuzeman, M. P. Lombardo and E. Pallante, Phys. Lett. B **670** (2008) 41.
[6] A. Deuzeman, M. P. Lombardo and E. Pallante, Phys. Rev. D **82** (2010) 074503.
[7] A. Hasenfratz, Phys. Rev. D **82** (2010) 014506.
[8] A. Hasenfratz, Phys. Rev. D **80** (2009) 034505.
[9] Z. Fodor, K. Holland, J. Kuti, D. Nogradi and C. Schroeder, Phys. Lett. **B703** (2011) 348-358.
[10] Z. Fodor, K. Holland, J. Kuti, D. Nogradi and C. Schroeder, Phys. Lett. B **681** (2009) 353.
[11] T. Karavirta, A. Mykkanen, J. Rantaharju, K. Rummukainen and K. Tuominen, JHEP **1106** (2011) 061; A. J. Hietanen, K. Rummukainen and K. Tuominen, Phys. Rev. D **80** (2009) 094504; A. J. Hietanen, J. Rantaharju, K. Rummukainen and K. Tuominen, JHEP **0905** (2009) 025.
[12] Y. Shamir, B. Svetitsky and E. Yurkovsky, Phys. Rev. D **83** (2011) 097502; O. Machtey and B. Svetitsky, Phys. Rev. D **81** (2010) 014501; Y. Shamir, B. Svetitsky and T. DeGrand, Phys. Rev. D **78** (2008) 031502.
[13] J. B. Kogut and D. K. Sinclair, Phys. Rev. D **81** (2010) 114507.
[14] Z. Fodor, K. Holland, J. Kuti, D. Nogradi and C. Schroeder, JHEP **0911** (2009) 103.
[15] T. Appelquist *et al.*, Phys. Rev. Lett. **104** (2010) 071601.
[16] T. Appelquist *et al.* [LSD Collaboration], Phys. Rev. Lett. **106** (2011) 231601.
[17] S. Catterall, J. Giedt, F. Sannino and J. Schneible, JHEP **0811** (2008) 009; S. Catterall and F. Sannino, Phys. Rev. D **76** (2007) 034504.
[18] L. Del Debbio, B. Lucini, A. Patella, C. Pica and A. Rago, Phys. Rev. D **82** (2010) 014510; **82** (2010) 014509.
[19] J. Braun, C. S. Fisher, H. Gies, Phys. Rev. **D84** (2011) 034045; J. Braun and H. Gies, JHEP **1005** (2010) 060; **0606** (2006) 024.
[20] W. E. Caswell, Phys. Rev. Lett. **33** (1974) 244.
[21] T. Banks and A. Zaks, Nucl. Phys. B **196** (1982) 189.
[22] T. Appelquist, J. Terning and L. C. R. Wijewardhana, Phys. Rev. Lett. **77** (1996) 1214; T. Appelquist, A. Ratnaweera, J. Terning and L. C. R. Wijewardhana, Phys. Rev. D **58** (1998) 105017.
[23] V. A. Miransky and K. Yamawaki, Phys. Rev. D **55** (1997) 5051 [Erratum-ibid. D **56** (1997) 3768].
[24] T. Appelquist, A. G. Cohen and M. Schmaltz, Phys. Rev. D **60** (1999) 045003.
[25] T. A. Ryttov and F. Sannino, Phys. Rev. D **78** (2008) 065001.
[26] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **229**, 381 (1983).
[27] D. D. Dietrich and F. Sannino, Phys. Rev. D **75** (2007) 085018.
[28] M. Velkovsky and E. V. Shuryak, Phys. Lett. B **437** (1998) 398.
[29] MILC Collaboration, http://www.physics.indiana.edu/~sg/milc.html
[30] C. Bernard *et al.*, Phys. Rev. D **75** (2007) 094505.
[31] M. Luscher and P. Weisz, Phys. Lett. B **158** (1985) 250; Commun. Math. Phys. **97** (1985) 59 [Erratum-ibid. **98** (1985) 433].
[32] M. A. Clark, PoS **LAT2006** (2006) 004.

[33] A. Kocic, J. B. Kogut and M. -P. Lombardo, Nucl. Phys. B **398** (1993) 376.
[34] G. P. Lepage, P. B. Mackenzie, Phys. Rev. D **48** (1993) 2250.
[35] M. Cheng, N. H. Christ, S. Datta, J. van der Heide, C. Jung, F. Karsch, O. Kaczmarek and E. Laermann *et al.*, Phys. Rev. D **77** (2008) 014511.
[36] S. Gupta, Phys. Rev. D **64** (2001) 034507.
[37] H. Kawai, R. Nakayama and K. Seo, Nucl. Phys. B **189** (1981) 40.
[38] J. Braun, Phys. Rev. **D81** (2010) 016008.