

# A left-right symmetric model with $SU(2)$ -triplet fermions

Pei-Hong Gu\*

Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

We consider an  $SU(3)_c \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$  left-right symmetric model with three Higgs scalars including an  $SU(2)_L$  doublet, an  $SU(2)_R$  doublet and an  $SU(2)_L \otimes SU(2)_R$  bidoublet. In addition to usual  $SU(2)$ -doublet fermions, our model contains  $SU(2)$ -triplet fermions with Majorana masses. The neutral components of the left-handed triplets can contribute a canonical seesaw while the neutral components of the right-handed triplets associated with the right-handed neutrinos can contribute a double/inverse-type seesaw. Our model can be embedded into an  $SO(10)$  grand unification theory where the triplets belong to the  $\mathbf{45} = (\mathbf{1}, \mathbf{3}, \mathbf{1}, 0) \oplus (\mathbf{1}, \mathbf{1}, \mathbf{3}, 0) \oplus \dots$  representations.

PACS numbers: 14.60.Pq, 12.60.Cn, 12.60.Fr

In the  $SU(3)_c \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$  left-right symmetric models [1], the standard model (SM) fermions plus the right-handed neutrinos are placed in the  $SU(2)_L$  or  $SU(2)_R$  doublets. With an appropriate Higgs content, we can generate the fermion masses by supplementing additional fermions or not. The minimal Higgs content seems to be an  $SU(2)_R$  doublet and an  $SU(2)_L$  doublet, which drive the left-right and electroweak symmetry breaking, respectively. With the  $SU(2)$ -doublet Higgs scalars, one can consider the universal [2] seesaw [3] scenario to generate the SM fermion masses and the neutrino masses by introducing  $SU(2)$ -singlet fermions. In the most popular version of the left-right symmetric models, the fermion fields only contain the usual  $SU(2)$  doublets while the Higgs fields contain two  $SU(2)$  triplets and one  $SU(2)_L \otimes SU(2)_R$  bidoublet. The right-handed Higgs triplet and the Higgs bidoublet are responsible for the left-right and electroweak symmetry breaking, respectively. The charged fermions can obtain their masses through the usual Yukawa interactions while the neutrinos can obtain their masses through the type-I [3] and II [4] seesaw mechanisms. In the original left-right symmetric model, the Higgs content contains two  $SU(2)$  doublets and one  $SU(2)_L \otimes SU(2)_R$  bidoublet. The right-handed Higgs doublet and the Higgs bidoublet account for the left-right and electroweak symmetry breaking, respectively. In this scenario, it is difficult to naturally explain the smallness of the neutrino masses. To solve this problem, one can introduce gauge-singlet fermions with Majorana masses to realize the linear [5] and double/inverse [6, 7] seesaw.

In this paper we shall extend the original left-right symmetric model by  $SU(2)$ -triplet fermions to generate the small neutrino masses in a nature way. This idea has been proposed in an early work [8]. However, the author explicitly broke the discrete left-right symmetry to protect the left-handed Higgs doublet from a nonzero vacuum expectation value (VEV). In consequence, a double/inverse-type seesaw induced by the neutral com-

ponents of the right-handed triplets associated with the right-handed neutrinos becomes the unique origin of the neutrino masses. We shall study the general scenario where the left-handed Higgs doublet has a nonzero VEV. Specifically, the neutrinos can also pick up their masses through a canonical seesaw induced by the neutral components of the left-handed triplets. The canonical seesaw even can dominate over the double/inverse-type seesaw. Furthermore, we will embed our model into the  $SO(10)$  grand unification theory (GUT). In the GUT context, the  $SU(2)$ -triplet fermions belong to the  $\mathbf{45}$  representations. A  $\mathbf{45}$  multiplet also contain a singlet fermion under the left-right gauge symmetry. The singlet fermions can result in a double/inverse-type seesaw and a linear seesaw.

We take the notations as follows:

$$\chi_L(\mathbf{1}, \mathbf{2}, \mathbf{1}, 1) = \begin{bmatrix} \chi_L^+ \\ \chi_L^0 \end{bmatrix}, \quad \chi_R(\mathbf{1}, \mathbf{1}, \mathbf{2}, 1) = \begin{bmatrix} \chi_R^+ \\ \chi_R^0 \end{bmatrix},$$

$$\phi(\mathbf{1}, \mathbf{2}, \mathbf{2}, 0) = \begin{bmatrix} \phi_1^0 & \phi_2^+ \\ \phi_1^- & \phi_2^0 \end{bmatrix} \quad (1)$$

are the Higgs scalars, while

$$q_L(\mathbf{3}, \mathbf{2}, \mathbf{1}, +\frac{1}{3}) = \begin{bmatrix} u_L \\ d_L \end{bmatrix}, \quad q_R(\mathbf{3}, \mathbf{1}, \mathbf{2}, +\frac{1}{3}) = \begin{bmatrix} u_R \\ d_R \end{bmatrix},$$

$$l_L(\mathbf{1}, \mathbf{2}, \mathbf{1}, -1) = \begin{bmatrix} \nu_L \\ e_L \end{bmatrix}, \quad l_R(\mathbf{1}, \mathbf{1}, \mathbf{2}, -1) = \begin{bmatrix} \nu_R \\ e_R \end{bmatrix},$$

$$T_L(\mathbf{1}, \mathbf{3}, \mathbf{1}, 0) = \begin{bmatrix} T_L^1 \\ T_L^2 \\ T_L^3 \end{bmatrix}, \quad T_R(\mathbf{1}, \mathbf{1}, \mathbf{3}, 0) = \begin{bmatrix} T_R^1 \\ T_R^2 \\ T_R^3 \end{bmatrix} \quad (2)$$

denote the fermions. It is convenient to rewrite the fermion triplets by the matrix representation,

$$T_{L,R} = \tau_1 T_{L,R}^1 + \tau_2 T_{L,R}^2 + \tau_3 T_{L,R}^3$$

$$= \begin{bmatrix} T_{L,R}^0 & \sqrt{2}T_{L,R}^+ \\ \sqrt{2}T_{L,R}^- & -T_{L,R}^0 \end{bmatrix} \quad (3)$$

\*Electronic address: peihong.gu@mpi-hd.mpg.de

with

$$T_{L,R}^{\pm} = \frac{T_{L,R}^1 \mp iT_{L,R}^2}{\sqrt{2}}, \quad T_{L,R}^0 = T_{L,R}^3. \quad (4)$$

The right-handed Higgs doublet  $\chi_R$  will develop a VEV

$$\langle \chi_R \rangle = \begin{bmatrix} 0 \\ v_R \end{bmatrix} \quad (5)$$

for the left-right symmetry breaking. Subsequently, the Higgs bidoublet will acquire a VEV

$$\langle \phi \rangle = \begin{bmatrix} v_1 & 0 \\ 0 & v_2 \end{bmatrix} \quad (6)$$

to break the electroweak symmetry. Because the left-handed Higgs doublet has the trilinear couplings with the right-handed Higgs doublet and the Higgs bidoublet, i.e.

$$V \supset \mu \chi_L^\dagger \phi \chi_R + \tilde{\mu} \chi_L^\dagger \tilde{\phi} \chi_R + \text{H.c.}, \quad (7)$$

it will pick up an induced VEV<sup>1</sup>,

$$\langle \chi_L \rangle = \begin{bmatrix} 0 \\ v_L \end{bmatrix} \quad \text{with} \quad v_L \simeq \frac{\mu v_R v_2 + \tilde{\mu} v_R v_1}{M_{\chi_L^0}^2} \sim \frac{\mu v_2 + \tilde{\mu} v_1}{v_R}, \quad (8)$$

which is a weak-scale value for  $\mu \sim \tilde{\mu} \sim v_R$  or a tiny value for  $\mu \sim \tilde{\mu} \ll v_R$ . The allowed Yukawa couplings should be

$$\begin{aligned} \mathcal{L}_Y = & -y_q \bar{q}_L \phi q_R - \tilde{y}_q \bar{q}_L \tilde{\phi} q_R - y_l \bar{l}_L \phi l_R - \tilde{y}_l \bar{l}_L \tilde{\phi} l_R \\ & - f_L \bar{l}_L^c i \tau_2 T_L \chi_L - f_R \bar{l}_R^c i \tau_2 T_R \chi_R + \text{H.c.} \end{aligned} \quad (9)$$

The fermion triplets further have the Majorana masses as below,

$$\mathcal{L}_M = -\frac{1}{4} M_L \text{Tr}(\bar{T}_L^c T_L) - \frac{1}{4} M_R \text{Tr}(\bar{T}_R^c T_R) + \text{H.c.} \quad (10)$$

Through the Yukawa couplings with the Higgs bidoublet, the charged fermions can obtain their masses, i.e.

$$\mathcal{L}_m \supset -m_d \bar{d}_L d_R - m_u \bar{u}_L u_R - m_e \bar{e}_L e_R + \text{H.c.}, \quad (11)$$

with

$$m_d = y_q v_2 + \tilde{y}_q v_1, \quad (12a)$$

$$m_u = y_q v_1 + \tilde{y}_q v_2, \quad (12b)$$

$$m_e = y_l v_2 + \tilde{y}_l v_1. \quad (12c)$$

As for the neutral fermions, including the left- and right-handed neutrinos as well as the neutral components of the left- and right-handed triplets, their mass term is given by

$$\begin{aligned} \mathcal{L}_m \supset & f_L v_L \bar{\nu}_L^c T_L^0 - \frac{1}{2} M_L (\bar{T}_L^0)^c T_L^0 - (y_l v_1 + \tilde{y}_l v_2) \bar{\nu}_L \nu_R \\ & + f_R v_R \bar{\nu}_R^c T_R^0 - \frac{1}{2} M_R (\bar{T}_R^0)^c T_R^0 + \text{H.c.} \\ = & -\frac{1}{2} [\bar{\nu}_L^c, (\bar{T}_L^0)^c] \begin{bmatrix} 0 & -f_L v_L \\ -f_L^T & M_L \end{bmatrix} \begin{bmatrix} \nu_L \\ T_L^0 \end{bmatrix} \\ & -\frac{1}{2} [\bar{\nu}_L^c, \bar{\nu}_R, (\bar{T}_R^0)^c] \\ & \times \begin{bmatrix} 0 & y_l^T v_1 + \tilde{y}_l^T v_2 & 0 \\ y_l v_1 + \tilde{y}_l v_2 & 0 & f_R^* v_R \\ 0 & f_R^\dagger v_R & M_R^\dagger \end{bmatrix} \begin{bmatrix} \nu_L \\ \nu_R^c \\ (T_R^0)^c \end{bmatrix} \\ & + \text{H.c.} \end{aligned} \quad (13)$$

It is straightforward to see the  $2 \times 2$  mass matrix can induce a canonical seesaw formula [3],

$$\mathcal{L}_m \supset -\frac{1}{2} m_\nu^L \bar{\nu}_L^c \nu_L + \text{H.c.} \quad \text{with} \quad m_\nu^L = -f_L \frac{v_L^2}{M_L} f_L^T, \quad (14)$$

like that in the type-I [3] or III [9] seesaw extension of the SM. On the other hand, the  $3 \times 3$  mass matrix can induce a double/inverse-type seesaw formula [6, 7],

$$\begin{aligned} \mathcal{L}_m \supset & -\frac{1}{2} m_\nu^R \bar{\nu}_L^c \nu_L + \text{H.c.} \quad \text{with} \\ m_\nu^R = & (y_l^T v_1 + \tilde{y}_l^T v_2) \frac{1}{f_R^\dagger v_R} M_R^\dagger \frac{1}{f_R^* v_R} (y_l v_1 + \tilde{y}_l v_2), \end{aligned} \quad (15)$$

which is the double seesaw [6] for  $M_R \gg f v_R$  or the inverse seesaw [7] for  $M_R \ll f v_R$ . In the double seesaw scenario, the right-handed neutrinos should obtain a Majorana mass term much smaller than  $M_R$ . For  $M_L = M_R$  (for the discrete left-right symmetry being a parity transformation) or  $M_L = M_R^\dagger$  (for the discrete left-right symmetry being a charge conjugation), the right-handed neutrinos should also be much lighter than the left-handed triplets. In the inverse seesaw scenario, the right-handed neutrinos and the neutral components of the right-handed triplets should form the pseudo-Dirac fermions with the masses of the order of  $f v_R$ . The left-handed triplets and the charged components of the right-handed triplets thus should be much lighter than the pseudo-Dirac fermions.

For an appropriate parameter choice, the canonical seesaw could dominate over the double/inverse-type seesaw. For example, we can obtain  $m_\nu^L \sim \mathcal{O}(0.1 \text{ eV}) \gg m_\nu^R$  by taking  $v_L \sim v_{1,2} = \mathcal{O}(100 \text{ GeV})$ ,  $M_L = M_R(M_R^\dagger) = \mathcal{O}(\text{TeV}) \ll f_R v_R$ ,  $f_L = f_R(f_R^*) \sim y_l \sim \tilde{y}_l \sim \mathcal{O}(10^{-6})$ .

<sup>1</sup> The electroweak symmetry breaking can be driven by the left-handed Higgs doublet. The Higgs bidoublet then can obtain a VEV at the weak scale through its trilinear couplings with the left- and right-handed Higgs doublets.

It is possible to detect the TeV-scale left-handed triplets at the LHC [10].

Our model can accommodate the leptogenesis [11] mechanism to explain the matter-antimatter asymmetry in the universe. In the double seesaw scenario, since the right-handed neutrinos are much lighter than the neutral components of the left-handed triplets, the leptogenesis should be realized by the decays of the right-handed neutrinos [11]. In the inverse seesaw scenario, the neutral components of the left-handed triplets are much lighter than the pseudo Dirac fermions, which are composed of the right-handed neutrinos and the neutral components of the right-handed triplets, so that their decays should account for the leptogenesis [12]. In the case of  $M_R \sim f v_R$ , the decays of the left- and right-handed neutral fermions may both contribute to the leptogenesis. Note that due to the trilinear couplings among the Higgs bidoublet and the left- and right-handed Higgs doublets, the left-handed Higgs doublet will mix to the Higgs doublets from the Higgs bidoublet after the left-right symmetry breaking. So, if the neutral components of the left-handed fermion triplets are lighter than the left-handed Higgs doublet, they should decay into the SM lepton and Higgs doublets as same as the right-handed neutrinos. Not only the right-handed neutrinos but also the neutral components of the left-handed triplets can mediate the loop corrections in the decays of the right-handed neutrinos [13]. The same story also exists in the decays of the neutral components of the left-handed triplets.

Our model can be embedded into an  $SO(10)$  GUT, where the Higgs doublets, the Higgs bidoublet, the fermion doublets and the fermion triplets belong to the following multiplets, respectively,

$$10_H = \phi(1, 2, 2, 0) \oplus (3, 1, 1, -\frac{1}{3}) \oplus (\bar{3}, 1, 1, \frac{1}{3}), \quad (16a)$$

$$16_H = \chi_L^*(1, 2, 1, -1) \oplus \chi_R(1, 1, 2, 1) \oplus (3, 2, 1, \frac{1}{3}) \oplus (\bar{3}, 1, 2, -\frac{1}{3}), \quad (16b)$$

$$16_F = l_L(1, 2, 1, -1) \oplus l_R^c(1, 1, 2, 1) \oplus q_L(3, 2, 1, \frac{1}{3}) \oplus q_R^c(\bar{3}, 1, 2, -\frac{1}{3}), \quad (16c)$$

$$45_F = S(1, 1, 1, 0) \oplus T_L(1, 3, 1, 0) \oplus T_R^c(1, 1, 3, 0) \oplus \Sigma(3, 1, 1, \frac{4}{3}) \oplus \Sigma^c(\bar{3}, 1, 1, -\frac{4}{3}) \oplus \Omega(3, 2, 2, \frac{2}{3}) \oplus \Omega^c(\bar{3}, 2, 2, -\frac{2}{3}) \oplus (8, 1, 1, 0). \quad (16d)$$

At the left-right level, the terms involving the singlets  $S$  should include

$$\mathcal{L} \supset -h_L \bar{l}_L \tilde{\chi}_L S - h_R \bar{l}_R \tilde{\chi}_R S^c - \frac{1}{2} M_S \bar{S} S^c + \text{H.c.} \quad (17)$$

By performing the seesaw mechanism on

$$\begin{aligned} \mathcal{L}_m &\supset -(y_l v_1 + \tilde{y}_l v_2) \bar{\nu}_L \nu_R - h_L v_L \bar{\nu}_L S - h_R v_R \bar{\nu}_R S^c \\ &\quad - \frac{1}{2} M_S \bar{S} S^c + \text{H.c.} \\ &= -\frac{1}{2} [\bar{\nu}_L, \bar{\nu}_R^c, S^c] \\ &\quad \times \begin{bmatrix} 0 & y_l^T v_1 + \tilde{y}_l^T v_2 & h_L v_L \\ y_l v_1 + \tilde{y}_l v_2 & 0 & h_R^* v_R \\ h_L^T v_L & h_R^\dagger v_R & M_S \end{bmatrix} \begin{bmatrix} \nu_L^c \\ \nu_R \\ S \end{bmatrix} \\ &\quad + \text{H.c.}, \end{aligned} \quad (18)$$

we can read the contribution from the singlets  $S$  to the neutrino masses,

$$\begin{aligned} \mathcal{L}_m &\supset -\frac{1}{2} m_\nu^S \bar{\nu}_L \nu_L^c + \text{H.c.} \quad \text{with} \\ m_\nu^S &= (y_l v_1 + \tilde{y}_l v_2) \frac{1}{h_R^\dagger v_R} M_S \frac{1}{h_R^* v_R} (y_l^T v_1 + \tilde{y}_l^T v_2) \\ &\quad - [(y_l v_1 + \tilde{y}_l v_2) + (y_l^T v_1 + \tilde{y}_l^T v_2)] \frac{v_L}{v_R}. \end{aligned} \quad (19)$$

The first term is a double/inverse-type seesaw [6, 7] while the the second term is a linear seesaw [5]. Therefore, we totally have two double/inverse-type seesaw (from the singlets  $S$  and the right-handed triplets  $T_R$ ), a canonical seesaw (from the left-handed triplets  $T_L$ ) and a linear seesaw (from the singlets  $S$ ). Only the double/inverse-type seesaw will survive while the canonical and linear seesaw will disappear if we choose the VEV  $\langle \chi_L \rangle$  to be zero as did in [14]. In the presence of a nonzero  $\langle \chi_L \rangle$ , the up- and down-type quarks can also obtain the seesaw induced masses by integrating out the  $(\Sigma, \Sigma^c)$  and the  $(\Omega, \Omega^c)$ , respectively. The VEV  $\langle \chi_R \rangle$  should be very large when it accounts for the symmetry breaking of the  $SU(2)_R \otimes U(1)_{B-L}$  down to the  $U(1)_Y$ . Accordingly, the  $45_F$  should be very heavy to guarantee the quark seesaw. Actually, the  $45_F$  could not be very light if they have the Yukawa couplings with the Higgs multiplets (such as a  $210_H$ ) which are responsible for breaking the  $SO(10)$ . In case the  $45_F$  are not very heavy, we may introduce a new  $45_H = (1, 3, 1, 0) \oplus (1, 1, 3, 0) \oplus \dots$  to break the  $SU(2)_R$  down to a  $U(1)_R$  [15]. Then the  $U(1)_R \otimes U(1)_{B-L}$  is broken down to the  $U(1)_Y$  by the  $16_H$ . We hence can take the  $\langle \chi_R \rangle$  at a lower scale such as the TeV.

In this paper we studied the left-right symmetric model with the  $SU(2)$ -triplet fermions. The neutral components of the  $SU(2)_L$ -triplet fermions lead to a canonical seesaw while the neutral components of the  $SU(2)_R$ -triplet fermions associated with the right-handed neutrinos result in a double/inverse-type seesaw. The neutrinos thus can naturally obtain the small masses. Meanwhile, the leptogenesis can work through the decays of the left- and/or right-handed neutral fermions. We further indicated that our model could be realized in an  $SO(10)$  GUT with the fermionic  $45$  multiplets.

**Acknowledgement:** This work is supported by the Alexander von Humboldt Foundation.

- 
- [1] J.C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974); R.N. Mohapatra and J.C. Pati, *ibid.* **11**, 566 (1975); R.N. Mohapatra and J.C. Pati, *ibid.* **11**, 2558 (1975); R.N. Mohapatra and G. Senjanović, *ibid.* **12**, 1502 (1975).
  - [2] Z.G. Berezhiani, Phys. Lett. B **129**, 99 (1983); D. Chang and R.N. Mohapatra, Phys. Rev. Lett. **58**, 1600 (1987); S. Rajpoot, Phys. Lett. B **191**, 122 (1987); A. Davidson and K.C. Wali, Phys. Rev. Lett. **59**, 393 (1987).
  - [3] P. Minkowski, Phys. Lett. B **67**, 421 (1977); T. Yanagida, in *Proc. of the Workshop on Unified Theory and the Baryon Number of the Universe*, ed. O. Sawada and A. Sugamoto (KEK, Tsukuba, 1979), p. 95; M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, ed. F. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979), p. 315; S.L. Glashow, in *Quarks and Leptons*, ed. M. Lévy *et al.* (Plenum, New York, 1980), p. 707; R.N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).
  - [4] M. Magg and C. Wetterich, Phys. Lett. B **94**, 61 (1980); J. Schechter and J.W.F. Valle, Phys. Rev. D **22**, 2227 (1980); T.P. Cheng and L.F. Li, Phys. Rev. D **22**, 2860 (1980); G. Lazarides, Q. Shafi, and C. Wetterich, Nucl. Phys. B **181**, 287 (1981); R.N. Mohapatra and G. Senjanović, Phys. Rev. D **23**, 165 (1981).
  - [5] S.M. Barr, Phys. Rev. Lett. **92**, 101601 (2004).
  - [6] R.N. Mohapatra, Phys. Rev. Lett. **56**, 561 (1986).
  - [7] R.N. Mohapatra and J.W.F. Valle, Phys. Rev. D **34**, 1642 (1986).
  - [8] P.F. Perez, JHEP **0903**, 142 (2009).
  - [9] R. Foot, H. Lew, X.G. He, and G.C. Joshi, Z. Phys. C **44**, 441 (1989).
  - [10] A. Arhrib *et al.*, Phys. Rev. D **82**, 053004 (2010).
  - [11] M. Fukugita and T. Yanagida, Phys. Lett. B **174**, 45 (1986).
  - [12] T. Hambye, Y. Lin, A. Notari, M. Papucci, and A. Strumia, Nucl. Phys. B **695**, 169 (2004).
  - [13] S. Blanchet and P.F. Perez, JCAP **0808**, 037 (2008).
  - [14] J. Chakraborty, S. Goswami, and A. Raychaudhuri, Phys. Lett. B **698**, 265 (2011).
  - [15] M. Malinsky, J.C. Romão, and J.W.F. Valle, Phys. Rev. Lett. **95**, 161801 (2005).