

Binary Icosahedral Flavor Symmetry for Four Generations of Quarks and Leptons

Chian-Shu Chen^{1,2a}, Thomas W. Kephart^{3b}, and Tzu-Chiang Yuan^{2,4c}

¹*Physics Division, National Center for Theoretical Sciences, Hsinchu, Taiwan 300*

²*Institute of Physics, Academia Sinica, Taipei, Taiwan 115*

³*Department of Physics and Astronomy, Vanderbilt University, Nashville, Tennessee 37235, USA*

⁴*Kavli Institute of Theoretical Physics China, CAS, Beijing 100190, China*

(Dated: February 18, 2022)

To include the quark sector, the $A_5 \equiv I$ (icosahedron) four generation lepton model is extended to a binary icosahedral symmetry I' flavor model. We find the masses of fermions, including the heavy sectors, can be accommodated. At leading order the CKM matrix is the identity and the PMNS matrix, resulting from same set of vacua, corresponds to tribimaximal mixings.

arXiv:1110.6233v2 [hep-ph] 8 Aug 2013

^a chianshu@phys.sinica.edu.tw

^b tom.kephart@gmail.com

^c tcyuan@phys.sinica.edu.tw

INTRODUCTION

The current version of the standard model (SM) consists of three generations of quarks and leptons. Recently we proposed [1] a four generation lepton model based on the non-abelian discrete symmetry $A_5 \equiv I$ (icosahedron), in which the best features of the three family $A_4 \equiv T$ (tetrahedral) model survive. Besides the new heavy degrees of freedom in the A_5 model, which satisfy the experimental constraints, we retain tribimaximal neutrino mixings, three light neutrino masses, and three SM charged lepton masses in the three light generation sector.

In this paper, we will explore a generalization of our A_5 model to include four generations of both quarks and leptons. But before launching into that discussion we must first discuss the viability of models with four generations given recent experimental developments. A fourth generation is now being constrained [2] by precision electroweak data [3], by flavor symmetries [4], and by the Higgs-like particle at 125 GeV recently reported at the LHC [5–7]. The new data provide an important step forward in distinguishing various four generation models, and in particular eliminating some from consideration. In particular, four sequential generation models are now highly disfavored [2]; however, it would be premature to dismiss all four generation models. While tension between four generation models and data has developed, a fourth generation is not excluded by the electroweak precision data [8], so the existence of a fourth generation is still a viable phenomenological possibility which can provide an explanation of the observed anomaly of CP asymmetries in the B meson system [9], and the baryon asymmetry of the universe [10], with additional mixings and CP phases. Also, there are a number of way to relieve this tension. For example two Higgs doublet models (See e.g., [11] and references therein) can accommodate a fourth generation of fermions and current data. For a comprehensive review see Ref. [12]. Typically these two Higgs doublet models are low energy effective field theories that require composite Higgses similar to top quark condensate models [13]. For some recent examples see Ref. [14]. Another possibility is to add electroweak doublets that are in color octets [15]. Further discussion can be found in Ref. [16]. While the model we will discuss has an extended Higgs structure, a full exploration of the possible composite nature of the scalar sector is beyond the scope of our present study.

To generalize our A_5 model to include four generations of quarks and leptons, we first recall the three family scenarios in which the binary tetrahedral group $T' \equiv SL_2(3)$ is capable of providing a model of both the quarks and leptons with tribimaximal mixings and a calculable Cabibbo angle [17]. The T' group is the double covering group of A_4 . It has four irreducible representations (irreps) with identical multiplication rules to those of A_4 , one triplet **3** and three singlets **1**, **1'**, and **1''**, plus three additional doublet irreps **2**, **2'**, and **2''**. The additional doublets allow the implementation of the $\mathbf{2} \oplus \mathbf{1}$ structure to the quark sector [18–22], thus the third family of quarks are treated differently and are assigned to a singlet. Hence they can acquire heavy masses [23, 24]. One should note that A_4 is not a subgroup of T' , therefore, the inclusion of quarks into the model is not strictly an extension of A_4 , but instead replaces it [25]. Based on the same philosophy, we study the model of four families of quarks and leptons by using the binary icosahedral group $I' \equiv SL_2(5)$. The relation between I' and A_5 is similar to that for T' and A_4 . The icosahedral group $A_5 \subset SO(3)$ has double-valued representations that are single-valued representations of the double icosahedral group $I' \subset SU(2)$. Hence, besides the irreps of I' that are coincident with those of A_5 , there are four additional spinor-like irreps $\mathbf{2}_s, \mathbf{2}'_s, \mathbf{4}_s$, and $\mathbf{6}_s$ of I' . We shall be able to assign quarks to the spinor-like representations, but to discuss model building using I' , we must first review our lepton model based on A_5 , which will remain essentially unchanged when generalized to I' . Some useful group theory details have been relegated to the Appendix.

THE LEPTONIC A_5 MODEL

The irreps of A_5 are one singlet **1**, two triplets **3** and **3'**, one quartet **4**, and one quintet **5**. The model is required to be invariant under the flavor symmetry of $A_5 \times Z_2 \times Z_3$ and the particle content is given by Table I.

Field	L_i	l_{R5}	l_{R3}^c	$l_{R1}^{(1),(2)}$	N_{R5}	$N^{(1)R}$	S_4	H_4	H'_4	Φ_3
$SU(2)_L$	2	1	1	1	1	1	1	2	2	2
A_5	4	5	3	1	5	1	4	4	4	3
Z_2	1	-1	-1	-1	1	1	1	1	-1	1
Z_3	ω	1	1	1	1	1	1	ω^2	ω^2	ω^2

TABLE I. Particle content of the lepton A_5 model.

Here $L_i = (\nu_i, l_i)^T$ is the left-handed $SU(2)_L$ doublet with generation index i , l_R 's and N_R 's are right-handed charged leptons and neutrinos respectively. H_4 , H'_4 and Φ_3 are $SU(2)_L$ doublet scalar fields, while S_4 is a singlet scalar. The representations of $SU(2)_L$ gauge symmetry should not be confused with the representations of the non-abelian discrete symmetry A_5 . The most general Yukawa interactions invariant under the symmetries can be expressed as

$$\begin{aligned}
L_{Y(\text{lepton})} = & \frac{1}{2} M_1 N_R^{(1)} N_R^{(1)} + \frac{1}{2} M_5 N_{R5} N_{R5} \\
& + Y_{S1}(S_4 N_{R5} N_{R5}) + Y_{S2}(S_4 (l_{R3}^-)^c l_{R5}^-) \\
& + Y_1(L_{L4} N_R^{(1)} H_4) + Y_2(L_{L4} N_{R5} H_4) \\
& + Y_3(L_{L4} N_{R5} \Phi_3) + Y_4(L_{L4} l_{R5} H'_4) \\
& + Y_5(L_{L4} l_{R1} H'_4) + Y_6(L_{L4} l_{R2} H'_4) + \text{H.c.}
\end{aligned} \tag{1}$$

If the scalar S_4 develops the vacuum expectation value (VEV) $\langle S_4 \rangle = (V_S, 0, 0, 0)$, then A_5 will break to A_4 causing the A_5 irreps to decompose as $\mathbf{1} \rightarrow \mathbf{1}$, $\mathbf{3} \rightarrow \mathbf{3}$, $\mathbf{3}' \rightarrow \mathbf{3}$, $\mathbf{4} \rightarrow \mathbf{1} \oplus \mathbf{3}$, and $\mathbf{5} \rightarrow \mathbf{1}' \oplus \mathbf{1}'' \oplus \mathbf{3}$. There is one vector-like $SU(2)_L$ singlet charged lepton field with the mass given by $\langle S_4 \rangle$. The masses of the four chiral generations of charged leptons are generated by scalar field H'_4 with an interesting result that the electron mass is predicted to be zero at tree level and induced through quantum corrections. We argued in [1] that there is enough freedom to fit the observed charged lepton masses. The canonical seesaw mechanism is responsible for the left-handed neutrino masses in the A_5 model¹, while the Dirac mass terms are provided by the scalars fields H_4 and Φ_3 . We are able to obtain one heavy neutrino and three SM light neutrino masses through arranging the VEVs of the two scalar fields without severe fine-tuning. Therefore, the mass structure of the three families under A_4 symmetry is retained. We refer the reader to Ref. [1] for the details of the model. Now we discuss the inclusion of the quark sector by extending A_5 to its double covering I' .

I' SYMMETRY AND THE QUARK SECTOR

The irreps of I' are one singlet $\mathbf{1}$, two triplets $\mathbf{3}$ and $\mathbf{3}'$, one quartet $\mathbf{4}$, and one quintet $\mathbf{5}$, which are also the irreps of A_5 , plus I' has four spinor-like irreps $\mathbf{2}_s, \mathbf{2}'_s, \mathbf{4}_s$, and $\mathbf{6}_s$. The characters and the multiplication rules of I' symmetry can be found in Table II and Table III of the Appendix.

In this work we confine ourselves to minimally extending the A_5 model, i.e., to include the four generations of quarks while minimizing the introduction of other new degrees of freedom. The assignment of the quark sector under $I' \times Z_2 \times Z_3$ is given as follows:

$$\underbrace{\begin{pmatrix} u \\ d \end{pmatrix}_L \begin{pmatrix} c \\ s \end{pmatrix}_L}_{U_{1L}(\mathbf{2}_s, +1, \omega)} \quad \text{and} \quad \underbrace{\begin{pmatrix} t \\ b \end{pmatrix}_L \begin{pmatrix} t' \\ b' \end{pmatrix}_L}_{U_{2L}(\mathbf{2}_s, +1, \omega^2)} \tag{2}$$

for the left-handed doublets, and

$$\underbrace{\underbrace{d_R, s_R}_{S_R} \underbrace{u_R, c_R}_{C_R}}_{D_{sR}(\mathbf{4}_s, +1, +1)} \quad , \quad \underbrace{b_R, b'_R}_{D_{bR}(\mathbf{2}'_s, -1, \omega^2)} \quad , \quad \underbrace{t_R, t'_R}_{D_{tR}(\mathbf{2}'_s, +1, \omega^2)} \tag{3}$$

for the right-handed singlets. Here the fields $(t', b')_L^T$, b'_R and t'_R denote the chiral fields of the fourth generation quarks and the Higgs sector is the same as the A_5 model. Thus, we can write the most general Yukawa interactions between quarks and scalar fields as

$$\begin{aligned}
L_{Y(\text{quark})} = & f_1(U_{1L} \otimes D_{sR}) \otimes \Phi_3 + f_2(U_{2L} \otimes D_{bR}) \otimes H'_4 \\
& + f_3(U_{2L} \otimes D_{tR}) \otimes H_4 + \text{H.c.}
\end{aligned} \tag{4}$$

Recall that in the A_5 model the first step of symmetry breaking $A_5 \rightarrow A_4$ was caused by the vacuum expectation values (VEVs) of the $SU(2)_L$ singlet scalars $\langle S_4 \rangle = (V_S, 0, 0, 0)$. Here the S_4 VEV breaks I' to T' symmetry. The

¹ Also see Ref. [26] for a general discussion of neutrino masses with four generations of fermions.

decomposition of the irreps at this stage of symmetry breaking $I' \rightarrow T'$ is given in Table IV in the Appendix. Therefore, the quark fields decompose as

$$\begin{aligned}
U_{1L}(\mathbf{2}_s, +1, \omega) &\rightarrow U_{1L}(\mathbf{2}, +1, \omega) , \\
U_{2L}(\mathbf{2}_s, +1, \omega^2) &\rightarrow U_{2L}(\mathbf{2}, +1, \omega^2) , \\
D_{sR}(\mathbf{4}_s, +1, +1) &\rightarrow S_R(\mathbf{2}', +1, +1) + C_R(\mathbf{2}'', +1, +1) , \\
D_{bR}(\mathbf{2}'_s, -1, \omega^2) &\rightarrow D_{bR}(\mathbf{2}, -1, \omega^2) , \\
D_{tR}(\mathbf{2}'_s, +1, \omega^2) &\rightarrow D_{tR}(\mathbf{2}, +1, \omega^2) ;
\end{aligned} \tag{5}$$

while for the scalars we have

$$\begin{aligned}
S_4(\mathbf{4}, +1, +1) &\rightarrow S_1(\mathbf{1}, +1, +1) + S_3(\mathbf{3}, +1, +1) , \\
H_4(\mathbf{4}, +1, \omega^2) &\rightarrow H_1(\mathbf{1}, +1, \omega^2) + H_3(\mathbf{3}, +1, \omega^2) , \\
H'_4(\mathbf{4}, -1, \omega^2) &\rightarrow H'_1(\mathbf{1}, -1, \omega^2) + H'_3(\mathbf{3}, -1, \omega^2) , \\
\Phi_3(\mathbf{3}, +1, \omega^2) &\rightarrow \Phi_3(\mathbf{3}, +1, \omega^2) .
\end{aligned} \tag{6}$$

The Yukawa interactions of Eq. (4) now reads

$$\begin{aligned}
&f_1[U_{1L}(\mathbf{2}) \otimes (S_R(\mathbf{2}') \oplus C_R(\mathbf{2}''))] \otimes \Phi_3 \\
&+ f_2[U_{2L}(\mathbf{2}) \otimes D_{bR}(\mathbf{2})] \otimes (H'_1 \oplus H'_3) \\
&+ f_3[U_{2L}(\mathbf{2}) \otimes D_{tR}(\mathbf{2})] \otimes (H_1 \oplus H_3) + \text{H.c.}
\end{aligned} \tag{7}$$

The Clebsch-Gordan coefficients of T' can be found in the review [27] and were already calculated in [19, 22, 28]. We adopt the coefficients shown in Ref. [27] and ignore the phases for simplicity. The Yukawa couplings are divided into terms involving the up-type quarks

$$\begin{aligned}
&f_1 \left[- \begin{pmatrix} u \\ d \end{pmatrix}_L u_R \Phi_{3_1} + \begin{pmatrix} c \\ s \end{pmatrix}_L c_R \Phi_{3_2} + \frac{1}{\sqrt{2}} \left(\begin{pmatrix} u \\ d \end{pmatrix}_L c_R + \begin{pmatrix} c \\ s \end{pmatrix}_L u_R \right) \Phi_{3_3} \right] \\
&+ \frac{f_3}{\sqrt{2}} \left[\begin{pmatrix} t \\ b \end{pmatrix}_L t'_R (H_1 + H_{3_1}) - \begin{pmatrix} t' \\ b' \end{pmatrix}_L t_R (H_1 - H_{3_1}) \right] \\
&+ f_3 \left[- \begin{pmatrix} t \\ b \end{pmatrix}_L t_R H_{3_2} + \begin{pmatrix} t' \\ b' \end{pmatrix}_L t'_R H_{3_3} \right]
\end{aligned} \tag{8}$$

and the down-type quarks

$$\begin{aligned}
&f_1 \left[\begin{pmatrix} c \\ s \end{pmatrix}_L s_R \Phi_{3_1} + \frac{1}{\sqrt{2}} \left(\begin{pmatrix} u \\ d \end{pmatrix}_L s_R + \begin{pmatrix} c \\ s \end{pmatrix}_L d_R \right) \Phi_{3_2} - \begin{pmatrix} u \\ d \end{pmatrix}_L d_R \Phi_{3_3} \right] \\
&+ \frac{f_2}{\sqrt{2}} \left[\begin{pmatrix} t \\ b \end{pmatrix}_L b'_R (H'_1 + H'_{3_1}) - \begin{pmatrix} t' \\ b' \end{pmatrix}_L b_R (H'_1 - H'_{3_1}) \right] \\
&+ f_2 \left[- \begin{pmatrix} t \\ b \end{pmatrix}_L b_R H'_{3_2} + \begin{pmatrix} t' \\ b' \end{pmatrix}_L b'_R H'_{3_3} \right]
\end{aligned} \tag{9}$$

respectively. Here we express the components of scalar fields as $\Phi_3 = (\Phi_{3_1}, \Phi_{3_2}, \Phi_{3_3})$, $H_4 = H_1 + H_3$ where $H_3 = (H_{3_1}, H_{3_2}, H_{3_3})$, and $H'_4 = H'_1 + H'_3$ where $H'_3 = (H'_{3_1}, H'_{3_2}, H'_{3_3})$ according to the breaking $I' \rightarrow T'$ shown in Eq. (6). The masses of up-type fermions are generated by the Φ_3 and H_4 VEVs, while down-type fermion masses are generated by the Φ_3 and H'_4 VEVs.

MASSES AND MIXINGS

We notice that each up- and down-type quark mass matrix is divided into two 2×2 block matrices and can be expressed as

$$M_U = \left(\begin{array}{c|c} M_{uc} & 0 \\ \hline 0 & M_{tt'} \end{array} \right) \quad \text{and} \quad M_D = \left(\begin{array}{c|c} M_{ds} & 0 \\ \hline 0 & M_{bb'} \end{array} \right). \tag{10}$$

This indicates the first two generations mix only with the third and fourth generations through higher order corrections. The 2×2 block mass matrices are given by

$$M_{uc} = f_1 \begin{pmatrix} -\langle \Phi_{31} \rangle & \frac{\langle \Phi_{33} \rangle}{\sqrt{2}} \\ \frac{\langle \Phi_{33} \rangle}{\sqrt{2}} & \langle \Phi_{32} \rangle \end{pmatrix}, \quad M_{ds} = f_1 \begin{pmatrix} -\langle \Phi_{33} \rangle & \frac{\langle \Phi_{32} \rangle}{\sqrt{2}} \\ \frac{\langle \Phi_{32} \rangle}{\sqrt{2}} & \langle \Phi_{31} \rangle \end{pmatrix}, \quad (11)$$

$$M_{tt'} = f_3 \begin{pmatrix} -\langle H_{32} \rangle & \frac{1}{\sqrt{2}}(\langle H_1 \rangle + \langle H_{31} \rangle) \\ \frac{1}{\sqrt{2}}(-\langle H_1 \rangle + \langle H_{31} \rangle) & \langle H_{33} \rangle \end{pmatrix}, \quad (12)$$

and

$$M_{bb'} = f_2 \begin{pmatrix} -\langle H'_{32} \rangle & \frac{1}{\sqrt{2}}(\langle H'_1 \rangle + \langle H'_{31} \rangle) \\ \frac{1}{\sqrt{2}}(-\langle H'_1 \rangle + \langle H'_{31} \rangle) & \langle H'_{33} \rangle \end{pmatrix}. \quad (13)$$

With the $\langle \Phi_3 \rangle = v(1, 1, 1)$ VEV, which is enforced by the requirement of tribimaximal mixings in neutrinos at leading order [1], we find that the masses of u, d, s, c are degenerate $m_u = m_d = m_s = m_c = \sqrt{\frac{3}{2}}f_1v$. Also, the mass matrices of both M_{uc} and M_{ds} take the same form. Therefore, the matrices $M_{uc}M_{uc}^\dagger$ and $M_{ds}M_{ds}^\dagger$ are diagonalized by using the same unitary matrix. Thus we conclude that, at first order, the CKM matrix is forced to be the identity², which is an acceptable first approximation to V_{CKM} . It is interesting that the light quark masses (u, d, s, c) and three SM light neutrino masses³ have the same origin; both come from the VEVs of Φ_3 field, and the unit CKM matrix and the tribimaximal PMNS mixings arise from the single subgroup Z_3 , which is the remnant symmetry left in the vacuum $\langle \Phi_3 \rangle = v(1, 1, 1)$. To correct the CKM mixings by the high-order effects, the relevant dimension-five and -six effective operators are $U_{2L}D_{sR}\Phi_3^2$, $U_{2L}D_{sR}\Phi_3H_4$, $U_{2L}D_{sR}H_4'^2$, $U_{2L}D_{sR}H_4^2$ and $U_{1L}D_{bR}H_4^2H_4'$, $U_{1L}D_{tR}H_4^3$, $U_{1L}D_{tR}H_4'^2H_4$ respectively. Also, as mentioned in Ref. [1], perturbations of the Φ_3 VEVs are needed to accommodate realistic neutrino masses. These high-order corrections will link together the derivations of the Cabibbo angle θ_C and θ_{13} in the quark and lepton mixing matrices respectively. A nonzero value of θ_{13} has recently been indicated by several experiments [29–32] and by global analyses [33]. If we consider perturbations of the VEVs by taking $\langle \Phi_3 \rangle = (v + \Delta_1, v + \Delta_2, v)$, the light quark masses are calculated to be

$$m_{u,c}^2 = \frac{f_1^2}{2} \left[3v^2 + 2v(\Delta_2 - \Delta_1) + (\Delta_1^2 + \Delta_2^2) \right. \\ \left. \mp \sqrt{(\Delta_1^2 + \Delta_2^2)^2(6v^2 - 4v(\Delta_1 + \Delta_2))} \right]$$

and

$$m_{d,s}^2 = \frac{f_1^2}{2} \left[3v^2 + 2v(\Delta_1 + \Delta_2) + (\Delta_1^2 + \Delta_2^2) \right. \\ \left. \mp \Delta_1 \sqrt{(2v + \Delta_1)^2 + 2(v + \Delta_2)^2} \right].$$

This indicates how the degeneracy of the light quark masses can be lifted.

Recall that in the lepton A_5 model, H_4 is responsible for the Dirac masses of neutrinos, and we require the condition $\langle H_1 \rangle \equiv V_1 \gg \langle H_{31,2,3} \rangle \equiv V_{31,2,3}$ in order to decouple the 4th generation neutrino from the three light SM neutrinos. Therefore, from $M_{t,t'}$, we obtain the masses of t and t' as $m_{t,t'}^2 \approx \left[V_1^2 + (V_{31}^2 + V_{32}^2 + V_{33}^2) \mp V_1 \sqrt{4V_{31}^2 + 2(V_{32} + V_{33})^2} \right] / 2$. For $M_{bb'}$, we also follow the lepton A_5 model by taking⁴ $\langle H'_4 \rangle = (V'_1, V', V', V')$, since this gives masses to charged leptons too. $m_{b,b'}^2$ are then given by $[V_1'^2 + 3V'^2 \mp 2\sqrt{3}V'_1V'] / 2$. In general, we have enough parameters to accommodate the heavy quark mass spectrum.

² The third and fourth generations can mix largely in principle.

³ Neutrino masses are generated through the seesaw mechanism, and therefore they are further suppressed by the lepton number breaking scale.

⁴ The four component VEVs, in general, can be different.




FIG. 1. Extended Dynkin diagram and irreducible representations of I' .

CONCLUSION

As mentioned in the introduction, the recent observations of a boson with a mass near 125 GeV [5, 6] have placed severe constraints on the standard model augmented by a sequential fourth generation of fermions. The CMS experiment has now excluded such a fourth generation of fermions with masses of up to 600 GeV [7]. Note that our I' model has three doublets and one singlet Higgs, all of which are in non-singlet irreps of I' . Hence it is not necessarily disfavored by the current experimental search. Indeed, such severe limits can be relaxed into the range of 400 \sim 600 GeV in a two Higgs doublets model with four generation of fermions, as discussed in Ref.[12]. It would be interesting to investigate whether our I' model (or one of its extensions) can be recast into a form designed in Ref.[12].

In summary, we construct a model of four fermion generations based on the binary icosahedral symmetry group I' . Many properties of the SM with three families are accommodated such as the mass spectrum, tribimaximal mixings in the neutrino sector, and an identity CKM matrix at leading order⁵. In addition, quarks and leptons relations are intimately connected as their masses are provided from the same set of scalars. We believe this makes the model both interesting and challenging. For example, one has to strike a balance between the result of tribimaximal mixings in the neutrino sector and derivation a realistic Cabibbo angle from perturbations in the quark sector. It is still not clear to us whether the higher order corrections will lead to a realistic Cabibbo angle or if we need extra degrees of freedom to realize it. We will leave it and other phenomenological aspects of this model to future work.

This work was supported in part by the US DOE grant DE-FG05-85ER40226, the National Science Council of Taiwan under Grant Numbers 98-2112-M-001-014-MY3, 101-2112-M-001-005-MY3, and the National Center for Theoretical Sciences of Taiwan (NCTS). TWK is grateful for the hospitality of the Physics Division of NCTS where this work was initiated. TCY thanks the hospitalities of KITPC (Beijing, China) and IMSc (Chennai, India) where progress of this work was made.

APPENDIX

Discrete Symmetry Groups I' and A_5

A_5 is the only simple finite discrete subgroup of $SO(3)$. Its 60 elements can be generated by products of the two generators a and b , which satisfy

$$a^2 = b^3 = (ab)^5 = e, \quad (14)$$

where e is the identity element.

I' is the double covering of A_5 , therefore it has 120 elements. The representation in terms of generators is similar to that of A_5 , namely

$$a^2 = b^3 = (ab)^5. \quad (15)$$

We note that a^2 is no longer the identity, but the negative of the identity, i.e., $a^4 = b^6 = e$. Any finite subgroup of $SU(2)$ must have (at least) one spinor doublet $\mathbf{2}_s$. By using the multiplication rules, the irreducible representations of the group [36, 37] can be visualized as an extended Dynkin diagrams depicted in Fig. 1.

⁵ Both A_5 and I' have also recently been used in the context of three generation models of fermion masses and mixings [35].

Recalling that the exact sequence relation between $SU(2)$ and $SO(3)$ is

$$1 \rightarrow Z_2 \rightarrow SU(2) \rightarrow SO(3) \rightarrow 1, \quad (16)$$

we can restrict to the discrete cases

$$1 \rightarrow Z_2 \rightarrow T' \rightarrow T \rightarrow 1 \quad (17)$$

and

$$1 \rightarrow Z_2 \rightarrow I' \rightarrow I \rightarrow 1 \quad (18)$$

to demonstrate the double coverings. As our interest is in I' we first reproduce its character table in Table II, from which we can easily calculate the multiplication table for I' irreps presented in Table III. The symmetry breaking to T' is also easily obtained, as given in Table IV.

	$C_1(1)$	$C_2(1)$	$C_3(12)$	$C_4(12)$	$C_5(12)$	$C_6(12)$	$C_7(30)$	$C_8(20)$	$C_9(20)$
1	1	1	1	1	1	1	1	1	1
3	3	3	$1 - \phi$	$1 - \phi$	ϕ	ϕ	-1	0	0
3'	3	3	ϕ	ϕ	$1 - \phi$	$1 - \phi$	-1	0	0
4	4	4	-1	-1	-1	-1	0	1	1
5	5	5	0	0	0	0	1	-1	-1
2_s	2	-2	$\phi - 1$	$1 - \phi$	$-\phi$	ϕ	0	-1	1
2'_s	2	-2	$-\phi$	ϕ	$\phi - 1$	$1 - \phi$	0	-1	1
4_s	4	-4	-1	1	-1	1	0	1	-1
6_s	6	-6	1	-1	1	-1	0	0	0

TABLE II. Character table of I' where $\phi = \frac{1+\sqrt{5}}{2}$ is the golden ratio.

\otimes	1	3	3'	4	5	2_s	2'_s	4_s	6_s
1	1	3	3'	4	5	2_s	2'_s	4_s	6_s
3	3	$1 \oplus 3 \oplus 5$	$4 \oplus 5$	$3' \oplus 4 \oplus 5$	$3 \oplus 3' \oplus 4 \oplus 5$	$2_s \oplus 4_s$	6_s	$2_s \oplus 4_s \oplus 6_s$	$2'_s \oplus 4_s \oplus 6_s \oplus 6_s$
3'	3'	$4 \oplus 5$	$1 \oplus 3' \oplus 5$	$3 \oplus 4 \oplus 5$	$3 \oplus 3' \oplus 4 \oplus 5$	6_s	$2'_s \oplus 4_s$	$2'_s \oplus 4_s \oplus 6_s$	$2_s \oplus 2'_s \oplus 4_s \oplus 6_s$
4	4	$3' \oplus 4 \oplus 5$	$3 \oplus 4 \oplus 5$	$1 \oplus 3 \oplus 3' \oplus 4 \oplus 5$	$3 \oplus 3' \oplus 4 \oplus 5 \oplus 5$	$2'_s \oplus 6_s$	$2_s \oplus 6_s$	$4_s \oplus 6_s \oplus 6_s$	$2_s \oplus 2'_s \oplus 4_s \oplus 4_s \oplus 6_s \oplus 6_s$
5	5	$3 \oplus 3' \oplus 4 \oplus 5$	$3 \oplus 3' \oplus 4 \oplus 5$	$3 \oplus 3' \oplus 4 \oplus 5 \oplus 5$	$1 \oplus 3 \oplus 3' \oplus 4 \oplus 4 \oplus 5 \oplus 5$	$4_s \oplus 6_s$	$4_s \oplus 6_s$	$2_s \oplus 2'_s \oplus 4_s \oplus 6_s \oplus 6_s$	$2_s \oplus 2'_s \oplus 4_s \oplus 4_s \oplus 6_s \oplus 6_s \oplus 6_s$
2_s	2_s	$2_s \oplus 4_s$	6_s	$2'_s \oplus 6_s$	$4_s \oplus 6_s$	$1 \oplus 3$	4	$3 \oplus 5$	$3' \oplus 4 \oplus 5$
2'_s	2'_s	6_s	$2'_s \oplus 4_s$	$2_s \oplus 6_s$	$4_s \oplus 6_s$	4	$1 \oplus 3'$	$3' \oplus 5$	$3 \oplus 4 \oplus 5$
4_s	4_s	$2_s \oplus 4_s \oplus 6_s$	$2'_s \oplus 4_s \oplus 6_s$	$4_s \oplus 6_s \oplus 6_s$	$2_s \oplus 2'_s \oplus 4_s \oplus 6_s \oplus 6_s$	$3 \oplus 5$	$3' \oplus 5$	$3' \oplus 4 \oplus 5$	$3 \oplus 3' \oplus 4 \oplus 4 \oplus 5 \oplus 5$
6_s	6_s	$2'_s \oplus 4_s \oplus 6_s \oplus 6_s$	$2_s \oplus 2'_s \oplus 4_s \oplus 6_s$	$2_s \oplus 2'_s \oplus 4_s \oplus 4_s \oplus 6_s \oplus 6_s$	$2_s \oplus 2'_s \oplus 4_s \oplus 4_s \oplus 6_s \oplus 6_s \oplus 6_s$	$3' \oplus 4 \oplus 5$	$3 \oplus 4 \oplus 5$	$3 \oplus 3' \oplus 4 \oplus 4 \oplus 5 \oplus 5$	$1 \oplus 3 \oplus 3' \oplus 3' \oplus 3' \oplus 4 \oplus 4 \oplus 5 \oplus 5 \oplus 5$

TABLE III. Multiplication rules for the binary icosahedral group I' .

I'	\rightarrow	T'	I'	\rightarrow	T'
1		1	2_s		2
3		3	2'_s		2
3'		3	4_s		2' + 2''
4		1 + 3	6_s		2 + 2' + 2''
5		1' + 1'' + 3			

TABLE IV. $I' \rightarrow T'$ symmetry breaking.

Higgs Potential

The most general form of the Higgs potential containing the scalar fields S_4 , H_4 , H'_4 and Φ_3 , invariant under the discrete $A_5 \times Z_2 \times Z_3$ symmetries is given by

$$\begin{aligned}
V = & V(S_4) + V(H_4) + V(H'_4) + V(\Phi_3) + V(S_4, H_4) \\
& + V(S_4, H'_4) + V(S_4, \Phi_3) + V(H_4, H'_4) \\
& + V(H_4, \Phi_3) + V(H'_4, \Phi_3) + V(H_4, H'_4, \Phi_3)
\end{aligned} \tag{19}$$

where the individual terms are written as

$$\begin{aligned}
V(S_4) &= \mu_{S_4}^2 S_4^2 + \mu_s (S_4^2)_4 S_4 + \lambda_\alpha^s (S_4^2)_\alpha (S_4^2)_\alpha, \\
V(H_4) &= \mu_H^2 (H_4^\dagger H_4)_1 + \lambda_\alpha^H (H_4^\dagger H_4)_\alpha (H_4^\dagger H_4)_\alpha, \\
V(H'_4) &= \mu_{H'}^2 (H_4'^\dagger H'_4)_1 + \lambda_\alpha^{H'} (H_4'^\dagger H'_4)_\alpha (H_4'^\dagger H'_4)_\alpha, \\
V(\Phi_3) &= \mu_\Phi^2 (\Phi_3^\dagger \Phi_3)_1 + \lambda_\beta^\Phi (\Phi_3^\dagger \Phi_3)_\beta (\Phi_3^\dagger \Phi_3)_\beta, \\
V(S_4, H_4) &= \delta^{HS} (H_4^\dagger H_4)_4 S_4 + \lambda_\alpha^{HS} (H_4^\dagger H_4)_\alpha (S_4^2)_\alpha, \\
V(S_4, H'_4) &= \delta^{H'S} (H_4'^\dagger H'_4)_4 S_4 + \lambda_\alpha^{H'S} (H_4'^\dagger H'_4)_\alpha (S_4^2)_\alpha, \\
V(S_4, \Phi_3) &= \delta^{\Phi S} (\Phi_3^\dagger \Phi_3)_4 S_4 + \lambda_\beta^{\Phi S} (\Phi_3^\dagger \Phi_3)_\beta (S_4^2)_\beta, \\
V(H_4, H'_4) &= \lambda_\alpha^{HH'} (H_4^\dagger H_4)_\alpha (H_4'^\dagger H'_4)_\alpha \\
& + \lambda_\alpha'^{HH'} (H_4^\dagger H'_4)_\alpha (H_4'^\dagger H_4)_\alpha \\
& + \left[\lambda_\alpha''^{HH'} (H_4^\dagger H'_4)_\alpha (H_4'^\dagger H_4)_\alpha + \text{H.c.} \right],
\end{aligned}$$

$$\begin{aligned}
V(H_4, \Phi_3) &= \lambda_\beta^{H\Phi} (H_4^\dagger H_4)_\beta (\Phi_3^\dagger \Phi_3)_\beta \\
& + \lambda_\gamma^{H\Phi} (H_4^\dagger \Phi_3)_\gamma (\Phi_3^\dagger H_4)_\gamma \\
& + \left[\lambda_\gamma''^{H\Phi} (H_4^\dagger \Phi_3)_\gamma (H_4^\dagger \Phi_3)_\gamma + \text{H.c.} \right], \\
V(H'_4, \Phi_3) &= \lambda_\beta^{H'\Phi} (H_4'^\dagger H'_4)_\beta (\Phi_3^\dagger \Phi_3)_\beta \\
& + \lambda_\gamma^{H'\Phi} (H_4'^\dagger \Phi_3)_\gamma (\Phi_3^\dagger H'_4)_\gamma \\
& + \left[\lambda_\gamma''^{H'\Phi} (H_4'^\dagger \Phi_3)_\gamma (H_4'^\dagger \Phi_3)_\gamma + \text{H.c.} \right], \\
V(H_4, H'_4, \Phi_3) &= \lambda_\gamma^{HH'\Phi} (H_4^\dagger \Phi_3)_\gamma (H_4'^\dagger H'_4)_\gamma \\
& + \lambda_\gamma'^{HH'\Phi} (H_4^\dagger \Phi_3)_\gamma (H_4'^\dagger H_4)_\gamma \\
& + \lambda_\gamma''^{HH'\Phi} (H_4^\dagger \Phi_3)_\gamma (H_4^\dagger H_4)_\gamma + \text{H.c.}
\end{aligned}$$

Here we have introduced the I' group representation indices $\alpha = \mathbf{1}, \mathbf{3}, \mathbf{3}', \mathbf{4}, \mathbf{5}$; $\beta = \mathbf{1}, \mathbf{3}, \mathbf{5}$; and $\gamma = \mathbf{3}', \mathbf{4}, \mathbf{5}$ respectively. The first stage of SSB takes us from the initial I' discrete symmetry to T' , and this is accomplished with a VEV for S_4 . Consider the pure S_4 sector of the Higgs potential

$$V_{S_4} = \mu_{S_4}^2 S_4 S_4 + \lambda (S_4 S_4)^2 + \lambda_1 (S_4 M_1 S_4)^2 + \lambda_2 (S_4 M_3 S_4)^2.$$

where M_1 and M_3 are I' group matrices. Since $(4 \times 4)_S$ in I' contains three terms there are three singlets in $[(4 \times 4)_S]^2$ and hence in V_{S_4} . There is also a potential cubic term that can be suppressed, either by imposing an additional Z_2

symmetry, or by making S_4 a complex field. For λ_1 and λ_2 sufficiently small and of the proper signs, the SSB is dominated by the λ term and we have $\langle S_4 \rangle = -\sqrt{\frac{\mu_{S_4}^2}{\lambda}}(1, 0, 0, 0)$ such that $I' \rightarrow T'$ with the SM gauge group unaffected and with the scalars decomposing as in Eq. (6).

The sector of the Higgs potential of the I' model that depends only on H_1 , $H_{1'}$, H_3 , and $H_{3'}$ is given in the appendix of [38] which only involves T' invariant terms and is identical to our form of the potential up to constraints due to residual I' relations on the coupling constants. The work of [38] demonstrates how T' is broken completely and how the light quark and lepton masses and mixings arise.

Our only other additional scalars are the I' triplet, SM doublet fields Φ_3 . Since $3 \rightarrow 3$ under $I' \rightarrow T'$, a Φ_3 breaks T' . The pure Φ_3 scalar sector can be rewritten as

$$V_{\Phi_3} = \mu_{\Phi_3}^2 \Phi_3^\dagger \Phi_3 + \lambda (\Phi_3^{a\dagger} \Phi_3^a)^2 + \lambda' (\Phi_3^{a\dagger} \Phi_3^b)^2$$

where we have included the triplet indices $a, b, \dots = 1, 2, 3$ and find two quartic terms due to the fact that $(3 \times 3)_S$ in I' contains two terms and hence $[(3 \times 3)_S]^2$ contains two singlets. Given the T' level VEVs for H_3 , and $H_{3'}$, there is no T' symmetry remaining to rotate the Φ_3 VEV. Hence the choice $\langle \Phi_3 \rangle = -v(1, 1, 1)$ is stable for $\lambda' \approx 0$ and of the form needed in the model.

A similar analysis can be applied to the investigation of the SSB in our A_5 model. Finally we note that typically, only an $O(10^{-1})$ fine tuning of scalar quartic coupling constants is needed to maintain the stability of such patterns of SSB.

-
- [1] C. -S. Chen, T. W. Kephart and T. -C. Yuan, JHEP **1104**, 015 (2011) [arXiv:1011.3199 [hep-ph]].
 - [2] O. Eberhardt, A. Lenz and J. Rohrwild, Phys. Rev. D **82**, 095006 (2010) [arXiv:1005.3505 [hep-ph]]; O. Eberhardt, G. Herbert, H. Lacker, A. Lenz, A. Menzel, U. Nierste and M. Wiebusch, Phys. Rev. D **86**, 013011 (2012) [arXiv:1204.3872 [hep-ph]]; O. Eberhardt, A. Lenz, A. Menzel, U. Nierste and M. Wiebusch, Phys. Rev. D **86**, 074014 (2012) [arXiv:1207.0438 [hep-ph]]; O. Eberhardt, G. Herbert, H. Lacker, A. Lenz, A. Menzel, U. Nierste and M. Wiebusch, Phys. Rev. Lett. **109**, 241802 (2012) [arXiv:1209.1101 [hep-ph]].
 - [3] C. Lazzeroni *et al.* [NA62 Collaboration], Phys. Lett. B **698**, 105 (2011) [arXiv:1101.4805 [hep-ex]].
 - [4] H. Lacker and A. Menzel, JHEP **1007**, 006 (2010) [arXiv:1003.4532 [hep-ph]].
 - [5] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012) [arXiv:1207.7214 [hep-ex]].
 - [6] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012) [arXiv:1207.7235 [hep-ex]].
 - [7] S. Chatrchyan *et al.* [CMS Collaboration], arXiv:1302.1764 [hep-ex].
 - [8] E. Gates and J. Terning, Phys. Rev. Lett. **67**, 1840 (1991); B. A. Kniehl and H. G. Kohrs, Phys. Rev. D **48**, 225 (1993); B. Holdom, Phys. Rev. D **54**, 721 (1996) [arXiv:hep-ph/9602248]; H. J. He, N. Polonsky and S. Su, Phys. Rev. D **64**, 053004 (2001) [arXiv:hep-ph/0102144]; B. Holdom, W. S. Hou, T. Hurth, M. L. Mangano, S. Sultansoy and G. Unel, PMC Phys. A **3**, 4 (2009) [arXiv:0904.4698 [hep-ph]].
 - [9] W. S. Hou, M. Nagashima and A. Soddu, Phys. Rev. Lett. **95**, 141601 (2005) [arXiv:hep-ph/0503072]; W. S. Hou, H. n. Li, S. Mishima and M. Nagashima, Phys. Rev. Lett. **98**, 131801 (2007) [arXiv:hep-ph/0611107]; A. Soni, A. K. Alok, A. Giri, R. Mohanta and S. Nandi, Phys. Lett. B **683**, 302 (2010) [arXiv:0807.1971 [hep-ph]].
 - [10] M. E. Shaposhnikov, Nucl. Phys. B **287** (1987) 757; W. S. Hou, Chin. J. Phys. **47**, 134 (2009) [arXiv:0803.1234 [hep-ph]].
 - [11] S. Bar-Shalom, S. Nandi and A. Soni, Phys. Rev. D **84**, 053009 (2011) [arXiv:1105.6095 [hep-ph]].
 - [12] S. Bar-Shalom, M. Geller, S. Nandi and A. Soni, arXiv:1208.3195 [hep-ph].
 - [13] W. A. Bardeen, C. T. Hill and M. Lindner, Phys. Rev. D **41**, 1647 (1990).
 - [14] P. Q. Hung and C. Xiong, Nucl. Phys. B **847**, 160 (2011) [arXiv:0911.3890 [hep-ph]]; P. Q. Hung and C. Xiong, Phys. Lett. B **694**, 430 (2011) [arXiv:0911.3892 [hep-ph]]; P. Q. Hung and C. Xiong, Nucl. Phys. B **848**, 288 (2011) [arXiv:1012.4479 [hep-ph]]; C. M. Ho, P. Q. Hung and T. W. Kephart, JHEP **1206**, 045 (2012) [arXiv:1102.3997 [hep-ph]].
 - [15] A. V. Manohar and M. B. Wise, Phys. Rev. D **74**, 035009 (2006) [hep-ph/0606172].
 - [16] X. -G. He and G. Valencia, Phys. Lett. B **707**, 381 (2012) [arXiv:1108.0222 [hep-ph]].
 - [17] P. H. Frampton, T. W. Kephart and S. Matsuzaki, Phys. Rev. D **78**, 073004 (2008) [arXiv:0807.4713 [hep-ph]].
 - [18] P. H. Frampton and T. W. Kephart, Int. J. Mod. Phys. A **10**, 4689 (1995) [arXiv:hep-ph/9409330].
 - [19] A. Aranda, C. D. Carone and R. F. Lebed, Phys. Lett. B **474**, 170 (2000) [arXiv:hep-ph/9910392] and Phys. Rev. D **62**, 016009 (2000) [arXiv:hep-ph/0002044].
 - [20] F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, Nucl. Phys. B **775**, 120 (2007) [Erratum-ibid. **836**, 127 (2010)] [arXiv:hep-ph/0702194].
 - [21] M. C. Chen and K. T. Mahanthappa, Phys. Lett. B **652**, 34 (2007) [arXiv:0705.0714 [hep-ph]].
 - [22] P. H. Frampton and T. W. Kephart, JHEP **0709**, 110 (2007) [arXiv:0706.1186 [hep-ph]].
 - [23] P. H. Frampton and T. W. Kephart, Phys. Rev. D **51**, 1 (1995) [arXiv:hep-ph/9409324].
 - [24] P. H. Frampton, *The third family is different*, in *Proceeding of the Fourth International Symposium on Particles, Strings and Cosmology*, K.C. Wali ed., World Scientific (1995), [arXiv:hep-ph/9409331] and *Treating top differently from charm and up*, in *Particle Theory and Phenomenology*, K.E. Lassila et al. eds., World Scientific (1996), [arXiv:hep-ph/9507351].

- [25] P. H. Frampton, T. W. Kephart and R. M. Rohm, Phys. Lett. B **679**, 478 (2009) [arXiv:0904.0420 [hep-ph]].
- [26] M. A. Schmidt and A. Y. Smirnov, Nucl. Phys. B **857**, 1 (2012) [arXiv:1110.0874 [hep-ph]].
- [27] H. Ishimori, T. Kobayashi, H. Ohki, H. Okada, Y. Shimizu and M. Tanimoto, Prog. Theor. Phys. Suppl. **183**, 1 (2010) [arXiv:1003.3552 [hep-th]].
- [28] W. M. Fairbairn, T. Fulton and W. H. Klink, J. Math. Phys. (N.Y.) **5**, 1038 (1964).
- [29] K. Abe *et al.* [T2K Collaboration], Phys. Rev. Lett. **107**, 041801 (2011). [arXiv:1106.2822 [hep-ex]].
- [30] Y. Abe *et al.* [DOUBLE-CHOOZ Collaboration], Phys. Rev. Lett. **108**, 131801 (2012) [arXiv:1112.6353 [hep-ex]].
- [31] F. P. An *et al.* [DAYA-BAY Collaboration], Phys. Rev. Lett. **108**, 171803 (2012) [arXiv:1203.1669 [hep-ex]].
- [32] J. K. Ahn *et al.* [RENO Collaboration], Phys. Rev. Lett. **108**, 191802 (2012) [arXiv:1204.0626 [hep-ex]].
- [33] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. M. Rotunno, Phys. Rev. D **84**, 053007 (2011) [arXiv:1106.6028 [hep-ph]]; T. Schwetz, M. Tortola and J. W. F. Valle, New J. Phys. **13**, 109401 (2011) [arXiv:1108.1376 [hep-ph]].
- [34] K. Nakamura *et al.* (Particle Data Group), J. Phys. G **37**, 075021 (2010).
- [35] L. L. Everett and A. J. Stuart, Phys. Rev. D **79**, 085005 (2009) [arXiv:0812.1057 [hep-ph]]; F. Feruglio and A. Paris, JHEP **1103**, 101 (2011) [arXiv:1101.0393 [hep-ph]]; G. -J. Ding, L. L. Everett and A. J. Stuart, Nucl. Phys. B **857**, 219 (2012) [arXiv:1110.1688 [hep-ph]]; L. L. Everett and A. J. Stuart, Phys. Lett. **B698**, 131-139 (2011) [arXiv:1011.4928 [hep-ph]].
- [36] C. Luhn, S. Nasri and P. Ramond, J. Math. Phys. **48**, 123519 (2007) [arXiv:0709.1447 [hep-th]].
- [37] P. Ramond, *Group Theory: A Physicist's Survey* (Cambridge University Press, 2010).
- [38] D. A. Eby and P. H. Frampton, Phys. Lett. B **713**, 249 (2012) [arXiv:1111.4938 [hep-ph]].