

Factorization of Heavy-to-Light Baryonic Transitions in SCET

Wei Wang *

Deutsches Elektronen-Synchrotron DESY, D-22607 Hamburg, Germany

(Dated: April 3, 2024)

Abstract

In the framework of the soft-collinear effective theory, we demonstrate that the leading-power heavy-to-light baryonic form factors at large recoil obey the heavy quark and large energy symmetries. Symmetry breaking effects are suppressed by Λ/m_b or Λ/E , where Λ is the hadronic scale, m_b is the b quark mass and $E \sim m_b$ is the energy of light baryon in the final state. At leading order, the leading power baryonic form factor $\xi_{\Lambda,p}(E)$, in which two hard-collinear gluons are exchanged in the baryon constituents, can factorize into the soft and collinear matrix elements convoluted with a hard-kernel of order α_s^2 . Including the energy release dependence, we derive the scaling law $\xi_{\Lambda,p}(E) \sim \Lambda^2/E^2$. We also find that this form factor $\xi_{\Lambda}(E)$ is numerically smaller than the form factor governed by soft processes, although the latter is formally power-suppressed.

Keywords: Heavy quark physics, QCD, b-physics

* Email:wei.wang@desy.de

I. INTRODUCTION

Precision test of the unitarity of the CKM matrix, allowing us to explore the SM description of the CP violation and reveal any physics beyond the SM, greatly depends on our knowledge of the nonperturbative matrix elements. Fortunately the calculation of the amplitudes of bottom meson decays is under control as the amplitudes can be expanded in terms of small ratios justified by both the large b -quark mass, and a large energy release in the decay. With this expansion, a number of theoretical predictions on different observables in various channels are found in global agreement with experimental measurements (see Ref. [1] for a review).

Decay processes of heavy baryons consisting of a bottom quark provide complementary information with the B meson and thereby are receiving growing attentions on both experimental and theoretical sides. Semileptonic decays, such as $\Lambda_b \rightarrow pl\bar{\nu}$, are simplest exclusive baryonic decays and governed by heavy-to-light form factors. In this retrospect, apart from the theoretical analysis based on the heavy quark effective theory [2–4], the simplification of baryonic form factors in the large energy limit is exploited [5, 6] (see Ref. [7, 8] for an earlier discussion), applying the method developed in the mesonic case [9, 10]. In the $\Lambda_b \rightarrow \Lambda$ transition, only one form factor is nonzero after the reduction and this universal function (soft form factor) is also calculated within the light-cone QCD sum rules in conjunction with the effective field theory [5].

Soft-collinear effective theory (SCET) [11–15] is a powerful tool to describe processes with particles having energy much larger than their mass. The heavy-to-light decay of heavy baryons, for instance $\Lambda_b \rightarrow pl\bar{\nu}$, is of this type. SCET makes use of the expansion in small ratios, in this case, $\lambda = \sqrt{\Lambda/m_b}$ with Λ as the hadronic scale and m_b as the b quark mass. One of the most important features of SCET is that the interaction between the soft and collinear sectors is taken into account, overcoming the shortcomings in the large energy effective theory [9, 10]. Therefore in SCET not only the reduction of the leading-power form factors is formulated on the QCD basis, but also the symmetry-breaking corrections can be systematically explored [16, 17].

In this work, we will analyze the baryonic form factors in SCET and follow the techniques developed in the $B \rightarrow \pi$ form factor which takes the following factorization form at the leading power [15, 18, 19]

$$F_i^{B \rightarrow \pi}(E) = C_i \xi_\pi(E) + \int d\tau C'_i(E, \tau) \Xi_\pi(\tau, E). \quad (1)$$

Here E is the energy of the final hadron and C_i and C'_i are the short-distance coefficients obtained by matching from QCD onto the effective field theory. The one-loop expressions for these coefficients can be found in Refs. [11, 13, 18–21]. In what follows we will adopt the ansatz that the final light particle is composed of collinear objects and thus hard-collinear gluon exchange is required to turn the soft spectators into energetic ones. In such picture, to the end we will show that the matrix elements parametrizing form factors, in the example of $\Lambda_b \rightarrow \Lambda$, are formally simple

$$\langle \Lambda(p') | \bar{s} \Gamma b | \Lambda_b(p) \rangle = C_i \xi_\Lambda(E) \bar{u}_\Lambda(p') \Gamma u_{\Lambda_b}(p) + \mathcal{O}(\lambda^2 \xi_\Lambda), \quad (2)$$

in which the spin indices are suppressed. For contributions dominated by soft processes which are not suppressed by α_s , please see Refs. [5, 6].

The remainder of this work is organized as follows. In Sec. II, we will present the form of the leading power and next-to-leading power heavy-to-light currents in SCET after integrating out the hard modes, and following Ref. [19] discuss their representations in the effective theory containing soft and collinear modes. In Sec. III, the transition form factors are directly calculated in QCD, and we show the correspondence with the SCET effective operators. Several implications from our analysis are given in Sec. IV, and a summary of our findings is presented in Sec. V.

II. SCET ANALYSIS

We use the position-space representation of SCET and closely follow the notations in Refs. [14, 19]. We work in the b-baryon rest frame and use the light-cone coordinate, in which a momentum p is decomposed as

$$p^\mu = (n_+ p) \frac{n_-^\mu}{2} + p_\perp^\mu + (n_- p) \frac{n_+^\mu}{2} \quad (3)$$

where $n_\pm$ are two light-like vectors: $n_+^2 = n_-^2 = 0$ and $n_+ \cdot n_- = 2$. The reference directions $n_\pm$ are chosen such that the energetic massless external lines in the recoiling system have $n_+ p$ of order m_b , while the magnitude of $n_- p$ is small. This type of momenta is collinear: $p_c = (n_+ p, p_\perp, n_- p) \sim (1, \lambda^2, \lambda^4)$. The slowly-moving degrees of freedom in the heavy baryon have soft momenta $q_s \sim (\lambda^2, \lambda^2, \lambda^2)$. For the heavy b quark, the statement of “soft” refers to the residual momentum after removing the large component which becomes a label of heavy quark. The hard-collinear mode, with $\mathcal{O}(m_b \Lambda)$ virtuality, arises from the interaction between soft and collinear sector: $p_{hc} \sim (1, \lambda, \lambda^2)$.

Power scalings of quark and gluon fields are determined by the configuration of their momenta. For the quark fields, we have

$$\begin{aligned} \xi_c &= \frac{\not{n}_- \not{n}_+}{4} \psi_c \sim \lambda^2, \quad \xi_{hc} = \frac{\not{n}_- \not{n}_+}{4} \psi_{hc} \sim \lambda, \\ q_s &\sim \lambda^3, \quad h_v = \frac{1 + \not{v}}{2} Q_v \sim \lambda^3. \end{aligned} \quad (4)$$

Here v is the velocity of the heavy quark. $\xi_{c,hc}$ and h_v are large components of the collinear, hard-collinear and heavy quark fields, respectively. Small components of the heavy quark field, H_v , and collinear quarks, η_{hc} and η_c , can be integrated out at tree level by solving the equation of motion. Scalings of gluon fields have a similar behavior with their momenta

$$\begin{aligned} n_+ A_c &\sim 1, \quad n_- A_c \sim \lambda^4, \quad A_{\perp c} \sim \lambda^2, \quad A_s \sim \lambda^2, \\ n_+ A_{hc} &\sim 1, \quad n_- A_{hc} \sim \lambda^2, \quad A_{\perp hc} \sim \lambda. \end{aligned} \quad (5)$$

From the relativistic normalization condition, we find that the baryonic states in the effective theory, taking the Λ_b and Λ as an example, have the scaling

$$|\Lambda_b\rangle \sim \lambda^{-3}, \quad |\Lambda\rangle \sim \lambda^{-2}, \quad (6)$$

where we did not specify the differences with the states in QCD. Presumably these differences may introduce more power corrections, but they are left out here, since the leading-power behavior is unlikely to change. Decay constants of baryons defined via [22, 23]

$$\begin{aligned}\epsilon^{ijk}\langle 0|(u^i C \gamma_5 d^j) h_v^k |\Lambda_b\rangle &= f_{\Lambda_b}^{(1)} u_{\Lambda_b}, \quad \epsilon^{ijk}\langle 0|(u^i C \gamma_5 \not{v} d^j) h_v^k |\Lambda_b\rangle = f_{\Lambda_b}^{(2)} u_{\Lambda_b}, \\ \epsilon^{ijk}\langle 0|(u^i C \gamma_5 \frac{\not{v}_+}{2} d^j) \frac{\not{v}_+}{2} s^k |\Lambda(p')\rangle &= f_{\Lambda} \frac{n_+ p'}{2} \frac{\not{v}_+}{2} u_{\Lambda},\end{aligned}$$

scale as $f_{\Lambda_b} \sim \lambda^6$ and $f_{\Lambda} \sim \lambda^4$ with f_{Λ_b} denoting both $f_{\Lambda_b}^{(1)}$ and $f_{\Lambda_b}^{(2)}$.

In SCET, integration of the fluctuations with large virtualities proceeds in two-steps [15, 19]. In the first step, hard scales, caused by the interaction between the collinear sector and heavy quark, and between two or more collinear sectors with different directions, are integrated out and thereby QCD is matched onto an intermediate effective theory, called SCET_I. In this effective theory gauge invariant operators are built out of fields of hard-collinear quarks or soft gluons and quarks. The leading-power and next-to-leading power terms having non-zero matrix elements between the baryonic transition are constructed as

$$\begin{aligned}O_j^A(s) &= (\bar{\xi}_{hc} W_{hc})_s \Gamma_j Y_s^\dagger h_v, \\ O_j^B(s_1, s_2) &= (\bar{\xi}_{hc} W_{hc})_{s_1} (W_{hc}^\dagger i D_{\perp \mu} W_{hc})_{s_2} \Gamma'_j Y_s^\dagger h_v, \\ O_j^C(s_1, s_2, s_3) &= (\bar{\xi}_{hc} W_{hc})_{s_1} (W_{hc}^\dagger i D_{\perp \mu_1} W_{hc})_{s_2} (W_{hc}^\dagger i D_{\perp \mu_2} W_{hc})_{s_3} \Gamma'_j Y_s^\dagger h_v, \\ O_j^D(s_1, s_2) &= (\bar{\xi}_{hc} W_{hc})_{s_1} (W_{hc}^\dagger i n_- D W_{hc} - i n_- D_s)_{s_2} \Gamma'_j Y_s^\dagger h_v, \\ O_j^E(s, t) &= (\bar{\xi}_{hc} W_{hc})_s (i D_s^\mu)_t \Gamma'_j Y_s^\dagger h_v,\end{aligned}\tag{7}$$

where the hard-collinear field with the subscript s is evaluated at $x + s n_+$, while the soft field with the subscript t is evaluated at $x + t n_-$, with x being the space coordinate from the QCD current. Γ'_j is one of the following gamma matrices

$$\Gamma'_j = (1, \gamma_5, \gamma_\perp, \gamma_\perp \gamma_5).\tag{8}$$

The W_{hc} and Y_s are hard-collinear and soft Wilson lines, respectively [15, 19].

Integration of the hard-collinear mode will result in the final SCET, named as SCET_{II} for convenience. In SCET_I, the generic power scalings of the operators in Eq. (7) are

$$O^A \sim \lambda^4, \quad Q^B \sim \lambda^5, \quad O^{C,D,E} \sim \lambda^6.\tag{9}$$

But none of them have the right quantum numbers with baryons in the initial and final state. Thus the matching of these operators from SCET_I onto SCET_{II} will induce additional power suppressions and one of our goals is to count these suppressions. To the end, we will demonstrate that the contribution from the O^A operator starts at the $\mathcal{O}(\lambda^9)$, while the other types of operators have the power λ^{11} .

A. General analysis in SCET_{II}

To represent the quantum numbers of the Λ_b and Λ baryon, at least the fields $q_s q_s h_v$ and three collinear quark fields are needed. In the light-cone gauge a most general form of an operator with

non-vanishing matrix elements can be taken as [19]

$$[\text{objects}] \times (\bar{\xi}_c \{1, \not{n}_+/2\} \Gamma'_k q_s) (\bar{\xi}_c \{1, \not{n}_+/2\} \Gamma'_l q_s) (\bar{\xi}_c \Gamma'_j h_v). \quad (10)$$

where the objects in the brackets are combinations of the building blocks:

$$\left| \begin{array}{c|c|c|c|c|c} (in_- \partial)^{-1} & n_-^\mu & \partial_\perp, A_{\perp c}, A_{\perp s} & n_- \partial, n_- A_c & \bar{q}_s \frac{\not{n}_+}{2} \Gamma'_m q_s & \bar{q}_s \Gamma''_m q_s \\ n_1 & n_3 & n_5 & n_7 & n_{9a} & n_{9c} \\ \hline (in_+ \partial)^{-1} & n_+^\mu & n_+ \partial, n_+ A_s & \bar{\xi}_c \frac{\not{n}_+}{2} \Gamma'_m \xi_c & \bar{q}_s \frac{\not{n}_-}{2} \Gamma'_m q_s & \\ n_2 & n_4 & n_6 & n_8 & n_{9b} & \end{array} \right|,$$

with the integers n_i being the number of occurrences of O_i in an operator. $\Gamma'_{j,k,l,m}$ take one of the forms in Eq. (8), while Γ''_m is a basis for the remaining eight boost-invariant Dirac structures. We, following Ref. [19], use the power scaling, boost invariance and the matching of mass dimensions to pick up the allowed forms. The notation for these symbols is used as: $[\lambda]_O = n$ means that O scales with λ^n , the “boost” label corresponds to the scaling α^n of O under boosts $n_- \rightarrow \alpha n_-$, $n_+ \rightarrow \alpha^{-1} n_+$; the mass dimension is denoted by $[d]_O$. Using the properties of these building blocks which are discussed in detail in table 2 of Ref. [19], we find an operator in the final effective theory has the scalings

$$\begin{aligned} [\lambda] &= 15 - 2n_1 + 2n_5 + 2n_6 + 4n_7 + 4n_8 + 6(n_{9a} + n_{9b} + n_{9c}), \\ [\alpha] &= 0 = -n_1 + n_2 + n_3 - n_4 - n_6 + n_7 - n_8 - n_{9a} + n_{9b}, \\ [d] &= 9 - n_1 - n_2 + n_5 + n_6 + n_7 + 3(n_8 + n_{9a} + n_{9b} + n_{9c}), \end{aligned} \quad (11)$$

from which we have

$$[\lambda] = 6 + [d] - n_3 + n_4 + n_5 + 2n_6 + 2n_7 + 2n_8 + 4n_{9a} + 2n_{9b} + 3n_{9c}. \quad (12)$$

In operators $O^{A,B,C,E}$ the only Lorentz structure having nonzero contraction with n_-^μ is $\epsilon_{\alpha\perp\beta\perp\mu\nu} n_-^\mu n_+^\nu$, and thus $n_3 \leq n_4$. For the O^A operator, $[d] = 3$ and there is only one nontrivial solution with $[\lambda] = 9$, $n_1 = n_2 = 3$, $n_3 = n_4$ and $n_i = 0 (i \geq 5)$. Since there is no free Lorentz index containing n_-^μ or n_+^μ in Γ'_j , the equality $n_3 = n_4$ rules out the possibility of $\not{n}_+/2$ in Eq. (10).

As for O^B operator, $[d] = 4$ thus $[\lambda] \geq 10$. Since n_1 is an integer, the leading contribution from this operator has the scaling $[\lambda] = 11$. One solution is $n_1 = 2$, $n_2 = 3$, $n_4 - n_3 = 1$, $n_i = 0 (i \geq 5)$ and the other is $n_3 = n_4$, $n_1 = n_2 = 3$, $n_5 = 1$, $n_i = 0 (i \geq 6)$. The latter one corresponds to the higher Fock state contribution, due to the presence of an extra soft or collinear gluon. The O^E also belongs to this type.

In the light-cone gauge $(W_{hc}^\dagger in_- DW_{hc} - in_- D_s)$ reduces to $n_- A_{hc}$. In the O^D operator, $[d] = 4$ and the factor n_-^μ contracts with the gluon field $A_{hc\mu}$. After the elimination of the hard-collinear fields, n_-^μ can not be a free Lorentz index and maybe it is contracted: with n_+ which is a constant or in the form of $\epsilon_{\alpha\perp\beta\perp\mu\nu} n_-^\mu n_+^\nu$; with a gamma matrix as $\bar{q}_s \frac{\not{n}_-}{2} \Gamma'_m q_s$; with a derivative to a soft field in the form of $n_- \partial$; or with a collinear gluon field as $n_- A_c$. In the first contraction, $n_4 \geq n_3$, and $[\lambda] \geq 10$. Due to the integer constraint on n_1 , this operator has the scaling $[\lambda] = 11$ and its solution is similar to the one in O^B . For the rest cases, n_3 may be larger than n_4 by one unit, but $n_7 > 0$ or $n_{9b} > 0$, causing more power suppressions and resulting in $[\lambda] > 11$.

For the operator O^C , $[d] = 5$ and $[\lambda] \geq 11$. The solution having the power $[\lambda] = 11$ is $n_1 = n_2 = 2$, $n_3 = n_4$ and $n_i = 0 (i \geq 5)$.

The above matching analysis indicates that the operator O^A is indeed dominant and others are λ^2 suppressed. Taking into account the power scalings of baryonic states, we obtain the scaling laws for operator matrix elements

$$\langle \Lambda | O^A | \Lambda_b \rangle \sim \lambda^4, \quad \langle \Lambda | O^{B,C,D,E} | \Lambda_b \rangle \sim \lambda^6. \quad (13)$$

B. Tree-level Matching

Now we will perform a tree-level matching from SCET_I to SCET_{II} , and identify various terms to different types of operators. In this procedure, the hard-collinear quark field is first expressed as a product of soft and collinear fields and the hard-collinear gluon fields. Then the hard-collinear gluons are integrated out by solving the equation of motion for the Yang-Mills fields and their expressions in terms of soft and collinear quarks and gluons will be substituted back into the hard-collinear quark field. For simplicity, we shall work in the light-cone gauge $n_+ A_{hc} = n_+ A_c = n_- A_s = 0$ and the gauge invariant form can be obtained by the field redefinition technique.

The QCD currents can be matched onto the effective currents in the SCET

$$J^{QCD} = [\bar{\psi}(x) \Gamma b](x) \rightarrow e^{-im_b v \cdot x} [\bar{\psi} \Gamma Q](x) \quad (14)$$

with

$$\begin{aligned} \psi &= \xi_c + \eta_c + \xi_{hc} + \eta_{hc} + q_s \\ &= \xi_c + \xi_{hc} + q_s - \frac{1}{in_+ D_s} \frac{\not{n}_+}{2} [(i\not{D}_\perp)(\xi_c + \xi_{hc}) + (g\mathcal{A}_{\perp c} + g\mathcal{A}_{\perp hc})q_s], \\ \mathcal{Q} &= \left(1 + \frac{i\not{D}_s}{2m_b}\right) h_v - \frac{1}{n_- v} \frac{\not{n}_-}{2m_b} (g\mathcal{A}_{\perp c} + g\mathcal{A}_{\perp hc}) h_v \\ &+ \frac{1}{2m_b n_- v} \left[\frac{1}{in_+ \partial} (g\mathcal{A}_{\perp c} + g\mathcal{A}_{\perp hc})(g\mathcal{A}_{\perp c} + g\mathcal{A}_{\perp hc}) \right] h_v \\ &- \left[\left\{ \frac{1}{m_b n_- v} \frac{\not{n}_- \not{n}_+}{4} - \frac{n_+ v}{n_- v in_+ \partial} \right\} (n_- A_{hc}) \right] h_v + \mathcal{O}(\lambda^4 h_v), \end{aligned} \quad (15)$$

with the derivative $1/n_+ \partial$ acting on the collinear field in the square bracket.

In the light-cone gauge, the collinear quark Lagrangian reads as

$$\mathcal{L} = \bar{\xi} \left(in_- D + [i\not{D}_\perp] \frac{1}{in_+ D_s} [i\not{D}_\perp] \right) \frac{\not{n}_+}{2} \xi + \dots, \quad (16)$$

with the ellipses standing for all other terms. Here ξ and the collinear gluon in the covariant derivative denote both collinear and hard-collinear field and will be substituted as $\xi \rightarrow \xi_c + \xi_{hc}$, $A_c \rightarrow A_c + A_{hc}$. With the use of the equation of motion, the ξ_{hc} can be integrated out and in particular, the solution (dropping the terms not satisfying momentum conservation)

$$\xi_{hc} \sim -\frac{1}{in_- \partial} \left(gn_- A_{hc} + i\not{D}_\perp \frac{1}{in_+ \partial} i\not{D}_\perp \right) \xi_c$$

contributes to $\psi^{(6)}$ as

$$\psi^{(6)} = -\frac{1}{in_- \partial} \left(gA_{\perp hc}^{(3)} \frac{1}{in_+ \partial} gA_{\perp hc}^{(3)} + gn_- A_{hc}^{(6)} \right) \xi_c + \dots, \quad (17)$$

in which the expressions of gluons will be specified below. The other useful pieces are [19]

$$\begin{aligned} \psi^{(2)} &= \xi_c, \\ \psi^{(5)} &= \frac{1}{in_+ \partial} gA_{\perp hc}^{(3)} \frac{\not{n}_+}{2} \xi_c - \frac{1}{in_- \partial} \left((i\not{D}_{\perp c} + gA_{\perp s}) \frac{1}{in_+ \partial} gA_{\perp hc}^{(3)} \right. \\ &\quad \left. + gA_{\perp hc}^{(3)} \frac{1}{in_+ \partial} (i\not{D}_{\perp c} + gA_{\perp s}) \right) \xi_c - \frac{1}{in_- \partial} gn_- A_{hc}^{(5)} \xi_c + \dots, \end{aligned} \quad (18)$$

where the first term in $\psi^{(5)}$ is from the small component of the hard-collinear quark field $\eta_{hc}^{(5)}$ and the ones in the large parentheses are from $\xi_{hc}^{(5)}$. The relevant hard-collinear gluon field is expanded as [19]

$$\begin{aligned} A_{\perp hc}^{(3)} &= gT^A \frac{1}{in_+ \partial in_- \partial} \{ \bar{q}_s \gamma_{\perp} T^A \xi_c + h.c. \}, \\ n_- A_{hc}^{(5)} &= -\frac{2}{(in_+ \partial)^2} \left\{ i\not{D}^{\mu\perp} [in_+ \partial A_{\mu\perp hc}^{(3)}] - g \left[in_+ \partial A_c^{\mu\perp}, A_{\mu\perp hc}^{(3)} \right] \right. \\ &\quad \left. - 2gT^A \left\{ \bar{\xi}_c T^A \left(\frac{\not{n}_+}{2} - \frac{1}{in_- \partial} gA_{\perp c} \right) q_s + h.c. \right\} \right\}, \\ n_- A_{hc}^{(6)} &= -\frac{2}{(in_+ \partial)^2} \left[-2[in_+ \partial A_{\perp hc}^{(3)\mu}, A_{\mu\perp hc}^{(3)}] + 2gT^A \left\{ \bar{\xi}_c T^A \left(\frac{1}{in_- \partial} gA_{\perp hc}^{(3)} \right) q_s + h.c. \right\} \right]. \end{aligned} \quad (19)$$

with the covariant derivative $i\not{D}^{\mu}\mathcal{O} = i\partial^{\mu}\mathcal{O} + g[A_c^{\mu} + A_s^{\mu}, \mathcal{O}]$.

Before substituting the hard-collinear fields into the currents, we first count the collinear quark numbers. The final baryonic state contains three quarks, and has collinear quark number +3. In order to have nonzero matrix elements, the effective currents in the SCET must have the collinear quark number 3 as well. Let us recall that the gluon field $A_{\perp hc}^{(3)}$ contains one collinear quark (or antiquark depending on the interaction form in the effective theory), while $n_- A_{hc}^{(6)}$ may contain two collinear quarks. For the expression of $\psi^{(n)}$, we note that $\psi^{(2)}$ (and also $\psi^{(4)}$) has collinear quark number -1, while $\psi^{(3)}$ has collinear quark number 0. The most nontrivial terms are: $\psi^{(5)}$ which has a collinear quark number -2 or 0, and $\psi^{(6)}$ with collinear quark number -3 (or ± 1). The combinations having the leading and next-to-leading power scalings indeed take the forms as O^A , $O^{B,C,D}$ and O^E .

Substituting $A_{\perp hc}^{(3)}$, $n_- A_{hc}^{(6)}$ and $\psi^{(6)}$ into the effective currents, we have the leading term in the expansion

$$J^{(9)} = -\bar{\xi}_c \left(gA_{\perp hc}^{(3)} \frac{1}{in_+ \partial} gA_{\perp hc}^{(3)} + gn_- A_{hc}^{(6)} \right) \frac{1}{in_- \partial} \Gamma h_v. \quad (20)$$

The first term in the above equation contains two hard-collinear gluons emitted from the hard-collinear quark, and is depicted as the first diagram in Fig. 1. In this figure, the dashed lines denote the collinear quarks, while the solid lines are soft spectators. The thick lines represent the




FIG. 1: Tree matching diagrams for the heavy-to-light baryonic form factors. The dashed lines denote the collinear quarks, while the solid lines are soft spectators. The thick lines represent the heavy bottom quark. The spring lines denote a collinear gluon n_-A_{hc} while the spring+solid lines denote the $A_{\perp hc}$. The hard modes have been integrated out and shrunk to the black point.

heavy bottom quark. The spring lines denote a collinear gluon n_-A_{hc} while spring+solid lines denote the $A_{\perp hc}$. In the $n_-A_{hc}^{(6)}$, the trigluon term, corresponding to Fig (1c), vanishes and it can be understood as follows. The three quarks have antisymmetric colors in both initial and final baryons, and thus the color rearrangement factor in this diagram is zero

$$\epsilon^{ijk}\epsilon^{i'j'k'}T_{ii'}^AT_{jj'}^BT_{kk'}^Cf^{ABC} = \epsilon^{ikj}\epsilon^{i'k'j'}T_{ii'}^AT_{kk'}^BT_{jj'}^Cf^{ABC} = 0. \quad (21)$$

The current $J^{(9)}$ originates from the large component of the hard-collinear quark field $\xi_{hc}^{(6)}$ as shown in Eq. (17) and thereby the Lorentz structure is reduced:

$$J^{(9)} \sim \frac{\not{n}_+\not{n}_-}{4}\Gamma\frac{1+\not{v}}{2} \rightarrow \Gamma'_j, \quad (22)$$

as expected in the large recoil limit.

The other combinations of operators start from λ^{11}

$$\begin{aligned} J^{(11)} = & -\frac{1}{n_-v}\bar{\psi}^{(5)}\Gamma\frac{\not{n}_-}{2m_b}gA_{\perp hc}^{(3)}h_v + \frac{1}{2m_bn_-v}\bar{\psi}^{(2)}\Gamma\frac{1}{in_+\partial}gA_{\perp hc}^{(3)}gA_{\perp hc}^{(3)}h_v \\ & -\bar{\psi}^{(2)}\Gamma\left[\left\{\frac{1}{m_bn_-v}\frac{\not{n}_-\not{n}_+}{4} - \frac{n_+v}{n_-vin_+\partial}\right\}(n_-A_{hc}^{(6)})\right]h_v + \bar{\psi}^{(6)}\Gamma\frac{i\not{D}_s}{2m_b}h_v + \dots, \end{aligned} \quad (23)$$

where these four pieces can be incorporated into the operators $O^{B,C,D,E}$ respectively. It should be noted that except the second term, the other terms can have different Lorentz structures with the reduced form as in Eq. (22). For instance, the fourth term is from the small component of the heavy bottom quark, which has the Lorentz structure $\frac{\not{n}_+\not{n}_-}{4}\Gamma\frac{1-\not{v}}{2}$.

We also show the tree-level matching diagrams for the $O^{B,C,D}$ operators in Fig. 1. However the higher Fock state contributions, either from O^E having the similar structure with O^A except that one additional soft gluon is emitted from the hard vertex, or from the operator O^B , are not

depicted. Graphically speaking the dominance of O^A can be understood as follows. In the three diagrams (a,d,e) one commonality is that the two gluons interact with a soft quark from the initial state and a collinear quark in the final external state, and thereby these two vertices have the same power scaling. However in the first diagram the quark propagator next to the weak vertex has the form $1/(n-p) \sim 1/\lambda^2$ while the rest quark propagators are of order λ^0 , leading to the enhancement of the first diagram.

III. ANALYSIS OF THE TRANSITION DIAGRAMS IN QCD

In this section, we will analyze the leading power behaviors of the baryonic transition form factors in QCD, whose Feynman diagrams are depicted in Fig. 2. We adopt the ansatz that the fast-moving baryon is composed of three collinear constituents, therefore at least two gluons are exchanged and these gluons must be far off-shell. We will not include the contributions involving higher Fock states, as at least one more gluon is needed. As we have already shown, the trigluon diagrams give vanishing contributions and thereby will not be considered either. There are seven diagrams shown in Fig. 2: three of them (a,b,c) containing the momentum exchange by two gluons between the spectator quark system and the energetic light quark connecting the the electroweak vertex; the same number of diagrams (e,f,g) having two gluons emitted from the heavy quark; the rest diagram (d) in which the light spectator system receives momentum exchange from both the energetic quark and the heavy quark. The inclusion of the flavor index will give another seven diagrams, but only leads to the exchange of momentum fractions of the light spectator quarks.

The leading twist LCDA of a light baryon, such as Λ , is [6, 23]

$$\epsilon^{ijk} \frac{n+p'}{8} \frac{1}{6} (C \not{n} - \gamma_5)_{\beta\alpha} (\bar{u}_\Lambda)_\gamma, \quad (24)$$

with i, j, k being the color indices and α, β, γ being the spinor indices. For the heavy baryon, several types of LCDAs emerge [22]

$$\begin{aligned} & \frac{1}{48} \epsilon^{ijk} (\not{n} + \gamma_5 C)_{\alpha\beta} (u_{\Lambda_b})_\gamma, \quad \frac{1}{48} \epsilon^{ijk} (\not{n} - \gamma_5 C)_{\alpha\beta} (u_{\Lambda_b})_\gamma, \\ & \frac{1}{48} \epsilon^{ijk} (\gamma_5 C)_{\alpha\beta} (u_{\Lambda_b})_\gamma, \quad \frac{1}{48} \epsilon^{ijk} (\not{n} + \not{n}' - \gamma_5 C)_{\alpha\beta} (u_{\Lambda_b})_\gamma. \end{aligned} \quad (25)$$

In the leading power matrix elements, only the first type of LCDA contributes. We choose the momentum fractions of the three collinear quarks in the light baryon as y_1, y_2, y_3 and the momentum fractions of the soft spectator quarks (in the direction n_+) in the initial state as x_2 and x_3 . The corresponding momenta will be denoted as p'_1, p'_2, p'_3 for the collinear quarks, and p_2, p_3 for the soft quarks.

The calculation will be simplified by the following two observations.

- If both vertices of a hard-collinear gluon are attached to collinear quarks, only the transverse component of this gluon contributes.
- In the light spectator system (usually called a diquark), only the diagrams with even number of gluon transverse indices are nonzero. For instance, as shown in Fig 2(b,e,f), if only one




FIG. 2: Feynman Diagrams for heavy-to-light baryonic form factors in QCD. Trigluon diagrams having wrong color factors are not shown.

hard-collinear gluon is emitted from the light quark or the heavy quark, this gluon has to be in the form $n_- A_{hc}$ or $n_+ A_{hc}$.

The first observation can be proved by writing the amplitudes as

$$[\bar{q}_1 \gamma_\mu \dots] \times [\bar{q}_2 \gamma^\mu \dots] = [\bar{q}_1 \gamma_{\perp \mu} \dots] \times [\bar{q}_2 \gamma^{\perp \mu} \dots], \quad (26)$$

with q_1 and q_2 being the two collinear quarks attached to the gluon. The second one is based on the fact that the two-spectator system technically forms a trace in the spinor space. There is no transverse index from the external wave functions, and thereby the internal ones from the exchanged gluons must be even.

The leading power contributions from Fig. 2(a,b) can be matched onto the O^A operator. In Fig. 2(a), using the first observation, one of the two gluons (the right one) is connected to two collinear quarks, and only the transverse component is left. With the second observation, the other gluon must take the transverse component as well. In the numerator of the quark propagator between the two gluons, the collinear momentum $\not{p}'_1 + \not{p}'_3$ does not contribute since it is next to the light spinor: $\bar{u}_\Lambda \gamma_\perp (\not{p}'_1 + \not{p}'_3) = 0$. This propagator is simplified as

$$\frac{i(-\not{p}_3 + \not{p}'_1 + \not{p}'_3)}{(-p_3 + p'_1 + p'_3)^2} \simeq \frac{i\not{p}_+}{2(y_1 + y_3)n_+ p'}, \quad (27)$$

which scales as λ^0 . The other quark propagator is reduced to

$$\frac{i(-\not{p}_3 - \not{p}_2 + \not{p}')}{(-p_2 - p_3 + p')^2} \simeq -\frac{i\not{p}_-}{2(x_2 + x_3)m_{\Lambda_b} n_- v}, \quad (28)$$

which has the scaling $1/\lambda^2$. Here we have used $x_2, x_3 \sim \Lambda/m_{\Lambda_b}$ for the soft momentum fraction. Combining these pieces, this diagram has the form

$$\begin{aligned}
F^{(a)} &= C_N g_s^4 \int dy_2 dy_3 dx_2 dx_3 f_{\Lambda_b} f_{\Lambda} \Phi_{\Lambda_b}(x_1, x_2, x_3) \Phi_{\Lambda}(y_1, y_2, y_3) \\
&\times \frac{i}{y_3 x_3 m_{\Lambda_b} n_+ p' n_- v} \frac{i}{y_2 x_2 m_{\Lambda_b} n_+ p' n_- v} \\
&\times \bar{u}_{\Lambda} \gamma_{\perp}^{\mu} \frac{i \not{p}_+}{2(y_1 + y_3) n_+ p'} \gamma_{\perp}^{\nu} \frac{-i \not{p}_-}{2(x_2 + x_3) m_{\Lambda_b} n_- v} \Gamma u_{\Lambda_b} \\
&\times \frac{n_+ p'}{64} (C \not{p}_- \gamma_5)_{\alpha\beta} (\gamma_{\nu})_{\alpha\alpha'} (\gamma_{\mu})_{\beta\beta'} (\not{p}_+ \gamma_5 C)_{\beta'\alpha'} \\
&\propto \lambda^{10}/\lambda^6 \sim \lambda^4,
\end{aligned} \tag{29}$$

where the scaling λ^{10} is from decay constants and $1/\lambda^6$ comes from the two gluons propagators and the propagator in Eq. (28). C_N is the color factor

$$C_N = \frac{1}{36} \epsilon^{ijk} \epsilon^{i'j'k'} (T^b T^a)^{ii'} (T^b)^{jj'} (T^a)^{kk'} = \frac{2}{27}. \tag{30}$$

Eq. (29) confirms our power counting analysis given in the previous section. Furthermore as indicated in the third line of the above equation, the light spectator (diquark) system is proportional to $g^{\perp\mu\nu}$ which results in the Lorentz structure

$$F^{(a)} \propto \bar{u}_{\Lambda} \frac{\not{p}_+ \not{p}_-}{4} \Gamma \frac{1 + \not{p}}{2} u_{\Lambda_b}, \tag{31}$$

where the large energy and heavy quark symmetries are manifestly demonstrated again.

In Fig. 2(b), the upper gluon vertex is replaced by $\not{p}_+/2$ and the quark propagator next to the electroweak vertex is of the form $\not{p}_-/2$. Therefore this diagram has the same structure:

$$\begin{aligned}
F^{(b)} &= C_N g_s^4 \int dy_2 dy_3 dx_2 dx_3 f_{\Lambda_b} f_{\Lambda} \Phi_{\Lambda_b}(x_1, x_2, x_3) \Phi_{\Lambda}(y_1, y_2, y_3) \\
&\times \frac{i}{y_3 x_3 m_{\Lambda_b} n_+ p' n_- v} \frac{i}{(y_2 + y_3)(x_2 + x_3) m_{\Lambda_b} n_+ p' n_- v} \times \bar{u}_{\Lambda} \frac{\not{p}_+}{2} \frac{-i \not{p}_-}{2(x_2 + x_3) n_- v m_{\Lambda_b}} \Gamma u_{\Lambda_b} \\
&\times \frac{n_+ p'}{2} (C \not{p}_- \gamma_5)_{\alpha\beta} (\gamma_{\perp\mu} \frac{i \not{p}_+}{2(y_2 + y_3) n_+ p'} \not{p}_-)_{\alpha\alpha'} (\gamma_{\perp}^{\mu})_{\beta\beta'} (\not{p}_+ \gamma_5 C)_{\beta'\alpha'} \\
&\propto \lambda^{10}/\lambda^6 \sim \lambda^4.
\end{aligned} \tag{32}$$

In Fig. 2(c), based on the first observation, the upper gluon is transverse; thus there are either one or three transverse indices in the light spectator system, leading to vanishing contribution.

In Fig. 2(d), both gluons can only contain transverse components and this diagram can be matched onto the operator O^B . Both the heavy quark and light quark propagators scale as λ^0 and thus

$$\begin{aligned}
F^{(d)} &= C_N g_s^4 \int dy_2 dy_3 dx_2 dx_3 f_{\Lambda_b} f_{\Lambda} \Phi_{\Lambda_b}(x_1, x_2, x_3) \Phi_{\Lambda}(y_1, y_2, y_3) \frac{i}{y_3 x_3 m_{\Lambda_b} n_+ p' n_- v} \frac{i}{x_2 y_2 m_{\Lambda_b} n_+ p' n_- v} \\
&\times \bar{u}_{\Lambda} \gamma_{\perp}^{\nu} \frac{i \not{p}_+}{2(y_1 + y_3) n_+ p'} \Gamma \frac{i \not{p}_-}{2 m_{\Lambda_b} n_- v} \gamma_{\perp}^{\mu} u_{\Lambda_b} \times \frac{n_+ p'}{2} (C \not{p}_- \gamma_5)_{\alpha\beta} (\gamma_{\perp\mu})_{\alpha\alpha'} (\gamma_{\perp\nu})_{\beta\beta'} (\not{p}_+ \gamma_5 C)_{\beta'\alpha'} \\
&\propto \lambda^{10}/\lambda^4 \sim \lambda^6.
\end{aligned} \tag{33}$$

with again λ^{10} from decay constants and $1/\lambda^4$ from the two gluon propagators. Of particular interest is that the Lorentz structure in this diagram has the form

$$F^{(d)} \propto \bar{u}_\Lambda \gamma_\perp^\nu \frac{i\not{p}_+}{2(y_1 + y_3)n_+p'} \Gamma \frac{i\not{p}_-}{2m_{\Lambda_b}n_-v} \gamma_\perp^\mu u_{\Lambda_b}, \quad (34)$$

which manifestly breaks the large recoil symmetries.

In Fig. 2(e), the gluon attaching to the two light quarks is transverse while the component n_-A_{hc} contributes at the heavy quark propagator. This diagram corresponds to the operator O^D . Using the first observation, the light quark propagator scales as λ^0 and thus

$$\begin{aligned} F^{(e)} &= C_N g_s^4 \int dy_2 dy_3 dx_2 dx_3 f_{\Lambda_b} f_\Lambda \Phi_{\Lambda_b}(x_1, x_2, x_3) \Phi_\Lambda(y_1, y_2, y_3) \\ &\quad \times \frac{i}{y_3 x_3 m_{\Lambda_b} n_+ p' n_- v} \frac{i}{(y_2 + y_3)(x_2 + x_3) m_{\Lambda_b} n_+ p' n_- v} \\ &\quad \times \bar{u}_\Lambda \Gamma \left[\frac{-in_+v}{(y_2 + y_3)n_+ p' n_- v} + \frac{\not{p}_- \not{p}_+}{4} \frac{i}{n_- v m_{\Lambda_b}} \right] u_{\Lambda_b} \\ &\quad \times \frac{n_+ p'}{2} (C \not{p}_- \gamma_5)_{\alpha\beta} (\gamma_{\perp\mu} \frac{i\not{p}_+}{2(y_2 + y_3)n_+ p'} \not{p}_-)_{\alpha\alpha'} (\gamma_\perp^\mu)_{\beta\beta'} (\not{p}_+ \gamma_5 C)_{\beta'\alpha'} \\ &\propto \lambda^{10}/\lambda^4 \sim \lambda^6. \end{aligned} \quad (35)$$

The first term in the square bracket obeys the large recoil symmetries, but the integral over $y_2 + y_3$ in it is divergent. It is worthwhile to point out that in the SCET solution for the operator O^E in the previous section, the number of the occurrence of $1/(in_+\partial)$ is found to be $n_2 = 3$, which means the momentum fractions for the light baryon can appear only three times. The additional momentum fraction arises from the short-distance coefficients, for instance at tree-level shown in Eq. (15).

In Fig. (2f), the gluon attaching to the b quark can not contribute with the transverse component based on the second observation. The n_+A_{hc} component can be absorbed into the Wilson line, one necessary piece in the gauge invariant definition of the SCET operators. Thus this diagram is incorporated into the operator O^A and its scaling is

$$\begin{aligned} F^{(f)} &= C_N g_s^4 \int dy_2 dy_3 dx_2 dx_3 f_{\Lambda_b} f_\Lambda \Phi_{\Lambda_b}(x_1, x_2, x_3) \Phi_\Lambda(y_1, y_2, y_3) \\ &\quad \times \frac{i}{y_3 x_3 m_{\Lambda_b} n_+ p' n_- v} \frac{i}{(y_2 + y_3)(x_2 + x_3) m_{\Lambda_b} n_+ p' n_- v} \times \bar{u}_\Lambda \Gamma \frac{-i}{(y_2 + y_3)n_+ p'} u_{\Lambda_b} \\ &\quad \times \frac{n_+ p'}{64} (C \not{p}_- \gamma_5)_{\alpha\beta} (\not{p}_+ \frac{-i\not{p}_-}{2(x_2 + x_3)m_{\Lambda_b}n_-v} \gamma_{\perp\mu})_{\alpha\alpha'} (\gamma_\perp^\mu)_{\beta\beta'} (\not{p}_+ \gamma_5 C)_{\beta'\alpha'} \\ &\propto \lambda^{10}/\lambda^6 \sim \lambda^4. \end{aligned} \quad (36)$$

In particular this contribution cancels the one from Fig. (2b).

In Fig. 2(g), the two heavy quark propagators have the offshellness of order m_b^2 and can be shrunk to one point. Suppose that the two gluons are transverse, and then it is incorporated into

O^C and its power scaling is

$$\begin{aligned}
F^{(g1)} &= C_N g_s^4 \int dy_2 dy_3 dx_2 dx_3 f_{\Lambda_b} f_{\Lambda} \Phi_{\Lambda_b}(x_1, x_2, x_3) \Phi_{\Lambda}(y_1, y_2, y_3) \frac{i}{y_3 x_3 m_{\Lambda_b} n_+ p' n_- v} \frac{i}{x_2 y_2 m_{\Lambda_b} n_+ p' n_- v} \\
&\quad \times \bar{u}_{\Lambda} \Gamma \frac{-1}{(y_2 + y_3) n_+ p'} \gamma_{\perp \mu} \gamma_{\perp \nu} u_{\Lambda_b} \times \frac{n_+ p'}{2} (C \not{\epsilon} - \gamma_5)_{\alpha \beta} (\gamma_{\perp \mu})_{\alpha \alpha'} (\gamma_{\perp}^{\mu})_{\beta \beta'} (\not{\epsilon} + \gamma_5 C)_{\beta' \alpha'} \\
&\propto \lambda^{10} / \lambda^4 \sim \lambda^6,
\end{aligned} \tag{37}$$

where the momentum fraction $1/(y_2 + y_3)$ in the second line comes from the Wilson coefficient for the operator O^C . If the two gluons take the $n_+ A_{hc}$ component for the vertices attaching to the heavy quark, this diagram can be matched onto operator O^A . The contribution is λ^2 suppressed compared to the leading power terms

$$\begin{aligned}
F^{(g2)} &= C_N g_s^4 \int dy_2 dy_3 dx_2 dx_3 f_{\Lambda_b} f_{\Lambda} \Phi_{\Lambda_b}(x_1, x_2, x_3) \Phi_{\Lambda}(y_1, y_2, y_3) \times \frac{i}{y_3 x_3 m_b n_+ p' n_- v} \frac{i}{x_2 y_2 m_b n_+ p' n_- v} \\
&\quad \times \bar{u}_{\Lambda} \Gamma \frac{i}{(y_2 + y_3) n_+ p'} \frac{i}{y_3 n_+ p'} u_{\Lambda_b} \times \frac{n_+ p'}{2} (C \not{\epsilon} - \gamma_5)_{\alpha \beta} (\not{\epsilon}_+)_{\alpha \alpha'} (\not{\epsilon}_+)_{\beta \beta'} (\not{\epsilon} - \gamma_5 C)_{\alpha' \beta'} \\
&\propto \lambda^{10} / \lambda^4 \sim \lambda^6,
\end{aligned} \tag{38}$$

and the integration in this term does not converge.

IV. DISCUSSIONS

As we have shown, in the dominant contribution from the O^A the inverse of derivatives to both collinear fields and soft fields appear three times. In the momentum space these factors will be converted to the inverse of momenta. Let them act on the collinear fields, we obtain the factor $1/(n_+ p')^3$. The energy dependence of a quark field can be read from the propagators

$$\langle 0 | \xi_c(x) \bar{\xi}_c(0) | 0 \rangle = \int \frac{d^4 p'}{(2\pi)^4} e^{-i n_+ p' n_- x/2} \frac{n_+ p'}{p'^2} \frac{\not{\epsilon}_-}{2}. \tag{39}$$

The measure $d^4 p'$ and p'^2 are Lorentz invariant, and thus $\xi_c \sim \sqrt{n_+ p'}$. Then the leading power baryonic transition matrix element scales as

$$\langle \Lambda(p') | O^A(0) | \Lambda_b(p) \rangle \sim \frac{(n_+ p')^{3/2}}{n_+ p'^3} = (n_+ p')^{-3/2},$$

where we have employed the energy independence of baryon states. Using $\bar{u}_{\Lambda} \sim \sqrt{n_+ p'}$ and the definition of the soft form factor in SCET

$$\langle \Lambda(p') | O^A(0) | \Lambda_b(p) \rangle = \xi_{\Lambda}(E) \bar{u}_{\Lambda}(p') \Gamma u_{\Lambda_b}(p),$$

and restoring correct mass dimensions, we obtain the momentum dependence

$$\xi_{\Lambda}(E) \sim \frac{\Lambda^2}{(n_+ p')^2}. \tag{40}$$

This behavior can also be read from the QCD calculation as shown in Eq. (29). But it should be noticed that the above scaling law is different with the results derived in different versions of QCD

light-cone sum rules [5, 6] in which the form factor is dominated by soft processes. To have the power counting, we represent the form factor as an overlap integral of the wave functions in both longitudinal and transverse momentum space

$$\xi_\Lambda(E) = \int \frac{dx_2 d^2\vec{k}_{2\perp} dx_3 d^2\vec{k}_{3\perp}}{(16\pi^3)^2} \psi_{\Lambda_b}(x_2, x_3, \vec{k}_{2\perp}, \vec{k}_{3\perp}) \psi_\Lambda(y_2(x_2), y_3(x_3), \vec{k}_{2\perp}, \vec{k}_{3\perp}), \quad (41)$$

with $y_2(x_2)$ and $y_3(x_3)$ to be fixed by kinematics. From the normalizations of the b-baryon state, we have

$$\int \frac{dx_2 d^2\vec{k}_{2\perp} dx_3 d^2\vec{k}_{3\perp}}{(16\pi^3)^2} |\psi_{\Lambda_b}(x_2, x_3, \vec{k}_{2\perp}, \vec{k}_{3\perp})|^2 = 1, \quad (42)$$

implying that $\psi_{\Lambda_b}(x_2, x_3, \vec{k}_{2\perp}, \vec{k}_{3\perp}) \sim \lambda^{-6}$ since $x_{2,3} \sim \lambda^2$ and $k_{2\perp,3\perp} \sim \lambda^2$. For the light particles, the momentum fraction in the normalization is of order 1, therefore for generic values of $y_{2,3}$, $\psi_\Lambda(y_2, y_3, \vec{k}_{2\perp}, \vec{k}_{3\perp}) \sim \lambda^{-4}$. However the dominance of soft processes leads to the phase suppression and in particular the scalings of the momentum fractions $y_2(x_2) \sim \lambda^2$ and $y_3(x_3) \sim \lambda^2$ result in $\psi_\Lambda(y_2(x_2), y_3(x_3), \vec{k}_{2\perp}, \vec{k}_{3\perp}) \sim 1$. Substituting the scalings for the wave-functions, we obtain

$$\xi_\Lambda(E) \sim \lambda^6, \quad (43)$$

from which we can see the contribution from the soft process is formally λ^2 -suppressed compared to the leading power contribution from the operator O^A .

As a comparison, it is also instructive to recapture the energy dependence of the $B \rightarrow \pi$ form factor in the SCET. Ref. [19] finds that when matching onto SCET_{II} the leading power contribution, from the operator $\bar{\xi}_{hc}\Gamma h_v$, has two powers of $1/(in_+\partial)$. Together with the scalings from the two collinear quark fields, the soft form factor, parametrized via

$$\langle \pi(p') | \bar{\xi}_{hc} h_v | \bar{B}(p) \rangle = 2E \xi_\pi(E)$$

with $E = n_- v n_+ p' / 2 = (m_B^2 - q^2) / (2m_B)$, behaves as

$$\xi_\pi(E) \sim \frac{\Lambda^{3/2} \sqrt{m_b}}{(n_+ p')^2} \sim \frac{(\Lambda/m_B)^{3/2}}{1 - q^2/m_B^2}. \quad (44)$$

When matching to SCET_{II}, the derivatives $1/(in_+\partial)$ and $1/(in_-\partial)$ also contain the momentum fractions: x_2, x_3 or $x_2 + x_3$ for the initial heavy baryon, y_1, y_3, y_3 or some linear combinations depending on the fields acting on. For example, the tree-level factorization formula from Fig. 2(a) has the following integration form as shown in Eq. (29)

$$\int dy_2 dy_3 dx_2 dx_3 \Phi_{\Lambda_b}(x_1, x_2, x_3) \Phi_\Lambda(y_1, y_2, y_3) \frac{1}{x_2 x_3 y_2 y_3 (y_1 + y_3) (x_2 + x_3)}. \quad (45)$$

With the assumption that $\Phi \sim x_2 x_3$ in the limit of $x_2, x_3 \rightarrow 0$ [22, 23], where Φ denotes the LCDA of Λ_b or Λ , the integration is convergent which is different with the mesonic transition form factor ξ_π . In Fig. 2(b) and Fig. 2(f), the involved integral

$$\int dy_2 dy_3 dx_2 dx_3 \Phi_{\Lambda_b}(x_1, x_2, x_3) \Phi_\Lambda(y_1, y_2, y_3) \frac{1}{y_3 x_3 (y_2 + y_3)^2 (x_2 + x_3)^2} \quad (46)$$

is finite as well. The absence of the divergences leads to the factorization of ξ_Λ

$$\xi_\Lambda = f_{\Lambda_b} \Phi_{\Lambda_b}(x_i) \otimes J(x_i, y_i) \otimes f_\Lambda \Phi_\Lambda(y_i), \quad (47)$$

in which $\otimes$ denotes the convolution over momentum fractions x_i and y_i , and the jet function is given as

$$J(x_i, y_i) = -\frac{1}{4} C_N g_s^4 \frac{1}{x_2 x_3 (x_2 + x_3)} \frac{1}{y_2 y_3 (y_1 + y_3)} \frac{1}{(m_{\Lambda_b}^2 - q^2)^2 m_{\Lambda_b}} + (x_2 \leftrightarrow x_3, y_2 \leftrightarrow y_3). \quad (48)$$

It should be cautious that although this formula is valid at tree-level (order α_s^2), whether it can be extended to all orders remains unknown to us and requires further analysis.

On the contrary, the subleading power corrections can not be factorized, for instance the second term from the diagram shown in Fig. 2(g), has the form

$$\int dy_2 dy_3 \frac{1}{y_2 y_3^2 (y_2 + y_3)} \Phi_\Lambda(y_1, y_2, y_3) \sim \log(y_3),$$

which is divergent when y_3 is approaching zero.

To have some numerical estimate, we use the QCD sum rule calculation of the f_{Λ_b} (next-to-leading order in α_s) [24] and f_Λ [23]

$$f_{\Lambda_b} = (0.032 \pm 0.004) \text{GeV}^3, \quad f_\Lambda = (6.0 \pm 0.3) \times 10^{-3} \text{GeV}^2, \quad (49)$$

together with the asymptotic form of Φ_Λ [23] and the parametrized model for Φ_{Λ_b} [22]

$$\Phi_{\Lambda_b}(x_1, x_2, x_3) = x_2 x_3 \left[\frac{m_{\Lambda_b}^4}{\epsilon_0^4} e^{-(x_2+x_3)m_{\Lambda_b}/\epsilon_0} + a_2 C_2^{3/2} (2u-1) \frac{m_{\Lambda_b}^4}{\epsilon_1^4} e^{-(x_2+x_3)m_{\Lambda_b}/\epsilon_1} \right],$$

$$\Phi_\Lambda(y_1, y_2, y_3) = 120 y_1 y_2 y_3, \quad (50)$$

where $\omega = (x_2 + x_3)m_{\Lambda_b}$, $u = x_2/(x_2 + x_3)$, $\epsilon_0 = (200_{-60}^{+130})$ MeV, $\epsilon_1 = (650_{-300}^{+650})$ MeV and $a_2 = 0.333_{-0.333}^{+0.250}$ [22]. With these inputs and the strong coupling constant at the scale $\mu \sim 2$ GeV: $\alpha_s \simeq 0.3$, we calculate the form factor as

$$\xi_\Lambda(q^2 = 0) = -0.012_{-0.023}^{+0.009}, \quad (51)$$

where the displayed uncertainties are from ϵ_0 . For comparison, we quote the soft form factor ξ_Λ computed in the SCET sum rules [5]

$$\xi_\Lambda(q^2 = 0) = 0.38, \quad (52)$$

which is larger by about one order of magnitude.

V. CONCLUSIONS

Weak decays of heavy baryons provide an ideal ground for the extraction of the helicity structure of the electroweak interaction, thanks to the spin correlation and polarization embedded in decay

amplitudes. In the heavy-to-light transition, the most important ingredients incorporating the QCD dynamics are form factors. Due to the variety in the Lorentz structures, the amplitude is governed by a number of form factors. The development of the effective field theory allows us to simplify the form factors and pick up the terms of great importance.

In this work we have analyzed the factorization properties and power scalings of heavy-to-light baryonic form factors at large recoil. Using the soft-collinear effective theory, we proved that the form factors are greatly simplified by the heavy quark and large energy symmetries at leading power in $1/m_b$. This finding indicates that only one function is necessary to parametrize the transition of $\Lambda_b \rightarrow p$ or $\Lambda_b \rightarrow \Lambda$. A general power counting analysis indicates the form factors are of the order Λ^2/E^2 . In contrast to the mesonic case, the leading power form factor can factorize into a convolution of a hard-scattering kernel of order α_s^2 and light-cone distribution amplitudes without encountering any divergence. Using the inputs mainly from QCD sum rules, we calculate the form factor $\xi_\Lambda(E)$ and find it is numerically smaller than the one governed by soft processes, although the latter is formally power-suppressed. We have also discussed the origins for symmetry breaking effects which are suppressed by powers of Λ/m_b and/or Λ/E .

Acknowledgement

The author is very grateful to Ahmed Ali for valuable discussions and carefully reading this manuscript, and to Thorsten Feldmann for pointing out an error in the earlier version of this manuscript. He also thanks Yu-Ming Wang and De-Shan Yang for useful discussions. This work is supported by the Alexander von Humboldt foundation.

-
- [1] M. Artuso, D. M. Asner, P. Ball, E. Baracchini, G. Bell, M. Beneke, J. Berryhill and A. Bevan *et al.*, Eur. Phys. J. C **57**, 309 (2008) [arXiv:0801.1833 [hep-ph]].
 - [2] N. Isgur and M. B. Wise, Nucl. Phys. B **348**, 276 (1991).
 - [3] T. Mannel, W. Roberts, Z. Ryzak, Nucl. Phys. **B355**, 38-53 (1991).
 - [4] T. Mannel and S. Recksiegel, J. Phys. GG **24**, 979 (1998) [hep-ph/9701399].
 - [5] T. Feldmann, M. W. Y. Yip, [arXiv:1111.1844 [hep-ph]].
 - [6] T. Mannel, Y. -M. Wang, [arXiv:1111.1849 [hep-ph]].
 - [7] G. Hiller, A. Kagan, Phys. Rev. **D65**, 074038 (2002) [hep-ph/0108074].
 - [8] G. Hiller, M. Knecht, F. Legger and T. Schietinger, Phys. Lett. B **649**, 152 (2007) [hep-ph/0702191].
 - [9] M. J. Dugan and B. Grinstein, Phys. Lett. B **255**, 583 (1991).
 - [10] J. Charles, A. Le Yaouanc, L. Oliver, O. Pene and J. C. Raynal, Phys. Rev. D **60**, 014001 (1999) [hep-ph/9812358].
 - [11] C. W. Bauer, S. Fleming, D. Pirjol, I. W. Stewart, Phys. Rev. **D63**, 114020 (2001) [hep-ph/0011336].
 - [12] C. W. Bauer, D. Pirjol, I. W. Stewart, Phys. Rev. Lett. **87**, 201806 (2001) [hep-ph/0107002].
 - [13] C. W. Bauer, D. Pirjol and I. W. Stewart, Phys. Rev. D **65**, 054022 (2002) [hep-ph/0109045].
 - [14] M. Beneke, A. P. Chapovsky, M. Diehl, T. Feldmann, Nucl. Phys. **B643**, 431-476 (2002) [arXiv:hep-ph/0206152 [hep-ph]].
 - [15] C. W. Bauer, D. Pirjol and I. W. Stewart, Phys. Rev. D **67**, 071502 (2003) [hep-ph/0211069].

- [16] R. J. Hill and M. Neubert, Nucl. Phys. B **657**, 229 (2003) [hep-ph/0211018].
- [17] M. Beneke and D. Yang, Nucl. Phys. B **736**, 34 (2006) [hep-ph/0508250].
- [18] M. Beneke and T. Feldmann, Nucl. Phys. B **592**, 3 (2001) [hep-ph/0008255].
- [19] M. Beneke, T. Feldmann, Nucl. Phys. **B685**, 249-296 (2004) [hep-ph/0311335].
- [20] M. Beneke, Y. Kiyo and D. s. Yang, Nucl. Phys. B **692**, 232 (2004) [hep-ph/0402241].
- [21] T. Becher and R. J. Hill, JHEP **0410**, 055 (2004) [hep-ph/0408344].
- [22] P. Ball, V. M. Braun and E. Gardi, Phys. Lett. B **665**, 197 (2008) [arXiv:0804.2424 [hep-ph]].
- [23] Y. -L. Liu and M. -Q. Huang, Nucl. Phys. A **821**, 80 (2009) [arXiv:0811.1812 [hep-ph]].
- [24] S. Groote, J. G. Korner and O. I. Yakovlev, Phys. Rev. D **56**, 3943 (1997) [hep-ph/9705447].