arXiv:1112.6017v2 [math.DS] 13 Mar 2012

ENTROPY AND EXACT DEVANEY CHAOS ON TOTALLY REGULAR CONTINUA

VLADIMÍR ŠPITALSKÝ

Abstract. We study topological entropy of exactly Devaney chaotic maps on totally regular continua, i.e. on (topologically) rectifiable curves. After introducing the so-called P-Lipschitz maps (where P is a finite invariant set) we give an upper bound for their topological entropy. We prove that if a non-degenerate totally regular continuum X contains a free arc which does not disconnect X or if X contains arbitrarily large generalized stars then X admits an exactly Devaney chaotic map with arbitrarily small entropy. A possible application for further study of the best lower bounds of topological entropies of transitive/Devaney chaotic maps is indicated.

1. Introduction

A *(discrete) dynamical system* is a pair (X, f) where X is a compact metric space and $f : X \to X$ is a continuous map. For $n \in \mathbb{N}$ we denote the composition $f \circ f \circ \cdots \circ f$ (n-times) by f^n. A point $x \in X$ is a *periodic point* of f if $f^n(x) = x$ for some $n \in \mathbb{N}$. We say that a dynamical system (X, f) is *(topologically) transitive* if for every nonempty open sets $U, V \subseteq X$ there is $n \in \mathbb{N}$ such that $f^n(U) \cap V \neq \emptyset$. If every iterate f^n ($n \geq 1$) is transitive we say that f is *totally transitive*. If $(X \times X, f \times f)$ is transitive then (X, f) is called *(topologically) weakly mixing*. A system (X, f) is *(topologically) mixing* provided for every nonempty open sets $U, V \subseteq X$ there is $n_0 \in \mathbb{N}$ such that $f^n(U) \cap V \neq \emptyset$ for every $n \geq n_0$. Further, (X, f) is *(topologically) exact* or *locally eventually onto* if for every nonempty open subset U of X there is $n \in \mathbb{N}$ such that $f^n(U) = X$. Finally, a system (X, f) is *Devaney chaotic* (*totally Devaney chaotic*, *exactly Devaney chaotic*) provided X is infinite, f is transitive (totally transitive, exact, respectively) and has dense set of periodic points.

One of the main quantitative characteristics of a dynamical system (X, f) is the *topological entropy*, denoted by $h(f)$. A challenging topic in discrete dynamics is the study of relationships between entropy and qualitative properties of dynamical systems, such as transitivity, exactness and (exact) Devaney chaos. This goes back to 1982 when Blokh in [8] showed that every transitive map on the unit interval $[0, 1]$ has entropy at least $(1/2) \log 2$ and that this is the best possible

2010 *Mathematics Subject Classification.* Primary 37B05, 37B20, 37B40; Secondary 54H20.

Key words and phrases. Exact Devaney chaos, topological entropy, Lipschitz map, totally regular continuum, continuum of finite length, rectifiable curve.

The author wishes to express his thanks to Ľubomír Snoha for his help with the preparation of the paper. The author was supported by the Slovak Research and Development Agency under the contract No. APVV-0134-10 and by the Slovak Grant Agency under the grant number VEGA 1/0978/11.

bound, i.e. $I^{\mathrm{T}}([0,1]) = (1/2)\log 2$, where

$$I^{\mathrm{T}}(X) = \inf\{h(f) : \quad f : X \to X \text{ is transitive}\}.$$

Moreover, this infimum is in fact the minimum, i.e. the infimum is attainable.

Instead of $I^{\mathrm{T}}(X)$ one can study infima/minima of entropies of more restrictive classes of dynamical systems on X. Denote by $I^{\mathrm{D}}(X)$ and $I^{\mathrm{ED}}(X)$ the infima of entropies of Devaney chaotic and exactly Devaney chaotic systems on X, respectively. We see at once that $I^{\mathrm{T}}(X) \le I^{\mathrm{D}}(X) \le I^{\mathrm{ED}}(X)$. For (connected) *graphs*, these infima and the existence of corresponding minima were studied e.g. in [10, 2, 1, 4, 3, 30, 5, 27, 22, 18]; see Table 1 for some of the results. (For the definition of classes $\mathcal{P}(i)$ of trees see [30].) Every transitive graph map which is not conjugate to an irrational rotation of the circle has dense periodic points [9], so $I^{\mathrm{D}}(X) = I^{\mathrm{T}}(X)$ whenever X is a graph. Notice also that any exact map on a non-degenerate space X has positive entropy, see e.g. [22]; hence the infimum is never attained if $I^{\mathrm{ED}}(X) = 0$.

The space X	$I^{\mathrm{T}}(X)$ / $I^{\mathrm{D}}(X)$	Attainable?	$I^{\mathrm{ED}}(X)$	Att.?
interval	$\frac{1}{2}\log 2$	yes	$\frac{1}{2}\log 2$	no
circle	0	yes/no	0	no
n-star	$\frac{1}{n}\log 2$	yes	$\frac{1}{n}\log 2$	?
tree with n ends	$\ge \frac{1}{n}\log 2$	depends on X	$\ge \frac{1}{n}\log 2$	?
tree $\in \mathcal{P}(i)$, n ends	$\le \frac{1}{n-i}\log 2$	depends on X	$\le \frac{1}{n-i}\log 2$	?
graph $\neq$ tree, circle	0	no	0	no

TABLE 1. Infima of entropies on graphs

In the present paper we study $I^{\mathrm{ED}}(X)$ for totally regular continua X. Recall that a *continuum* is a connected compact metric space. A continuum X is *totally regular* [26] if for every point $x \in X$ and every countable set $P \subseteq X$ there is a basis of neighborhoods of x with finite boundary not intersecting P. Equivalently one can define totally regular continua as those continua X which admit a compatible metric d such that (X, d) is a *rectifiable curve* (i.e. (X, d) has finite one-dimensional Hausdorff measure $\mathcal{H}^1_d(X)$). Every *graph* and even every *local dendrite* (a locally connected continuum containing only finitely many simple closed curves) is totally regular. As another example one can mention the Hawaiian earrings (an infinite wedge of circles), which is a totally regular continuum but not a local dendrite.

Recall that an *arc* is a homeomorphic image of $[0,1]$. A *free arc* in X is an arc $A \subseteq X$ such that every non-end point of A is an interior point of A (in the topology of X). We say that a free arc $A \subseteq X$ *does not disconnect* X if no non-end point of A disconnects X; for locally connected continua (hence also for totally regular ones) this is equivalent to the fact that A is a subset of a simple closed curve in X, see e.g. [29, IV.9.3]. We say that X *contains arbitrarily large generalized stars* if for any $k \in \mathbb{N}$ there are a point $a \in X$ and k components of $X \setminus \{a\}$ such that the closures of the components are homeomorphic relative to a (i.e. the corresponding homeomorphisms fix the point a).

In [28] we introduced the so-called length-expanding Lipschitz maps (briefly LEL-maps) and we showed that for every two non-degenerate totally regular continua X, Y there exists a LEL-map $f : X \to Y$; see Section 2.6 for details. LEL-maps form "building blocks" in our constructions of exactly Devaney chaotic systems with small entropy. Our main results are summarized in the following theorem.

Main Theorem. *For every non-degenerate totally regular continuum X it holds that $I^{\mathrm{ED}}(X) < \infty$. Moreover, if X contains arbitrarily large generalized stars or X contains a free arc which does not disconnect X, then X admits*

- *a Devaney chaotic map which is not totally Devaney chaotic, and*
- *an exactly Devaney chaotic map,*

both with arbitrarily small positive entropy; hence $I^{\mathrm{ED}}(X) = 0$.

Thus, in particular, $I^{\mathrm{ED}}(X)$ and hence also $I^{\mathrm{D}}(X)$ and $I^{\mathrm{T}}(X)$ are zero for many dendrites, e.g. for the ω-star or the Wazewski's universal dendrite. Still, the following general question is open.

Question (Baldwin [5])**.** *Is it true that $I^{\mathrm{T}}(X) = 0$ for every dendrite X which is not a tree?*

Our result gives $I^{\mathrm{ED}}(X) = 0$ also for many spaces X which are not dendrites, say for the Hawaiian earrings. Since every graph which is not a tree contains a free non-disconnecting arc, Main Theorem is a generalization of [4, Theorems 3.7 and 4.1], cf. the last row of Table 1. Let us note that $I^{\mathrm{ED}}(X) = 0$ is true also for other subclasses of the class of totally regular continua, see e.g. Lemma 18 and Remark 19. We even believe that the following conjecture is true.

Conjecture. *$I^{\mathrm{ED}}(X) = 0$ for every totally regular continuum X which is not a tree.*

A potential application of our results lies in the fact that the estimates of $I^{\mathrm{T}}(X)$, $I^{\mathrm{D}}(X)$ and $I^{\mathrm{ED}}(X)$ for one-dimensional continua can be used to estimate the corresponding quantities for some higher-dimensional continua. To do that, one needs two tools. The first one is an appropriate extension theorem. For instance, in [2] and [20] it was proved that a transitive map $f : X \to X$ on a compact metric space X without isolated points can be extended to a transitive map F on $X \times [0, 1]$ without increasing the entropy. An analogous result is true for Devaney chaotic maps [6]. The second tool are procedures enabling to go from $X \times [0, 1]$ to the cone or the suspension over X, or to the space $X \times \mathbb{S}^1$, similarly as in [6]. However, it is not the topic of the present paper to continue in this direction.

Finally let us notice that the study of infima of entropies of transitive/Devaney chaotic maps on a given space is naturally connected with the question whether the space admits a *zero entropy* transitive/Devaney chaotic map. Using techniques developed in [6] one can easily construct a Devaney chaotic map on the Cantor fan (the cone over the Cantor set) which has zero entropy. On the other hand, if X is a compact metric space containing a free arc and X is not a union of finitely many disjoint circles, then every transitive map on X has positive entropy [14]. However, the following question is still open.

Question (Baldwin [5])**.** *Is there a dendrite admitting a transitive map with zero entropy?*

The paper is organized as follows. In Section 2 we recall all the needed definitions and facts. In Section 3 we introduce the so-called P-Lipschitz maps and we give an upper bound for their topological entropy. Finally, in Section 4 we prove Main Theorem.

2. Preliminaries

Here we briefly recall all the notions and results which will be needed in the rest of the paper. The terminology is taken mainly from [21, 25, 23, 15].

If M is a set, its cardinality is denoted by $\#M$. The cardinality of infinite countable sets is denoted by $\aleph_0$. If M is a singleton set we often identify it with its only point. We write $\mathbb{N}$ for the set of positive integers $\{1, 2, 3, \dots\}$, $\mathbb{R}$ for the set of reals and I for the unit interval $[0, 1]$. By an interval we mean any nonempty connected subset of $\mathbb{R}$ (possibly degenerate to a point). For intervals J, J' we write $J \le J'$ if $t \le s$ for every $t \in J$, $s \in J'$.

By a *space* we mean any nonempty metric space. A space is called *degenerate* provided it has only one point; otherwise it is called *non-degenerate*. If E is a subset of a space $X = (X, d)$ we denote the closure, the interior and the boundary of E by $\overline{E}$, $\operatorname{int}(E)$ and ∂E, respectively, and we write $d(E)$ for the diameter of E. We say that two sets $E, F \subseteq X$ are *non-overlapping* if they have disjoint interiors. For $x \in X$ and $r > 0$ we denote the closed ball with the center x and radius r by $B(x, r)$.

Let (X, f) be a dynamical system. A set $A \subseteq X$ is called *f-invariant* if $f(A) \subseteq A$. For the definitions of (total) transitivity, (weak, strong) mixing, exactness and (exact) Devaney chaos see Section 1.

2.1. Minkowski dimension and topological entropy. A subset S of a metric space (X, d) is called *ε-separated* if $d(x, y) > \varepsilon$ for every points $x \neq y$ from S. A subset R of X is said to *ε-span* X if for every $x \in X$ there is $y \in R$ with $d(x, y) \le \varepsilon$. We will denote by $s(X, \varepsilon)$ and $r(X, \varepsilon)$ the maximal cardinality of ε-separated set and the minimal cardinality of ε-spanning set, respectively. We see at once that $r(X, \varepsilon) \le s(X, \varepsilon) \le r(X, \varepsilon/2)$. The *upper* and *lower Minkowski (ball) dimensions* of X are defined by (see e.g [24, 5.3] or [19, Definition 3.2.7])

$$\overline{\dim}(X) = \limsup_{\varepsilon \to 0} \frac{\log r(X, \varepsilon)}{-\log \varepsilon} \qquad \text{and} \qquad \underline{\dim}(X) = \liminf_{\varepsilon \to 0} \frac{\log r(X, \varepsilon)}{-\log \varepsilon}\,.$$

Let (X, f) be a dynamical system. For $n \in \mathbb{N}$ define a metric d_n on X by $d_n(x, y) = \max_{0 \le i < n} d(f^i(x), f^i(y))$. We say that a set $S \subseteq X$ is *(n, ε)-separated* if it is an ε-separated subset of (X, d_n), i.e. if for every distinct $x, y \in S$ there is $i < n$ with $d(f^i(x), f^i(y)) > \varepsilon$. A set $R \subseteq X$ is said to *(n, ε)-span* X if it ε-spans (X, d_n). Let $s_n(X, \varepsilon, f)$ and $r_n(X, \varepsilon, f)$ denote the maximal cardinality of (n, ε)-separated subsets and the minimal cardinality of (n, ε)-spanning subsets of X, respectively. Then, by [11] and [13], the *topological entropy* $h(f)$ of the system (X, f) is given by

$$h(f) = \lim_{\varepsilon \to 0} \limsup_{n \to \infty} \frac{1}{n} \log r_n(X, \varepsilon, f) = \lim_{\varepsilon \to 0} \limsup_{n \to \infty} \frac{1}{n} \log s_n(X, \varepsilon, f)\,. \tag{2.1}$$

2.2. Continua. A *continuum* is a connected compact metric space. A *cut point* (or a *separating point*) of a continuum X is any point $x \in X$ such that $X \setminus \{x\}$ is disconnected. A point x of a continuum X is called a *local separating point* of X if there is a connected neighborhood U of x such that $U \setminus \{x\}$ is not connected.

If a, b are points of X then any cut point of X such that a, b belong to different components of $X \setminus \{x\}$ is said to *separate* a, b. The set of all such points is denoted by $\mathrm{Cut}(a, b)$ or $\mathrm{Cut}_X(a, b)$. If $a = b$ then obviously $\mathrm{Cut}(a, b) = \emptyset$.

Let X be a continuum. A metric d on X is said to be *convex* provided for every distinct $x, y \in X$ there is $z \in X$ such that $d(x, z) = d(z, y) = d(x, y)/2$. By [7, Theorem 8] every locally connected continuum admits a compatible convex metric.

If X is a continuum, we introduce the following notion. A *splitting* of X is any system $\mathcal{A} = \{X_1, \dots, X_n\}$ of non-degenerate subcontinua covering X such that $P_{\mathcal{A}} = \bigcup_{i \neq j} X_i \cap X_j$ is finite. In a splitting $\mathcal{A}$, every $X_i \in \mathcal{A}$ is *regular closed* (i.e. X_i is closed and $\mathrm{int}(X_i)$ is dense in it), $\mathrm{int}(X_i) = X_i \setminus \left(\bigcup_{j \neq i} X_j\right) \supseteq X_i \setminus P_{\mathcal{A}}$ and $\partial X_i \subseteq P_{\mathcal{A}}$.

Lemma 1. *Let X be a continuum and $\mathcal{A} = \{X_1, \dots, X_n\}$ be a splitting of X. For $i = 1, \dots, n$ let d_i be a convex metric on X_i satisfying $d_i(x, y) = d_j(x, y)$ whenever $i \neq j$ and $x, y \in X_i \cap X_j$. For $x, y \in X$ put*

$$d(x, y) = \inf \left\{ \sum_{k=1}^{m} d_{i_k}(x_{k-1}, x_k) : \ m \geq 1, \ x_0 = x, \ x_m = y \right. \\ \left. \text{and } x_{k-1}, x_k \in X_{i_k} \text{ for } k = 1, \dots, m \right\}. \tag{2.2}$$

Then d is a convex metric on X equivalent with the original one and the infimum in (2.2) is in fact the minimum. Moreover, if $d(x, y) = d_i(x, y)$ whenever $x, y \in P_{\mathcal{A}} \cap X_i$, then

$$d|_{X_i \times X_i} = d_i.$$

Proof. Using convexity of every d_i we may assume that, in the infimum from (2.2), $i_k \neq i_{k+1}$, $x_k \in P_{\mathcal{A}}$ for $1 \leq k < m$ and $x_k \neq x_l$ for every $k \neq l$. Since $P_{\mathcal{A}}$ is finite we have that, for any fixed x, y, the infimum is in fact the minimum and there are $m \geq 1$, $i_1, \dots, i_m$, $x_0 = x, x_1, \dots, x_m = y$ such that $x_0, \dots, x_m$ are pairwise distinct, $x_0 \in X_{i_1}$, $x_m \in X_{i_m}$,

$$d(x, y) = \sum_{k=1}^{m} d_{i_k}(x_{k-1}, x_k) \quad \text{and} \quad x_k \in X_{i_k} \cap X_{i_{k+1}} \cap P_{\mathcal{A}} \quad \text{for } 1 \leq k < m. \tag{2.3}$$

The fact that d is a convex pseudometric is straightforward and, by (2.3), $d(x, y) = 0$ implies $x = y$. To show that d is equivalent with the original metric, fix a sequence $(x_k)_k$ and a point x; since X is compact we only need to prove that $x_k \to x$ implies $d(x_k, x) \to 0$. It is no loss of generality in assuming that, for some i, $x_k \in X_i$ for every k. Since $x_k \to x$, also $x \in X_i$ and $d(x_k, x) \leq d_i(x_k, x) \to 0$.

Now fix i and assume that $d(x, y) = d_i(x, y)$ for every $x, y \in P_{\mathcal{A}} \cap X_i$. Take arbitrary $x, y \in X_i$. Then $d(x, y) \leq d_i(x, y)$ by (2.2). To show the opposite inequality take $i_1, \dots, i_m$ and $x_0 = x, x_1, \dots, x_m = y$ satisfying (2.3); we can assume that $i_1 = i_m = i$ (so possibly $x_0 = x_1$ or $x_{m-1} = x_m$). Since $x_1, x_{m-1} \in P_{\mathcal{A}} \cap X_i$, we get $d(x_1, x_{m-1}) = d_i(x_1, x_{m-1})$ and

$$d_i(x, y) \leq d_i(x, x_1) + d_i(x_1, x_{m-1}) + d_i(x_{m-1}, y) \leq \sum_{k=1}^{m} d_{i_k}(x_{k-1}, x_k) = d(x, y)\,.$$

Thus $d(x, y) = d_i(x, y)$ for every $x, y \in X_i$. □

2.3. Hausdorff one-dimensional measure. For a Borel subset B of a metric space (X,d) the *one-dimensional Hausdorff measure* of B is defined by

$$\mathcal{H}^1_d(B)=\lim_{\delta\to 0}\mathcal{H}^1_{d,\delta}(B),\quad \mathcal{H}^1_{d,\delta}(B)=\inf\left\{\sum_{i=1}^{\infty}d(E_i):\ B\subseteq\bigcup_{i=1}^{\infty}E_i,\ d(E_i)<\delta\right\}.$$

We say that (X,d) has *finite length* if $\mathcal{H}^1_d(X)<\infty$. By e.g. [17, Proposition 4A],

$$\mathcal{H}^1_d(C)\geq d(C)\qquad\text{whenever } C \text{ is a connected Borel subset of } X. \tag{2.4}$$

If $A\subseteq X$ is an arc then $\mathcal{H}^1_d(A)$ is equal to the length of A [15, Lemma 3.2]. In the case when (X,d) is the Euclidean real line $\mathbb{R}$ and $J\subset\mathbb{R}$ is an interval, $\mathcal{H}^1_d(J)$ is equal to the length of J and we denote it simply by $|J|$.

If (X,d) is a continuum of finite length endowed with a convex metric d, then it has the so-called *geodesic property* (see e.g. [17, Corollary 4E]): for every distinct $x,y\in X$ there is an arc A with end points x,y such that $d(x,y)=\mathcal{H}^1_d(A)$; any such arc A is called a *geodesic arc* or shortly a *geodesic*. Every subarc of a geodesic is again a geodesic. If x,y are the end points of a geodesic A and $z\in A$ then $d(x,y)=d(x,z)+d(z,y)$.

2.4. Lipschitz maps. A map $f:(X,d)\to(Y,\varrho)$ between metric spaces is called *Lipschitz* with a Lipschitz constant $L\geq 0$, shortly *Lipschitz-L*, provided $\varrho(f(x),f(x'))\leq L\cdot d(x,x')$ for every $x,x'\in X$; the smallest such L is denoted by $\operatorname{Lip}(f)$ and is called the *Lipschitz constant* of f. If $f:X\to Y$ is Lipschitz-L then $\mathcal{H}^1_\varrho(f(B))\leq L\cdot\mathcal{H}^1_d(B)$ for every Borel set $B\subset X$ such that $f(B)$ is Borel-measurable [15, p. 10]. We will use the following simple fact.

Lemma 2. *Let X be a continuum with a convex metric d and let $\mathcal{A}=\{X_1,\dots,X_n\}$ be a splitting of X. Let (Y,ϱ) be a metric space. Then any map $f:(X,d)\to(Y,\varrho)$ with Lipschitz-L restrictions $f|_{X_i}$ $(i=1,\dots,n)$ is Lipschitz-L.*

Proof. By Lemma 1, for any $x,y\in X$ there are $x_0=x,x_1,\dots,x_{m-1},x_m=y$ such that $x_{k-1},x_k\in X_{i_k}$ $(k=1,\dots,m)$ and $d(x,y)=\sum_{k=1}^m d(x_{k-1},x_k)$. So

$$\varrho(f(x),f(y))\leq\sum_{k=1}^m\varrho(f(x_{k-1}),f(x_k))\leq L\cdot\sum_{k=1}^m d(x_{k-1},x_k)=L\cdot d(x,y).$$

□

2.5. Totally regular continua. By e.g. [21, 26], a continuum X is called

- a *dendrite* if it is locally connected and contains no simple closed curve;
- a *local dendrite* if it is locally connected and contains at most finitely many simple closed curves;
- *completely regular* if it contains no non-degenerate nowhere dense subcontinuum;
- *totally regular* if for every $x\in X$ and every countable set $P\subseteq X$ there is a basis of neighborhoods of x with finite boundary not intersecting P;
- *regular* if every $x\in X$ has a basis of neighborhoods with finite boundary;
- *hereditarily locally connected* if every subcontinuum of X is locally connected;
- a *curve* if it is one-dimensional.

Notice that (local) dendrites as well as completely regular continua are totally regular and (totally) regular continua are hereditarily locally connected, hence they are locally connected curves.

We will need the following simple estimate of the maximal cardinality of ε-separated sets in totally regular continua X. Notice that an analogous estimate follows from the fact that the 1-dimensional Minkowski content is equal to $\mathcal{H}^1(X)$ for continua $X \subseteq \mathbb{R}^n$ of finite length ([16, Theorem 3.2.39], see also [24, p. 75]).

Lemma 3. *Let $X = (X, d)$ be a non-degenerate totally regular continuum. Then*

$$\underline{\dim}(X) = \overline{\dim}(X) = 1 \quad \textit{and} \quad s(X,\varepsilon) \le \frac{2 \cdot \mathcal{H}^1(X)}{\varepsilon} \quad \textit{for every } 0 < \varepsilon < d(X)\,.$$

Proof. Let us first prove the inequality for $s(X, \varepsilon)$. There is no loss of generality in assuming $\mathcal{H}^1(X) < \infty$. By (2.4),

$$\mathcal{H}^1(B(x,r)) \ge r \qquad \text{for every } x \in X \text{ and } 0 < r < d(X)/2\,. \tag{2.5}$$

Take $0 < \varepsilon < d(X)$ and an ε-separated set $S = \{x_1, \dots, x_k\}$ of maximal cardinality $k = s(X, \varepsilon)$. Since $d(x_i, x_j) > \varepsilon$ for every $i \ne j$, the closed balls $B(x_i, \varepsilon/2)$ are disjoint. Applying (2.5) we conclude that $\mathcal{H}^1(X) \ge \sum_i \mathcal{H}^1(B(x_i, \varepsilon/2)) \ge k\varepsilon/2$, so $s(X, \varepsilon) \le 2\mathcal{H}^1(X)/\varepsilon$.

The equality $\underline{\dim}(X) = \overline{\dim}(X) = 1$ is a special case of [16, Theorem 3.2.39]. (Notice that $\overline{\dim}(X) \le 1$ immediately follows from the upper estimate for $s(X, \varepsilon)$.) □

The following lemma will be used in the proof of Lemma 15.

Lemma 4. *If X is a totally regular continuum which is not a dendrite, then there are points $a, b \in X$ and subcontinua X_0, X_1 of X such that $X_0 \cup X_1 = X$, $X_0 \cap X_1 = \{a, b\}$ and $\mathrm{Cut}_{X_i}(a, b)$ is uncountable for $i = 0, 1$.*

Proof. By [12] we can write X as the inverse limit $\varprojlim(X_n, f_n)$, where X_n are graphs and $f_n : X_{n+1} \to X_n$ are monotone; let $\pi_n : X \to X_n$ $(n \in \mathbb{N})$ be the natural projections. Since X is not a dendrite, by [25, Theorem 10.36] there is m such that X_m contains a circle S. Let C be the set of all points $x_m \in S$ which are not vertices of X_m and are such that $\pi_m^{-1}(x_m)$ are singletons. Then $S \setminus C$ is countable since every system of disjoint non-degenerate subcontinua of X is countable. Take $a_m \ne b_m$ from C and put $a = \pi_m^{-1}(a_m)$, $b = \pi_m^{-1}(b_m)$. Write X_m as the union $X_{m,0} \cup X_{m,1}$ of two non-degenerate subgraphs with $X_{m,0} \cap X_{m,1} = \{a_m, b_m\}$. Then $X_i = \pi_m^{-1}(X_{m,i})$ $(i = 0, 1)$, being inverse limits of continua [23, Proposition 2.1.17], are subcontinua of X covering X and $X_0 \cap X_1 = \pi_m^{-1}(X_{m,0} \cap X_{m,1}) = \{a, b\}$. □

2.6. Length-expanding Lipschitz maps on totally regular continua. Here we recall the main results of [28]. We say that a family $\mathcal{C}$ of non-degenerate subcontinua of X is *dense* if every nonempty open set in X contains a member of $\mathcal{C}$. By $\mathcal{C}_I$ we denote the system of all non-degenerate closed subintervals of I; the Euclidean metric on I is denoted by d_I.

Definition 5 ([28])**.** Let $X = (X, d)$, $X' = (X', d')$ be non-degenerate (totally regular) continua of finite length and let $\mathcal{C}, \mathcal{C}'$ be dense systems of subcontinua of X, X', respectively. We say that a continuous map $f : X \to X'$ is *length-expanding*

with respect to $\mathcal{C}, \mathcal{C}'$ if there exists $\varrho > 1$ (called *length-expansivity constant* of f) such that, for every $C \in \mathcal{C}$, $f(C) \in \mathcal{C}'$ and

$$\text{if} \quad f(C) \neq X' \qquad \text{then} \quad \mathcal{H}^1_{d'}(f(C)) \geq \varrho \cdot \mathcal{H}^1_d(C). \tag{2.6}$$

Moreover, if f is surjective and Lipschitz-L we say that $f : (X, d, \mathcal{C}) \to (X', d', \mathcal{C}')$ is (ϱ, L)-*length-expanding Lipschitz*. Sometimes we briefly say that f is (ϱ, L)-*LEL* or only *LEL*.

Proposition 6 ([28])**.** *Let $f : (X, d, \mathcal{C}) \to (X, d, \mathcal{C})$ be a LEL map. Then f is exact and has finite positive entropy. Moreover, if f is the composition $\varphi \circ \psi$ of some maps $\psi : X \to I$ and $\varphi : I \to X$, then f has the specification property and so it is exactly Devaney chaotic.*

For $k \geq 1$ let $f_k : I \to I$ be the piecewise linear continuous map which fixes 0 and maps every $[(i-1)/k, i/k]$ onto I.

Proposition 7 ([28])**.** *For every non-degenerate totally regular continuum X and every $a, b \in X$ we can find a convex metric $d = d_{X,a,b}$ on X and Lipschitz surjections $\varphi_{X,a,b} : I \to X$, $\psi_{X,a,b} : X \to I$ with the following properties:*

(a) $\mathcal{H}^1_d(X) = 1$*;*
(b) *the system $\mathcal{C} = \mathcal{C}_{X,a,b} = \{\varphi_{X,a,b}(J) : \ J \text{ is a closed subinterval of } I\}$ is a dense system of subcontinua of X;*
(c) *for every $\varrho > 1$ there are a constant L_ϱ (depending only on ϱ) and (ϱ, L_ϱ)-LEL maps*

$$\varphi : (I, d_I, \mathcal{C}_I) \to (X, d, \mathcal{C}) \quad \text{and} \quad \psi : (X, d, \mathcal{C}) \to (I, d_I, \mathcal{C}_I)$$

with $\varphi(0) = a$, $\varphi(1) = b$, $\psi(a) = 0$ and such that $\varphi = \varphi_{X,a,b} \circ f_k$, $\psi = f_l \circ \psi_{X,a,b}$ for some $k, l \geq 3$.

Moreover, if $\mathrm{Cut}_X(a, b)$ is uncountable, d, φ, ψ can be assumed to satisfy

(d) *$d(a, b) > 1/2$ and $\psi(b) = 1$.*

Proposition 8 ([28])**.** *Keeping the notation from Proposition 7, for every $\varrho > 1$, every non-degenerate totally regular continua X, X' and every points $a, b \in X$, $a', b' \in X'$ there are a constant L_ϱ (depending only on ϱ) and (ϱ, L_ϱ)-LEL map*

$$f : (X, d_{X,a,b}, \mathcal{C}_{X,a,b}) \to (X', d_{X',a',b'}, \mathcal{C}_{X',a',b'})$$

with $f(a) = a'$ and, provided $\mathrm{Cut}_X(a, b)$ is uncountable, $f(b) = b'$. Moreover, f is equal to the composition $\varphi \circ \psi$ of two LEL-maps $\psi : X \to I$ and $\varphi : I \to X'$.

We will need Proposition 9, which is a simple generalization of Proposition 8. Before giving the formulation of it we need to introduce the following notation. If $p \geq 1$, X_i $(i = 1, \dots, p)$ are non-degenerate subcontinua of X and a_i $(i = 0, \dots, p)$ are points of X such that $a_0 \in X_1, a_p \in X_p$, $a_i \in X_i \cap X_{i+1}$ for $1 \leq i < p$ and $\mathcal{C}_i = \mathcal{C}_{X_i, a_{i-1}, a_i}$ for $1 \leq i \leq p$, we write

$$(X, d, \mathcal{C}) = \coprod_{i=1}^{p} (X_i, d_i, \mathcal{C}_i)$$

if $X = \bigcup_i X_i$, d is a convex metric on X such that $d|_{X_i \times X_i} = d_{X_i, a_{i-1}, a_i}$ for $i = 1, \dots, p$ and $\mathcal{C}$ is the system of all unions $C_i \cup X_{i+1} \cup \dots \cup X_{j-1} \cup C_j$ where $i \leq j$, $C_i \in \mathcal{C}_i$, $C_j \in \mathcal{C}_j$, and, if $i < j$, $C_i = \varphi_{X_i, a_{i-1}, a_i}([s, 1])$ and $C_j = \varphi_{X_j, a_{j-1}, a_j}([0, t])$ for some $s, t \in I$. Particularly, if $p = 1$ then $d = d_1$ and $\mathcal{C} = \mathcal{C}_{X, a_0, a_1}$.

Proposition 9. *Let X, X' be non-degenerate totally regular continua. Let $a, b \in X$ and $d = d_{X,a,b}$, $\mathcal{C} = \mathcal{C}_{X,a,b}$. Let $a'_0, a'_1, \dots, a'_p$ be points of X' such that*

$$(X', d', \mathcal{C}') = \prod_{i=1}^{p} (X'_i, d_{X'_i, a'_{i-1}, a'_i}, \mathcal{C}_{X'_i, a'_{i-1}, a'_i}).$$

Then for every $\varrho > 1$ there is a map $f : X \to X'$ such that:

(a1) *$f : (X, d) \to (X', d')$ is a Lipschitz-$\tilde{L}$ surjection, where $\tilde{L} = \tilde{L}_{p,\varrho}$ depends only on p and ϱ;*

(a2) *for every $C \in \mathcal{C}$ we have $f(C) \in \mathcal{C}'$ and*

$$\text{if} \quad f(C) \not\supseteq X'_i \quad \text{for every } i \qquad \text{then} \quad \mathcal{H}^1_{d'}(f(C)) \geq \rho \cdot \mathcal{H}^1_d(C);$$

(b) *f is the composition $\varphi \circ \psi$ of two Lipschitz-$\tilde{L}$ surjections $\psi : (X, d) \to I$ and $\varphi : I \to (X', d')$ such that $\psi(a) = 0$, $\varphi(0) = a'_0$, $\varphi(1) = a'_p$ and, provided $\mathrm{Cut}_X(a, b)$ is uncountable, $\psi(b) = 1$.*

Notice that, by (b), $f(a) = a'_0$ and, provided $\mathrm{Cut}_X(a, b)$ is uncountable, $f(b) = a'_p$;

Proof. We may assume that $\varrho \geq 2$. For $i = 1, \dots, p$ put $d'_i = d'_{X'_i, a'_{i-1}, a'_i}$ and $\mathcal{C}'_i = \mathcal{C}'_{X'_i, a'_{i-1}, a'_i}$. Let L_ϱ be as in Proposition 8 and let $\psi : (X, d, \mathcal{C}) \to (I, d_I, \mathcal{C}_I)$, $\varphi'_i = \varphi_{X'_i, a'_{i-1}, a'_i} \circ f_{k_i} : (I, d_I, \mathcal{C}_I) \to (X'_i, d'_i, \mathcal{C}'_i)$ $(i = 1, \dots, p)$ be (ϱ, L_ϱ)-LEL maps with $\varphi'_i(0) = a'_{i-1}$, $\varphi'_i(1) = a'_i$, $\psi(a) = 0$ and, provided $\mathrm{Cut}_X(a, b)$ is uncountable, also $\psi(b) = 1$.

For $i = 1, \dots, p$ let $I_i = [(i-1)/p,\, i/p\,]$ and $g_i : I_i \to I$ be the increasing linear surjection. Define $f : X \to X'$ by $f = \varphi' \circ \psi$, where $\varphi' : I \to X'$ is given by $\varphi'|_{I_i} = \varphi'_i \circ g_i$ for $i = 1, \dots, p$. Then $\varphi'(i/p) = a'_i$ for $0 \leq i \leq p$, φ' is continuous and Lipschitz with $\mathrm{Lip}(\varphi') \leq pL_\varrho$ by Lemma 2. Since $\mathrm{Lip}(f) \leq pL^2_\varrho$, f satisfies (a1) and (b) with $\tilde{L} = pL^2_\varrho$.

To show that f satisfies also (a2) fix any $C \in \mathcal{C}$ and put $L = \psi(C)$ and $L_i = L \cap I_i$ for $i = 1, \dots, p$; notice that L and nonempty L_i's are closed intervals. The fact that $f(C) = \varphi'(L) \in \mathcal{C}'$ immediately follows from the definitions of f and $\mathcal{C}'$. If L contains some I_i then $f(C) \supseteq X'_i$. Otherwise $|L| \geq \varrho \cdot \mathcal{H}^1_d(C)$ since $L \neq I$ and there is i such that $L = L_i \cup L_{i+1}$. We may assume that $|L_i| \geq |L|/2$. Then

$$\mathcal{H}^1_{d'}(f(C)) \geq \mathcal{H}^1_{d'}(\varphi'_i \circ g_i(L_i)) \geq \varrho p \cdot |L_i| \geq \frac{\varrho^2 p}{2} \cdot \mathcal{H}^1_d(C) \geq \varrho \cdot \mathcal{H}^1_d(C)$$

since $\varrho \geq 2$. So f satisfies also (a2) and the proof is finished. □

3. P-Lipschitz maps and an entropy bound

In this section we introduce a special class of maps called P-Lipschitz maps (do not confuse with Lipschitz-L maps) and we show a relatively easy way of obtaining an upper bound for their entropy (see Proposition 12), which will be used in Section 4. For the definition of a splitting see Section 2.2.

Definition 10 (P-Lipschitz maps)**.** Let X be a non-degenerate continuum and $f : X \to X$ be a continuous map. Let P be a finite f-invariant subset of X, $\mathcal{A}$ be a splitting of X with $P_\mathcal{A} \subseteq P$ and $(L_A)_{A \in \mathcal{A}}$ be positive constants. Then we say that f is a *P-Lipschitz map (w.r.t. the splitting $\mathcal{A}$ and the constants $(L_A)_{A \in \mathcal{A}}$)* if, for every $A \in \mathcal{A}$, $f(A)$ is non-degenerate and $\mathrm{Lip}(f|_A) \leq L_A$.

Roughly speaking, P-Lipschitz maps are "piecewise-Lipschitz" maps with the "pieces" being subcontinua intersecting only in a finite invariant subset P. Trivially, every non-constant Lipschitz map is P-Lipschitz with $P = \emptyset$ and $\mathcal{A} = \{X\}$. On the other hand, if the metric d of X is convex then every P-Lipschitz map is Lipschitz with $\mathrm{Lip}(f) \le \max_{A\in\mathcal{A}} L_A$ (see Lemma 2).

Let f be a P-Lipschitz map w.r.t. $\mathcal{A}$. For $A, B \in \mathcal{A}$ we write $A \to B$ or, more precisely, $A \xrightarrow{f} B$ provided $f(A)$ intersects the interior of B. So if $A \to B$ then $f(A)\cap B$ is a finite union of subcontinua of X at least one of which is non-degenerate; the number of components of $f(A) \cap B$ is smaller than or equal to the cardinality of ∂B. Since every $f(A)$ is non-degenerate, for every $A \in \mathcal{A}$ there is $B \in \mathcal{A}$ with $A \to B$ and, provided f is surjective, also $C \in \mathcal{A}$ with $C \to A$.

The *P-transition graph G_f of f* is the directed graph the vertices of which are the sets $A \in \mathcal{A}$ and the edges of which correspond to $A \to B$. The corresponding 01-transition matrix will be called the *P-transition matrix* and will be denoted by M_f. For an integer $n \ge 1$ put

$$\mathcal{A}^n = \{\mathbb{A} = (A_0, A_1, \dots, A_{n-1}) :\ A_i \in \mathcal{A},\ A_0 \to A_1 \to \cdots \to A_{n-1}\}.$$

So $\mathcal{A}^n$ is the set of all paths of length $n-1$ in G_f.

Lemma 11. *Let f be a P-Lipschitz map w.r.t. $\mathcal{A}$ and let $x \in X$. Then for every integer $n \ge 1$ there is $\mathbb{A} = (A_0, A_1, \dots, A_{n-1}) \in \mathcal{A}^n$ such that*

$$f^i(x) \in A_i \qquad \text{for every } i = 0, 1, \dots, n-1.$$

Proof. We prove the lemma by induction. If $n = 1$ then the assertion follows from the fact that $\mathcal{A} = \mathcal{A}^1$ covers X. Assume now that the assertion of the lemma is true for some $n \ge 1$; we are going to show that it is true also for $n+1$. To this end take any point $x \in X$; we want to find $\mathbb{A} \in \mathcal{A}^{n+1}$ with $f^i(x) \in A_i$ for $i \le n$. By the induction hypothesis there is $(A_0, A_1, \dots, A_{n-1}) \in \mathcal{A}^n$ such that $f^i(x) \in A_i$ for every $i \le n-1$. Put $y = f^{n-1}(x) \in A_{n-1}$. If $f(y) \notin P$ then $f(y) \in \mathrm{int}(A_n)$ for some $A_n \in \mathcal{A}$; hence $A_{n-1} \to A_n$ and we can put $\mathbb{A} = (A_0, A_1, \dots, A_{n-1}, A_n) \in \mathcal{A}^{n+1}$. Now assume that $f(y) \in P$. Let $B_1, \dots, B_k$ $(k \ge 1)$ be the collection of all sets from $\mathcal{A}$ containing $f(y)$. Since $f(A_{n-1})$ has no isolated point (it is a non-degenerate continuum) and contains $f(y)$, it must intersect the interior of some B_i. If we put $\mathbb{A} = (A_0, A_1, \dots, A_{n-1}, B_i)$, we are done. □

Let X be a continuum with a convex metric d and let f be a P-Lipschitz map on X w.r.t. $\mathcal{A}$ and $(L_A)_{A\in\mathcal{A}}$. For every nonempty subset $\mathcal{B}$ of $\mathcal{A}$ put

$$L_{\mathcal{B}} = \max_{A\in\mathcal{B}} L_A.$$

Since any P-Lipschitz map f has $\mathrm{Lip}(f) \le L_{\mathcal{A}}$, we immediately have (see e.g. [19, Theorem 3.2.9])

$$h(f) \le \overline{\dim}(X) \cdot \log^+ L_{\mathcal{A}}$$

where $\log^+ L = \max\{\log L, 0\}$.

Assume now that X is a non-degenerate totally regular continuum and the metric d of X is convex with $\mathcal{H}^1_d(X) < \infty$. Then $\overline{\dim}(X) = 1$ (see Lemma 3), hence $h(f) \le \log^+ L_{\mathcal{A}}$. But this upper bound is often too pessimistic. Consider e.g. the following example. Let f and $\mathcal{A} = \{X_0, \dots, X_{k-1}\}$ be such that $f(X_i) \subseteq X_{(i+1) \bmod k}$ and that $L_{X_0} = L > 1$, $L_{X_i} \le 1$ for $i \ge 1$. Then $L_{\mathcal{A}} = L$ and we have $h(f) \le \log L$. But since X_i's are f^k-invariant and $\mathrm{Lip}(f^k|_{X_i}) \le L$, we have a much better estimate

$h(f) \le (1/k)\log L$. The key point here is the fact that the sets $A \in \mathcal{A}$ with "large" L_A occurs "rarely" in paths $\mathbb{A} \in \mathcal{A}^n$ of f.

To formalize this idea, for nonempty $\mathcal{B} \subseteq \mathcal{A}$ put

$$\theta_{\mathcal{B}} = \limsup_{n\to\infty} \frac{k_n^{\mathcal{B}}}{n} \qquad \text{where} \quad k_n^{\mathcal{B}} = \max_{\mathbb{A}\in\mathcal{A}^n} \#\{k = 0,\dots,n-1 : A_k \notin \mathcal{B}\}\,. \tag{3.1}$$

This quantity measures the maximal "asymptotic frequency" of occurrences of $A \in \mathcal{A}\setminus\mathcal{B}$ in paths of f. The following proposition asserts that if $L_{\mathcal{B}}$ is close to 1 and $\theta_{\mathcal{B}}$ is sufficiently small, then the entropy of f can be small even for large $L_{\mathcal{A}}$.

Proposition 12 (Upper bound for the entropy of P-Lipschitz maps)**.** *Let X be a non-degenerate totally regular continuum endowed with a convex metric d such that $\mathcal{H}^1_d(X) < \infty$. Let $f : X \to X$ be P-Lipschitz w.r.t. $\mathcal{A}$ and $(L_A)_{A\in\mathcal{A}}$. Then, for every nonempty subsystem $\mathcal{B}$ of $\mathcal{A}$,*

$$h(f) \le \log^+ L_{\mathcal{B}} + 2\theta_{\mathcal{B}} \log^+ L_{\mathcal{A}}.$$

Proof. Since the assertion of the proposition does not change if we replace every L_A by $\max\{L_A, 1\}$, we may assume that $L_A \ge 1$ for every $A \in \mathcal{A}$. Fix any $0 < \varepsilon < d(X)$ such that $\varepsilon < \operatorname{dist}(A,B)$ for every disjoint $A, B \in \mathcal{A}$ and $\varepsilon < d(x,y)$ for every distinct $x, y \in P$.

Let $n \in \mathbb{N}$ and let S be an (n,ε)-separated subset of X of maximal cardinality $\#S = s_n(X,\varepsilon,f)$. Take any $x \ne y$ from S. By Lemma 11 there are $\mathbb{A}^x = (A_i^x)_{i<n}, \mathbb{A}^y = (A_i^y)_{i<n} \in \mathcal{A}^n$ such that $x_i = f^i(x) \in A_i^x$ and $y_i = f^i(y) \in A_i^y$ for $i < n$. Put $d_i = d(x_i, y_i)$ for $i < n$ and $j = \min\{i : d_i > \varepsilon\}$; then $0 \le j \le n-1$.

Fix any $0 \le i < j$. Let B be a geodesic arc from x_i to y_i. Then, by (2.4), $d(B) \le \mathcal{H}^1(B) = d(x_i, y_i) \le \varepsilon$. By the choice of ε the set B contains at most one point from P. If $B \cap P = \emptyset$ then B does not intersect the boundaries of A_i^x, A_i^y and so $B \subseteq \operatorname{int}(A_i^x) \cap \operatorname{int}(A_i^y)$. Hence $A_i^x = A_i^y$ and

$$d_{i+1} \le L_{A_i^x} \cdot d_i. \tag{3.2}$$

Otherwise $B \cap P = \{z\}$ is a singleton. We claim that $z \in A_i^x$. Indeed, if $z \ne x_i$ then x_i belongs to the interior of A_i^x. Since B is connected, intersects the interior of A_i^x and $\partial A_i^x \subseteq P$, the subarc $x_i z$ of B is contained in A_i^x and so $z \in A_i^x$. If $z = x_i$ then again $z \in A_i^x$. Analogously $z \in A_i^y$ and thus $z \in A_i^x \cap A_i^y$. Using the facts that $f|_A$ is Lipschitz on every $A \in \mathcal{A}$ and that B is geodesic (hence $d(x_i,z) + d(z,y_i) = d_i$) we obtain

$$\begin{aligned} d_{i+1} &\le d(x_{i+1}, f(z)) + d(f(z), y_{i+1}) \le L_{A_i^x} \cdot d(x_i, z) + L_{A_i^y} \cdot d(z, y_i) \\ &\le \max\{L_{A_i^x}, L_{A_i^y}\} \cdot d_i\,. \end{aligned} \tag{3.3}$$

Combination of (3.2) and (3.3) for $i < j$ gives

$$d_j \le d_0 \cdot \prod_{i<j} \max\{L_{A_i^x}, L_{A_i^y}\} \le d_0 \cdot \prod_{i<n} \max\{L_{A_i^x}, L_{A_i^y}\}\,.$$

Thus, by the definition (3.1) of $k_n = k_n^{\mathcal{B}}$,

$$\varepsilon < d_j \le d_0 \cdot L_{\mathcal{A}}^{2k_n} \cdot L_{\mathcal{B}}^{n-2k_n} \le d_0 \cdot L_{\mathcal{A}}^{2k_n} \cdot L_{\mathcal{B}}^{n}\,.$$

So the set S is ε'-separated with $\varepsilon' = \varepsilon/(L_{\mathcal{A}}^{2k_n} \cdot L_{\mathcal{B}}^n)$ and Lemma 3 gives

$$s_n(X,\varepsilon,f) \le s(X,\varepsilon') \le \frac{2\mathcal{H}^1(X)}{\varepsilon} \cdot L_{\mathcal{A}}^{2k_n} \cdot L_{\mathcal{B}}^n\,.$$

From this and (2.1), $h(f) \le 2\theta_{\mathcal{B}} \log L_{\mathcal{A}} + \log L_{\mathcal{B}}$. □

In the following example we show how Proposition 12 can be used to obtain a "good" upper estimate of the topological entropy.

Example 13. Consider a graph X which is not a tree. We can write $X = I \cup G$, where I is a free arc in X identified with $[0,1]$ and G is a subgraph of X such that $I \cap G = \{0,1\}$. Let $n \geq 3$ and $g = g_n : X \to X$ be the map constructed in the proof of [4, Theorem 4.1]. Recall that if we put $X_i = [i/n, (i+1)/n]$ for $0 \leq i < n$, $i \neq 1$ and $X_1^- = \left[\frac{1}{n}, \frac{3}{2n}\right]$, $X_1^+ = \left[\frac{3}{2n}, \frac{2}{n}\right]$, then g maps linearly X_0 onto $X_1^- \cup X_1^+$, X_1^- onto X_1^+, X_1^+ onto $X_1^+ \cup X_2$ and X_i onto X_{i+1} for $i = 2, \dots, n-2$. Moreover, $g|_{X_{n-1}} : X_{n-1} \to G$ and $g|_G : G \to X_0$ are piecewise linear maps with the number of pieces not depending on n.

Put $\lambda = \sqrt[n-2]{2}$ and define the convex metric $d = d_n$ on X in such a way that $d(0,1) = d\left(0, \frac{1}{n}\right) = 1$, $d\left(\frac{1}{n}, \frac{3}{2n}\right) = d\left(\frac{3}{2n}, \frac{2}{n}\right) = \frac{1}{2}$, $d\left(\frac{i}{n}, \frac{i+1}{n}\right) = \frac{1}{2}\lambda^{i-1}$ for $i = 2, \dots, n-1$, and the Lipschitz constants of $g|_{X_{n-1}}$, $g|_G$ are bounded from above by a constant L not depending on n. Then g is P-Lipschitz with $P = \left\{\frac{i}{n},\ i = 0, \dots, n\right\} \cup \left\{\frac{3}{2n}\right\}$, $\mathcal{A} = \{X_0, X_1^-, X_1^+, X_2, \dots, X_{n-1}, G\}$ and $L_{X_0} = L_{X_1^-} = 1$, $L_{X_1^+} = L_{X_2} = \dots = L_{X_{n-2}} = \lambda$, $L_{X_{n-1}} = L_G = L$. Put $\mathcal{B} = \mathcal{A} \setminus \{G, X_{n-1}\}$. Then $L_{\mathcal{A}} = L$, $L_{\mathcal{B}} = \lambda$ and an easy computation gives that $\theta_{\mathcal{B}} \leq \frac{2}{n}$. According to Proposition 12 we have

$$h(g_n) \leq \log \lambda + \frac{2}{n} \log L \leq \frac{\log(2L^2)}{n-2},$$

and thus $\limsup_n h(g_n) = 0$.

We finish this section with a simple lemma giving a sufficient condition for transitivity and exactness of P-Lipschitz maps. Recall that a 01-square matrix M is called *irreducible* if for every indices i, j there is $n > 0$ with $M_{ij}^n > 0$; i.e. the corresponding directed graph G_M is *strongly connected* (there is a path from each vertex to every other vertex). The *period* p_M of an irreducible matrix M is the greatest common divisor of those $n > 0$ for which $M_{ii}^n > 0$ for some i. Equivalently, p_M is the greatest common divisor of the lengths of cycles (closed paths) in G_M. An irreducible matrix M and the corresponding graph G_M are called *primitive* provided $p_M = 1$.

Lemma 14. *Let X be a non-degenerate totally regular continuum endowed with a convex metric d such that $\mathcal{H}_d^1(X) < \infty$. Let a map $f : X \to X$ be P-Lipschitz w.r.t. $\mathcal{A}$ such that $A \to B$ implies $f(A) \supseteq B$ for $A, B \in \mathcal{A}$. Assume that there is a dense system $\mathcal{D}$ of subcontinua of X such that for every $D \in \mathcal{D}$ there are $n \in \mathbb{N}$, $A \in \mathcal{A}$ with $f^n(D) \supseteq A$. Then the following hold:*

(a) *f is transitive but not totally transitive if and only if M_f is irreducible but not primitive;*
(b) *f is exact if and only if M_f is primitive.*

Proof. If M_f is not irreducible there are $A, B \in \mathcal{A}$ such that there is no path from A to B. Hence $f^n(\operatorname{int}(A)) \cap \operatorname{int}(B) = \emptyset$ for every $n > 0$, which contradicts transitivity of f. If M_f is irreducible with the period $p > 1$, then there are $A, B \in \mathcal{A}$ such that there is no path from A to B of length np. I.e. f^p is not transitive and thus f is not totally transitive.

Assume now that M_f is irreducible and take any nonempty open sets U, V. Since $\bigcup_{A \in \mathcal{A}} \operatorname{int}(A) \supseteq X \setminus P$ is dense in X, there is $B \in \mathcal{A}$ intersecting V. By the assumption on $\mathcal{D}$ there are $D \in \mathcal{D}$, $n \in \mathbb{N}$ and $A \in \mathcal{A}$ such that $D \subseteq U$ and $f^n(D) \supseteq A$. Since M_f is irreducible there is m with $f^m(A) \supseteq B$. Hence

$f^{n+m}(D) \supseteq B$ and so $f^{n+m}(U)$ intersects V. Thus f is transitive. Moreover, if M_f is primitive there is p such that, for every $C \in \mathcal{A}$, there is a path of length p from A to C. So $f^{n+p}(U) \supseteq f^{n+p}(D) \supseteq f^p(A) = X$ and f is exact. □

4. Proof of Main Theorem

In this section we prove Main Theorem. We use the notation from Section 2.6.

4.1. Lipschitz, exactly Devaney chaotic maps.

Lemma 15. *There is $L > 0$ such that any non-degenerate totally regular continuum X admits a convex metric d with $\mathcal{H}^1_d(X) < \infty$ and Lipschitz-L, Devaney chaotic maps $f, g : (X, d) \to (X, d)$ with positive entropies such that f is not totally transitive and g is exact.*

Proof. The existence of a map $g : (X, d) \to (X, d)$ was shown in [28], so we only need to construct f. If X contains a circle, there are points $a \neq b$ such that X is the union of two subcontinua X_0, X_1 with $X_0 \cap X_1 = \{a, b\}$ and $\mathrm{Cut}_{X_i}(a, b)$ is uncountable for $i = 0, 1$. (Such points a, b can be chosen as follows. By [12] we can write $X = \varprojlim(X_n, f_n)$, where X_n are graphs and $f_n : X_{n+1} \to X_n$ are monotone; let $\pi_n : X \to X_n$ $(n \in \mathbb{N})$ be the natural projections. Since X is not a dendrite, by [25, Theorem 10.36] there is m such that X_m contains a circle S. Let C be the set of all points $x_m \in S$ which are not vertices of X_m and are such that $\pi_m^{-1}(x_m)$ are singletons. Then $S \setminus C$ is countable since every system of disjoint non-degenerate subcontinua of X is countable. Take $a_m \neq b_m$ from C and put $a = \pi_m^{-1}(a_m)$, $b = \pi_m^{-1}(b_m)$. Write X_m as the union $X_{m,0} \cup X_{m,1}$ of two non-degenerate subgraphs with $X_{m,0} \cap X_{m,1} = \{a_m, b_m\}$. Then $X_i = \pi_m^{-1}(X_{m,i})$ $(i = 0, 1)$, being inverse limits of continua [23, Proposition 2.1.17], are non-degenerate subcontinua of X covering X and $X_0 \cap X_1 = \pi_m^{-1}(X_{m,0} \cap X_{m,1}) = \{a, b\}$.)

Without loss of generality we may assume that $d_{X_0,a,b}(a, b) = d_{X_1,a,b}(a, b)$ (otherwise we replace one of the metrics by a constant multiple of it). Let d be the convex metric on X defined by (2.2) for the splitting $\{X_0, X_1\}$ and metrics $d_0 = d_{X_0,a,b}$, $d_1 = d_{X_1,a,b}$; by Lemma 1, d coincides with $d_{X_i,a,b}$ on X_i for $i = 0, 1$. Let $f_0 : X_0 \to X_1$, $f_1 : X_1 \to X_0$ be maps from Proposition 9 (with $\varrho > 1$, $p = 1$) fixing points a and b. Define $f : X \to X$ by $f(x) = f_i(x)$ if $x \in X_i$. Then, for $i = 0, 1$, $f^2|_{X_i} : X_i \to X_i$ is LEL and so, by Proposition 6, $f^2|_{X_i}$ is exactly Devaney chaotic. Thus f is Devaney chaotic and has positive entropy. Since f^2 is not transitive, f is not totally transitive.

If X does not contain a circle, then X is a dendrite. Take any non-end point $a \in X$ and write $X = X_0 \cup X_1$, where X_i's are non-degenerate subdendrites with $X_0 \cap X_1 = \{a\}$. Fix $b_i \in X_i \setminus \{a\}$ and notice that, since X_i's are dendrites, the sets $\mathrm{Cut}_{X_i}(a, b_i)$ are uncountable. Now we can proceed analogously as in the non-dendrite case; the only change is that instead of $f_i(b) = b$ we require $f_0(b_0) = b_1$ and $f_1(b_1) = b_0$. □

Corollary 16. *Every non-degenerate totally regular continuum admits a (not totally) Devaney chaotic, as well as an exactly Devaney chaotic map with finite positive entropy. Every compact metric space which is the disjoint union of finitely many non-degenerate totally regular continua admits a Devaney chaotic map with finite positive entropy.*

4.2. Exactly Devaney chaotic maps with arbitrarily small entropy. In what follows we show that under some conditions a totally regular continuum X admits (exactly) Devaney chaotic maps with arbitrarily small entropy. We consider two subclasses of totally regular continua: those containing arbitrarily large generalized stars and those containing a non-disconnecting free arc. The constructions are modifications of those for stars [2, Theorem 1.2] and for graphs which are not trees [4, Theorems 3.7 and 4.1]. The key difference is that instead of constructing piecewise linear Markov maps we construct P-Lipschitz maps using Proposition 9.

Let X be a continuum and $k \geq 2$ be an integer. We say that X *contains a generalized k-star* if there are a point $a \in X$ and k components $C_1, \dots, C_k$ of $X \setminus \{a\}$ such that the closures $X_i = \overline{C_i} = C_i \cup \{a\}$ $(i = 1, \dots, k)$ are homeomorphic relative to a; i.e. for every i, j there is a homeomorphism $h_{ij} : X_i \to X_j$ fixing the point a (see Figure 1). Recall that X *contains arbitrarily large generalized stars* if it contains a generalized k-star for every $k \geq 2$.

Lemma 17. *There is a constant $c > 0$ such that any totally regular continuum X containing a generalized k-star ($k \geq 2$) admits a (not totally) Devaney chaotic map f as well as an exactly Devaney chaotic map g with positive topological entropies bounded from above by c/k. Consequently, if X contains arbitrarily large generalized stars then $I^{\mathrm{ED}}(X) = 0$.*




FIGURE 1. A continuum containing a generalized k-star

Proof. Since X contains a generalized k-star, there is $a \in X$ such that we can write $X = X_0 \cup X_1 \cup \cdots \cup X_k$, where X_i's are subcontinua of X, $X_i \cap X_j = \{a\}$ for every $i \neq j$ and X_i, X_j are homeomorphic relative to a for $i, j \geq 1$. Further, $X_1, \dots, X_k$ are non-degenerate and, by replacing k with $k-1$ if necessary, we may assume that also X_0 is such. For $i = 0, \dots, k$ put $d_i = d_{X_i,a,a}$ and $\mathcal{C}_i = \mathcal{C}_{X_i,a,a}$.

We may assume that the metrics d_i are such that for every $1 \leq i < k$ there is an isometry $f_i : (X_i, d_i) \to (X_{i+1}, d_{i+1})$ fixing a. Since $\bigcup_{i \neq j} X_i \cap X_j = \{a\}$ is a singleton, by Lemma 1 there is a convex metric d on X such that $d|_{X_i \times X_i} = d_i$ for every i. Let $f_k : X_k \to X_0$, $f_0 : X_0 \to X_1$ be maps from Proposition 9 (with $p = 1$, $\varrho > 1$) such that $f_k(a) = f_0(a) = a$. Without loss of generality we may assume that if $C \in \mathcal{C}_i$ $(i = 1, \dots, k-1)$ then $f_i(C) \in \mathcal{C}_{i+1}$. We define the map $f : X \to X$ by $f(x) = f_i(x)$ for $x \in X_i$, $i = 0, \dots, k$. Proposition 6

implies that $f^{k+1}|_{X_0} : X_0 \to X_0$, being LEL, is exactly Devaney chaotic. Hence $h(f) > 0$, f is a (not totally) Devaney chaotic map, and, since d is convex, f is Lipschitz with $\mathrm{Lip}(f) \le \tilde{L}$ by Lemma 2. Moreover, since $\mathrm{Lip}(f^{k+1}) \le \tilde{L}^2$ we have $h(f) \le \tilde{L}^2/(k+1) \le c/k$ for some constant c depending only on $\tilde{L}$.

To construct an exactly Devaney chaotic map $g : X \to X$ we take the metric d on X as in the previous case and we define $g|_{X_i} = f_i$ for $i = 1, \dots, k$. The only difference is the definition of g on X_0: we put $g|_{X_0} = g_0$, where $g_0 : X_0 \to (X_1 \cup X_2)$ is a map obtained from Proposition 9 with $p = 2$, $\varrho' > 2$, $X' = X_1 \cup X_2$ and $a'_i = a$ for $i = 0, 1, 2$. Then $g_0(a) = a$. By Proposition 9 we have that, for $C \in \mathcal{C}_0$, if $g_0(C)$ contains neither X_1 nor X_2, then $g_0(C) = C_1 \cup C_2$, where $C_1 \in \mathcal{C}_1$ and $C_2 \in \mathcal{C}_2$ are continua and $\mathcal{H}^1(g_0(C)) = \mathcal{H}^1(C_1) + \mathcal{H}^1(C_2) \ge \varrho' \cdot \mathcal{H}^1(C)$. Hence

$$\mathcal{H}^1(C') \ge \frac{\varrho'}{2} \cdot \mathcal{H}^1(C) \qquad \text{for some} \quad C' \in \mathcal{C}_1 \cup \mathcal{C}_2,\ C' \subseteq g(C). \tag{4.1}$$

Notice that $\mathrm{Lip}(g)$ is bounded from above by an absolute constant L, see Lemma 2.

We are going to show the entropy bound for g using Proposition 12. Put $P = \{a\}$, $\mathcal{A} = \{X_0, \dots, X_k\}$, $\mathcal{B} = \{X_1, \dots, X_{k-1}\}$, $L_{X_i} = 1$ for $1 \le i < k$ and $L_{X_0} = L_{X_k} = L$. Then g is P-Lipschitz w.r.t. $\mathcal{A}$ and $(L_{X_i})_i$, $L_{\mathcal{A}} = L$ and $L_{\mathcal{B}} = 1$. Moreover, $X_i \to X_j$ if and only if $j = (i+1) \bmod (k+1)$ or $i = 0$, $j = 2$. Fix any $n \in \mathbb{N}$ and $\mathbb{A} = (A_0, \dots, A_{n-1}) \in \mathcal{A}^n$ with exactly $k_n = k_n^{\mathcal{B}}$ members from $\mathcal{A} \setminus \mathcal{B}$, see (3.1). Let $l_1 < l_2 < \cdots < l_{k_n}$ be the indices l with $A_l \notin \mathcal{B}$, i.e. with $A_l \in \{X_0, X_k\}$. Then, for any l_j, $1 \le i \le k-1$ and $l_j + i < n$, it holds that $A_{l_j+i} \in \{X_{i-1}, X_i, X_{i+1}\}$ and thus $A_{l_j+2}, \dots, A_{l_j+k-2} \in \mathcal{B}$. So $l_{j+2} - l_j \ge k-1$ for every $j \le k_n - 2$. We see at once that $n - 1 \ge l_{k_n} \ge (k_n - 1)(k-1)/2$, i.e. $\theta_{\mathcal{B}} = \limsup_n k_n/n \le 2/(k-1)$. By Proposition 12, $h(g) \le \log^+ L_{\mathcal{B}} + 2\theta_{\mathcal{B}} \log^+ L_{\mathcal{A}} \le (4/(k-1)) \log L \le c/k$ for some absolute constant c.

To prove that g is exact we use Lemma 14 with $\mathcal{D} = \cup_i \mathcal{C}_i$. To this end take any i and $C \in \mathcal{C}_i$ and suppose that for every n the set $g^n(C)$ does not contain any X_i. Then Proposition 9 and (4.1) give that, for every $n \ge 1$,

$$\mathcal{H}^1(g^n(C)) \ge q^n \cdot \mathcal{H}^1(C) \qquad \text{with} \quad q = \min\{\varrho, \varrho'/2\} > 1.$$

(The inequality can be shown by induction as follows. Trivially it holds for $n = 0$. Put $C_0 = C$ and assume that, for some $n \ge 0$, there is i such that the set $g^n(C)$ contains some $C_n \in \mathcal{C}_i$ with $\mathcal{H}^1(C_n) \ge q^n \cdot \mathcal{H}^1(C)$. If $i \ne 0$ then put $C_{n+1} = g(C_n) \in \mathcal{C}_{(i+1) \bmod (k+1)}$ and use Proposition 9. If $i = 0$, put $C_{n+1} = C'$, where C' is the continuum from (4.1).) But this contradicts the fact that $\mathcal{H}^1(X) < \infty$. Hence $\mathcal{D}$ satisfies the assumption from Lemma 14. Notice also that $A \to B$ implies $g(A) \supseteq B$. And since the P-transition graph of g contains cycles $X_0 \to X_1 \to \cdots \to X_k \to X_0$ and $X_0 \to X_2 \to X_3 \to \cdots \to X_k \to X_0$ of lengths $k+1$ and k, the P-transition matrix of g is primitive. Hence, by Lemma 14, g is exact.

To finish it suffices to show that g has dense periodic points. Let $\tilde{X} = X \cup A$ be a totally regular continuum obtained from X by adding to it an arc A such that $A \cap X = \{a\}$. Take a convex metric $\tilde{d}$ on $\tilde{X}$ such that it coincides with d on X and the length of A is 1. Then there is an isometry from I onto A mapping 0 to a. So, by the choice of g_0, there are Lipschitz maps $\psi : X_0 \to A$ and $\varphi : A \to X_1 \cup X_2$

with $\varphi \circ \psi = g_0$ and $\psi(a) = \varphi(a) = a$. Define the map $\tilde{g} : \tilde{X} \to \tilde{X}$ by

$$\tilde{g}(x) = \begin{cases} g(x) & \text{if } x \in X_1 \cup \cdots \cup X_k; \\ \psi(x) & \text{if } x \in X_0; \\ \varphi(x) & \text{if } x \in A. \end{cases}$$

Analogously as before, $\tilde{g}$ is exact. Since $\tilde{X}$ contains a disconnecting interval (an open set homeomorphic to $(0,1)$ such that any point of it disconnects X into exactly two components), $\tilde{g}$ has dense periodic points by [2, Theorem 1.1]. Trivially, any periodic point of $\tilde{g}$ in X is also a periodic point of g. So also the periodic points of g are dense. □

Lemma 18. *There is $c > 0$ with the following property. Let X be a totally regular continuum which can be written as the union of non-degenerate subcontinua*

$$X = X_0 \cup X_1 \cup \cdots \cup X_k,$$

where $k \geq 2$ and the following hold:

(a) *there are distinct points $a_0, \ldots, a_k$ with $X_i \cap X_{(i+1) \bmod (k+1)} = \{a_i\}$ for $i = 0, \ldots, k$;*
(b) *$X_i \cap X_j = \emptyset$ whenever $2 \leq |i - j| < k$;*
(c) *for $i = 1, \ldots, k-1$ there is a homeomorphism $h_i : X_i \to X_{i+1}$ such that $h_i(a_{i-1}) = a_i$ and $h_i(a_i) = a_{i+1}$;*
(d) *$\mathrm{Cut}_{X_1}(a_0, a_1)$ is uncountable.*

Then X admits a (not totally) Devaney chaotic map f and an exactly Devaney chaotic map g with positive topological entropies bounded from above by c/k.

See Figure 2 for an illustration of a continuum satisfying the assumptions of the lemma.




FIGURE 2. A continuum from Lemma 18

Proof. We may assume that $\mathrm{Cut}_{X_0}(a_k, a_0)$ is uncountable, for if not, we replace k by $k' = k - 1$ and X_{k-1} by $X'_{k-1} = X_{k-1} \cup X_k$ (if $k' = 1$ we simply apply Lemma 15).

Consider the metrics $d_i = d_{X_i,a_{i-1},a_i}$ for $i = 0,\dots,k$ (with $a_{-1} := a_k$); in view of (c) we may assume that $f_i : (X_i, d_i) \to (X_{i+1}, d_{i+1})$ are isometries for $1 \le i < k$. Let d be the convex metric on X defined by (2.2). Since $\bigcup_{i\neq j} X_i \cap X_j = \{a_i : \ i = 0,\dots,k\}$, $1 > d_i(a_{i-1}, a_i) > 1/2$ and $X_i \cap X_j$ are empty or singletons for $i \neq j$, Lemma 1 gives that $d|_{X_i\times X_i} = d_i$ for every i.

Let $f_k : X_k \to X_0$, $f_0 : X_0 \to X_1$ and $g_0 : X_0 \to (X_1 \cup X_2)$ be maps from Proposition 9 with $(\varrho > 2)$ such that $f_0(a_k) = g_0(a_k) = a_0$, $f_0(a_0) = g_0(a_0) = a_1$, $f_k(a_{k-1}) = a_k$ and $f_k(a_k) = a_0$. Define $f, g : X \to X$ by

$$f(x) = \begin{cases} f_0(x) & \text{if } x \in X_0; \\ h_i(x) & \text{if } x \in X_i,\ 1 \le i < k; \\ f_k(x) & \text{if } x \in X_k; \end{cases} \qquad g(x) = \begin{cases} g_0(x) & \text{if } x \in X_0; \\ h_i(x) & \text{if } x \in X_i,\ 1 \le i < k; \\ f_k(x) & \text{if } x \in X_k. \end{cases}$$

Analogously as in the proof of Lemma 17 we can show that f is (not totally) Devaney chaotic, g is exact and the entropies of f, g are positive and bounded from above by c/k, where c is an absolute constant. What is left is to prove that g has dense periodic points.

To this end put $P = \{a_i : \ i = 0,\dots,k\}$ and realize that $g(P) = P$. Let $\mathcal{D}$ be the decomposition of X into P and singletons $\{x\}$, $x \notin P$. Let $X' = X/\mathcal{D}$ (i.e. we collapse P into a point), $\pi : X \to X'$ be the natural projection and $\{a'\} = \pi(P)$, $X'_i = \pi(X_i)$. Then X' is a compact metric space since the decomposition $\mathcal{D}$ is upper semicontinuous. The map $g : X \to X$ induces $g' : X' \to X'$ which fixes a' and, being a factor of g, is exact. Notice that $g'|_{X'_0} : X'_0 \to X'_1 \cup X'_2$ can be written as the composition $\varphi \circ \psi$ with continuous $\psi : X'_0 \to I$ and $\varphi : I \to X'_1 \cup X'_2$. So, analogously as in the proof of Lemma 17, g' has dense periodic points. Hence also g has dense periodic points, since whenever $x' \neq a'$ is a periodic point of g', then its only π-preimage x is a periodic point of g. □

Finally, we are ready to prove our main result.

Main Theorem. *For every non-degenerate totally regular continuum X it holds that $I^{\mathrm{ED}}(X) < \infty$. Moreover, if X contains arbitrarily large generalized stars or X contains a free arc which does not disconnect X, then X admits*

- *a Devaney chaotic map which is not totally Devaney chaotic, and*
- *an exactly Devaney chaotic map,*

both with arbitrarily small positive entropy; hence $I^{\mathrm{ED}}(X) = 0$.

Proof. Note that if X contains a non-disconnecting free arc A then, for every $k \ge 2$, X can be written as in Lemma 18 with $X_1,\dots,X_k$ being subarcs of A. So the theorem immediately follows from Corollary 16 and Lemmas 17, 18. □

Remark 19. Lemma 18 gives $I^{\mathrm{ED}}(X) = 0$ also for some continua containing no non-disconnecting free arc. For example if, for some $m \in \{3, 4, \dots, \aleph_0\}$, X contains the universal dendrite D_m of order m with two point boundary and with connected $X \setminus \operatorname{int}(D_m)$, we again have $I^{\mathrm{ED}}(X) = 0$. (Recall that D_m is the topologically unique dendrite such that the branch points of it are dense and all have the order m.)

References

[1] L. Alsedà, S. Baldwin, J. Llibre and M. Misiurewicz, *Entropy of transitive tree maps*, Topology **36** (1997), no. 2, 519–532.

[2] L. Alsedà, S. Kolyada, J. Llibre and Ľ. Snoha, *Entropy and periodic points for transitive maps*, Trans. Amer. Math. Soc. **351** (1999), no. 4, 1551–1573.
[3] Ll. Alsedà, J. Llibre and M. Misiurewicz, *Combinatorial Dynamics and Entropy in Dimension One*, Second Edition, Advanced Series in Nonlinear Dynamics 5, World Scientific, Singapore, 2000.
[4] L. Alsedà, M. A. del Río and J. A. Rodríguez, *A splitting theorem for transitive maps*, J. Math. Anal. Appl. **232** (1999), no. 2, 359–375.
[5] S. Baldwin, *Entropy estimates for transitive maps on trees*, Topology **40** (2001), no. 3, 551–569.
[6] F. Balibrea and Ľ. Snoha, *Topological entropy of Devaney chaotic maps*, Topology Appl. **133** (2003), no. 3, 225–239.
[7] R. H. Bing, *Partitioning a set*, Bull. Amer. Math. Soc. **55** (1949), 1101–1110.
[8] A. Blokh, *On sensitive mappings of the interval*, Russian Math. Surveys **37** (1982), 203–204.
[9] A. Blokh, *On transitive mappings of one-dimensional branched manifolds. (Russian)*, Differential-difference equations and problems of mathematical physics (Russian), 3–9, 131, Akad. Nauk Ukrain. SSR, Inst. Mat., Kiev, 1984.
[10] A. Blokh, *On the connection between entropy and transitivity for one-dimensional mappings*, Russ. Math. Surv. **42** (1987), no. 5, 165–166.
[11] R. Bowen, *Entropy for group endomorphisms and homogeneous spaces*, Trans. Amer. Math. Soc. **153** 1971, 401–414.
[12] R. D. Buskirk, J. Nikiel and E. D. Tymchatyn, *Totally regular curves as inverse limits*, Houston J. Math. **18** (1992), no. 3, 319–327.
[13] E. I. Dinaburg, *A connection between various entropy characterizations of dynamical systems*, (Russian) Izv. Akad. Nauk SSSR Ser. Mat. **35** (1971), 324–366.
[14] M. Dirbák, Ľ. Snoha and V. Špitalský, *Dynamical consequences of a free interval: minimality, transitivity, mixing and topological entropy*, `arXiv:1112.1191v1 [math.DS]` (2011).
[15] K. J. Falconer, *The geometry of fractal sets*, Cambridge University Press, Cambridge, 1986.
[16] H. Federer, *Geometric measure theory*, Springer-Verlag New York Inc., New York 1969.
[17] D. H. Fremlin, *Spaces of finite length*, Proc. London Math. Soc. (3) **64** (1992), 449–486.
[18] G. Harańczyk, D. Kwietniak and P. Oprocha, *Topological structure and entropy of mixing graph maps*, `arXiv:1111.0566v1 [math.DS]` (2011).
[19] A. Katok and B. Hasselblatt, *Introduction to the modern theory of dynamical systems*, Cambridge University Press, Cambridge, 1995.
[20] S. Kolyada and M. Matviichuk, *On extensions of transitive maps*, Discrete Contin. Dyn. Syst. **30** (2011), no. 3, 767–777.
[21] K. Kuratowski, *Topology, vol. 2*, Academic Press and PWN, Warszawa, 1968.
[22] D. Kwietniak and M. Misiurewicz, *Exact Devaney chaos and entropy*, Qual. Theory Dyn. Syst. **6** (2005), no. 1, 169–179.
[23] S. Macías, *Topics on continua*, Chapman & Hall/CRC, Boca Raton, FL, 2005.
[24] P. Mattila, *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability*, Cambridge University Press, Cambridge, 1995.
[25] S. B. Nadler, *Continuum theory. An introduction*, Monographs and Textbooks in Pure and Applied Mathematics, 158, Marcel Dekker, Inc., New York, 1992.
[26] J. Nikiel, *Locally connected curves viewed as inverse limits*, Fund. Math. **133** (1989), no. 2, 125–134.
[27] S. Ruette, *Chaos for continuous interval maps — a survey of relationship between the various sorts of chaos*, preprint, available at `http://www.math.u-psud.fr/~ruette/articles/chaos-int.pdf`.
[28] V. Špitalský, *Length-expanding Lipschitz maps on totally regular continua*, `arXiv:1203.2352v1 [math.DS]` (2012).
[29] G. T. Whyburn, *Analytic topology*, American Mathematical Society, New York, 1942.
[30] X. Ye, *Topological entropy of transitive maps of a tree*, Ergodic Theory Dynam. Systems **20** (2000), no. 1, 289–314.

Department of Mathematics, Faculty of Natural Sciences, Matej Bel University, Tajovského 40, 974 01 Banská Bystrica, Slovakia
E-mail address: `vladimir.spitalsky@umb.sk`