

# Some further notes on the Kruskal - Szekeres completion.

Kayll Lake

Department of Physics, Queen's University, Kingston, Ontario, Canada, K7L 3N6

(Dated: April 3, 2024)

In this pedagogical note, the Kruskal - Szekeres completion of the Schwarzschild spacetime is obtained explicitly in a few steps.

PACS numbers: 04.20.Cv, 04.20.Jb, 04.20.Gz

A discussion of the Kruskal [1] - Szekeres [2] completion of the Schwarzschild spacetime is a cornerstone of any modern introduction to general relativity. In a previous note [3] I reviewed some history of these coordinates and showed how they can be obtained from Israel coordinates [4] and also from an integration of Einstein's equations. The purpose of the present note is to point out a very brief but direct and explicit derivation of the Kruskal - Szekeres completion. We use geometrical units and a signature of +2.

We start with the standard form of the Schwarzschild spacetime

$$ds^2 = -\left(1 - \frac{2m}{r}\right) dt^2 + \frac{dr^2}{1 - \frac{2m}{r}} + r^2 d\Omega_2^2, \quad (1)$$

where  $d\Omega_2^2$  is the metric of a unit two-sphere and  $m$  is a constant  $> 0$ . The radial null geodesic equations of (1) are

$$\frac{dr}{dt} = \pm \left(1 - \frac{2m}{r}\right). \quad (2)$$

The solutions to (2) can be given as

$$r = 2m(\mathcal{L}(\Psi) + 1) \quad (3)$$

and

$$r = 2m(\mathcal{L}(\Phi) + 1). \quad (4)$$

Here  $\mathcal{L}(x)$  is the Lambert W function [5], defined by  $\mathcal{L}(x) \exp(\mathcal{L}(x)) = x$  [6],

$$\Psi^2 \equiv \left(\frac{u^2}{e}\right)^2 \exp\left(\frac{t}{m}\right), \quad (5)$$

$$\Phi^2 \equiv \left(\frac{v^2}{e}\right)^2 \exp\left(-\frac{t}{m}\right), \quad (6)$$

and  $u$  and  $v$  label the radial null geodesics. Now at any two - sphere  $(u, v)$ , where the branches of the radial null geodesics cross,  $\Psi^2 = \Phi^2$  and so from (5) and (6) we find

$$\exp\left(\frac{t}{m}\right) = \left(\frac{v}{u}\right)^2 \quad (7)$$

and so

$$\Psi^2 = \Phi^2 = \left(\frac{uv}{e}\right)^2. \quad (8)$$

Now in taking the appropriate roots to (7) and (8) we must orientate the  $u-v$  axes. Noting that  $\mathcal{L}(-1/e) = -1$  and  $\mathcal{L}(0) = 0$  we take  $uv > 0$  over the range  $0 < r < 2m$  and so  $\Psi = \Phi = -\frac{uv}{e}$ . We have arrived at [7]

$$d\tilde{s}^2 = \frac{-(4m)^2}{(1 + \mathcal{L}) \exp(1 + \mathcal{L})} dudv + (2m)^2 (1 + \mathcal{L})^2 d\Omega_2^2, \quad (9)$$

where we have now written  $\mathcal{L} \equiv \mathcal{L}(-uv/e)$  and take  $\exp(t/2m) = v/u$  for  $uv > 0$  and  $\exp(t/2m) = -v/u$  for  $uv < 0$ . This is the null form of the Kruskal - Szekeres completion.

We convert our results into standard notation in the Appendix.

## ACKNOWLEDGMENTS

I thank Bill Unhru for useful criticism of an earlier draft. This work was supported by a grant from the Natural Sciences and Engineering Research Council of Canada. Portions of this work were made possible by use of *GRTensorII* [9].

- 
- [1] M. Kruskal, Phys. Rev., **119**, 1743 (1960).
  - [2] G. Szekeres, Gen. Rel. Grav., **34**, 2001 (2002) (Reprinted from Publicationes Mathematicae Debrecen **7**, 285 (1960)).
  - [3] K. Lake, Class. Quantum Grav., **27**, 097001 (2010), [arXiv:gr-qc/1002.3600].
  - [4] W. Israel, Phys. Rev., **143**, 1016 (1966).
  - [5] See, for example, R. M. Corless, G. H. Gonnet, D. E.

- G. Hare, D. J. Jeffrey, and D. E. Knuth, Advances in Computational Mathematics **5**, 329 (1996).
- [6] Note that  $x \geq -1/e$ . We require  $\mathcal{L}(x) \geq -1$  so that  $\mathcal{L}(x)$  is single valued.
- [7] Note that  $d\mathcal{L}/dz = \mathcal{L}/z(1 + \mathcal{L})$ .
- [8] C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation* (W. H. Freeman, San Francisco, 1973).
- [9] This package runs within Maple. The GRTensorII software

and documentation is distributed freely from the address  
<http://grtensor.org>

## APPENDIX

Following, for example [8], define

$$V \equiv \frac{v+u}{2}, \quad U \equiv \frac{v-u}{2} \quad (10)$$

so that

$$v = V + U, \quad u = V - U. \quad (11)$$

Since

$$r = 2m(1 + \mathcal{L}(-\frac{uv}{e})), \quad \mathcal{L} \exp(\mathcal{L}) = -\frac{uv}{e}$$

we have

$$\left(\frac{r}{2m} - 1\right) \exp\left(\frac{r}{2m}\right) = U^2 - V^2, \quad (12)$$

the usual implicit relation for  $r$ . We now have [7]

$$ds^2 = -\frac{32m^3}{r} \exp\left(-\frac{r}{2m}\right) (dV^2 - dU^2) + r^2 d\Omega_2^2, \quad (13)$$

the usual form of the Kruskal - Szekeres metric.

Finally, for  $uv > 0$  we have  $v/u = \exp(t/2m)$  and so

$$\frac{U}{V} = \tanh\left(\frac{t}{4m}\right), \quad V^2 > U^2 \quad (14)$$

and for  $uv < 0$  we have  $v/u = -\exp(t/2m)$  and so

$$\frac{V}{U} = \tanh\left(\frac{t}{4m}\right), \quad V^2 < U^2. \quad (15)$$

The usual procedure is to introduce (12), (14) and (15) as coordinate transformations to change (1) to (13). The purpose of this note is to point out that these seemingly mysterious coordinate transformations need never be introduced.