

ISOSPECTRAL FLOWS ON A CLASS OF FINITE-DIMENSIONAL JACOBI MATRICES

TOBIAS SUTTER, DEBASISH CHATTERJEE, FEDERICO A. RAMPONI,
AND JOHN LYGEROS

ABSTRACT. We present a new matrix-valued isospectral ordinary differential equation that asymptotically block-diagonalizes $n \times n$ zero-diagonal Jacobi matrices employed as its initial condition. This o.d.e. features a right-hand side with a nested commutator of matrices, and structurally resembles the double-bracket o.d.e. studied by R.W. Brockett in 1991. We prove that its solutions converge asymptotically, that the limit is block-diagonal, and above all, that the limit matrix is defined uniquely as follows: For n even, a block-diagonal matrix containing 2×2 blocks, such that the super-diagonal entries are sorted by strictly increasing absolute value. Furthermore, the off-diagonal entries in these 2×2 blocks have the same sign as the respective entries in the matrix employed as initial condition. For n odd, there is one additional 1×1 block containing a zero that is the top left entry of the limit matrix. The results presented here extend some early work of Kac and van Moerbeke.

§ 1. INTRODUCTION AND MAIN RESULT

The tasks of sorting a list, diagonalizing a matrix, and solving a linear programming problem are traditionally solved with computer science algorithms, for example the quicksort algorithm for sorting or the simplex method for solving linear programs. Brockett [Brockett, 1991] showed that solutions to such problems can also be obtained by means of a smooth dynamical system, in particular as the limit of solutions to certain matrix-valued ordinary differential equations (o.d.e.'s). A classical problem from linear algebra is therefore solvable by calculus. Motivated by Brockett's work, new problems, conventionally tackled by algebraic methods, have been assigned to calculus. For instance, [Faybusovich, 1992] proposed an ordinary differential equation (structurally similar to the one proposed by Brockett) as the starting point in a general approach to interior point methods for linear programming.

By a *Jacobi matrix* we mean a symmetric tridiagonal matrix (in general, infinite) with real entries and distinct eigenvalues. In this article we present a matrix-valued ordinary differential equation which asymptotically block-diagonalizes a finite-dimensional zero-diagonal Jacobi matrix taken as its initial condition. Jacobi matrices arise in a variety of applications, for example in solid state physics to characterize the Toda lattice, which is a simple model for a one dimensional

Key words and phrases. Isospectral flows; Kac-van Moerbeke flow; Jacobi matrices; Linear algebra; Krasovskij-LaSalle invariance principle.

This research was partially supported by the European Commission under the project MoVeS (FP7-ICT-2009.3.5), and the HYCON2 Network of Excellence (FP7-ICT-2009-5).

crystal—see e.g. [Moser, 1975], [Helmke and Moore, 1994, pp. 59-60] for a detailed study. There is also a strong connection between Brockett’s double bracket flow [Brockett, 1991] and the Toda lattice equation, which was first observed by [Bloch, 1990].

We offer a second motivation here that has intrinsic appeal and interest, relating to the computation of the roots of certain polynomials. Orthogonal polynomials on the real line corresponding to a Borel probability measure μ have considerable applications in mathematical physics and engineering [Simon, 2005]. Let $\langle \cdot, \cdot \rangle$ denote the standard inner product on the Hilbert space $L_2(\mathbb{R}, \mu)$. Then a sequence of monic orthogonal polynomials on the real line is defined recursively [Szegő, 1959] by

$$\begin{aligned} P_{n+1}(x) &= xP_n(x) - a_n^2 P_{n-1}(x) - b_{n+1} P_n(x), \quad n \in \mathbb{N}, \\ P_{-1}(\cdot) &= 0, \quad P_0(\cdot) = 1, \end{aligned}$$

where

$$a_n := \frac{\|P_n\|}{\|P_{n-1}\|}, \text{ for } n \in \mathbb{N} \text{ and } b_{n+1} := \frac{\langle xP_n, P_n \rangle}{\|P_n\|^2} \text{ for } n \in \mathbb{N}_0.$$

To a given measure μ , there corresponds a (generally infinite) Jacobi matrix

$$(1.1) \quad \mathcal{J} := \begin{pmatrix} b_1 & a_1 & 0 & 0 & \cdots \\ a_1 & b_2 & a_2 & 0 & \cdots \\ 0 & a_2 & b_3 & a_3 & \cdots \\ 0 & 0 & a_3 & b_4 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

with strict positive off-diagonal entries derived from its orthogonal polynomials, and it is well-known that the zeros of these orthogonal polynomials are precisely the eigenvalues of finite truncations of this Jacobi matrix. Conversely, Favard’s Theorem [Favard, 1935] shows that to every finite-dimensional symmetric tridiagonal matrix with strictly positive off-diagonal entries there corresponds a finitely supported measure. (The uniqueness of this measure for an infinite-dimensional Jacobi matrix is an issue that relates to the solvability of the “moment problem” [Akhiezer, 1965], with which we shall not deal here.) Since the space of square integrable functions corresponding to this measure is finite dimensional (the measure itself being finitely supported), it is enough that from the orthogonal polynomials it is possible to recover the measure, and this leads to the problem of finding the roots of such polynomials. The latter, in general, is known to be a difficult task. As mentioned above, for our applications it suffices to determine the eigenvalues of the finite truncations of the corresponding Jacobi matrix, which we shall do in this article with the aid of an appropriate matrix-valued ordinary differential equation. For this we further specialize the measures to non-negative linear combinations of finitely many Dirac measures on the real line placed symmetrically around 0. These measures give rise to zero-diagonal Jacobi matrices, and our main result to the problem of finding the roots of the orthogonal polynomials corresponding to these measures.

As a concrete application we consider the Gaussian quadrature method. For a given positive weight function $w(x)$ on an interval $[a, b]$ of the real line, the n -point

Gaussian quadrature rule approximates an integral by

$$\sum_{i=1}^n w_i f(x_i) \approx \int_a^b f(x) w(x) dx.$$

It is known (see [Simon, 2005, p. 17], [Szegő, 1959, p. 351], [Watkins, 2005, p. 21]) that the optimal points x_1, \dots, x_n coincide with the zeros of the orthogonal polynomial $P_n(x)$ introduced above, when we use the probability measure $\mu = \frac{w(x) dx}{\int_a^b w(x) dx}$, with dx being the Lebesgue measure. As mentioned, these zeros are precisely the eigenvalues of a certain $n \times n$ truncation of the Jacobi matrix in (1.1). Furthermore, if the interval $[a, b]$ is symmetric about 0 and the weight function $w(x)$ is even, the resulting Jacobi matrix has zero diagonal entries.

In this article, we treat the problem of obtaining the eigenvalues of zero-diagonal finite dimensional Jacobi matrices from the asymptotic limit of a smooth dynamical system. Preparatory to stating our main result, we need some preliminary notation: We let $\text{Sym}(n)$ and $\text{Skew}(n)$ denote respectively the set of symmetric and skew-symmetric $n \times n$ matrices with real entries. We define $\text{Jac}_0(n)$ as the set of all $n \times n$ Jacobi matrices with real entries and zeros on its diagonal. For a matrix A , $\|A\|$ is the Frobenius norm defined as $\|A\| = \sqrt{\text{tr}(AA^\top)}$. Let $\mathcal{C}(\mathbb{R}_{\geq 0}, \mathbb{R})$ denote the set continuous functions from $\mathbb{R}_{\geq 0}$ to \mathbb{R} . The bracket $[\cdot, \cdot]$ is the usual matrix commutator $[A, B] = AB - BA$. For $H_0 \in \text{Sym}(n)$ we let $M(H_0) := \{\Theta^\top H_0 \Theta \mid \Theta \in \mathcal{O}(n, \mathbb{R})\}$ denote the set of all real matrices orthogonally similar to H_0 [Moore et al., 1994]. For $H_0 \in \text{Jac}_0(n)$ we define $M_J(H_0)$ to be the set of all zero-diagonal Jacobi matrices that are isospectral to H_0 (that is, they have the same eigenvalues). Moreover, for $p = 0, 1, \dots, n-1$, we let

$$\mathcal{D}_{u,p}(a_1, a_2, \dots, a_{n-p}) := \begin{pmatrix} 0 & \dots & 0 & a_1 & & \\ & 0 & \dots & 0 & a_2 & \\ & & \ddots & & & \ddots \\ & & & 0 & a_{n-p} & \\ & & & & 0 & \\ & & & & & \vdots \\ & & & & & & 0 \end{pmatrix} \in \mathbb{R}^{n \times n}$$

and

$$\mathcal{D}_p(a_1, a_2, \dots, a_{n-p}) := \mathcal{D}_{u,p}(a_1, a_2, \dots, a_{n-p}) + \mathcal{D}_{u,p}(a_1, a_2, \dots, a_{n-p})^\top.$$

The following is our main result:

(1.2). **Theorem.** *Let n be a positive integer. Consider the zero-diagonal Jacobi matrix*

$$(1.3) \quad H = \begin{pmatrix} 0 & a_1 & & & \\ a_1 & 0 & a_2 & & \\ & a_2 & & \ddots & \\ & & & & a_{n-2} & \\ & & & a_{n-2} & 0 & a_{n-1} \\ & & & & 0 & a_{n-1} & 0 \end{pmatrix} \in \text{Jac}_0(n),$$

and the skew-symmetric matrix

$$(1.4) \quad K(H) := \mathcal{D}_{u,2}(a_1 a_2, a_2 a_3, \dots, a_{n-2} a_{n-1}) \\ - \mathcal{D}_{u,2}(a_1 a_2, a_2 a_3, \dots, a_{n-2} a_{n-1})^\top,$$

derived from H . Consider the matrix-valued o.d.e.

$$(1.5) \quad \frac{d}{dt} H(t) = [H(t), K(H(t))], \quad H(0) := H_0 \in \text{Jac}_0(n).$$

- (i) (1.5) defines an isospectral flow on the set of all Jacobi matrices with zero diagonal entries.
- (ii) The solutions $H(t)$ of (1.5) exist for all $t \geq 0$ and approach asymptotically the set of equilibrium points of (1.5), which is given by

$$\{\bar{H} \in \text{M}_J(H_0) \mid K(\bar{H}) = 0\}.$$

- (iii) $\lim_{t \rightarrow \infty} H(t)$ exists.
- (iv) If n is even, then for all initial conditions $H_0 \in \text{Jac}_0(n)$ other than the equilibria, where $H_0 = \mathcal{D}_1(a_1, \dots, a_{n-1})$ and $\sigma(H_0) = \{\pm \lambda_1, \dots, \pm \lambda_{\frac{n}{2}}\}$ with $|\lambda_1| < |\lambda_2| < \dots < |\lambda_{\frac{n}{2}}|$, the solution $H(t)$ of (1.5) converges to

$$\lim_{t \rightarrow \infty} H(t) = \mathcal{D}_1(\text{sgn}(a_1)|\lambda_1|, 0, \text{sgn}(a_3)|\lambda_2|, 0, \dots, \text{sgn}(a_{n-1})|\lambda_{\frac{n}{2}}|).$$

If n is odd, then for all initial conditions $H_0 \in \text{Jac}_0(n)$ other than the equilibria, where $H_0 = \mathcal{D}_1(a_1, \dots, a_{n-1})$ and $\sigma(H_0) = \{0, \pm \lambda_1, \dots, \pm \lambda_{\frac{n-1}{2}}\}$ with $0 < |\lambda_1| < |\lambda_2| < \dots < |\lambda_{\frac{n-1}{2}}|$, the solution $H(t)$ of (1.5) converges to

$$\lim_{t \rightarrow \infty} H(t) = \mathcal{D}_1\left(0, \text{sgn}(a_2)|\lambda_1|, 0, \text{sgn}(a_4)|\lambda_2|, 0, \dots, \text{sgn}(a_{n-1})|\lambda_{\frac{n-1}{2}}|\right).$$

Remarks and contributions.

1. The assertions (i) – (iii) imply that the o.d.e. (1.5) evolves on the set of zero-diagonal Jacobi matrices with a fixed spectrum determined by its initial condition. Solutions exist for all $t \geq 0$, attain a limit as $t \rightarrow \infty$ and approach the set of equilibrium points (having cardinality greater than 1 for $n > 2$) asymptotically. For all initial conditions $H_0 \in \text{Jac}_0(n)$ other than the equilibria, however, property (iv) defines the limit matrix uniquely as follows: For n even, a block-diagonal matrix containing 2×2 blocks, such that the super-diagonal entries are sorted by strictly increasing absolute value. Furthermore, the off-diagonal entries in these 2×2 blocks have the same sign as the respective entries in the matrix employed as initial condition. For n odd, there is one additional 1×1 block containing a zero that is the top left entry of the limit matrix.
2. [Brockett, 1991] studied the o.d.e. (1.5) in which the map K on the right-hand side of (1.5) was set to $K(H) = [H, N]$, where N is a constant symmetric matrix. In our case, however, $K(H) = [H, N(H)]$, where N is a linear function of H , not a constant. As an important consequence of the definition of K as in (1.4), all equilibrium points of (1.5) are non-hyperbolic. In contrast, for $K(H) = [H, N]$, where N is a constant symmetric matrix, all the equilibrium points are known to be hyperbolic [Brockett, 1991]. In particular, the proof techniques in [Brockett, 1991] do not carry over, and in order to prove the sorting property (iv) in Theorem (1.2)—the stable manifold theorems [Helmke

and Moore, 1994, p. 362ff] cannot be employed. As such, the analysis of (1.5) requires new tools.

3. It is possible to treat N as a parameter in [Brockett, 1991], and can therefore be employed, e.g., to sort the eigenvalues of $\lim_{t \rightarrow \infty} H(t)$ according to a particular order by selecting an appropriate matrix N . Such variations in our case are not readily available since K is a fixed function defined by (1.4).
4. The dynamical system $\frac{d}{dt}H(t) = [K(H(t)), H(t)]$ was studied first in [Kac and van Moerbeke, 1975]. This article studied several properties of (1.5) by considering the dynamics of the individual components, and the techniques relied on properties of the orthogonal polynomials associated to Jacobi matrices. In contrast, our technical tools are system theoretic. The analysis of the properties of (1.5) from a double bracket perspective, to our knowledge, has been carried out here for the first time. In addition, the sorting property (iv) in Theorem (1.2) is an entirely new observation.

Outline of the article: §2 presents the proof of the Theorem, which we illustrate with some numerical examples in §3. We conclude in §4 with a summary of our work and comment on possible subjects of further research.

§ 2. PROOF OF THEOREM (1.2)

Some preliminaries are needed in order to prove Theorem (1.2). We begin with the following classical result, which will play a key role behind proving that the solutions to (1.5) are isospectral.

(2.1). **Proposition** ([Lax, 1968]). *Let $\varphi : \text{Sym}(n) \rightarrow \text{Skew}(n)$ be a smooth mapping, and suppose that $\gamma : \mathbb{R}_{\geq 0} \rightarrow \text{Sym}(n)$ is a curve satisfying*

$$(2.2) \quad \frac{d\gamma}{dt}(t) = [\varphi \circ \gamma(t), \gamma(t)], \quad t \geq 0.$$

Then there exists a smooth family of unitary matrices $(U(t))_{t \geq 0}$ with $U(0) = I_n$ such that

$$\gamma(t) = U(t)^{-1} \gamma(0) U(t), \quad t \geq 0.$$

The family $(\gamma(t))_{t \geq 0}$ is thus isospectral.

Next we define the mapping

$$(2.3) \quad \text{Sym}(n) \ni A := \begin{pmatrix} a_{11} & a_{12} & \dots & & \\ a_{12} & a_{22} & & & \\ \vdots & & \ddots & & \\ & & & a_{n-1,n-1} & a_{n-1,n} \\ & & & a_{n-1,n} & a_{nn} \end{pmatrix} \mapsto$$

$$N(A) := \mathcal{D}_1(-a_{12}, 0, a_{34}, 2a_{45}, \dots, (n-3)a_{n-1,n}) \in \text{Sym}(n).$$

The mapping $N(\cdot)$ is linear, and can be written as

$$(2.4) \quad N(A) = \sum_{i=1}^{n-1} (i-2) (E_i A E_{i+1} + E_{i+1} A E_i),$$

where $A \in \text{Sym}(n)$ and E_i is the matrix with 1 at its (i, i) -th entry and zeros elsewhere.

(2.5). **Proposition.** *Let*

$$(2.6) \quad H_n = \begin{pmatrix} 0 & a_1 & 0 & & & \\ a_1 & 0 & a_2 & & & \\ 0 & a_2 & & & & \\ & & \ddots & & & \\ & & & a_{n-2} & 0 & \\ & & & a_{n-2} & 0 & a_{n-1} \\ & & & 0 & a_{n-1} & 0 \end{pmatrix} \in \text{Jac}_0(n).$$

Then the commutator of H_n and $N(H_n)$ is given by

$$(2.7) \quad \begin{aligned} K(H_n) &:= [H_n, N(H_n)] \\ &= \mathcal{D}_{u,2}(a_1 a_2, \dots, a_{n-2} a_{n-1}) - \mathcal{D}_{u,2}(a_1 a_2, \dots, a_{n-2} a_{n-1})^\top, \end{aligned}$$

where $N(\cdot)$ is the linear mapping defined in (2.3).

Proof. The proof proceeds by induction. Observe that $\mathcal{D}_{u,2} = 0$ if $n = 1$. For $n = 1$, we have $H_n = 0$, and therefore $K(H_n) = 0$. For $n = 2$, we have $H_n = \begin{pmatrix} 0 & a_1 \\ a_1 & 0 \end{pmatrix}$ and $N(H_n) = \begin{pmatrix} 0 & -a_1 \\ -a_1 & 0 \end{pmatrix}$. Since H_n and $N(H_n)$ commute, $K(H_n) = 0$.

We consider the induction step:

$$(2.8) \quad H_{n+1} = \begin{pmatrix} & & & & 0 & 0 \\ & & & & \vdots & \vdots \\ & \boxed{H_{n-1}} & & & 0 & 0 \\ & & & & a_{n-1} & 0 \\ 0 & \dots & 0 & a_{n-1} & 0 & a_n \\ 0 & \dots & 0 & 0 & a_n & 0 \end{pmatrix}$$

and

$$(2.9) \quad N(H_{n+1}) = \begin{pmatrix} & & & & 0 & 0 \\ & & & & \vdots & \vdots \\ & \boxed{N(H_{n-1})} & & & 0 & 0 \\ & & & & (n-3)a_{n-1} & 0 \\ 0 & \dots & 0 & (n-3)a_{n-1} & 0 & (n-2)a_n \\ 0 & \dots & 0 & 0 & (n-2)a_n & 0 \end{pmatrix}.$$

Simple matrix multiplications lead to
(2.10)

$$H_{n+1}N(H_{n+1}) = \begin{pmatrix} & & 0 & 0 \\ & \boxed{H_{n-1}N(H_{n-1})} & \vdots & \vdots \\ & & 0 & 0 \\ & & (n-3)a_{n-1}a_{n-2} & 0 \\ & & 0 & \textcircled{3} \\ 0 & \dots & 0 & (n-4)a_{n-1}a_{n-2} & 0 & \textcircled{4} & 0 \\ 0 & \dots & 0 & 0 & \textcircled{1} & 0 & \textcircled{2} \end{pmatrix},$$

where

$$\begin{aligned} \textcircled{1} &= (n-3)a_n a_{n-1}, \\ \textcircled{2} &= (n-2)a_n^2, \\ \textcircled{3} &= (n-2)a_n a_{n-1}, \\ \textcircled{4} &= (n-3)a_{n-1}^2 + (n-2)a_n^2, \end{aligned}$$

and
(2.11)

$$N(H_{n+1})H_{n+1} = \begin{pmatrix} & & 0 & 0 \\ & \boxed{N(H_{n-1})H_{n-1}} & \vdots & \vdots \\ & & 0 & 0 \\ & & (n-4)a_{n-1}a_{n-2} & 0 \\ & & 0 & \textcircled{6} \\ 0 & \dots & 0 & (n-3)a_{n-1}a_{n-2} & 0 & \textcircled{4} & 0 \\ 0 & \dots & 0 & 0 & \textcircled{5} & 0 & \textcircled{2} \end{pmatrix},$$

where

$$\begin{aligned} \textcircled{5} &= (n-2)a_n a_{n-1}, \\ \textcircled{6} &= (n-3)a_n a_{n-1}. \end{aligned}$$

By the induction hypothesis,

$$[H_{n-1}, N(H_{n-1})] = H_{n-1}N(H_{n-1}) - N(H_{n-1})H_{n-1} = K(H_{n-1}).$$

Therefore, by (2.10) and (2.11) we get

$$\begin{aligned} [H_{n+1}, N(H_{n+1})] &= \begin{pmatrix} & & 0 & 0 \\ & \boxed{K(H_{n-1})} & \vdots & \vdots \\ & & 0 & 0 \\ & & a_{n-1}a_{n-2} & 0 \\ & & 0 & a_n a_{n-1} \\ 0 & \dots & 0 & -a_{n-1}a_{n-2} & 0 & 0 & 0 \\ 0 & \dots & 0 & 0 & -a_n a_{n-1} & 0 & 0 \end{pmatrix} \\ &= K(H_{n+1}). \end{aligned} \quad \square$$

Note, that in view of Proposition (2.5), the modified Kac-van Moerbeke equation (1.5) can be represented as the double bracket o.d.e.

$$(2.12) \quad \frac{d}{dt}H(t) = [H(t), [H(t), N(H(t))]], \quad H_0 := H(0) \in \text{Jac}_0(n).$$

We shall employ the following auxiliary lemma in the proof of Theorem (1.2).

(2.13). **Lemma.** *If $A = \mathcal{D}_1(a_1, a_2, \dots, a_{n-1})$ with $a_i \in \mathbb{R}$ for all $i = 1, \dots, n-1$, then*

$$(2.14) \quad [A, [A, N(A)]] = \mathcal{D}_1(-a_1a_2^2, -a_2a_3^2 + a_1^2a_2, \dots, -a_{n-2}a_{n-1}^2 + a_{n-3}^2a_{n-2}, a_{n-2}^2a_{n-1}),$$

where the mapping $N(\cdot)$ is defined in (2.3).

Proof. We have already shown in Proposition (2.5) that $K(A) = [A, N(A)]$. Let A_1 denote the upper triangular part and A_2 the lower triangular part of A , such that $A = A_1 + A_2$. Using an analogous decomposition for $K(A)$ we get $K_1(A) + K_2(A) = K(A)$. We abbreviate and simply write K_i for $K_i(A)$, $i \in \{1, 2\}$. We decompose

$$(2.15) \quad \begin{aligned} [A, K(A)] &= [A_1 + A_2, K_1 + K_2] \\ &= [A_1, K_1] + [A_1, K_2] + [A_2, K_1] + [A_2, K_2]. \end{aligned}$$

We first show by induction that $[A_1, K_1] = 0$. Indeed, if we denote with a superscript the size of a matrix and consider $A^n = \mathcal{D}_1(a_1, a_2, \dots, a_{n-1})$, i.e., $A_1^n = \mathcal{D}_{u,1}(a_1, a_2, \dots, a_{n-1})$ and $A_2^n = \mathcal{D}_{u,1}(a_1, a_2, \dots, a_{n-1})^\top$, we observe that for $n = 1$ we have $[A_1^n, K_1^n] = 0$. The induction step can be done as follows:

$$\begin{aligned} &[A_1^{n+1}, K_1^{n+1}] \\ &= A_1^{n+1}K_1^{n+1} - K_1^{n+1}A_1^{n+1} \\ &= \begin{pmatrix} & 0 \\ \boxed{A_1^n} & \vdots \\ & 0 \\ 0 & \dots & 0 & a_n \end{pmatrix} \begin{pmatrix} & 0 \\ \boxed{K_1^n} & \vdots \\ & a_{n-1}a_n \\ 0 & \dots & 0 & 0 \end{pmatrix} \\ &\quad - \begin{pmatrix} & 0 \\ \boxed{K_1^n} & \vdots \\ & a_{n-1}a_n \\ 0 & \dots & 0 & 0 \end{pmatrix} \begin{pmatrix} & 0 \\ \boxed{A_1^n} & \vdots \\ & 0 \\ 0 & \dots & 0 & a_n \end{pmatrix} \\ &= \begin{pmatrix} & 0 \\ \boxed{A_1^n K_1^n} & \vdots \\ & a_{n-2}a_{n-1}a_n \\ 0 & \dots & 0 & 0 \end{pmatrix} - \begin{pmatrix} & 0 \\ \boxed{K_1^n A_1^n} & \vdots \\ & a_{n-2}a_{n-1}a_n \\ 0 & \dots & 0 & 0 \end{pmatrix} \\ &= 0. \end{aligned}$$

Moreover, we have $A_1^\top = A_2$ and $K_1^\top = -K_2$. Therefore,

$$[A_2, K_2] = A_2 K_2 - K_2 A_2 = -A_1^\top K_1^\top + K_1^\top A_1^\top = (A_1 K_1 - K_1 A_1)^\top = [A_1, K_1]^\top.$$

Thus, $[A_2, K_2] = 0$ as well, and it remains to show (again by induction) that

$$\begin{aligned} & [A_2, K_1] + [A_1, K_2] \\ &= \mathcal{D}_1(-a_1 a_2^2, -a_2 a_3^2 + a_1^2 a_2, \dots, -a_{n-2} a_{n-1}^2 + a_{n-3}^2 a_{n-2}, a_{n-2}^2 a_{n-1}). \end{aligned}$$

As the next step, we claim that

$$[A_2^n, K_1^n] = \mathcal{D}_{u,1}(-a_1 a_2^2, -a_2 a_3^2 + a_1^2 a_2, \dots, -a_{n-2} a_{n-1}^2 + a_{n-3}^2 a_{n-2}, a_{n-2}^2 a_{n-1}).$$

The induction base is trivial. Then we have:

$$\begin{aligned} & [A_2^{n+1}, K_1^{n+1}] \\ &= A_2^{n+1} K_1^{n+1} - K_1^{n+1} A_2^{n+1} \\ &= \begin{pmatrix} & 0 \\ \boxed{A_2^n} & \vdots \\ & 0 \\ & 0 \\ 0 & \dots & a_n & 0 \end{pmatrix} \begin{pmatrix} & 0 \\ \boxed{K_1^n} & \vdots \\ & a_{n-1} a_n \\ & 0 \\ 0 & \dots & 0 & 0 \end{pmatrix} \\ &\quad - \begin{pmatrix} & 0 \\ \boxed{K_1^n} & \vdots \\ & a_{n-1} a_n \\ & 0 \\ 0 & \dots & 0 & 0 \end{pmatrix} \begin{pmatrix} & 0 \\ \boxed{A_2^n} & \vdots \\ & 0 \\ & 0 \\ 0 & \dots & a_n & 0 \end{pmatrix} \\ &= \begin{pmatrix} & 0 \\ \boxed{A_2^n K_1^n} & \vdots \\ & 0 \\ & 0 & a_{n-1}^2 a_n \\ 0 & \dots & 0 & 0 \end{pmatrix} \\ &\quad - \begin{pmatrix} & 0 \\ \boxed{K_1^n A_2^n} & \vdots \\ & 0 \\ & 0 \\ 0 & \dots & 0 & 0 \end{pmatrix} - \begin{pmatrix} & 0 \\ & \vdots \\ & 0 \\ & 0 \\ 0 & \dots & 0 & 0 \end{pmatrix} \\ &= \mathcal{D}_{u,1}(-a_1 a_2^2, -a_2 a_3^2 + a_1^2 a_2, \dots, -a_{n-1} a_n^2 + a_{n-2}^2 a_{n-1}, a_{n-1}^2 a_n). \end{aligned}$$

Again, in view of $A_1^\top = A_2$ and $K_1^\top = -K_2$, we obtain

$$[A_1, K_2] = A_1 K_2 - K_2 A_1 = -A_2^\top K_1^\top + K_1^\top A_2^\top = (A_2 K_1 - K_1 A_2)^\top = [A_2, K_1]^\top.$$

Thus,

$$[A_1^n, K_2^n] = \mathcal{D}_{u,1}(a_1 a_2^2, a_2 a_3^2 + a_1^2 a_2, \dots, -a_{n-2} a_{n-1}^2 + a_{n-3}^2 a_{n-2}, a_{n-2}^2 a_{n-1})^\top.$$

According to (2.15), for $A = \mathcal{D}_1(a_1, a_2, \dots, a_{n-1})$ we have

$$[A, K(A)] = \mathcal{D}_1(-a_1 a_2^2, -a_2 a_3^2 + a_1^2 a_2, \dots, -a_{n-2} a_{n-1}^2 + a_{n-3}^2 a_{n-2}, a_{n-2}^2 a_{n-1}),$$

which completes the proof. \square

(2.16). **Lemma.** For $A, B \in \text{Sym}(n)$,

$$[A, [A, B]] = 0 \text{ if and only if } [A, B] = 0.$$

Proof. The “if” part is trivial. To prove the “only if” part, suppose that $[A, [A, B]] = 0$. This implies $B[A, [A, B]] = 0$. Using the techniques in [Helmke and Moore, 1994, p. 49], we compute

$$\begin{aligned} 0 &= \text{tr}(B[A, [A, B]]) \\ &= \text{tr}(B(A^2B - 2ABA + BA^2)) \\ &= \text{tr}(BA^2B - 2BABA + B^2A^2) \\ &= \text{tr}(BA^2B - BABA - ABAB + AB^2A) \\ &= \text{tr}((BA - AB)(AB - BA)) \\ &= \text{tr}([B, A][A, B]) \\ &= \text{tr}([A, B]^\top [A, B]) \\ &= \|[A, B]\|^2, \end{aligned}$$

which immediately gives $[A, B] = 0$. \square

(2.17). **Lemma.** Consider the continuous function

$$(2.18) \quad M(H_0) \ni H \mapsto f(H) := -\frac{1}{4}\|H - N(H)\|^2 + \frac{1}{4}\|N(H)\|^2 \in \mathbb{R}.$$

With respect to the o.d.e.

$$(2.19) \quad \frac{d}{dt}H(t) = [H(t), [H(t), N(H(t))]], \quad H(0) = H_0 \in \text{Sym}(n),$$

the time derivative of $f(H(\cdot))$ is given by

$$(2.20) \quad \frac{d}{dt}f(H(t)) = \|[H(t), N(H(t))]\|^2.$$

Proof. We start by simplifying the function

$$\begin{aligned} f(H(t)) &= -\frac{1}{4}\|H(t) - N(H(t))\|^2 + \frac{1}{4}\|N(H(t))\|^2 \\ &= -\frac{1}{4}\text{tr}(H(t)H(t) - H(t)N(H(t)) - N(H(t))H(t) + N(H(t))N(H(t))) \\ &\quad + \frac{1}{4}\|N(H(t))\|^2 \\ &= -\frac{1}{4}\|H(t)\|^2 + \frac{1}{2}\text{tr}(N(H(t))H(t)) - \frac{1}{4}\|N(H(t))\|^2 + \frac{1}{4}\|N(H(t))\|^2 \\ &= -\frac{1}{4}\|H(t)\|^2 + \frac{1}{2}\text{tr}(N(H(t))H(t)). \end{aligned}$$

Since $H(t) \in M(H_0)$ by Proposition (2.1), $\|H(t)\|$ is constant for all $t \geq 0$. We calculate the derivative of f along the trajectories of (2.19) as follows:

$$\begin{aligned} \frac{d}{dt}f(H(t)) &= \frac{1}{2}\text{tr}\left(\left(\frac{d}{dt}N(H(t))\right)H(t) + N(H(t))\dot{H}(t)\right) \\ &= \frac{1}{2}\text{tr}\left(\left(\frac{d}{dt}N(H(t))\right)H(t)\right) \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2} \operatorname{tr} (N(H(t)) [H(t), [H(t), N(H(t))]]) \\
& = \frac{1}{2} \operatorname{tr} \left(\left(\frac{d}{dt} N(H(t)) \right) H(t) \right) \\
& \quad + \frac{1}{2} \operatorname{tr} \left(\underbrace{[N(H(t)), H(t)]}_{[H(t), N(H(t))]^\top} [H(t), N(H(t))] \right) \\
(2.21) \quad & = \frac{1}{2} \operatorname{tr} \left(\left(\frac{d}{dt} N(H(t)) \right) H(t) \right) + \frac{1}{2} \|[H(t), N(H(t))]\|^2,
\end{aligned}$$

where, at the third equality, we employed the fact [Bernstein, 2009, p. 162] that for $A, B, C \in \mathbb{R}^{n \times n}$, $\operatorname{tr}(A[B, C]) = \operatorname{tr}([A, B]C)$. Therefore, it remains to show that

$$\operatorname{tr} \left(\left(\frac{d}{dt} N(H(t)) \right) H(t) \right) = \|[H(t), N(H(t))]\|^2.$$

Note that

$$\frac{d}{dt} N(H(t)) = N(\dot{H}(t))$$

since N is linear, and since from (2.21) it follows that $\operatorname{tr}(N(H(t))\dot{H}(t)) = \|[H(t), N(H(t))]\|^2$, our proof will be complete if we show that

$$(2.22) \quad \operatorname{tr}(N(\dot{H}(t))H(t)) = \operatorname{tr}(N(H(t))\dot{H}(t)).$$

To this end, employing the expansion of N in (2.4), we see that

$$\begin{aligned}
\operatorname{tr}(N(\dot{H}(t))H(t)) &= \operatorname{tr} \left(\sum_{i=1}^{n-1} (i-2) (E_i \dot{H}(t) E_{i+1} + E_{i+1} \dot{H}(t) E_i) H(t) \right) \\
&= \sum_{i=1}^{n-1} (i-2) \operatorname{tr} ((E_i \dot{H}(t) E_{i+1} + E_{i+1} \dot{H}(t) E_i) H(t)) \\
&= \sum_{i=1}^{n-1} (i-2) (\operatorname{tr}(E_i \dot{H}(t) E_{i+1} H(t)) + \operatorname{tr}(E_{i+1} \dot{H}(t) E_i H(t))) \\
&= \sum_{i=1}^{n-1} (i-2) (\operatorname{tr}(E_{i+1} H(t) E_i \dot{H}(t)) + \operatorname{tr}(E_i H(t) E_{i+1} \dot{H}(t))) \\
&= \sum_{i=1}^{n-1} (i-2) \operatorname{tr} ((E_i H(t) E_{i+1} + E_{i+1} H(t) E_i) \dot{H}(t)) \\
&= \operatorname{tr}(N(H(t))\dot{H}(t)),
\end{aligned}$$

which establishes (2.22), and completes the proof. \square

(2.23). **Lemma.** *The o.d.e.*

$$(2.24) \quad \frac{d}{dt} H(t) = [H(t), [H(t), N(H(t))]], \quad H(0) = H_0 \in \operatorname{Jac}_0(n),$$

has a finite number of equilibrium points on $\operatorname{Jac}_0(n)$ that are isospectral to H_0 .

Proof. In view of Proposition (2.1) and Lemma (2.13), $H(t) \in M_J(H_0)$ for all solutions of (2.24) and for all $t \geq 0$. Moreover, for $H_0 \in \text{Jac}_0(n)$ we have $M_J(H_0) \subseteq \text{Jac}_0(n)$. By Lemma (2.16),

$$(2.25) \quad \begin{aligned} E &:= \{H(t) \in M_J(H_0) \mid \|[H(t), N(H(t))]\|^2 = 0\} \\ &= \{H(t) \in M_J(H_0) \mid [H(t), N(H(t))] = 0\} \end{aligned}$$

is the set of all equilibrium points of (2.24) on $\text{Jac}_0(n)$ that are isospectral to H_0 . Let

$$\tilde{H} = \mathcal{D}_1(a_1, a_2, a_3, a_4, \dots, a_{n-2}, a_{n-1}) \in \text{Jac}_0(n).$$

At this point it is crucial to recall that according to our definition Jacobi matrices have distinct eigenvalues. We treat the case of n even and n odd separately:

n even

Consider the set of matrices

$$\tilde{E} := \{\tilde{H} \in \text{Jac}_0(n) \mid [\tilde{H}, N(\tilde{H})] = 0\}.$$

In view of (2.7), the only possibility for \tilde{H} to lie in \tilde{E} is if $a_i = 0$ for all i even and $a_i \neq 0$, for all $i \in \{1, 3, \dots, n-1\}$ with $a_i \neq a_j$ for all $i \neq j$, such that

$$\tilde{H}_{\tilde{E}} = \mathcal{D}_1(a_1, 0, a_3, 0, \dots, 0, a_{n-1}).$$

Note that $\tilde{H}_{\tilde{E}}$ has the spectrum $\sigma(\tilde{H}_{\tilde{E}}) = \{\pm a_i \mid i \in \{1, 3, \dots, n-1\}\}$ containing only distinct eigenvalues. Now E as defined in (2.25) is a subset of \tilde{E} satisfying the isospectral conditions; it is the restriction of \tilde{E} to the set of zero-diagonal Jacobi matrices isospectral to H_0 , i.e., $E = \tilde{E}|_{M_J(H_0)}$. Considering all the possible permutations of the a_i for $i \in \{1, 3, \dots, n-1\}$, the set E contains $(\frac{n}{2})!$ equilibrium points on $\text{Jac}_0(n)$ that are isospectral to H_0 .

n odd

First of all, since $\tilde{H} = \mathcal{D}_1(a_1, a_2, a_3, a_4, \dots, a_{n-2}, a_{n-1})$ is a (zero-diagonal) Jacobi matrix, we need to evoke the fact [Penskoi, 2008] that its spectrum has the form

$$(2.26) \quad \sigma(\tilde{H}) = \{0, \pm\lambda_1, \pm\lambda_2, \dots, \pm\lambda_{\frac{n-1}{2}}\}, \quad \text{where } \lambda_i \neq \lambda_j \text{ for all } i \neq j.$$

Moreover, if we consider

$$\tilde{E} := \{\tilde{H} \in \text{Jac}_0(n) \mid [\tilde{H}, N(\tilde{H})] = 0\},$$

in view of (2.7), \tilde{H} must satisfy

$$(2.27) \quad a_i a_{i+1} = 0 \quad \text{for all } i = 1, \dots, n-2$$

in order to lie in \tilde{E} . Furthermore, in view of (2.26) and (2.27) \tilde{H} has to be a block-diagonal matrix containing $\frac{n-1}{2}$ blocks of the form $\begin{pmatrix} 0 & a_j \\ a_j & 0 \end{pmatrix}$ with $a_j \neq 0$ and one 1×1 block containing a zero, where the block entries are distinct (since the eigenvalues of \tilde{H} need to be distinct). Accordingly, there are $\frac{n+1}{2}$ possibilities to place the 1×1 block in \tilde{H} . As above $E = \tilde{E}|_{M_J(H_0)}$ and considering all possible permutations of the blocks we get that E contains $(\frac{n+1}{2})(\frac{n-1}{2})!$ equilibrium points on $\text{Jac}_0(n)$ that are isospectral to H_0 . \square

(2.28). **Lemma.** Let $g \in \mathcal{C}(\mathbb{R}_{\geq 0}, \mathbb{R})$ and suppose that $\lim_{t \rightarrow \infty} g(t)$ exists. If

$$\lim_{t \rightarrow \infty} |g(t)| = \eta, \quad \text{for some } \eta > 0$$

and

$$\lim_{t \rightarrow \infty} \int_0^t g(s) ds = -\infty,$$

then

$$\lim_{t \rightarrow \infty} g(t) < 0.$$

Proof. First, note that $\lim_{t \rightarrow \infty} |g(t)| = \eta > 0$ implies that there exists $T_0 > 0$ such that for all $t > T_0$, $g(t) \neq 0$. Therefore, by continuity, the sign of $g(t)$ is the same for all $t > T_0$. Suppose $\lim_{t \rightarrow \infty} g(t) \geq 0$. If the limit is equal to 0, then its absolute value would have to converge to 0 as well, which is a contradiction. Therefore, $\lim_{t \rightarrow \infty} g(t) > 0$. But then $g(t) = |g(t)|$ for all $t > T_0$ and $\lim_{t \rightarrow \infty} g(t) = \eta$. Then, there exists $T_1 > 0$ such that for all $t > T_1$, $g(t) > \frac{\eta}{2}$. Without loss of generality we assume that $T_1 > T_0$ (otherwise put $T'_1 := \max\{T_0, T_1\}$). By continuity, g is bounded on the interval $[0, T_1]$ by some constant C . Therefore,

$$-\infty < -T_1 C \leq \int_0^{T_1} g(s) ds \leq T_1 C < \infty.$$

Now we see that for $t > T_1$

$$\int_0^t g(s) ds = \int_0^{T_1} g(s) ds + \int_{T_1}^t g(s) ds \geq -T_1 C + \int_{T_1}^t \frac{\eta}{2} ds \rightarrow \infty - T_1 C = \infty$$

for $t \rightarrow \infty$. Therefore, $\lim_{t \rightarrow \infty} \int_0^t g(s) ds = \infty$ which is a contradiction. \square

Proof of Theorem (1.2). In view of Lemma (2.13) it follows that the right-hand side of (1.5) is a symmetric tridiagonal matrix with zero diagonal entries given by (2.14). The fact that (1.5) is isospectral is an immediate consequence of Proposition (2.1). Therefore, the flow of (1.5) evolves on the set of zero-diagonal Jacobi matrices isospectral to H_0 , i.e., $H(t) \in M_J(H_0)$ for all $t \geq 0$. This settles the claim in (i).

In order to show (ii), note that in view of (i), $H(t) \in M_J(H_0)$ for all $t \geq 0$. Since $M_J(H_0)$ is known to be a compact manifold [Penskoi, 2008, Proposition 1.2], $H(t)$ exists for all $t \geq 0$. By Lemma (2.16) we see that the set of equilibrium points \bar{H} of (1.5) is given by $\{\bar{H} \in M_J(H_0) \mid [\bar{H}, N(\bar{H})] = 0\}$. To show that $H(t)$ approaches the set of equilibrium points, consider the function

$$(2.29) \quad M_J(H_0) \ni H \mapsto f(H) := -\frac{1}{4} \|H(t) - N(H(t))\|^2 + \frac{1}{4} \|N(H(t))\|^2 \in \mathbb{R}.$$

According to Lemma (2.17),

$$(2.30) \quad \frac{d}{dt} f(H(t)) = \|[H(t), N(H(t))]\|^2 \geq 0.$$

We invoke the Krasovskij-LaSalle's Invariance Principle [Khalil, 2002, Theorem 4.4], [Vidyasagar, 2002, p. 178]: First, we define the set

$$E := \left\{ H \in M_J(H_0) \mid [H, N(H)] = 0 \right\}.$$

By Lemma (2.16), E coincides with the set of all equilibrium points of (1.5). Therefore, E is an invariant set with respect to (1.5). Second, recall that $M_J(H_0)$ is a compact set. Therefore, by the Krasovskij-LaSalle's Invariance Principle, every solution $(H(t))_{t \geq 0}$ starting in $M_J(H_0)$ approaches the set of equilibrium points E asymptotically, which proves the claim in (ii).

To show property (iii) note that we have already shown in part (ii) that every solution of (1.5) approaches the set of equilibrium points asymptotically. However, according to Lemma (2.23), the number of equilibrium points is finite. By continuity of trajectories, therefore, $(H(t))_{t \geq 0}$ converges to a single equilibrium point, i.e., $\lim_{t \rightarrow \infty} H(t)$ exists, which proves the claim in (iii).

To prove (iv), consider first the o.d.e.

$$(2.31) \quad \frac{dx(t)}{dt} = x(t)g(t), \quad x(0) = x_0,$$

where $x(t) \in \mathbb{R}$, $g \in \mathcal{C}(\mathbb{R}_{\geq 0}, \mathbb{R})$. Suppose that $x_0 \neq 0$. The unique solution ([Khalil, 2002]) to (2.31) is given by

$$x(t) = x_0 \exp \left(\int_0^t g(s) ds \right), \quad t \geq 0,$$

and it follows at once that

$$(2.32) \quad \lim_{t \rightarrow \infty} x(t) = 0 \quad \text{implies} \quad \lim_{t \rightarrow \infty} \int_0^t g(s) ds = -\infty.$$

We consider the components of the o.d.e. (1.5):

$$\begin{aligned} \frac{da_1(t)}{dt} &= -a_1(t)a_2^2(t), \\ \frac{da_i(t)}{dt} &= a_i(t) (a_{i-1}^2(t) - a_{i+1}^2(t)), \quad i = 2, \dots, n-2, \\ \frac{da_{n-1}(t)}{dt} &= a_{n-1}(t)a_{n-2}^2(t), \end{aligned}$$

and distinguish two different cases depending on the parity of n :

n even

For each $i = 1, \dots, \frac{n}{2} - 1$, define the function $g_{2i} \in \mathcal{C}(\mathbb{R}_{\geq 0}, \mathbb{R})$ by $g_{2i}(t) := a_{2i-1}^2(t) - a_{2i+1}^2(t)$. With the definition of g_{2i} , the even-numbered components of the o.d.e. (1.5) can be represented as

$$\frac{da_{2i}(t)}{dt} = a_{2i}(t) (a_{2i-1}^2(t) - a_{2i+1}^2(t)) = a_{2i}(t)g_{2i}(t).$$

This o.d.e. is of the form (2.31), and by Lemma (2.23) and properties (ii) and (iii) of Theorem (1.2), it follows that $\lim_{t \rightarrow \infty} a_{2i}(t)$ exists and is equal to zero. By (2.32),

$$(2.33) \quad \lim_{t \rightarrow \infty} \int_0^t g_{2i}(s) ds = -\infty.$$

Independently of the preceding steps, recall that the eigenvalues of any Jacobi matrix are distinct. From the definition of g_{2i} it now follows that

$$(2.34) \quad \text{there exists } \eta_{2i} > 0 \text{ such that } \lim_{t \rightarrow \infty} |g_{2i}(t)| = \eta_{2i}.$$

In view of (2.33) and (2.34), Lemma (2.28) and property (iii) of Theorem (1.2) lead to

$$(2.35) \quad \lim_{t \rightarrow \infty} a_{2i-1}^2(t) < \lim_{t \rightarrow \infty} a_{2i+1}^2(t) \quad \text{for all } i = 1, \dots, \frac{n}{2} - 1.$$

n odd

We introduce the function $f(t) := a_{n-1}^2(t)$, $t \geq 0$, where $a_{n-1}(\cdot)$ is the solution of the

final component of (1.5). The derivative of f given by $\frac{df(t)}{dt} = 2a_{n-1}^2(t)a_{n-2}^2(t) \geq 0$ shows that f is monotonically non-decreasing. Since $a_{n-1}(0) \neq 0$, it follows from the fact that f is monotonically non-decreasing, that $\lim_{t \rightarrow \infty} a_{n-1}(t) \neq 0$. In view of Lemma (2.23), and properties (ii) and (iii) of Theorem (1.2), it follows that $\lim_{t \rightarrow \infty} a_{2i-1}(t)$ exists and is equal to zero for all $i = 1, \dots, \frac{n-1}{2}$. As in the case of n even, we define, for each $i = 2, \dots, \frac{n-1}{2}$, a function $g_{2i-1} \in \mathcal{C}(\mathbb{R}_{\geq 0}, \mathbb{R})$ by $g_{2i-1}(t) := a_{2i-2}^2(t) - a_{2i}^2(t)$. With this definition of g_{2i-1} , we see that the odd-numbered components (greater than one) of the o.d.e. (1.5) can be represented as

$$\frac{da_{2i-1}(t)}{dt} = a_{2i-1}(t) (a_{2i-2}^2(t) - a_{2i}^2(t)) = a_{2i-1}(t)g_{2i-1}(t).$$

This o.d.e. is of the form (2.31), and since we know that $\lim_{t \rightarrow \infty} a_{2i-1}(t) = 0$, it follows from (2.32) that

$$(2.36) \quad \lim_{t \rightarrow \infty} \int_0^t g_{2i-1}(s) ds = -\infty.$$

Since the eigenvalues of any Jacobi matrix are distinct, by definition of g_{2i-1} we see that

$$(2.37) \quad \text{there exists } \eta_{2i-1} > 0 \text{ such that } \lim_{t \rightarrow \infty} |g_{2i-1}(t)| = \eta_{2i-1}.$$

In view of (2.36) and (2.37), Lemma (2.28) and property (iii) of Theorem (1.2) lead to

$$(2.38) \quad \lim_{t \rightarrow \infty} a_{2i-2}^2(t) < \lim_{t \rightarrow \infty} a_{2i}^2(t), \quad \text{for all } i = 2, \dots, \frac{n-1}{2}.$$

Finally, note that independently of the parity of n , for each component of the o.d.e. (1.5), zero is always an equilibrium point. Therefore, if $a_i(0) \neq 0$, then $a_i(\cdot)$ cannot take the value 0, and this in turn implies that if $\lim_{t \rightarrow \infty} a_i(t) \neq 0$, then $\text{sgn}(a_i(0)) = \text{sgn}(\lim_{t \rightarrow \infty} a_i(t))$ for all $i = 1, \dots, n-1$. This completes the proof. \square

§ 3. EXAMPLES

We illustrate Theorem (1.2) with three numerical examples. We solve the o.d.e. (1.5) numerically and plot the upper-diagonal entries of $H(t)$ as functions of time for a chosen initial condition. The figures show that the solution of (1.5) converges rather quickly to an equilibrium.

(3.1). *Example.* We start with a 4×4 example. Consider an initial condition

$$(3.2) \quad H(0) = H_0 = \mathcal{D}_1(5, -6, -2)$$

with the spectrum

$$\sigma(H_0) = \{ \pm 1.26, \pm 7.96 \}.$$

By solving the o.d.e. (1.5) numerically with this initial condition we see again that the transient behavior vanishes well before 1 second of simulation. At $T = 1$ s we have

$$H(T) = \mathcal{D}_1(1.26, 0, -7.96).$$

Figure 1 shows the evolution of the super-diagonal components of $H(t)$ against time t .




FIGURE 1. Flow of o.d.e. (1.5) for the initial condition (3.2).

(3.3). *Example.* Consider the following initial condition of size 10×10

$$(3.4) \quad H(0) = H_0 = \mathcal{D}_1(-3, 10, 1, -2, -6, -11, 5, 6, 12),$$

having the spectrum $\sigma(H_0) = \{\pm 0.21, \pm 2.71, \pm 10.48, \pm 12.34, \pm 14.36\}$. By solving the o.d.e. (1.5) numerically, we see that the transient behavior vanishes well before 1 second of simulation. At $T = 1s$ we have

$$H(T) = \mathcal{D}_1(-0.21, 0, 2.71, 0, -10.48, 0, 12.34, 0, 14.36).$$

Figure 2 shows the evolution of the super-diagonal components of $H(t)$ against time t .

(3.5). *Example.* Consider the following initial condition of size 29×29

$$(3.6) \quad H(0) = H_0 = \mathcal{D}_1(-6, 7, -8, 2, -13, 7, -12, 7, -2, 9, 2, -4, 2, \\ 4, 6, -15, -7, 11, -7, 9, 9, 15, 1, 5, -3, 11, -1, -3),$$

having the spectrum $\sigma(H_0) = \{0, \pm 2.81, \pm 2.98, \pm 4.17, \pm 4.66, \pm 4.84, \pm 6.26, \pm 9.29, \pm 10.84, \pm 11.53, \pm 11.83, \pm 12.48, \pm 17.11, \pm 17.98, \pm 18.85\}$. By solving the o.d.e. (1.5) numerically, we see that the transient behavior vanishes well before 1 second of simulation. At $T = 1s$ we have

$$H(T) = \mathcal{D}_1(0, 2.81, 0, 2.98, 0, 4.17, 0, 4.66, 0, 4.84, 0, -6.26, 0, 9.29, 0, -10.84, 0, \\ 11.53, 0, 11.83, 0, 12.48, 0, 17.11, 0, 17.98, 0, -18.85).$$

Figure 2 shows the evolution of the super-diagonal components of $H(t)$ against time t .




FIGURE 2. Flow of o.d.e. (1.5) for the initial condition (3.4).




FIGURE 3. Flow of o.d.e. (1.5) for the initial condition (3.6).

§ 4. CONCLUSIONS AND FUTURE DIRECTION

We presented a matrix-valued isospectral ordinary differential equation that asymptotically block-diagonalizes a finite-dimensional zero-diagonal Jacobi matrix employed as its initial condition. We demonstrated that this o.d.e. can be represented as a double bracket equation, thus establishing a connection to [Brockett, 1991], and we have proved certain new key properties of this o.d.e. by system-theoretic techniques. In particular that the limit is block-diagonal and the blocks of the limit point are sorted by increasing magnitude of the corresponding eigenvalue.

The domain of the o.d.e. (1.5) can be expanded to the set of real symmetric matrices $\text{Sym}(n)$. Since $M(H_0)$ for $H_0 \in \text{Sym}(n)$ is again a compact manifold [Helmke and Moore, 1994], assertions (i) and (ii) of Theorem (1.2) hold also for the symmetric case, and the proof proceeds analogously. Extensive simulations lead us to conjecture that the solutions converge asymptotically to block diagonal matrices, as in the case of zero-diagonal Jacobi matrices employed as initial conditions. However, a proof for this conjecture is still an open problem; the primary technical difficulty arises from the fact that in contrast to the case of zero-diagonal Jacobi matrices, in this case there exist infinitely many equilibrium points.

REFERENCES

- Akhiezer, N. I. [1965], *The Classical Moment Problem and Some Related Questions in Analysis*, Hafner Publishing Co., New York. Translated by N. Kemmer.
- Bernstein, D. S. [2009], *Matrix Mathematics*, 2 edn, Princeton University Press.
- Bloch, A. [1990], *Mathematical Developments Arising from Linear Programming*, Contemporary Mathematics, American Mathematical Society, Providence, R.I.
- Brockett, R. W. [1991], ‘Dynamical systems that sort lists, diagonalize matrices, and solve linear programming problems’, *Linear Algebra and its Applications* **146**, 79 – 91.
- Favard, J. [1935], ‘Sur les polynomes de Tchebicheff’, *Comptes Rendus de l’Académie des Sciences* **200**, 2052–2053.
- Faybusovich, L. [1992], ‘Interior-point methods and entropy’, *IEEE Decision and Control* **3**, 2094 – 2095.
- Helmke, U. and Moore, J. B. [1994], *Optimization and Dynamical Systems*, Communications and Control Engineering Series, Springer-Verlag London Ltd., London. With a foreword by R. Brockett.
- Kac, M. and van Moerbeke, P. [1975], ‘On an explicitly soluble system of nonlinear differential equations related to certain Toda lattices’, *Advances in Mathematics* **16**, 160–169.
- Khalil, H. K. [2002], *Nonlinear Systems*, 3 edn, Prentice Hall, Upper Saddle River, New Jersey.
- Lax, P. D. [1968], ‘Integrals of nonlinear equations of evolution and solitary waves’, *Communications on Pure and Applied Mathematics* **21**, 467–490.
- Moore, J. B., Mahony, R. E. and Helmke, U. [1994], ‘Numerical gradient algorithms for eigenvalue and singular value calculations’, *SIAM Journal on Matrix Analysis and Applications* **15**, 881–902.

- Moser, J. [1975], ‘Finitely many mass points on the line under the influence of an exponential potential – an integrable system’, *Dynamical Systems, Theory and Applications* **38**, 467–497.
- Penskoi, A. [2008], ‘Integrable systems and the topology of isospectral manifolds’, *Theoretical and Mathematical Physics* **155**, 627–632.
- Simon, B. [2005], *Orthogonal Polynomials on the Unit Circle, Part 1: Classical Theory*, Vol. 54.1 of *Colloquium Publication*, American Mathematical Society, Providence, R.I.
- Szegő, G. [1959], *Orthogonal Polynomials*, American Mathematical Society Colloquium Publications, Vol. 23. Revised ed, American Mathematical Society, Providence, R.I.
- Vidyasagar, M. [2002], *Nonlinear Systems Analysis*, Vol. 42, SIAM Classics in Applied Mathematics, Philadelphia.
- Watkins, D. S. [2005], ‘Product eigenvalue problems’, *SIAM Review* **47**, 3–40.

(T. Sutter, and J. Lygeros) AUTOMATIC CONTROL LABORATORY, ETL I28, ETH ZÜRICH, PHYSIKSTRASSE 3, 8092 ZÜRICH, SWITZERLAND

E-mail address, T. Sutter and J. Lygeros: sutttert@student.ethz.ch, jlygeros@control.ee.ethz.ch

URL: <http://control.ee.ethz.ch>

(D. Chatterjee) SYSTEMS & CONTROL ENGINEERING, IIT-BOMBAY, POWAI, 400 076 MUMBAI, INDIA

E-mail address, D. Chatterjee: chatterjee@sc.iitb.ac.in

URL: <http://www.sc.iitb.ac.in/~chatterjee>

(F. A. Ramponi) UNIVERSITÀ DEGLI STUDI DI BRESCIA, DIPARTIMENTO DI INGEGNERIA DELL’INFORMAZIONE, VIA BRANZE 38, 25123 BRESCIA, ITALY

E-mail address, F. A. Ramponi: federico.ramponi@ing.unibs.it