

THE CHERN CHARACTER OF CERTAIN INFINITE RANK BUNDLES ARISING IN GAUGE THEORY

JOUKO MICKELSSON

Department of Mathematics and Statistics, University of
Helsinki, and Royal Institute of Technology, Stockholm

ABSTRACT: A cocycle $\Omega : P \times G \rightarrow H$ taking values in a Lie group H for a free right action of G on P defines a principal bundle Q with the structure group H over $X = P/G$. The Chern character of a vector bundle associated to Q defines then characteristic classes on X . This observation becomes useful in the case of infinite dimensional groups. It typically happens that a representation of G is not given by linear operators which differ from the identity by a trace-class operator. For this reason the Chern character of a vector bundle associated to the principal fibration $P \rightarrow P/G$ is ill-defined. But it may happen that the Lie algebra representations of the group H are given in terms of trace-class operators and therefore the Chern character is well-defined; this observation is useful especially if the map $g \mapsto \Omega(p; g)$ is a homotopy equivalence on the image for any $p \in P$. We apply this method to the case $P = \mathcal{A}$, the space of gauge connections in a finite-dimensional vector bundle, and $G = \mathcal{G}$ is the group of (based) gauge transformations. The method for constructing the appropriate cocycle Ω comes from ideas in quantum field theory, used to define the renormalized gauge currents in a Fock space.

1. INTRODUCTION

In gauge theory one is led to study infinite rank vector bundles where the structure group is the group U^p , resp. U_p , (or its subgroup), consisting of unitary operators g in a complex Hilbert space H such that $g - 1$ is in L_p in the even case, resp. $[\epsilon, g] \in L_p$ in the odd case, [MR]. Here ϵ defines a polarization in H , and even/odd refers to the parity of the dimension of the underlying compact physical space M . L_p is the Schatten ideal of bounded operators T such that $|T|^p$ is trace-class, $p \geq 1$. The index p depends on the dimensionality of M , $p > n = \dim M$. The reason for the choice of p is related to the fact that the Green's function D^{-1} of a Dirac operator D is in L_p for $p > n$.

Starting from the even case, one can pose the more general question how to define the Chern character for U^p bundles over some space X . Typically, X is the moduli space of gauge connections. The problem in the infinite rank setting is that the powers \mathcal{F}^k of the curvature are not always trace-class operators. That would be the case in the even case if $k \geq p$.

On the other hand, all the groups U^p are homotopy equivalent to U^1 , due to a theorem by Richard Palais [P]. Concretely, the U^p bundles arise from principal gauge group bundles through the embedding of the group of gauge transformations \mathcal{G} to U^p . The group U^p can be replaced by U^1 using a cocycle $\Omega : P \times \mathcal{G} \rightarrow U^1$. For each point $A \in P$ the map $\mathcal{G} \rightarrow U^1$ defined by $g \mapsto \Omega(A; g)$ is homotopic to the embedding $\mathcal{G} \subset U^p$, in the topology of the larger group U^p . This method has been previously used in the quantization of chiral fermions in background gauge fields and in computation of hamiltonian anomalies (Schwinger terms), [M].

In order to define the Chern character one needs a connection. Normally, when a vector bundle is associated to a principal bundle P over X the connection form acting on sections of the vector bundle is obtained from the principal bundle connection by the action of the Lie algebra representation ρ in the model fiber of the vector bundle. However, in the case when the associated vector bundle is defined by a cocycle Ω there is a complication. There are two candidates for a connection. The first comes from the canonical 1-form defined by the derivative ω of Ω with respect to the second argument, and computing $\omega(p; \theta)$ with θ a connection 1-form for the bundle $P \rightarrow X$, is a natural candidate to replace ρ , but this does not quite define a connection. It does not transform properly under the right translations by \mathcal{G} . The second candidate is determined from the 'renormalization process' defining the cocycle Ω . That does define a covariant differentiation in the associated vector bundle. However, the connection form takes values in a larger Lie algebra than $\mathfrak{u}^1 = Lie(U^1)$. It turns out that we can still use this latter 1-form to compute the characteristic classes. The reason for this is that the gauge variations of the corresponding Chern-Simons forms are given by traces of trace class operator valued forms, due to the fact that the transition functions take values in \mathfrak{u}^1 . Pairing Chern classes with singular homology classes and using Stokes' theorem the integral of a Chern class against a singular cycle becomes a sum of integrals of the gauge variations of Chern-Simons form over the faces of the individual singular simplices and thus are well-defined.

2. CHERN CHARACTER FOR U^1 BUNDLES AND FAMILIES OF DIRAC OPERATORS

Let M be a compact oriented connected spin manifold and G a compact Lie group and \mathfrak{g} its Lie algebra. Denote by \mathcal{A} the space of smooth \mathfrak{g} valued 1-forms on M , the gauge potentials in a trivial G bundle over M . In the following we could generalize the discussion to nontrivial G bundles, but I want to avoid unnecessary technical complications which actually are irrelevant for the main ideas below. Denote by \mathcal{G} the group of gauge transformations (bundle automorphisms which fix the base), which in the case of a trivial bundle can be identified as the group of smooth maps from M to G .

Let us first consider the case when the dimension of M is odd. In quantum field theory one wants to associate a bundle of fermionic Fock spaces to \mathcal{A} , such that the quantized Dirac operators \hat{D}_A with $A \in \mathcal{A}$, transform covariantly under the group of gauge transformations. It turns out that in the case of chiral Dirac operators

(acting on massless Weyl fermions) there is generically an obstruction to this, in the form of a *commutator anomaly* (Schwinger terms) which means that instead of \mathcal{G} one has to construct certain group extension $\hat{\mathcal{G}}$ which acts on sections of the Fock bundle, [M]. There is a technical difficulty in this construction: The group $\hat{\mathcal{G}}$ has an unitary representation in the Fock space only when $\dim M = 1$.

When the dimension is higher than one the representations have to be understood in a generalized sense. Instead of a group homomorphism $\hat{\mathcal{G}} \rightarrow U(\mathcal{H})$, where \mathcal{H} is the fermionic Fock space, one has to construct first a cocycle $\Omega : \mathcal{A} \times \mathcal{G} \rightarrow U(H)$, where H is the fermionic 1-particle space, such that the values of Ω are in the restricted unitary group $U_{res}(H) \subset U(H)$. The restricted unitary group consists of unitaries such that the off-diagonal blocks with respect to a fixed polarization of H are Hilbert-Schmidt operators. In that case the operators $\Omega(A; g)$ can be represented by unitaries $\hat{\Omega}(A; g) \in U(\mathcal{H})$ which then define a group extension of \mathcal{G} .

In the present article I want to consider the problems arising in the case of an even dimensional manifold M from the point of view of families index theory. In short, the problem is the following: Consider \mathcal{A} as a principal bundle over the moduli space \mathcal{A}/\mathcal{G} with fiber \mathcal{G} . As \mathcal{G} I take now the group of smooth *based maps* from M to G . Based means that we fix a point $x_0 \in M$ and consider maps g such that $g(x_0) = 1$, the neutral element in G . This guarantees that $X = \mathcal{A}/\mathcal{G}$ has no singularities, it is a Frechet manifold. Given a finite dimensional representation $\rho : G \rightarrow \text{Aut}(V)$ we can construct a Hilbert bundle \mathcal{E} over \mathcal{A}/\mathcal{G} through the natural action of \mathcal{G} on sections of $S(M) \otimes E$ where E is a (trivial) complex vector bundle over M with fiber V and $S(M)$ is the spin bundle over M .

Any Hilbert bundle with structure group $U(H)$ (in the norm topology) is topologically trivial by Kuiper's theorem, [K]. However, in our case there is additional geometrical structure. The bundle has \mathcal{G} as its structure group and we can define a connection on \mathcal{E} taking values in the Lie algebra of \mathcal{G} . The problem next is to define the Chern character of the bundle in the usual way as $\text{tr} e^{\mathcal{F}/2\pi i}$, where \mathcal{F} is the curvature of the connection. As it stands, the Chern character is ill-defined since the trace of the curvature diverges. One way to avoid the divergence is to replace the curvature by the curvature of an appropriate Quillen superconnection, [Q]. This method was worked through in detail by J. - M. Bismut, [B], for the derivation of the local Atiyah-Singer index theorem for families of Dirac operators, [AS1]. In this article I want to propose a different solution, very close to the idea of renormalization in [M] in the odd dimensional case.

Let us first assume that the Dirac index of D_A is zero. Then we can fix an invertible hermitean operator D_0 such that $D_A - D_0$ is bounded for any $A \in \mathcal{A}$. Let $\epsilon = D_0/|D_0|$. Perturbing D_A by a suitable finite rank operator q one obtains an invertible operator $D'_A = D_A + q$ which anticommutes with the grading operator Γ and so also $F_A = D'_A/|D'_A|$ anticommutes with Γ .

Using the method in [LM] one can construct a family T_A of unitary operators commuting with Γ such that $T_A^{-1}F_A T_A - \epsilon$ is a trace-class operator. The operators T_A are not uniquely defined, if $R_A \in U_1$ (with respect to the polarization defined by the sign ϵ) then also $R_A T_A$ satisfies the condition. In order to be self contained, we give the (somewhat simplified) proof adapted to the present discussion.

Theorem 1. *For all interactions A in $D_A = D_0 + A$ with A a pseudodifferential*

operator of order zero, there is an unitary pseudodifferential operator T_A , commuting with the spinor grading operator Γ , such that the transformed interaction A' satisfies $[\epsilon, A'] \in L_1$ and $T_A = 1$ for $A = 0$. The symbol of T_A depends smoothly on the symbol of A .

Proof of Theorem. We first prove the following key lemma providing the recipe for constructing T_A :

Lemma 1. *Let A be a pseudodifferential operator such that $[\epsilon, A]$ is of order $k \leq 0$. Then A' , defined by $D_0 + A' = T_A^{-1}(D_0 + A)T_A$ with the unitary operator*

$$T_A = e^\alpha, \quad \alpha = -\frac{1}{8} (|D_0|^{-1}[\epsilon, A] + [\epsilon, A]|D_0|^{-1})$$

has the property $[\epsilon, A']$ is of order $k - 1$

Proof of Lemma. Denote the space of pseudodifferential operators of order k by O_k ; this is in fact an algebra if $k \leq 0$. We write $A' = A'_1 + A'_2$ where

$$A'_1 = A + [D_0, \alpha]$$

is the leading term in the asymptotic expansion, and

$$A'_2 = T_A^{-1}[D_0, T_A - \alpha - 1] + T_A^{-1}[A, T_A - 1]$$

is the rest.

Since obviously α is of order $k - 1$, and the fact that A'_2 is of order $k - 1$ trivially follows from $\alpha D_0, D_0 \alpha \in O_k$ and

$$T_A - 1 - \alpha = \frac{1}{2}\alpha^2 + \frac{1}{6}\alpha^3 + \dots \in O_{2k-2} \subset O_{k-2}.$$

The nontrivial part thus is to show that $[\epsilon, A'_1] \in O_{k-1}$. This can be seen by the following calculation,

$$\begin{aligned} [\epsilon, A'_1] &= [\epsilon, A] - \frac{1}{8} [\epsilon, [D_0, |D_0|^{-1}[\epsilon, A] + [\epsilon, A]|D_0|^{-1}]] \\ &= \frac{1}{8} (8[\epsilon, A] - [\epsilon, \epsilon[\epsilon, A] - |D_0|^{-1}[\epsilon, A]|D_0|\epsilon + \epsilon|D_0|[\epsilon, A]|D_0|^{-1} - [\epsilon, A]\epsilon]) \\ &= \frac{1}{8} (8[\epsilon, A] - 4[\epsilon, A] - 2|D_0|[\epsilon, A]|D_0|^{-1} - 2|D_0|^{-1}[\epsilon, A]|D_0|) \\ &= \frac{1}{4} (|D_0|^{-1} [[D_0], [\epsilon, A]] - [[D_0], [\epsilon, A]] |D_0|^{-1}) \end{aligned}$$

where we used $[\epsilon, A] = -\epsilon[\epsilon, A]\epsilon$ and $\epsilon D_0^{\pm 1} = D_0^{\pm 1}\epsilon = |D_0^{\pm 1}|$. Thus

$$[\epsilon, A'_1] = \frac{1}{4} [[D_0|^{-1}, [[D_0|, [\epsilon, A]]]$$

which is in O_{k-1} since $[[D_0], O_p] \subset O_p$. If we replace D_0^{-1} by $D_0(D_0^2 + \lambda)^{-1}$ a similar calculation leads to the same conclusion,

$$[\epsilon, A'_1] = \frac{1}{4} [[D_0|(D_0^2 + \lambda)^{-1}, [[D_0], [\epsilon, A]]] + \frac{1}{2} \lambda ((D_0^2 + \lambda)^{-1}[\epsilon, A] + [\epsilon, A](D_0^2 + \lambda)^{-1}).$$

This proves our Lemma.

We can apply this method successively: Starting from some interaction $A_0 = A$ such that $[\epsilon, A] \in O_0$ we get a new interaction $A_1 = A'$ using the conjugation T_A , with $[\epsilon, A_1] \in O_{-1}$. We can then insert A_1 as an argument to T and obtain an unitary operator T_{A_1} . This defines again a new interaction $A_2 = A'_1$ such that $[\epsilon, A_2] \in O_{-2}$. Continuing this way we obtain, after p steps, an unitary operator $T_A^p = T_{A_{p-1}} \dots T_{A_0}$ such that the interaction A_p in $T_A^p(D_0 + A)(T_A^p)^{-1} = D_0 + A_p$ has the property $[\epsilon, A_p] \in L_1$. The number p of iterations depends on the dimension of M , one must take $p > \dim M$. \square

Remark Actually, in quantum field theory applications (and also in Section 3 below) it is sufficient that $[\epsilon, A']$ is Hilbert-Schmidt; in practice, this means that one needs to perform one less iteration in the algorithm.

Since T_A commutes with Γ , the operator $T_A^{-1}F_A T_A$ anticommutes with Γ and we can write

$$T_A^{-1}F_A T_A = \begin{pmatrix} 0 & g \\ g^* & 0 \end{pmatrix}$$

where the blocks are defined with respect to the grading Γ . Here $g : S_- \rightarrow S_+$ is a unitary operator from the left chirality spinors to right chirality spinors. Likewise,

$$\epsilon = \begin{pmatrix} 0 & g_0 \\ g_0^* & 0 \end{pmatrix}.$$

with $g - g_0$ a trace-class operator. Thus we can write $g_0^{-1}g = h_A$ and $h_A - 1$ is trace-class. If T'_A is another unitary operator which satisfies the condition $T_A^{-1}F_A T_A - \epsilon \in L_1$ then we can write (by the ideal property of L_1) $T'_A = T_A S_A$ where $S_A \in U_1$.

Since T_A and the gauge transformations $h \in \mathcal{G}$ commute with Γ , we can write

$$(2.1) \quad T_A^{-1}h T_A^h = \Omega(A; h) = \begin{pmatrix} \Omega_+(A; h) & 0 \\ 0 & \Omega_-(A; h) \end{pmatrix}$$

where Ω satisfies the cocycle relation

$$(2.2) \quad \Omega(A; h_1)\Omega(A^{h_1}; h_2) = \Omega(A; h_1 h_2).$$

The change $T_A \mapsto T'_A = T_A S_A$ acts on the cocycle as

$$(2.3) \quad \Omega'(A; h) = S_A^{-1}\Omega(A; h)S_A^h.$$

Lemma 2. *The cocycle $\Omega = (\Omega_+, \Omega_-)$ satisfies*

$$\Omega_+ - g_0 \Omega_- g_0^{-1} \in L_1.$$

Proof. By $T_{A^h}^{-1} F_{A^h} T_{A^h} - \epsilon \in L_1$ and $F_{A^h} = h^{-1} F_A h + \text{finite rank operator}$, we obtain by

$$h T_{A^h} \epsilon T_{A^h}^{-1} h^{-1} - F_A \in L_1$$

and subtracting from $T_A \epsilon T_A^{-1} - F_A \in L_1$ the relation

$$h T_{A^h} \epsilon T_{A^h}^{-1} h^{-1} - T_A \epsilon T_A^{-1} \in L_1.$$

Finally, multiplying this relation from the left by T_A^{-1} and from the right by $h T_{A^h}$ gives

$$[\epsilon, T_A^{-1} h T_{A^h}] = [\epsilon, \Omega(A; h)] \in L_1,$$

which can be written as a pair of relations

$$g_0 \Omega_- - \Omega_+ g_0 \in L_1 \text{ and } g_0^{-1} \Omega_+ - \Omega_- g_0^{-1} \in L_1,$$

which actually are equivalent. The claim follows immediately by multiplying the second relation by g_0 from the left and from the ideal property of L_1 . \square

Corollary 1. *The conditional supertrace $\text{str}_c(\Omega) = \frac{1}{2} \text{tr}(\Gamma \Omega + \epsilon \Gamma \Omega \epsilon)$ is absolutely converging.*

If Ω is trace-class then the conditional supertrace above is just the supertrace $\text{tr} \Gamma \Omega$.

The set of unitary matrices $\Omega = \begin{pmatrix} \Omega_+ & 0 \\ 0 & \Omega_- \end{pmatrix}$ satisfying the property in Lemma 2 is an infinite dimensional Lie group $UU^1(H)$ which is homotopy equivalent to U^1 . Namely, its elements are parametrized by the unitaries Ω_- acting in the Hilbert space of left handed spinor fields and the unitary matrix $z = (g_0 \Omega_-)^{-1} \Omega_+ g_0$. By repeated use of Lemma 2 the operator z acting on left-handed spinors differs from the identity by a trace-class operator.

Remark Since the space of gauge connections is contractible, $(A, s) \rightarrow sA$ with $0 \leq s \leq 1$ is a contraction, the cocycle $\Omega(A; h)$ is contractible to the tautological cocycle $(A, h) \mapsto h = \Omega(sA; h)$ for $s = 0$. However, this homotopy is defined only in the framework of $U(H)$ valued cocycles, not in cohomology with values in $UU^1(H)$.

The above construction can be extended to the case when $\text{ind} D_A \neq 0$. Now we fix ϵ such that the parameter $g_0 : S_- \rightarrow S_+$ has Fredholm index equal to the Dirac index of D_A . In addition, we can require that g_0 is an isomorphism from the orthogonal complement of its kernel to the orthogonal complement of the cokernel. Likewise, one can choose ϵ_A such that it is equal to the sign of D_A in the orthogonal complement of $\ker D_A$ and is equal to zero on $\ker D_A$. Then ϵ_A anticommutes with Γ . The construction of T_A is essentially the same as in the index zero case: Instead of D_0^{-1} in the formula for T_A one chooses an operator R_0 such that $R_0 D_0$ is equal to the identity on the orthogonal complement of $\ker D_0$. This does not fix R_0 uniquely,

only up to a finite rank operator, but that is sufficient for the construction of the cocycle Ω in the Lemma.

The cocycle Ω defines a principal bundle P over X with structure group $UU^1(H)$. The total space of P is

$$P = \mathcal{A} \times_{\Omega} UU^1(H,)$$

the set of equivalence classes of pairs $(A, g) \in \mathcal{A} \times UU^1(H)$ where the equivalence relation is $(A, g) \equiv (A^{h^{-1}}, \Omega(A; h)g)$ where $h \in \mathcal{G}$. The right action of $UU^1(H)$ on P is given by the right multiplication on the second component.

Given a system $\psi_{\alpha} : U_{\alpha} \rightarrow \mathcal{A}$ of local trivializations of the bundle $\mathcal{A} \rightarrow X$ one has local trivializations $\tilde{\psi}_{\alpha} : U_{\alpha} \rightarrow P$, $\tilde{\psi}_{\alpha}(x) = (\psi_{\alpha}(x), 1)$. The transition functions are $\tilde{g}_{\alpha\beta}(x) = \Omega(\psi_{\alpha}(x), g_{\alpha\beta}(x))$ and they satisfy the usual cocycle relation

$$\tilde{g}_{\alpha\beta}\tilde{g}_{\beta\gamma} = \tilde{g}_{\alpha\gamma}$$

as a consequence of the twisted cocycle relation for Ω .

Following [AS2], a connection in the principal bundle $\mathcal{A} \rightarrow \mathcal{A}/\mathcal{G}$ is constructed as follows. For $A \in \mathcal{A}$ the horizontal subspace H_A in the tangent space of \mathcal{A} consists of all $B \in \mathcal{A}$ such that $\sum \partial_i B^i + [A_i, B^i] = 0$. Here $B^i = \sum g^{ij} A_j$ where g^{ij} is the inverse of the metric tensor g_{ij} on M . If $B \in H_A \cap V_A$ then on one hand $B_i = D_i^A X$ for some $X \in C^{\infty}(M, \mathfrak{g})$ but on the other hand $D_i^A B^i = \Delta_A X = 0$. The harmonic functions X satisfy

$$(2.4) \quad 0 = \int_M \langle X, \Delta_A X \rangle dvol_M = \int_M \langle D_i^A X, D^{A,i} X \rangle,$$

where $\langle \cdot, \cdot \rangle$ is an invariant inner product on \mathfrak{g} . Thus $D_i^A X = 0$ for each coordinate index i . On a connected manifold M a covariantly constant function X is completely fixed by its value at a base point x_0 . Thus for based infinitesimal gauge transformations we must have $X = 0$ everywhere and so $V_A \cap H_A = \{0\}$.

The connection form θ corresponding to the horizontal distribution H_A can be given fairly explicitly, [AS2]. Since the only harmonic function vanishing at the base point is zero, the form

$$(2.5) \quad \theta_A(B) = \Delta_A^{-1}(D_i^A B^i)$$

is uniquely defined, with values in the Lie algebra $Map_0(M, \mathfrak{g})$. It vanishes on horizontal forms B and is tautological on the vertical tangent vectors $B_i = D_i^A X$.

Given a principal bundle $\pi : P \rightarrow M$ with structure group G and a cocycle $\Omega : P \times G \rightarrow K$ where K is a group of invertible bounded operators in a Hilbert space V , one can construct an associated vector bundle $E = P \times_{\Omega} V$. The elements of E are equivalence classes of pairs $(p, v) \in P \times V$ with the equivalence relation defined by $(p, v) \sim (pg^{-1}, \Omega(p; g)v)$. As in the standard case, when the cocycle Ω is simply a homomorphism $\rho(g) = \Omega(p; g)$, a connection in the principal bundle P should define a connection in the associated vector bundle. The construction is slightly more complicated in the more general situation. A section ξ of E can be identified as a map $\xi : P \rightarrow V$ such that $\xi(pg) = \Omega(p; g)^{-1}\xi(p)$ for all $(p, g) \in P \times G$.

Fixing a local section ψ of P , a section of E is given by a local vector valued function $f = \xi \circ \psi$. Denote by ω the derivative of $\Omega(p; g)$ with respect to the second argument at $g = 1$.

Returning to the case when $P = \mathcal{A}$ is the space of gauge connections and Ω is the cocycle (2.1) we have

Lemma 3. *A change $T'_A = S_A T_A$ induces a gauge transformation*

$$\omega'(A; \theta(v)) = S_A \omega(A; \theta(v)) S_A^{-1} - (\mathcal{L}_{\theta(v)} S_A) S_A^{-1}$$

on the form ω .

Proof. Differentiating (2.3) with respect to parameter t at $t = 0$ for $h = e^{t\theta(v)}$ we obtain

$$\begin{aligned} \omega'(; \theta(v)) &= \left. \frac{d}{dt} \right|_{t=0} \Omega'(A; e^{t\theta(v)}) = \left. \frac{d}{dt} \right|_{t=0} S_A \Omega(A; e^{t\theta(v)}) S_A^{-1} \\ &= S_A \omega(A; \theta(v)) S_A^{-1} - (\mathcal{L}_{\theta(v)} S_A) S_A^{-1} \end{aligned}$$

where v is a tangent vector on \mathcal{A} at the point A and $\mathcal{L}_{\theta(v)}$ is the Lie derivative in the direction of the infinitesimal gauge transformation $\theta(v)$. \square

The natural form $\omega(p; \theta)$ taking values in the Lie algebra of K is not quite a connection form as the following proposition shows:

Proposition 1. *Let ω_α be the pull-back of the form $\omega(p; \theta)$ with respect to the local section ψ_α of the bundle $P \rightarrow M$. Then*

$$(2.6) \quad \omega_\beta = \Omega(\psi_\alpha; g_{\alpha\beta})^{-1} \omega_\alpha \Omega(\psi_\alpha; g_{\alpha\beta}) + \Omega(\psi_\alpha; g_{\alpha\beta})^{-1} D^{vert,1} \Omega(\psi_\alpha; g_{\alpha\beta})$$

where $D^{vert,1}$ means the differentiation taken in the vertical directions in the first argument,

$$D_v^{vert,1} \Omega(\psi; g) = \left. \frac{d}{dt} \right|_{t=0} \Omega(\psi \cdot e^{t\theta_\alpha(v)}; g) + D_v^{(2)} \Omega(\psi_\alpha; g),$$

where $D_v^{(2)}$ is the differentiation applied to the second argument, in the direction of the vector v , and $\theta_\alpha = \psi_\alpha^* \theta$ is the pull-back to U_α .

Proof. Using the cocycle identity $\Omega(p; g_0) \Omega(p \cdot g_0; g_0^{-1} g) = \Omega(p; g)$ we obtain first

$$D_v^{(2)} \Omega(p; g) = D_v^{(2)} \Omega(p; g_0 g_0^{-1} g) = \Omega(p; g_0) D_v^{(2)} \Omega(p \cdot g_0; g_0^{-1} g)$$

where $D_v^{(2)}$ (resp. $D_v^{(1)}$) denotes the derivative with respect to the second (resp. first) argument in the direction of the vector v on M ; here g_0 is a constant and $g = g(z)$ is a group valued function on M .

Taking now $g = g_{\alpha\beta}(z)$ and $p = \psi_\alpha(z)$, and after taking the derivatives set $g_0 = g_{\alpha\beta}(z)$, we obtain

$$(2.7) \quad \omega(\psi_\alpha(z) \cdot g_{\alpha\beta}(z); g_{\alpha\beta}^{-1} D_v g_{\alpha\beta}) = \Omega(\psi_\alpha(z); g_{\alpha\beta}(z))^{-1} D_v^{(2)} \Omega(\psi_\alpha(z); g_{\alpha\beta}(z)).$$

Next, using again the cocycle property in the second equality below, we obtain

(2.8)

$$\begin{aligned}
& \omega(\psi_\beta(z); g_{\alpha\beta}(z)^{-1} \theta_\alpha(v) g_{\alpha\beta}(z)) \\
&= \frac{d}{dt} \Big|_{t=0} \Omega(\psi_\alpha(z) \cdot g_{\alpha\beta}(z); g_{\alpha\beta}(z)^{-1} e^{t\theta_\alpha(v)} g_{\alpha\beta}(z)) \\
&= \frac{d}{dt} \Big|_{t=0} \Omega(\psi_\alpha(z) \cdot g_{\alpha\beta}(z); g_{\alpha\beta}(z)^{-1}) \Omega(\psi_\alpha(z); e^{t\theta_\alpha(v)} g_{\alpha\beta}(z)) \\
&= \frac{d}{dt} \Big|_{t=0} \Omega(\psi_\alpha(z); g_{\alpha\beta}(z))^{-1} \Omega(\psi_\alpha(z); e^{t\theta_\alpha(v)} g_{\alpha\beta}(z)) \\
&= \frac{d}{dt} \Big|_{t=0} \Omega(\psi_\alpha(z); g_{\alpha\beta}(z))^{-1} \Omega(\psi_\alpha(z); e^{t\theta_\alpha(v)}) \Omega(\psi_\alpha \cdot e^{t\theta_\alpha(v)}; g_{\alpha\beta}(z)) \\
&= \Omega(\psi_\alpha; g_{\alpha\beta})^{-1} \omega(\psi_\alpha; \theta_\alpha(v)) \Omega(\psi_\alpha; g_{\alpha\beta}) + \Omega(\psi_\alpha; g_{\alpha\beta})^{-1} D_v^{(1,vert)} \Omega(\psi_\alpha; g_{\alpha\beta}).
\end{aligned}$$

Adding equation (2.8) to (2.7) we arrive at (2.6) \square

Thus the defect in $\omega(p; \theta)$, for being a connection form, is the missing differentiation in the horizontal direction in the first argument on the right-hand-side in (2.6). A true connection form θ_Q on the principal K bundle $Q = P \times_\Omega K$ comes from a \mathfrak{k} valued 1-form θ_P on P as $\theta_Q = k^{-1} \theta_P k + k^{-1} dk$ where $k \in K$ if and only if θ_P satisfies the following conditions:

- (1) $\theta_P^g = \Omega(\cdot; g)^{-1} \theta_P \Omega(\cdot; g) - \Omega(\cdot; g)^{-1} d_P \Omega(\cdot; g)$
- (2) $\theta_P(p; \hat{X}) = \omega(p; X)$

where \hat{X} denotes the vertical vector field on P generated by the element X in the Lie algebra of G . These two conditions guarantee that 1) the form θ_Q on $P \times K$ indeed descends to $P \times_\Omega K$, 2) it is tautological in vertical K directions, and 3) is equivariant with respect to the combined right and adjoint action of K . In the more standard situation, when Ω does not depend on the argument $p \in P$, the conditions above are the usual properties of a connection form.

A true connection exists always in the finite dimensional case. In an infinite dimensional case, for a non-Hilbert manifold, one has to prove the existence of a partition of unity subordinate to an open trivializing cover of the base manifold. In the present case when $P = \mathcal{A}$ and $X = \mathcal{A}/\mathcal{G}$ such a partition of unity exists as proven in [CMM]; the proof is based a theorem of John Milnor [Mi]. So given an open cover $\{U_\alpha\}$ of X with local trivializations ψ_α and transition functions $\tilde{g}_{\alpha\beta}$ the standard construction applies, the local connection forms can be defined as

$$- \sum_{\beta} \rho_{\beta} d\tilde{g}_{\alpha\beta} \tilde{g}_{\alpha\beta}^{-1}.$$

However, this definition is not very user friendly for the calculation of characteristic classes because of the necessity to pick up the partition of unity.

In the case when $P = \mathcal{A}$ and $\Omega(A; g) = T_A^{-1} g T_{A^g}$ we can construct a connection as a covariant derivative acting on sections of the associated vector bundle through the global 1-form $\tilde{\theta} = T_A^{-1} \theta T_A + T_A^{-1} dT_A$. Although this form does not take values in the Lie algebra of the group UU^1 the covariant differentiation of a section f of the vector bundle can nevertheless be defined as

$$\nabla f_\alpha = df_\alpha + \tilde{\theta}_\alpha f.$$

We observe from the formula $\Omega(A; g) = T_A^{-1} g T_{A^g}$ that actually $\tilde{\theta}$ is equal to $\omega(A; \theta)$ *along the vertical directions* of the fibration $\mathcal{A} \rightarrow \mathcal{A}/\mathcal{G} = X$. Along horizontal directions the latter form is zero whereas the former is nonvanishing. It might sound strange that the true connection is nonvanishing along horizontal directions but one should bear in mind that the horizontal directions in the fibration \mathcal{A}/\mathcal{G} are not the same as the horizontal directions in the associated UU^1 principal bundle over \mathcal{A}/\mathcal{G} . This fact is due to the nonvanishing of the derivative of Ω with respect to the first argument in the horizontal directions of the fibration $\mathcal{A} \rightarrow X$.

Although the curvature \mathcal{F} constructed from the connection form $\tilde{\theta}$ does not take values in the space of trace class operators, one can still define the Chern character via the secondary characteristic classes. The essential reason for this is that along vertical directions in $\pi : \mathcal{A} \rightarrow \mathcal{A}/\mathcal{G}$ the connection $\tilde{\theta}$ agrees with $\omega(\cdot; \theta)$ and the latter takes values in the Lie algebra of U^1 . For example, a spherical cycle $S^{2k} \subset \mathcal{A}/\mathcal{G}$ can be lifted to \mathcal{A} after removing one point x from S^{2k} . The inverse image $\pi^{-1}(x) \subset \mathcal{A}$ is then a cycle of dimension $2k - 1$ inside of a gauge orbit. The integral of the ill-defined form $\text{tr } \mathcal{F}^k$ over the cycle S^{2k} becomes then an integral of the corresponding Chern-Simons form over the cycle in the fiber $\pi^{-1}(x)$. But this integration involves only the vertical directions in the connection and curvature, and are thus well-defined because of the trace class property of the connection along vertical directions. This process can be generalized to arbitrary homology cycles as explained below.

Recall that in the trace class case, on contractible sets $c^{2n} := \text{tr } \mathcal{F}^n = d \text{tr } C^{2n-1} = dc^{2n-1}$ where C^{2n-1} is a differential polynomial of the connection form. Explicitly, in a general vector bundle with a local connection 1-form A and the corresponding curvature form F the Chern-Simons form c^{2n-1} is given by the formula

$$(2.9) \quad c^{2n-1} = n \int_0^1 dt \text{tr } A_t F_t^{n-1}$$

where $A_t = tA$ and $F_t = t dA + \frac{t^2}{2} [A, A]$.

The integral of c^{2n} over a simplicial cycle c_{2n} of dimension $2n$ is then given as the sum of integrals of c^{2n-1} over all the faces $c_{2n-1, \alpha\beta}$ in the singular simplex $c_{2n} = \sum c_{2n, \alpha}$. The convention is that on the face $c_{2n+1, \alpha\beta}$ one integrates the pull-back form c_α^{2n-1} (defined by the local section ψ_α) over the boundary component $c_{2n-1, \alpha\beta} = c_{2n, \alpha} \cap c_{2n, \beta}$ with the orientation defined by the outward normal to $c_{2n, \alpha}$ and the fixed orientation on c_{2n} . For a given index pair α, β one has the pair of integrals

$$\int_{c_{2n-1, \alpha\beta}} c^{2n-1, \alpha} \quad \text{and} \quad \int_{c_{2n-1, \beta\alpha}} c^{2n-1, \beta}.$$

Now the form $c_{2n-1, \alpha\beta}$ is a gauge transform of $c_{2n-1, \beta\alpha}$ with respect to the change of local trivialization $\psi_\beta = \psi_\alpha \cdot g_{\alpha\beta}$ so, taking into account the orientations, the above pair gives the contribution

$$\int_{c_{2n-1, \alpha\beta}} (c^{2n-1, \alpha} - g_{\alpha\beta}^* c^{2n-1, \alpha}) = \int_{c_{2n-1, \alpha\beta}} c^{2n-1, \alpha\beta}.$$

The difference under the integral is a trace of a differential polynomial in the local connection form $\psi_\alpha^* \tilde{\theta}$ and the Maurer-Cartan 1-form $g_{\alpha\beta}^{-1} dg_{\alpha\beta}$; using (2.9) one observes that each term at least of degree one in the latter. When the Maurer-Cartan form takes values in the Lie algebra of U^1 , that is, in the ideal of trace-class operators, all the traces actually converge.

Thus we can conclude, using Corollary 1 and Lemma 3,

Theorem 2. *Let \mathcal{F} be the curvature corresponding to the connection ∇ , defined from the 1-form $\tilde{\theta}$ on \mathcal{A} . The formal Chern character $\text{str}_c e^{\mathcal{F}/2\pi i}$ defines a cohomology class on the base X as a pairing with singular cycles, evaluated using the descent equations in the Čech - de Rham double complex up to the cocycle in $H^{(*,1)}$ of Čech degree one. The class does not depend on the choice of the regularization T_A .*

Proof. What remains to show is that the Chern character in cohomology does not depend on the choice of the family T_A . A different choice T'_A is related to the first by $T'_A = T_A S_A$ with $S_A \in U_1$. Since the space \mathcal{A} of smooth connections is flat, we can choose a homotopy $T_A(t) = T_A S_A(t)$ with $S_A(0) = 1$ and $S_A(1) = S_A$. This gives a homotopy $\nabla(t)$ of the connections, connecting ∇ to the connection ∇' defined by T'_A . The rest is then the standard argument relating the Chern characters defined by a pair of connections in a vector bundle, the difference being the exterior derivative of a Chern-Simons form.

One should also note that the choice of the base point ϵ is not essential. Any other base point ϵ' (a self-adjoint operator with $\epsilon'^2 = 1$ and essential spectrum at ± 1) is written as $\epsilon' = W\epsilon W^{-1}$ for some unitary operator W . The connection $W\tilde{\theta}W^{-1}$ satisfies then the same conditional trace-class properties (in the vertical directions) with respect to ϵ' as does $\tilde{\theta}$ with respect to ϵ and so the Chern character defined by the pair $(\epsilon, \tilde{\theta})$ is equal to the Chern character defined by $(\epsilon', \tilde{\theta}')$.

□

3. THE ODD CASE: CHERN CHARACTER FOR U_p BUNDLES

When M is an odd dimensional spin manifold and D a family of Dirac operators acting on sections of the tensor product of the spin bundle and a (trivial) vector bundle over M , the Chern character takes values in the odd cohomology groups of the parameter space X . In the following I want to describe how the renormalization method in Section 2, with the operators like T_A , can be used to compute the Chern character on $X = \mathcal{A}/\mathcal{G}$.

The first step is to replace the family of Dirac operators D_A by a family of operators $\epsilon + B$ where B is a bounded operator (which depends on A) such that $[\epsilon, B] \in L_2$ and $\epsilon B + B\epsilon \in L_1$. First, choose a family of operators T_A such that $T_A^{-1}\epsilon T_A - F_A$ is trace-class, [LM]; equivalently, $\epsilon - T_A^{-1}F_A T_A$ is trace-class. Actually, since the operators D_A might have zero modes we have to replace the sign operator F_A by an approximate sign operator, say $F_A = D_A(|D_A| + \exp(-D_A^2))^{-1}$. Since these operators transform equivariantly with respect to gauge transforma-

tions, they descend to a family of bounded operators parametrized by $X = \mathcal{A}/\mathcal{G}$. We denote $T_A F_A T_A^{-1} = \epsilon + B(A)$.

As in the even case, we have a cocycle $\Omega : \mathcal{A} \times \mathcal{G} \rightarrow U_1$. Since in the odd case we do not have the grading Γ , there is no further refinement of this cocycle to a U^1 valued cocycle. The cocycle ω defines a U_1 bundle P over the moduli space X . The Atiyah-Singer connection in the principal bundle $\mathcal{A} \rightarrow X$ defines again a connection ∇ in the U_1 bundle (and in the associated Hilbert bundle) over X . We can then define a *superconnection*

$$\Theta = \delta + B + \nabla.$$

Here δ is an exterior differentiation in the sense of noncommutative geometry, [Co]: It is acting on 'k-forms' of the type $\Phi = \alpha_0 \delta \alpha_1 \dots \delta \alpha_k$ as $\delta \Phi = \delta \alpha_0 \delta \alpha_1 \dots \delta \alpha_k$ where the α_i 's are bounded operators in H such that $\delta \alpha = [\epsilon, \alpha] \in L_p$. One can check that when k is odd $\delta \Phi = \epsilon \Phi + \Phi \epsilon$ whereas when k is even $\delta \Phi = [\epsilon, \Phi]$. In this sense, B is a 1-form. We have a double complex of forms $\Omega^{k,l}$ where k gives the δ -degree and l the form degree related to the exterior differentiation on the base X . Finally, we modify the the action $\delta \mapsto (-1)^l \delta$ in order that $\delta d + d \delta = 0$.

Choosing an open cover $\{U_\alpha\}$ of X with local trivializations of the bundle P and transition functions $\tilde{g}_{\alpha\beta} : U_{\alpha\beta} \rightarrow U_1$, the local operator families B_α are related by

$$B_\beta = \tilde{g}_{\alpha\beta}^{-1} B_\alpha \tilde{g}_{\alpha\beta} + \tilde{g}_{\alpha\beta}^{-1} [\epsilon, \tilde{g}_{\alpha\beta}]$$

and the local 1-forms η_α on X which define the connection ∇ locally as $\nabla = d + \tilde{\theta}_\alpha$ are related by

$$\tilde{\theta}_\beta = \tilde{g}_{\alpha\beta}^{-1} \tilde{\theta}_\alpha \tilde{g}_{\alpha\beta} + \tilde{g}_{\alpha\beta}^{-1} d \tilde{g}_{\alpha\beta}$$

In this way, the superconnection transforms covariantly,

$$\Theta_\beta = \tilde{g}_{\alpha\beta}^{-1} \Theta_\alpha \tilde{g}_{\alpha\beta}.$$

When the U_1 bundle arises as an associated bundle to the gauge bundle $\mathcal{A} \rightarrow \mathcal{A}/\mathcal{G}$ through the cocycle Ω a superconnection can be defined with the help of local operator families $\epsilon + B_\alpha(A) = F_A$ defined using a local section $\psi_\alpha : \mathcal{A}/\mathcal{G} \rightarrow \mathcal{A}$.

The square of the superconnection gives the graded curvature,

$$\Theta^2 = \mathcal{F}^{(2,0)} + \mathcal{F}^{(1,1)} + \mathcal{F}^{(0,2)}$$

with $\mathcal{F}^{(2,0)} = \epsilon B + B \epsilon + B^2$, $\mathcal{F}^{(1,1)} = [\epsilon, \tilde{\theta}] + dB + [B, \tilde{\theta}]$ and $\mathcal{F}^{(0,2)} = d\tilde{\theta} + \tilde{\theta}^2$.

Since the classifying space $BU_1 = U(\infty)$ has odd homotopy type, the characteristic classes of U_1 bundles are located in in odd cohomology of the base space. The lowest nontrivial form is in $H^1(X)$. In a similar way as in the even case, we have:

Theorem 3. *The terms $tr_c (\mathcal{F}^n)^{(1,2n-1)}$ in the formal Chern character on X define odd closed index forms on $X = \mathcal{A}/\mathcal{G}$, through the terms of Čech degree one in the corresponding Čech - de Rham complex. They do not depend on the choice of the regularization family T_A .*

As in the even case, if one chooses a connection form $\tilde{\theta}$ such that $[\epsilon, \tilde{\theta}]$ is trace-class then also the forms $\text{tr } \mathcal{F}^n$ are well-defined directly.

Let us consider two simple examples, where we can in fact choose $T_A \equiv 1$.

Example 1 For $X = S^1$ the nontrivial U_1 bundles are classified by the homotopy type of the transition function $g \in U_1$ at the ends of the interval $0 \leq t \leq 2\pi$ parametrizing the circle $e^{it} \in S^1$. The group consists of disconnected components labelled by the Fredholm index of the a block in

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

The superconnection is given by $\Theta = \delta + d + \frac{t}{2\pi}g^{-1}[\epsilon, g]$ and $\tilde{\theta} = 0$. The representative for the index form in $H^1(S^1)$ is then $\frac{1}{2\pi}dt \text{tr } g^{-1}[\epsilon, g] = \text{tr } \mathcal{F}^{(1,1)}$. The integral of this form over the circle S^1 is equal to twice the Fredholm index of a .

Remark Actually, it is sufficient that the off-diagonal blocks of all components $\mathcal{F}^{(a,b)}$ are Hilbert-Schmidt (so we can take as the structure group of the bundle U_2 instead of U_1) and in addition the diagonal blocks of $\mathcal{F}^{(1,1)}$ are trace-class. Then all the odd forms on X arising in $\text{tr } \mathcal{F}^k$ are conditionally converging traces.

Example 2 The space \mathcal{A} of smooth vector potentials on the circle S^1 with values in the Lie algebra \mathfrak{g} of the gauge group $G = SU(2)$ is a principal ΩG bundle over G . The group ΩG of based smooth loops acts freely, as gauge transformations, on \mathcal{A} . The group ΩG acts in the Hilbert space H of square integrable functions on the circle with values in \mathbb{C}^2 by point-wise multiplication. The space $H = H_+ \oplus H_-$ is polarized to nonnegative and negative Fourier modes. It is known that ΩG embeds to $U_2(H)$, [PS]. (In fact, also $\Omega G \subset U_1$). Thus the ΩG bundle over $SU(2)$ extends naturally to a U_2 bundle. A superconnection is defined as follows. On the equator $S^2 \subset S^3 = SU(2)$ we have the transition function $g : S^2 \rightarrow \Omega G$ (we extend the equator to a small open neighborhood thereof and extend g as a constant to the normal direction). Explicitly, g can be given as

$$g(\mathbf{n}) = \exp(if(\phi)\mathbf{n} \cdot \sigma)$$

where $\mathbf{n} = (n_1, n_2, n_3)$ is a unit vector in \mathbb{R}^3 parametrizing the sphere S^2 and $\sigma = (\sigma_1, \sigma_2, \sigma_3)$ is the set of complex hermitean trace-less 2×2 Pauli matrices, $\sigma_k^2 = 1$ and $\sigma_1\sigma_2 = i\sigma_3$, and cyclic permutations of this relation. $f(\phi)$ is a smooth function on the interval $[0, 2\pi]$ with $f(0) = 0$ and $f(2\pi) = n \cdot 2\pi$, $n \in \mathbb{Z}$. We set $B = 0 = \tilde{\theta}$ on the half sphere $x_3 > 0$ and $B = \frac{t}{\pi}[\epsilon, g]g^{-1}$, $\tilde{\theta} = \frac{t}{\pi}g^{-1}dg$ on the other hemisphere, where $0 \leq t \leq \pi$ is the distance from the 'South Pole' on S^3 . This defines the superconnection Θ on the Southern hemisphere whereas $\Theta = \delta + d$ on the Northern hemisphere. The $(1, 3)$ component of the square of the supercurvature on the Southern hemisphere is now

$$\text{tr}_H (\mathcal{F}^2)^{(1,3)} = \frac{dt}{\pi} \left(-\frac{t}{\pi} + \left(\frac{t}{\pi}\right)^2 \right) \text{tr}_H g^{-1}[\epsilon, g](g^{-1}dg)^2$$

and the integral of the square of the curvature over the whole 3-sphere becomes

$$I(g) = \frac{i}{12\pi} \int_{0 \leq \phi \leq 2\pi} \int_{S^2} \text{tr}_{\mathbb{C}^2} (g'g^{-1})(dgg^{-1})^2$$

where g' is the derivative with respect to ϕ and d stands for the derivatives to the S^2 directions. We have used the formula

$$\mathrm{tr}_H[\epsilon, X]Y = \frac{1}{2\pi i} \int_{S^1} \mathrm{tr}_{\mathbb{C}^2} X'Y d\phi$$

for matrix valued functions on the unit circle. The value of the integral $I(g)$ is equal to $2\pi i$ times the winding number of the map $\tilde{g} : S^3 \rightarrow G$, where \tilde{g} is defined as $\tilde{g}(\phi, \mathbf{n}) = g(\mathbf{n})(\phi)$ with $\mathbf{n} \in S^2$ and ϕ parametrizes S^1 ; since g is a map from S^2 to the *based loop group* ΩG , this function on $S^2 \times S^1$ lifts to a map on S^3 . For the particular choice g above, the winding number is $2n$. With a minor modification one can get also the odd winding numbers: For that purpose one defines $g(\mathbf{n})(\phi)$ for $0 \leq \phi \leq \pi$ as above, but for $\pi \leq \phi \leq 2\pi$ one sets $g(\mathbf{n})(\phi) = h(\phi)$ where h is a fixed path in $SU(2)$ with end points $\pm \mathbf{1}$. Now the winding number of \tilde{g} is equal to n .

REFERENCES

- [AS1] M.F. Atiyah and I. M. Singer: The index of elliptic operators. IV. *Ann. of Math.* (2) **93** (1971), 119 - 138.
- [AS2] M.F. Atiyah and I.M. Singer: Dirac operators coupled to vector potentials. *Proc. Nat. Acad. Sci. U.S.A.* **81** (1984), no. 8, Phys. Sci., 2597 - 2600.
- [B] J.-M. Bismut: The Atiyah-Singer index theorem for families of Dirac operators: two heat equation proofs. *Invent. Math.* **83** (1985), no. 1, 91 - 151.
- [CMM] A. Carey, J. Mickelsson, and M. Murray: Index theory, gerbes, and Hamiltonian quantization. *Comm. Math. Phys.* **183** (1997), no. 3, 707 - 722.
- [C] Alain Connes: *Noncommutative Geometry*. Academic Press (1994); also <http://www.alainconnes.org/docs/book94bigpdf.pdf>
- [K] N. Kuiper: The homotopy type of the unitary group of Hilbert space. *Topology* **3** (1965), 19 - 30.
- [LM] E. Langmann and J. Mickelsson: Scattering matrix in external field problems. *J. Math. Phys.* **37** (1996), no. 8, 3933 - 3953.
- [M] J. Mickelsson: Wodzicki residue and anomalies of current algebras. In: *Integrable models and strings* (Espoo, 1993), 123 - 135, *Lecture Notes in Phys.*, **436**, Springer, Berlin, (1994).
- [Mi] John Milnor: On infinite dimensional Lie groups. Preprint, Institute for Advanced Studies, Princeton (1982)
- [MR] J. Mickelsson and S. Rajeev: Current algebras in d+1-dimensions and determinant bundles over infinite-dimensional Grassmannians. *Comm. Math. Phys.* **116** (1988), no. 3, 365 - 400.
- [P] Richard Palais: On the homotopy type of certain groups of operators. *Topology* **3** (1965), 271 - 279.
- [PS] A. Pressley and G. Segal: *Loop Groups*. Clarendon Press, Oxford (1986)
- [Q] Daniel Quillen: Superconnections and the Chern character. *Topology* **24** (1985), no. 1, 89 - 95.