Adrian Dumitrescu^{*} Minghui Jiang[†]

February 19, 2022

Abstract

For a planar point-set P, let D(P) be the minimum number of pairwise-disjoint empty disks such that each point in P lies on the boundary of some disk. Further define D(n) as the maximum of D(P) over all *n*-element point sets. Hosono and Urabe recently conjectured that $D(n) = \lceil n/2 \rceil$. Here we show that $D(n) \ge n/2 + n/236 - O(\sqrt{n})$ and thereby disprove this conjecture.

Keywords: Empty disk, point-circle incidence, graph, connected component.

Mathematics Subject Classification (MSC): 52C10 Erdős problems and related topics of discrete geometry.

1 Introduction

Hosono and Urabe [7] have recently considered two families of disks "incident" to a point set, as follows. For a given planar point set P, let B(P) be the minimum number of empty disks such that each point in P lies on the boundary of some disk. Further define the *bubble number* B(n) as the maximum of B(P) over all *n*-element point sets.

Analogously, let D(P) be the minimum number of pairwise-disjoint empty disks such that each point in P lies on the boundary of some disk. Further define the *disjoint bubble number* D(n) as the maximum of D(P) over all n-element point sets.

A point is said to *support* an empty disk if it lies on the boundary of the disk. Conversely, we say that the disk is *supported* by the point; note that a disk can be supported by multiple points. A point set is said to *support* a set of empty disks if each point supports at least one (empty) disk. Conversely, we say that a set of empty disks is *supported* by the point-set. Such a set of empty disks is also referred to as a *bubble set* for the given set of points.

Consider a set of points P. Since any set of pairwise-disjoint empty disks supporting P is simply a set of empty disks supporting P, we have $B(P) \leq D(P)$ for every P, and consequently

$$B(n) \le D(n), \text{ for every } n.$$
 (1)

Hosono and Urabe [7] have shown the following upper and lower bounds for B(n): for every $n \ge 14$, $\lceil n/2 \rceil \le B(n) \le \lfloor (2n-2)/3 \rfloor$. They also conjectured that $B(n) = D(n) = \lceil n/2 \rceil$. Here we show that out of three implied equalities, B(n) = D(n), $D(n) = \lceil n/2 \rceil$, and $B(n) = \lceil n/2 \rceil$, only the third holds, while the first two are false.

^{*}Department of Computer Science, University of Wisconsin-Milwaukee, USA. Email: dumitres@uwm.edu. Supported in part by NSF grant DMS-1001667.

[†]Department of Computer Science, Utah State University, Logan, USA. Email: mjiang@cc.usu.edu.

Dillencourt [6] proved that every Delaunay triangulation of n points (n even) has a perfect matching. By definition, for every edge in a Delaunay triangulation, there is an empty disk with the two endpoints of this edge on its boundary. Consequently, we have the following.

Proposition 1. For every $n \ge 1$, $B(n) = \lceil n/2 \rceil$.

In contrast, we show that $D(n) > \lfloor n/2 \rfloor$ for even n as small as 174.

Theorem 1. For every even $n \ge 174$, $D(n) \ge n/2 + 1$.

Proposition 1 and Theorem 1 together imply that for every even $n \ge 174$, $B(n) \ne D(n)$ and $D(n) > \lceil n/2 \rceil$, which disprove the conjecture of Hosono and Urabe [7] that $B(n) = D(n) = \lceil n/2 \rceil$. Our main result is the following sharper bound that holds for every n.

Theorem 2. $D(n) \ge n/2 + n/236 - O(\sqrt{n}).$

In a related work, Ábrego et al. [1, 2] consider the problem of matching a given set of n points (n even) by using n/2 disks with exactly two points in each disk. When the disks could overlap, such a perfect matching always exists if the n points are in general position. Similar to Proposition 1, this follows from the result of Dillencourt [6] that every Delaunay triangulation of n points (n even) has a perfect matching. Further, we can shrink the n/2 disks until each disk is empty and has the two points on its boundary.

On the other hand, when the disks are required to be pairwise-disjoint, a perfect matching need not exist. Ábrego et al. [1, 2] used the same circular construction as in our proof of Theorem 1 (details in Section 2) to prove that for even $n \ge 74$, that particular *n*-element point set does not admit any perfect matching with n/2 pairwise-disjoint empty disks. On the positive side, they showed that any *n* points in the plane (in general position) admit a matching of n/4 - O(1) points with pairwise-disjoint empty disks. One can verify that for any *n* points in the plane (not necessarily in general position), their proof implies that there exists a set of n/8 - O(1) disjoint empty disks incident to at least n/4 - O(1) points such that each disk is supported by at least two points. As a consequence we have $D(n) \le n/8 + 3n/4 + O(1) = 7n/8 + O(1)$. This leads to the following upper bound complementing the lower bound in Theorem 2.

Proposition 2. $D(n) \le 7n/8 + O(1)$.

To put our constructions in context, we mention some constructions with disks and squares which are build, similar to ours, by using the same principle of "exerting pressure". For illustration, the reader is referred to the articles [3, 4, 5, 8] for other constructions that we see related in this spirit.

Preliminaries. In this paper we restrict ourselves to disjoint bubbles, in particular to the estimation of D(n). Note that since any two disks supported by a common point must be interior-disjoint, each point supports at most two disks. Since all disks we consider are empty, such a disk Q is incident to a point p if and only if its boundary, ∂Q , is incident to p. Before presenting our lower bound construction, we briefly review the two constructions from [7] yielding the lower bound $D(n) \ge n/2$. In fact, our constructions yielding Theorems 1 and 2 were suggested by them.

1. Let P be a collinear set of n points; refer to Figure 1 (left). Any point must support at least one empty disk, but no empty disk can be supported by more than two points. This point set yields $B(P) \ge |P|/2$, thus $B(n) \ge B(P) \ge |P|/2 = n/2$, and by (1), $D(n) \ge B(n) \ge n/2$.




Figure 1: Two constructions of Hosono and Urabe. Left: a set of n = 11 points on a line. Right: a set of n = 10 points consisting of the center and the vertices of a regular (n - 1)-gon.

2. Assume that n is even. Let P consist of the vertices of a regular (n-1)-gon together with its center (i.e., the center of the circumscribed circle of the regular (n-1)-gon); refer to Figure 1 (right). The circle through any three vertices of the regular (n-1)-gon must be its circumscribed circle, which contains its center and hence is not empty. It follows that any empty disk can be supported by at most two (consecutive) vertices of the regular (n-1)-gon. Thus for every even $n \ge 4$, $B(n) \ge B(P) \ge \lceil (n-1)/2 \rceil = n/2$, and by (1), $D(n) \ge B(n) \ge n/2$.

Notation. An empty disk incident to only one point of P is called a *singleton disk*. In turn, the unique point on a singleton disk is called a *singleton point*.

Given a disk Q, a horizontal line ℓ , so that $Q \cap \ell \neq \emptyset$, let $\xi(Q, \ell)$ denote the point of intersection between ℓ and the vertical diameter of Q. For convenience, points on horizontal lines are frequently identified with their x-coordinates when there is no danger of confusion.

Organization of the paper. In Section 2 we use a circular construction showing that the conjectured equality $D(n) = \lceil n/2 \rceil$ is already false for n as small as 174. We then use this construction to obtain a lower bound of $D(n) \ge n/2 + n/257 - O(1)$ for every n. In Section 3 we present a linear construction and its grid-generalization which achieves our record lower bound of $D(n) \ge n/2 + n/236 - O(\sqrt{n})$ for every n.

2 Circular constructions

For any $n \ge 4$, define an *n*-gadget G_n as a set of *n* points consisting of the center *p* of a unit-radius circle *C* and the vertices q_1, \ldots, q_{n-1} of a regular (n-1)-gon inscribed in the circle. We call *p* the center point and q_1, \ldots, q_{n-1} the boundary points. We refer to Figure 2 for a 12-gadget.

Hosono and Urabe noted that for any even $n \ge 4$, the bubble number $B(G_n)$ of the *n*-gadget G_n is at least n/2. To see this, observe that the circle through any three points on the boundary of C must be C itself, which contains its center p and hence is not empty. It follows that any empty disk can be supported by at most two (consecutive) points on the boundary of C. Thus for every even $n \ge 4$, the n-1 boundary points must support at least $\lceil (n-1)/2 \rceil = n/2$ empty disks.

Hosono and Urabe then conjectured that for every n, $D(n) = \lceil n/2 \rceil$. Our Theorem 1 shows that this conjecture is false, for even n as small as 174. Specifically, we prove that for every even

PSfrag replacements




Figure 2: An *n*-gadget with n = 12.

 $n \ge 174, D(G_n) \ge n/2 + 1.$

Proof of Theorem 1. Consider any bubble set Q of disjoint empty disks for G_n . We will show that $|Q| \ge n/2 + 1$. Recall that any empty disk can be supported by at most 2 boundary points. Thus the n-1 boundary points of G_n support at least $\lceil (n-1)/2 \rceil = n/2$ empty disks in Q. Let $Q \in Q$ be an empty disk supported by the center point p of G_n . If Q is not supported by any boundary point, then the total number of disks in Q is at least n/2 + 1. We next consider the two remaining cases:

- 1. Q is supported by the center point p and only one boundary point, say q_1 . We claim that at least one of the four points q_2, q_3, q_4, q_5 and at least one of the four points $q_{n-1}, q_{n-2}, q_{n-3}, q_{n-4}$ are singleton points.
- 2. Q is supported by the center point p and two consecutive boundary points, say q_1 and q_2 . We claim that at least one of the four points q_3, q_4, q_5, q_6 and at least one of the four points $q_{n-1}, q_{n-2}, q_{n-3}, q_{n-4}$ are singleton points.

If our claims in the two cases are valid, then, excluding the at most two boundary points supporting Q and two singleton points near Q, the remaining at least (n-1)-2-2=n-5 boundary points would support at least $\lceil (n-5)/2 \rceil = n/2 - 2$ empty disks. Adding back the disk Q and the two disks supported by the two singleton points, the total number of empty disks would thus be at least (n/2-2)+1+2=n/2+1, as desired.

We first consider the first part of the claim in case 1, that at least one of the four points q_2, q_3, q_4, q_5 is a singleton point. and prove it by contradiction. Suppose to the contrary that none of q_2, q_3, q_4, q_5 is a singleton point. Then there must exist a disk $Q_1 \in \mathcal{Q}$ supported by both q_2 and q_3 , and a disk $Q_2 \in \mathcal{Q}$ supported by both q_4 and q_5 . We will show that for every even $n \geq 174$, the two disks Q_1 and Q_2 intersect, which contradicts our assumption of disjoint empty disks in the bubble set.

Put $\theta = \pi/(n-1)$. Let r be the radius, and o the center, of the disk Q. For $i \in \{1, 2\}$, the radius r_i of the disk Q_i is uniquely determined by the distance d_i from the disk center o_i to p,

according to the cosine rule:

$$r_i^2 = 1^2 + d_i^2 - 2 \cdot 1 \cdot d_i \cdot \cos \theta = d_i^2 - 2d_i \cos \theta + 1.$$
(2)

Without loss of generality, we assume that both Q_1 and Q_2 are tangent to Q. Then the distance between o and o_i is exactly $r + r_i$. Thus again by the cosine rule, we have

$$(r+r_i)^2 = r^2 + d_i^2 - 2 \cdot r \cdot d_i \cdot \cos \angle opo_i.$$
(3)

Given r and $\angle opo_i$ (these parameters will be fixed later in the argument, in equations (5) and (6)), the distance d_i and the radius r_i for each $i \in \{1, 2\}$ are uniquely determined by the two equations (2) and (3). This can be seen by a morphing argument. Observe that as d_i changes continuously, the portion of the disk Q_i inside the circle C also changes continuously: the portion for a smaller d_i properly contains the portion for a larger d_i . Subsequently, given $\angle o_1 po_2$, the distance $|o_1 o_2|$ between o_1 and o_2 is uniquely determined by applying the cosine rule once more:

$$|o_1 o_2|^2 = d_1^2 + d_2^2 - 2 \cdot d_1 \cdot d_2 \cdot \cos \angle o_1 p o_2.$$
(4)

To verify that Q_1 and Q_2 intersect, it suffices to show $|o_1 o_2| < r_1 + r_2$ for every even $n \ge 174$.

Before we proceed with the calculation, observe that the second part of the claim in case 1, that at least one of the four points $q_{n-1}, q_{n-2}, q_{n-3}, q_{n-4}$ is a singleton point, can be proved by contradiction in a similar way, where Q_1 is a disk supported by both q_{n-1} and q_{n-2} , and Q_2 is a disk supported by both q_{n-3} and q_{n-4} . In particular, equations (2), (3), and (4) still apply. Moreover, the same argument also applies to case 2. Indeed case 2 is the bottleneck case since the disk Q in case 2 exerts less pressure on the four points near it on each side than the disk Q in case 1 does. Since the two parts of the claim in case 2 are symmetric, it suffices to do the calculation for the first part of the claim in case 2 that at least one of q_3, q_4, q_5, q_6 is a singleton point.

We now do the calculation for this part, where Q is supported by both q_1 and q_2 , Q_1 is supported by both q_3 and q_4 , and Q_2 is supported by both q_5 and q_6 . We clearly have

$$\angle opo_1 = 4\theta, \quad \angle opo_2 = 8\theta, \quad \angle o_1po_2 = 4\theta.$$
 (5)

Since $|pq_1| = |pq_2| = 1$, $|op| = |oq_1| = |oq_2| = r$, and $\angle opq_1 = \angle opq_2 = \theta$, we also have

$$r = 1/(2\cos\theta). \tag{6}$$

Plug the parameters in (5) and (6) into the three equations (2), (3), and (4). One can verify that for n = 174, the values in (7) satisfy the equations.

$$\theta = 1.0404...^{\circ}, \quad r = 0.5000...,$$

$$d_1 = 1.0762..., \quad d_2 = 1.0113..., \quad r_1 = 0.0785..., \quad r_2 = 0.0215..., \quad (7)$$

$$|o_1o_2| = 0.0997..., \quad r_1 + r_2 = 0.1000....$$

In particular, $|o_1o_2| < r_1 + r_2$ for n = 174, and hence for every even $n \ge 174$. Thus for every even $n \ge 174$, the two disks Q_1 and Q_2 intersect, giving the desired contradiction. This completes the proof.

Theorem 3. For every $n, D(n) \ge (1/2 + 1/257)n - O(1)$.




Figure 3: The set P consists of m 174-gadgets and m 340-gadgets in an alternating pattern along a horizontal line. Each bridge disk (three are shown here) is supported by two or more points from two consecutive gadgets.

Proof. Let $m = \lfloor n/(174+340) \rfloor$, and let $\hat{n} = (174+340)m$. Since D(n) is a non-decreasing function in n, it suffices to show that $D(\hat{n}) \ge (1/2 + 1/257)\hat{n} = \hat{n}/2 + 2m$. To this end, we will construct a set P of \hat{n} points, and then show that $D(P) \ge \hat{n}/2 + 2m$.

The set P of \hat{n} points are constructed as follows. Refer to Figure 3. Arrange m 174-gadgets and m 340-gadgets in an alternating pattern along a horizontal line, with a very large horizontal distance d between the center points of consecutive gadgets. For each gadget, place the center point and one boundary point on the x-axis, so that the remaining boundary points are symmetric about the x-axis (this is possible because the number of points in each gadget is even). Scale the m 174-gadgets very slightly such that all 2m gadgets are sandwiched between two horizontal lines symmetric about the x-axis, with each of the two lines containing exactly one boundary point from each gadget (again this is possible because the number of points in each gadget is even).

We proceed to show that $D(P) \ge \hat{n}/2 + 2m$. By construction, the scaling in particular, any disk supported by two points from two non-consecutive gadgets cannot be empty. Let Q be any bubble set of disjoint empty disks for P. Then each disk in Q is either a *local disk* supported by one or more points from the same gadget, or a *bridge disk* supported by two or more points from two consecutive gadgets, that is, a 174-gadget and a 340-gadget. We charge each bridge disk to the 174-gadget that supports it. By Theorem 1 (note that scaling does not affect the result), each 174-gadget supports at least 174/2 + 1 disks in Q (both local and bridge). It remains to show that each 340-gadget supports at least 340/2 + 1 local disks in Q.

Since the horizontal distance d between the center points of consecutive gadgets is very large, every bridge disk of radius $\Omega(d)$ appears as an approximately straight line near either gadget that supports it. Consequently each 340-gadget can support at most two disjoint bridge disks that appear as two approximately parallel lines. If a 340-gadget does not support any bridge disk, then by Theorem 1 it supports at least 340/2 + 1 local disks, as desired. We next consider a 340-gadget supporting either one bridge disk or two bridge disks.

First consider a 340-gadget supporting only one bridge disk. We approximate the bridge disk by a straight line ℓ as justified earlier. Refer to Figure 4. Without loss of generality, we orient the construction such that ℓ is horizontal and supports the gadgets from below. We consider two cases

PSfrag replacements




Figure 4: A 340-gadget supporting a bridge disk approximated by a line ℓ (the figure is not drawn in precise proportion; the angles of the circular arcs between consecutive boundary points are enlarged to show the details).

analogous to the two cases in the proof of Theorem 1:

- 1. ℓ is supported by only one boundary point, say q_1 . We claim that at least two of the four points q_2, q_3, q_4, q_5 and at least two of the four points $q_{339}, q_{338}, q_{337}, q_{336}$ are singleton points.
- 2. ℓ is supported by two consecutive boundary points, say q_1 and q_2 . We claim that at least two of the four points q_3, q_4, q_5, q_6 and at least two of the four points $q_{339}, q_{338}, q_{337}, q_{336}$ are singleton points.

As in the proof of Theorem 1, case 2 is the bottleneck case, and by symmetry we only need to show that if ℓ is supported by two consecutive boundary points q_1 and q_2 , then at least two of the four points q_3, q_4, q_5, q_6 are singleton points. To show this by contradiction, we use the same technique of two disks Q_1 and Q_2 . Suppose the contrary that at most one of the four points q_3, q_4, q_5, q_6 is a singleton point. Then we must have two disks Q_1 and Q_2 in one of the following three subcases:

- i. Q_1 is supported by q_3 and q_4 , and Q_2 supported by q_5 and q_6 ;
- ii. Q_1 is supported by q_3 and q_4 , and Q_2 is supported by q_6 and q_7 ;
- iii. Q_1 is supported by q_4 and q_5 , and Q_2 is supported by q_6 and q_7 .

Put k = 340, and $\theta = \pi/(k-1) = 0.5309...^{\circ}$. As in the proof of Theorem 1, let r be the radius, and o the center, of the disk Q. For $i \in \{1, 2\}$, let r_i be the radius, and o_i the center, of the disk Q_i , and let d_i be the distance from o_i to the center point p. Then the two equations (2) and (4) continue to hold.

In each subcase, we can assume that both Q_1 and Q_2 are tangent to ℓ , just as we assumed they are tangent to Q in the proof of Theorem 1. Let q be the midpoint of q_1 and q_2 . Then $|pq| = \cos \theta$, and we have the following equation analogous to (3):

$$d_i \cos \angle qpo_i + r_i = \cos \theta. \tag{8}$$

We also have the following three sets of parameters, analogous to the parameters in (5), for the three subcases respectively:

$$\angle qpo_1 = 4\theta, \quad \angle qpo_2 = 8\theta, \quad \angle o_1po_2 = 4\theta, \tag{9}$$

$$\angle qpo_1 = 4\theta, \quad \angle qpo_2 = 10\theta, \quad \angle o_1po_2 = 6\theta, \tag{10}$$

 $\angle qpo_1 = 6\theta, \quad \angle qpo_2 = 10\theta, \quad \angle o_1po_2 = 4\theta.$ (11)

Plug the parameters in (9), (10), (11), respectively, into the three equations (2), (8), (4) (this is analogous to plugging (5) into (2), (3), (4) in the proof of Theorem 1). One can now verify that the following three sets of values in (12), (13), (14), respectively, satisfy the equations.

$$d_1 = 0.9333..., \quad d_2 = 0.9854..., \quad r_1 = 0.0672..., \quad r_2 = 0.0172..., \\ |o_1 o_2| = 0.0631..., \quad r_1 + r_2 = 0.0845....$$
(12)

$$d_1 = 0.9333..., \quad d_2 = 0.9919..., \quad r_1 = 0.0672..., \quad r_2 = 0.0122..., \\ |o_1 o_2| = 0.0794..., \quad r_1 + r_2 = 0.0795....$$
(13)

$$d_1 = 0.9721\dots, \quad d_2 = 0.9919\dots, \quad r_1 = 0.0293\dots, \quad r_2 = 0.0122\dots, \\ |o_1 o_2| = 0.0414\dots, \quad r_1 + r_2 = 0.0415\dots$$
(14)

Note that $|o_1o_2| < r_1 + r_2$ for each of the three subcases. Thus Q_1 and Q_2 intersect, giving the desired contradiction.

We have shown that at least 4 of the 8 boundary points near ℓ are singleton points. Excluding the at most 2 boundary points on ℓ and 4 singleton points near ℓ , the remaining at least (k-1)-2-4 = k-7 boundary points support at least $\lceil (k-7)/2 \rceil = k/2 - 3$ empty disks. Adding back the 4 empty disks supported by the 4 singleton points, the total number of local disks is at least k/2 - 3 + 4 = k/2 + 1. That is, the 340-gadget supports at least 340/2 + 1 local disks.

Finally consider a 340-gadget supporting two bridge disks. Recall that the two bridge disks appear as two approximately parallel lines. Thus the 8 boundary points near one line are approximately opposite to (and hence disjoint from) the 8 boundary points near the other line. And thus the same analysis for the two lines show that there are altogether 8 singleton points among the 16 boundary points. Excluding the at most 4 boundary points on the two lines and 8 singleton points near the two lines, the remaining at least (k-1) - 4 - 8 = k - 13 boundary points support at least $\lceil (k-13)/2 \rceil = k/2 - 6$ empty disks. Adding back the 8 empty disks supported by the 8 singleton points, the total number of local disks is at least k/2 - 6 + 8 = k/2 + 2. That is, the 340-gadget supports at least 340/2 + 2 local disks, more than what we desired.

In summary, we have shown that each 174-gadget supports at least 174/2 + 1 (both local and bridge) disks in Q, and that each 340-gadget supports at least 340/2 + 1 local disks in Q. Summing up the number of disks over all m 174-gadgets and m 340-gadgets, we have $D(P) \ge \hat{n}/2 + 2m$. This completes the proof of Theorem 3.

3 Linear constructions

In Section 3.1 we outline a preliminary (simpler) construction that is easier to analyze. In Section 3.2 we present a refinement which yields a better lower bound.

3.1 Preliminary construction

For every n, we construct a set P of n points, with $D(P) \ge n/2 + n/966 - O(1)$. We first describe a construction for every $n = C_1 j + C_2$, j = 1, 2..., for some suitable integers $C_1, C_2 \ge 1$. We then extend the construction for every n.

To achieve this bound we enforce that $\Omega(n)$ points of P are singleton points. Place the n points on three parallel horizontal lines ℓ_1, ℓ_2, ℓ_3 , as illustrated in Figure 5. The distance between ℓ_1 and ℓ_3 is 4; the line in the middle, ℓ_2 , is equidistant from ℓ_1 and ℓ_3 .

Let $0 < \delta \le 1/50$, so that $4/\delta$ is an integer; this condition will be ensured by the choice of δ . Let $j \ge 1$ be a positive integer. We first place the points on the three lines in a regular fashion.




Figure 5: Construction and three bubbles: one supported by a point on ℓ_2 and a point on ℓ_3 , one supported by a point on ℓ_2 and two points on ℓ_1 , and one supported by a point on each of the three lines.

Then shift right very slightly by the same distance ε those on ℓ_3 , so that no four points of P are co-circular on an empty circle. Place $(4/\delta+1)j+1$ uniformly distributed points on ℓ_1 , with any two consecutive points at distance δ ; proceed in the same way for ℓ_3 . Place j+1 uniformly distributed points on ℓ_2 , labeled p_1, \ldots, p_{j+1} , with any two consecutive points at distance $4 + \delta$. The leftmost points on the three lines are aligned vertically, say at x = 0. The small perturbation (shift) does not affect the points on ℓ_1 and ℓ_2 .

We refer to ℓ_1 and ℓ_3 as the *dense lines* and to the middle line ℓ_2 as the *sparse line*. The total number of points (on the three lines) is

$$n = \left(\frac{4}{\delta} + \frac{4}{\delta} + 3\right)j + 3 = \left(\frac{8}{\delta} + 3\right)j + 3.$$
(15)

In an alternative view of this construction, the initial points (before perturbation) are placed on the boundary and on the horizontal line through the center of the rectangle $R = [0, (4 + \delta)j] \times$ [0, 4]. We refer to the j - 1 points of P in the interior of R, i.e., on ℓ_2 , as *interior points*. As yet another view, the initial points are placed on the boundaries of the j axis-aligned rectangles $[(4 + \delta)i, (4 + \delta)(i + 1)] \times [0, 4], i = 0, \ldots, j - 1.$

Let \mathcal{Q} be a set of pairwise-disjoint bubbles (disks) supported by P. Let \mathcal{Q}' be the subset of disks that are either incident to some interior point of P or are incident to exactly three points of P but not to p_1 or p_{j+1} . Observe that the total number of disks in \mathcal{Q} incident to p_1 or p_{j+1} is O(1).

First, observe that the radius of any empty disk incident to an interior point and some other point is at least 1. Second, observe also that if an empty disk whose center is in R intersects both ℓ_1 and ℓ_3 , then its radius is at least 2. We refer to such disks (of the above two types) as *large*, and to any other disk as *small*. Third, observe that the radius of any empty disk supported by an interior point is at most

$$\sqrt{16+\delta^2}/2 < 2+\delta/2.$$

Similarly, the radius of any empty disk supported by a point in ℓ_1 and a point in ℓ_3 is also bounded from above by the same quantity. It follows that no empty disk can be incident to two interior points (on ℓ_2).

The next two lemmas show that if an interior point $p \in P$ on the sparse line supports a disk that is also supported by another point of P, then sufficiently many points on one of the two dense lines must be singleton points. Moreover these "forced" points can be uniquely "assigned" to the corresponding disk incident to p. This suggests that in a minimum-size bubble set supported by P, each interior point must be a singleton point. A calculation making this intuition precise is at the end of this subsection. **Lemma 1.** Let p_1, p_2, p_3, p_4 be four consecutive points on the dense line ℓ_3 . Let $Q_{1,2}$ and $Q_{3,4}$ be two disks, the first supported by p_1 and p_2 , and the second supported by p_3 and p_4 . Denote by ℓ'_3 the horizontal line at distance $(1 - \sqrt{3}/2)\delta$ above ℓ_3 . If both disks lie strictly below ℓ'_3 , then they overlap in their interior (hence they are not disjoint).

Proof. Refer to Figure 6. Let $\omega_{1,2}$ be the disk supported by p_1 and p_2 and tangent to ℓ'_3 . Similarly, let $\omega_{3,4}$ be the disk supported by p_3 and p_4 and tangent to ℓ'_3 . Denote by π_3 the closed half-plane below ℓ_3 . We clearly have that $\omega_{1,2} \cap \pi_3 \subset Q_{1,2} \cap \pi_3$ and $\omega_{3,4} \cap \pi_3 \subset Q_{3,4} \cap \pi_3$. Therefore, it suffices to show that $\omega_{1,2}$ and $\omega_{3,4}$ are tangent to each other, since then, it follows that $Q_{1,2}$ and $Q_{3,4}$ are not pairwise-disjoint.




Figure 6: Proof of Lemma 1.

The two disks $\omega_{1,2}$ and $\omega_{3,4}$ are congruent, and since they are both tangent to ℓ'_3 , their centers lie on the same horizontal line. It suffices to show that their radius r is at least δ . It is well-known that the radius of a circle circumscribed to a triangle with side lengths a,b,c is equal to abc/(4S), where S stands for the triangle area. It follows that

$$r = \frac{\delta \left[\left(\frac{\delta}{2}\right)^2 + \left(\left(1 - \frac{\sqrt{3}}{2}\right)\delta \right)^2 \right]}{(2 - \sqrt{3})\delta^2} = \delta$$

Hence the two disks $\omega_{1,2}$ and $\omega_{3,4}$ are tangent to each other, as desired, and this completes the proof.

Lemma 2. Let $\ell \in {\ell_1, \ell_3}$. Let $Q \in Q'$ be a large empty disk centered at o that intersects ℓ . Then there exist two singleton points of P on ℓ at horizontal distance at most $7\delta/2$ from o, and that are not incident to Q. As such, these points can be uniquely assigned to Q (and to no other disk in Q).

Proof. Assume for simplicity that $\ell = \ell_3$, and put $u = \xi(Q, \ell_3)$. Then u is contained in the interval between two consecutive points, say q and q'. Refer to Figure 7 (left). For simplicity suppose that u is contained in the left half-interval of qq' (the other case is symmetric).

If u is the midpoint of qq', since Q is empty, Q is either incident to both q and q' or to neither of them. If u is not the midpoint of qq', then Q can be only incident to q, but not to q', if at all.

Let q_i , $i = 1, \ldots, 4$, be the first four points of $P \cap \ell_3$ right of u that are not incident to Q. Observe that the farthest is at distance at most $9\delta/2$ from u, hence the x-coordinate of the center of the farthest disk incident to some pair $q_i, q_{i+1}, i = 1, 2, 3$, is at most 4δ . Let r denote the radius of Q; obviously, $r \ge 1$. Let y be the vertical distance between the midpoint of q_3q_4 and Q. It suffices to show that $y \le 0.1337\delta$, since then, $y \le 0.1337\delta < (1 - \sqrt{3}/2)\delta$. Recall that $\delta \le 1/50$, so we have

$$y \le r - \sqrt{r^2 - 16\delta^2} = \frac{16\delta^2}{r + \sqrt{r^2 - 16\delta^2}} \le \frac{16\delta^2}{1 + \sqrt{1 - 16\delta^2}} \le \frac{16\delta^2}{1 + \sqrt{1 - 16\delta^2}} \le \frac{16\delta^2}{1.997} \le 8.02\delta^2.$$




Figure 7: Proof of Lemma 2.

Set now $\delta = 1/60$, which ensures that $y \le 8.02\delta^2 \le 0.1337\delta$. Moreover, $4/\delta = 240$ is an integer multiple of δ , as desired.

It remains to show the existence of two singleton points in the vicinity of u. We show that there is at least one on each side of u. According to our previous calculation, any disk incident to some pair $q_i, q_{i+1}, i = 1, 2, 3$, must lie below the line ℓ'_3 . By Lemma 1, it follows that at least one of the points $q_i, i = 1, \ldots, 4$, is a singleton point. Note that since only consecutive points can be "matched" on an empty small circle, this implies that at least one of the points $q_i, i = 1, \ldots, 3$, is a singleton point. The farthest of these three points is at distance at most $7\delta/2$ from u. Similarly, there is a singleton point left of u at distance at most $7\delta/2$ from u. Observe that the choice of these points may depend on the number of points on ℓ_3 (zero, one or two) incident to Q.

Recall that by construction, each disk in \mathcal{Q} is incident to at most three points in P. Moreover, no disk in \mathcal{Q} is incident to two interior points (on ℓ_2).

Lemma 3. Let $Q \in Q'$ be an empty disk. Then there exists a subset $Q_1 \subset Q$ of disks such that $Q \in Q_1$ and Q_1 is supported by exactly m points of P, where $|Q_1| \ge (m+1)/2$.

Proof. Assume that $Q \in Q'$ is incident to (exactly) *i* points of *P*. By the previous observation, we know that $1 \le i \le 3$. We distinguish two cases:

Case 1. Q is incident to at least one interior point p (on ℓ_2).

If i = 1, then set $Q_1 = \{Q\}$, and the claimed inequality holds with m = 1.

If i = 2, then Q is also incident to a point on $q \in \ell_3$ (the case $q \in \ell_1$ is symmetric). Then by Lemma 2 there exist two points q_1, q_2 in the neighborhood of q (left and right of q), each incident to exactly one disk in Q, say Q_1 and Q_2 . Then set $Q_1 = \{Q, Q_1, Q_2\}$, and the claimed inequality holds with m = 4: $|Q_1| = 3 \ge (4+1)/2$.

If i = 3, assume first that Q is also incident to two points $q, q' \in \ell_3$ (the case with two points on ℓ_1 is symmetric). Then by Lemma 2 there exist two points (left of q and right of q'), each incident to exactly one disk in Q, say Q_1 and Q_2 , and such that Q_1 and Q_2 are each incident to exactly one point in P. Then set $Q_1 = \{Q, Q_1, Q_2\}$, and the claimed inequality holds with m = 5: $|Q_1| = 3 \ge (5+1)/2$. Second, assume that Q is also incident to a point $q \in \ell_1$ and a point $q' \in \ell_3$. Again by Lemma 2, there exist two points on ℓ_1 (left and right of q) and two points on ℓ_3 (left and right of q'), each incident to exactly one disk in Q. Let Q_1, Q_2, Q_3, Q_4 be these four disks. Then set $Q_1 = \{Q, Q_1, Q_2, Q_3, Q_4\}$, and the claimed inequality holds with m = 7: $|Q_1| = 5 \ge (7+1)/2$.

Case 2. Q is not incident to any interior point (on ℓ_2). Then, by the definition of Q', Q is not incident to any point on ℓ_2 . It follows that Q is incident to three points on $\ell_1 \cup \ell_3$. By Lemma 2, there exist four other points, two on ℓ_1 and two on ℓ_3 , each incident to exactly one disk in Q. Let

 Q_1, Q_2, Q_3, Q_4 be these four disks. Then set $Q_1 = \{Q, Q_1, Q_2, Q_3, Q_4\}$, and the claimed inequality holds with m = 8: $|Q_1| = 5 \ge (8+1)/2$.

This completes our case analysis and the proof of the lemma.

We now analyze the construction. We have set $\delta = 1/60$, thus $4/\delta = 240$. Let $Q_2 \subset Q$ be the subset of disks constructed from Q' by taking the union of all disks in Q_1 from all cases in the analysis in the proof of Lemma 3. Clearly $Q' \subset \cup Q_1 = Q_2 \subset Q$. Put $d = |Q_2|$. Let $T \subset P$ be the set of points incident to disks in Q_2 ; write t = |T|. Observe that each interior point yields one inequality of the form $|Q_1| \ge (m+1)/2$, so there are at least j-1 such inequalities. The corresponding sets Q_1 are pairwise disjoint by construction, thus by adding up all these inequalities yields

$$|\mathcal{Q}_2| \ge \frac{t+j-1}{2}.$$

Apart from O(1) exceptions, each disk in $\mathcal{Q} \setminus \mathcal{Q}_2$ is incident to at most two points in P. Consequently,

$$|\mathcal{Q}| = |\mathcal{Q}_2| + |\mathcal{Q} \setminus \mathcal{Q}_2| \ge \frac{t+j-1}{2} + \frac{n-t}{2} - O(1) = \frac{n+j}{2} - O(1).$$
(16)

By substituting the value of j resulting from (15), it follows that the number of disjoint empty disks in any disjoint bubble set for P is at least

$$\frac{n}{2} + \frac{j}{2} - O(1) = \frac{n}{2} + \frac{n}{16/\delta + 6} - O(1) = \frac{n}{2} + \frac{n}{966} - O(1).$$
(17)

Write $C_1 = 8/\delta + 3$. To extend the construction for every n, we add a small cluster of at most $C_1 - 1$ collinear points on ℓ_3 , far away from the "main" construction above. Observe that there can be at most two disks incident to both the main part and the additional cluster. Recall that each disk is incident to at most three points (this condition can be maintained), and conclude that the previous analysis yields the same bound.

3.2 A refined approach

We modify the previous construction and refine its analysis. These changes are motivated as follows.

- 1. The lower bound can be raised by "replicating" the construction vertically: use k + 1 dense lines separated by k sparse lines in between. The analysis however becomes more involved. Our preliminary construction has k = 1.
- 2. In our previous analysis we only require that small disks incident to two consecutive points on ℓ_3 and near a large disk intersecting ℓ_3 lie below ℓ'_3 . Observe that such a triplet of disjoint disks can be enlarged until the three disks become pairwise tangent. Thus δ can be increased and this brings another improvement in the lower bound on D(n).

Lemma 4. Let $x \leq 1/10$. Let C be a disk of unit radius centered at (0,1). Let C_1 be a disk incident to the points (3x/2,0) and (5x/2,0). Let C_2 be a disk incident to the points (7x/2,0) and (9x/2,0). If the three disks are pairwise tangent then $x = 2\sqrt{\lambda} = 0.03486...$, where $\lambda = \frac{4}{945}(7 - 4\sqrt{3})$ is the smaller solution of the quadratic equation $893025\lambda^2 - 52920\lambda + 16 = 0$.

Proof. Let C_1 be centered at $(2x, -y_1)$, and C_2 be centered at $(4x, -y_2)$, where $0 < y_2 < y_1$. Denote by r_i the radius of C_i , i = 1, 2. We have five unknowns, x, r_1, r_2, y_1, y_2 , and five connecting equations: two expressing incidences and three expressing tangencies. Put x = 2z.

$$z^{2} + y_{1}^{2} = r_{1}^{2}$$

$$z^{2} + y_{2}^{2} = r_{2}^{2}$$

$$16z^{2} + (1 + y_{1})^{2} = (1 + r_{1})^{2}$$

$$64z^{2} + (1 + y_{2})^{2} = (1 + r_{2})^{2}$$

$$16z^{2} + (y_{1} - y_{2})^{2} = (r_{1} + r_{2})^{2}$$

After eliminating y_1 and y_2 the system becomes

$$16z^{2} + \left(1 + \sqrt{r_{1}^{2} - z^{2}}\right)^{2} = (1 + r_{1})^{2}$$
$$64z^{2} + \left(1 + \sqrt{r_{2}^{2} - z^{2}}\right)^{2} = (1 + r_{2})^{2}$$
$$16z^{2} + \left(\sqrt{r_{1}^{2} - z^{2}} - \sqrt{r_{2}^{2} - z^{2}}\right)^{2} = (r_{1} + r_{2})^{2}$$

Recall that $z \leq r_1, r_2$, and make the substitutions

$$r_1 = \frac{z}{\cos \alpha}, \quad r_2 = \frac{z}{\cos \beta}, \text{ where } \alpha, \beta \in (0, \pi/2).$$

The above system can be rewritten as

$$16z^{2} + (1+z\tan\alpha)^{2} = \left(1+\frac{z}{\cos\alpha}\right)^{2}.$$
$$64z^{2} + (1+z\tan\beta)^{2} = \left(1+\frac{z}{\cos\beta}\right)^{2}.$$
$$16z^{2} + (z\tan\alpha - z\tan\beta)^{2} = z^{2}\left(\frac{1}{\cos\alpha} + \frac{1}{\cos\beta}\right)^{2}.$$

Express $\cos \alpha$ and $\tan \alpha$ as functions of $\tan(\alpha/2)$:

$$\cos \alpha = \frac{1-s^2}{1+s^2}$$
, and $\tan \alpha = \frac{2s}{1-s^2}$, where $s = \tan \frac{\alpha}{2}$.

Similarly, express $\cos \beta$ and $\tan \beta$ as functions of $\tan(\beta/2)$:

$$\cos\beta = \frac{1-t^2}{1+t^2}$$
, and $\tan\beta = \frac{2t}{1-t^2}$, where $t = \tan\frac{\beta}{2}$

Now the first equation of our system can be rewritten as

$$16z^{2} + \left(1 + z\frac{2s}{1 - s^{2}}\right)^{2} = \left(1 + z\frac{1 + s^{2}}{1 - s^{2}}\right)^{2}.$$
 (18)

By rearranging terms, (18) is equivalent to

$$2(1-s^2)(1-s)^2 = 15z(1-s^2)^2.$$
(19)

The solution s = 1 is infeasible, thus (18) yields

$$s = \frac{2 - 15z}{2 + 15z}.\tag{20}$$

Similarly, the second equation of our system can be rewritten as

$$64z^{2} + \left(1 + z\frac{2t}{1 - t^{2}}\right)^{2} = \left(1 + z\frac{1 + t^{2}}{1 - t^{2}}\right)^{2}.$$
(21)

By rearranging terms, (21) is equivalent to

$$2(1-t^2)(1-t)^2 = 63z(1-t^2)^2.$$
(22)

The solution t = 1 is infeasible, thus (22) yields

$$t = \frac{2 - 63z}{2 + 63z}.\tag{23}$$

Recall that s, t > 0, thus (20) and (23) require z < 2/15 and z < 2/63; overall, z < 2/63. We can now express all trigonometric functions occurring in the third equation of our system as functions of z.

$$\tan \alpha = \frac{2s}{1-s^2} = \frac{4-225z^2}{60z},$$

$$\tan \beta = \frac{2t}{1-t^2} = \frac{4-3969z^2}{252z},$$

$$\cos \alpha = \frac{1-s^2}{1+s^2} = \frac{60z}{4+225z^2},$$

$$\cos \beta = \frac{1-t^2}{1+t^2} = \frac{252z}{4+3969z^2}.$$

Finally we substitute this in the third equation of the system

$$16 + (\tan \alpha - \tan \beta)^2 = \left(\frac{1}{\cos \alpha} + \frac{1}{\cos \beta}\right)^2.$$

and obtain an equation in z only:

$$16 + \left(\frac{4 - 225z^2}{60z} - \frac{4 - 3969z^2}{252z}\right)^2 = \left(\frac{4 + 225z^2}{60z} + \frac{4 + 3969z^2}{252z}\right)^2,$$
(24)

or equivalently,

 $16 \cdot 1260^2 z^2 + (64 + 15120z^2)^2 = (104 + 24570z^2)^2.$ ⁽²⁵⁾

Make the substitution $\lambda = z^2$ and (after simplifying by 42) the quartic equation in z (25) becomes a quadratic equation in λ :

$$893025\lambda^2 - 52920\lambda + 16 = 0. \tag{26}$$

Its solutions are

$$\lambda_{1,2} = \frac{4}{945} \left(7 \pm 4\sqrt{3}\right).$$

Correspondingly, we have

$$z_{1,2} = \sqrt{\frac{4}{945} \left(7 \pm 4\sqrt{3}\right)}.$$

The larger solution z_1 is infeasible, as it contradicts our assumption z < 2/63. Finally

$$x = 2z_2 = 2\sqrt{\frac{4}{945} \left(7 - 4\sqrt{3}\right)} = 0.03486\dots$$

Refined construction. Set $\delta = 1/29 < 2\sqrt{\lambda} = 0.03486...$, where $\lambda = \frac{4}{945}(7 - 4\sqrt{3})$. The points on ℓ_1 and ℓ_3 are (initially) placed as before, at multiples of δ . The points on ℓ_2 are (initially) placed as before, at multiples of $4 + \delta$. We now replicate the above construction vertically: use k + 1 dense lines separated by k sparse lines in between. In total there are 2k + 1 lines, labeled $\ell_1, \ldots, \ell_{2k+1}$. Write $\mathcal{L} = \{\ell_1, \ldots, \ell_{2k+1}\}$. Finally shift right the points on every other odd-numbered line $\ell_3, \ell_7, \ell_{11}, \ldots$ by a small ε so that no four points of P are co-circular on an empty circle. Our preliminary construction has k = 1. The total number of points on the 2k + 1 lines is

$$n = ((k+1)(4/\delta + 1) + k)j + (2k+1) = jk(4/\delta + 2) + (4/\delta + 1)j + (2k+1).$$
(27)

For a fixed dense line $\ell \in \mathcal{L}$, we commonly denote the points in P on ℓ by q_i , where $i \in \mathbb{N}$. Suppose a large disk intersects a dense line; since almost the entire disk lies either above or below the line, we say that the disk intersects ℓ from above or from below.

Observation 1. Let ℓ be one of the dense horizontal lines, and $I \subset \ell$ be an interval of length $|I| = 8\delta$. Let $\{Q_1, \ldots, Q_h\}$ be large disks in Q intersecting ℓ , so that $\xi(Q_i, \ell) \in I$, for each $i = 1, \ldots, h$. Then $h \leq 2$. Moreover, Q_1 and Q_2 intersect ℓ from different sides, one from above and one from below.

Observation 2. Let $Q \in \mathcal{Q}$ be a large disk incident to at least one point in P on a dense line ℓ . Let m_j denote the midpoint of the subdivision interval $[q_j, q_{j+1}], j \in \mathbb{N}$, on ℓ .

(i) If Q is incident to exactly one point on ℓ , namely q_i , then $|q_i m_{i+1}| \leq 2\delta$ and $|m_{i-2}q_i| \leq 2\delta$.

(ii) If Q is incident to exactly two points on ℓ , namely q_i and q_{i+1} , then $|m_i m_{i+2}| \leq 2\delta$ and $|m_{i-2}m_i| \leq 2\delta$.

Corollary 1. Let $Q \in Q$ be a disk incident to at least one point in P on a dense line ℓ .

(i) If Q is incident to exactly one point on ℓ , namely q_i , then there exists a singleton point among $\{q_{i-3}, q_{i-2}, q_{i-1}\}$, and one among $\{q_{i+1}, q_{i+2}, q_{i+3}\}$.

(ii) If Q is incident to exactly two points on ℓ , namely q_i and q_{i+1} , then there exists a singleton point among $\{q_{i-3}, q_{i-2}, q_{i-1}\}$, and one among $\{q_{i+2}, q_{i+3}, q_{i+4}\}$.

Proof. We first prove statement (ii). Let Q_j denote a disk incident to the points q_j and q_{j+1} . If there exist two disks Q_{i+2} and Q_{i+4} such that the three disks $Q = Q_i$, Q_{i+2} , and Q_{i+4} are pairwise disjoint, shrink Q until it becomes tangent to ℓ_3 (if not already tangent). The three disks remain pairwise disjoint. By Observation 2(ii), the horizontal distance between the centers of any two consecutive disks is at most 2δ . However, by the choice of δ in Lemma 4, the three disks must be pairwise tangent in the final configuration, a contradiction.

Statement (i) is proved by a similar argument, using the two inequalities in Observation 2(i).

For two disks $Q_1, Q_2 \in \mathcal{Q}$ intersecting a common line ℓ , denote by $f(Q_1, Q_2)$ the number of points of P in the interval $[\xi(Q_1, \ell), \xi(Q_2, \ell)]$ that are not incident to Q_1 or Q_2 .

Define an undirected graph G with vertex set the set of large disks in \mathcal{Q} incident to points on dense lines in \mathcal{L} . Two disks $Q_1, Q_2 \in \mathcal{Q}$ are adjacent in G if they are each incident to points of Pon a common dense line $\ell \in \mathcal{L}$, and $f(Q_1, Q_2) \in \{0, 1, 2, 3, 4, 5\}$. By Observation 1, if $Q_1, Q_2 \in \mathcal{Q}$ are adjacent in G then the two large disks intersect the common line one from below, and one from above.

Lemma 5. Let $a \leq 6$ be a positive integer, and $\delta \leq 1/29 < 2\sqrt{\lambda} = 0.03486...$ Define the function

$$g(a,\delta) = \frac{a^2\delta^2}{1+\sqrt{1-a^2\delta^2}}.$$

Then g() is an increasing function of a, and $g(3,\delta) + g(6,\delta) \le 0.9\delta$.

Proof. The first claim is easy to verify, so it remains to check the second claim. We have

$$g(3,\delta) + g(6,\delta) = \frac{9\delta^2}{1 + \sqrt{1 - 9\delta^2}} + \frac{36\delta^2}{1 + \sqrt{1 - 36\delta^2}}$$
$$\leq \frac{9\delta^2}{1.99} + \frac{36\delta^2}{1.97} \leq 23\delta^2 \leq 0.9\delta,$$

as required.

Lemma 6. Let $\ell \in \mathcal{L}$ be a dense horizontal line, and $Q_1, Q_2 \in \mathcal{Q}$ be two large disks intersecting ℓ (one from above and one from below, with the center of Q_1 left of the center of Q_2), such that the pair (Q_1, Q_2) is an edge in G. Then there exist 4 singleton points on ℓ uniquely associated with this edge of G.

Proof. Let $r_i \ge 1$ be the radius of Q_i , i = 1, 2. Let q_i be the leftmost point on ℓ incident to Q_1 and q_j be the rightmost point on ℓ incident to Q_2 ; clearly $i \le j$. Write $f := f(Q_1, Q_2)$ for simplicity. We distinguish 6 cases.

Case 0. f = 0. It suffices to show that q_{j+1} and q_{j+2} are singleton points. By a symmetric argument q_{i-1} and q_{i-2} are then also singleton points. We will show that a small disk incident to both q_{j+2} and q_{j+3} cannot be in Q. Obviously, the diameter of any such disk is at least δ . The same argument will show that a small disk incident to both q_{j+1} and q_{j+2} cannot be in Q.

Assume otherwise, for contradiction. Observe that the horizontal distance between $\xi(Q_2, \ell)$ and the midpoint of $[q_{j+2}, q_{j+3}]$ is at most 3δ . Similarly, the horizontal distance between $\xi(Q_1, \ell)$ and the same midpoint is at most 5δ . We now verify that the "sandwich" Q_1, Q_2 forces the vertical diameter of a disk incident to both q_{j+2} and q_{j+3} to be smaller than δ . Indeed, this vertical diameter is at most $y_1 + y_2$, where y_i , i = 1, 2, is the vertical distance between the midpoint of $[q_{j+2}, q_{j+3}]$ and the corresponding disk. We have

$$y_1 \le r_1 - \sqrt{r_1^2 - 25\delta^2} = \frac{25\delta^2}{r_1 + \sqrt{r_1^2 - 25\delta^2}} \le \frac{25\delta^2}{1 + \sqrt{1 - 25\delta^2}} = g(5, \delta),$$

and

$$y_2 \le r_2 - \sqrt{r_2^2 - 9\delta^2} = \frac{9\delta^2}{r_2 + \sqrt{r_2^2 - 9\delta^2}} \le \frac{9\delta^2}{1 + \sqrt{1 - 9\delta^2}} = g(3, \delta).$$

By Lemma 5,

$$y_1 + y_2 = g(5,\delta) + g(3,\delta) \le g(6,\delta) + g(3,\delta) \le 0.9\delta_2$$

which contradicts the minimum- δ requirement on the diameter. It follows that q_{j+1} and q_{j+2} are singleton points.

Case 1. f = 1. As in Case 0, it suffices to show that a small disk incident to both q_{j+2} and q_{j+3} cannot be in \mathcal{Q} . Assume otherwise, for contradiction. Observe that the horizontal distance between $\xi(Q_2, \ell)$ and the midpoint of $[q_{j+2}, q_{j+3}]$ is at most 3δ . Similarly, the horizontal distance between $\xi(Q_1, \ell)$ and the same midpoint is at most 6δ . As in Case 0, we check that $y_1 + y_2 \leq g(6, \delta) + g(3, \delta) \leq 0.9\delta$, which holds by Lemma 5.

Case 2. f = 2. By Corollary 1, q_{i-1} and q_{j+1} are singleton points. It suffices to show that the two points of P on ℓ in between Q_1 and Q_2 are singleton points. The relevant inequality is $g(2, \delta) + g(2, \delta) \leq 0.9\delta$, which holds by Lemma 5.

Case 3. f = 3. By Corollary 1, q_{i-1} and q_{j+1} are singleton points. It suffices to show that the three points of P on ℓ in between Q_1 and Q_2 are singleton points. The relevant inequality is $g(2, \delta) + g(3, \delta) \leq 0.9\delta$, which holds by Lemma 5.

Case 4. f = 4. It suffices to show that the four points of P on ℓ in between Q_1 and Q_2 are singleton points. The relevant inequalities are $g(2, \delta) + g(4, \delta) \leq 0.9\delta$, and $g(3, \delta) + g(3, \delta) \leq 0.9\delta$, which hold by Lemma 5.

Case 5. f = 5. It suffices to show that the five points of P on ℓ in between Q_1 and Q_2 are singleton points. The relevant inequalities are $g(2,\delta) + g(5,\delta) \leq 0.9\delta$, and $g(3,\delta) + g(4,\delta) \leq 0.9\delta$, which hold by Lemma 5.

The end of our case analysis concludes the proof of the lemma.

The connected components of G are vertex-disjoint paths; see Figure 8 for an example.




Figure 8: Construction and a 3-vertex path in G.

Lemma 7. Let C be a connected component of G with i vertices. Then there exists a subset $Q_1 \subset Q$ of disks such that $C \subset Q_1$ and Q_1 is supported by exactly m points of P, where $|Q_1| \ge (m+i)/2$.

Proof. Recall that C is a path of i vertices in G, for some $i \ge 1$. If $i \ge 2$, each of the i-1 edges of this path is uniquely associated with a dense line in \mathcal{L} . By Lemma 6, for each edge in C there exist 4 singleton points uniquely associated with this edge. Let \mathcal{Q}_1 consist of the *i* large disks (in C) and the 4(i-1) singleton disks associated with the i-1 edges of C. Thus $|\mathcal{Q}_1| = i + 4(i-1) = 5i - 4$. Since each of the *i* large disks in C is incident to at most 3 points in P, we have $m \le 3i + 4(i-1) = 7i - 4$. It is now easy to verify that $|\mathcal{Q}_1| = 5i - 4 \ge 4i - 2 = ((7i - 4) + i)/2 \ge (m + i)/2$, as required. Note that equality is possible when i = 2.

We can now finalize the proof of Theorem 2 as follows. We have set $\delta = 1/29$, thus $4/\delta = 116$. Consider a connected component of G, say involving i large disks incident to at most i sparse points. (Recall that no large disk can be incident to two sparse points.) Let $Q_2 \subset Q$ be the subset of disks constructed from Q' by taking the union of all disks in Q_1 in the proof of Lemma 7. Clearly $Q' \subset \cup Q_1 = Q_2 \subset Q$. Put $d = |Q_2|$. Let $T \subset P$ be the set of points incident to disks in Q_2 ; write t = |T|. Observe that each connected component yields one inequality of the form $|Q_1| \ge (m+i)/2$, so each interior point generates a "surplus" of 1/2. The corresponding sets Q_1 are pairwise disjoint by construction, thus by adding up all these inequalities yields

$$|\mathcal{Q}_2| \ge \frac{t+(j-1)k}{2}.$$

Apart from O(k) exceptions, each disk in $\mathcal{Q} \setminus \mathcal{Q}_2$ is incident to at most two points in P. Consequently,

$$|\mathcal{Q}| = |\mathcal{Q}_2| + |\mathcal{Q} \setminus \mathcal{Q}_2| \ge \frac{t + (j-1)k}{2} + \frac{n-t}{2} - O(k) = \frac{n + (j-1)k}{2} - O(k).$$
(28)

Analysis. We let $j = \Theta(\sqrt{n})$, and $k = \Theta(\sqrt{n})$, satisfying (27). Analogous to (16) and (17), by substituting the value of (j-1)k resulting from (27), it follows that the number of disjoint empty disks in a minimizing bubble set is at least

$$\frac{n}{2} + \frac{(j-1)k}{2} - O(k) = \frac{n}{2} + \frac{n}{8/\delta + 4} - O(j+k) = \frac{n}{2} + \frac{n}{236} - O(\sqrt{n})$$

This construction can be also extended for every n, by adding a small cluster of at most $O(\sqrt{n})$ collinear points on one of the lines. The above lower bound is preserved and this ends the proof of Theorem 2.

4 Concluding remarks

Via suitable slight perturbations, our lower bound constructions can be realized with points in general position, for instance with no three on a line and no four on a circle. However, the analysis needs to be adapted, and we leave the details to the reader.

In our linear construction points are distributed uniformly on the dense lines. One can reduce the number of points on these lines by making a non-uniform distribution, with points farther away from each other when x is close to odd multiples of $2 + \delta/2$. The reason is that the radius of a disk in Q incident to an interior point $p \in \ell_2$ and intersecting ℓ_1 or ℓ_3 farther away horizontally from p must be significantly larger than 1, a value which was used for simplicity in our calculation. We however opted to leave out the non-uniform distribution due to its complicated analysis.

While we believe it is possible to further raise the lower bound $D(n) \ge n/2 + n/236 - O(\sqrt{n})$, e.g., replace 236 by 190 by making this and other changes, making progress beyond this bound probably requires new ideas. From the other direction, it is very likely that the upper bound 7n/8 + O(1) can be substantially reduced.

Acknowledgement. We thank an anonymous source for bringing the articles of Dillencourt [6] and Ábrego et al. [1, 2] to our attention and for indicating that Proposition 1 follows from the result of Dillencourt [6].

References

- B. M. Abrego, E. M. Arkin, S. Fernández-Merchant, F. Hurtado, M. Kano, J. S. B. Mitchell, and J. Urrutia, Matching points with circles and squares, *Proceedings of the Japan Conference* on Discrete and Computational Geometry (JCDCG 2004), vol. 3742 of LNCS, 2005, pp. 1–15.
- [2] B. M. Abrego, E. M. Arkin, S. Fernández-Merchant, F. Hurtado, M. Kano, J. S. B. Mitchell, and J. Urrutia, Matching points with squares, *Discrete & Computational Geometry*, **41** (2009), 77–95.
- [3] M. Ajtai, The solution of a problem of T. Rado, Bulletin de l'Académie Polonaise des Sciences, Série des Sciences Mathématiques, Astronomiques et Physiques, 21 (1973), 61–63.

- [4] S. Bereg, A. Dumitrescu, and M. Jiang, Maximum area independent set in disk intersection graphs, *International Journal of Computational Geometry & Applications*, **20** (2010), 105–118.
- [5] S. Bereg, A. Dumitrescu, and M. Jiang, On covering problems of Rado, Algorithmica, 57 (2010), 538–561.
- [6] M. B. Dillencourt, Toughness and Delaunay triangulations, Discrete & Computational Geometry, 5 (1990), 575–601.
- [7] K. Hosono and M. Urabe, Empty disks supported by a point set, Proceedings of the XIV Spanish Meeting on Computational Geometry (EGC 2011), June 2011, pp. 99–100.
- [8] J. Pach and G. Tóth, On the independence number of coin graphs, *Geombinatorics*, 6 (1996), 30–33.