BIRMAN-MURAKAMI-WENZL TYPE ALGEBRAS FOR GENERAL COXETER GROUPS

Zhi Chen

Abstract

We introduce a BMW type algebra for every Coxeter group, These new algebras are introduced as deformations of the Brauer type algebras introduced by the author, they have the corresponding Hecke algebras as quotients.

1 introduction

The Birman-Murakami-Wenzl algebras $B_n(\tau, l)$ (BMW algebras)introduced by Birman,Wenzl in [BW] are natural deformations of Brauer's centralizer algebras $B_n(\tau)$. These two kinds of algebras were studied extensively during last two decades. For an explanation of their backgrounds and their relation to three dimensional Topology, consult[BW] [Mu].

In 2001, Häring-Oldenberg [Ha] introduced the Cyclotomic BMW algebras and Cyclotomic Brauer algebras associated with G(m, 1, n) type pseudo reflection groups. The Cyclotomic BMW algebras were studied intensively in recent years. In [RX] Rui and Xu classified irreducible representations of these algebras. In [2] Goodman proved they support cellular structures. In [Yu] Yu studied freeness of them for certain parameter ring, and also proved cellularity of these algebras. In [GH] Goodman and Hauschild introduced Affine BMW algebras from topological considerations and give them a presentation. In 2005, Cohen, Gijsbers and Wales [CGW1] introduced a BMW type algebra and a Brauer type algebra for every simply laced Coxeter group. Then they calculated dimension, and established the semisimplicity and cellularity of these algebras (of finite type) in [1], [CW2]. In this paper we present a BMW type algebra for every Coxeter group, as the deformations of the generalized Brauer type algebras are isomorphic to the algebras from [CGW1](Proposition 5.2).

Since the present paper is very closely related to the simply laced BMW algebras, we write down the presentation of them according to [CGW1]. Let Γ be a simply laced Dynkin diagram. Here 'simply laced' means containing no multiple bonds. For two nodes i, j of Γ , if i and j are connected by a bond, we write $i \sim j$, otherwise we write $i \sim j$.

Definition 1.1 ([CGW1]). Let Γ be a rank n simply laced Dynkin diagram. The type Γ Brauer algebra $B_{\Gamma}(\tau)$ over $\mathbb{Q}(\tau)$ and the type Γ BMW algebra B_{Γ} over $\mathbb{Q}(l, x)$ are

	$B_{\Gamma}(\tau)$	$B_{\Gamma}(\tau, l)$
Generators	$s_i(i \in I); e_i(i \in I)$	$X_{\mathfrak{i}}(\mathfrak{i}\in I)$; $E_{\mathfrak{i}}(\mathfrak{i}\in I)$
Relations	$s_i s_j s_i = s_j s_i s_j$, if $i \sim j$;	$X_iX_jX_i = X_jX_iX_j$, if $i \sim j$;
	$s_i s_j e_i = e_j s_i s_j$ if $i \sim j$;	$X_{\mathfrak{i}}X_{\mathfrak{j}}E_{\mathfrak{i}}=E_{\mathfrak{j}}X_{\mathfrak{i}}X_{\mathfrak{j}} \text{ if } \mathfrak{i}\sim \mathfrak{j};$
	$s_i s_j = s_j s_i$,if $i \not\sim j;$	$X_i X_j = X_j X_i$, if $i \nsim j$;
	$s_i^2 = 1$, for all i;	$l(X_i^2 + \nu X_i - 1) = \nu E_i, \text{ for all } i;$
	$s_i e_i = e_i$, for all i;	$X_i E_i = l^{-1} E_i$, for all i;
	$e_i s_j e_i = e_i$, if $i \sim j$;	$E_i X_j E_i = l E_i$, ifi ~ j;
	$s_i e_j = e_j s_i$, if $i \not\sim j$;	$X_i E_j = E_j X_i$, ifi $\nsim j$;
	$e_i^2 = \tau e_i$, for all i.	$E_i^2 = \tau E_i$.

TABLE 1. Presentation for $B_{\Gamma}(\tau)$ and $B_{\Gamma}(\tau, l)$.

Where $v = \frac{l-l^{-1}}{l-\tau}$. When the Coxeter group is of A_{n-1} type,that is, being the n-th symmetric group, the corresponding simply laced BMW algebra and Brauer algebra are just $B_n(\tau, l)$ and $B_n(\tau)$ respectively. In [CH] the author introduced a Brauer type algebra(see Definition 2.1, Definition 2.2 of the present paper) for every Coxeter group and every pseudo reflection group as candidates for generalized Brauer type algebras. In that construction, certain KZ connections and generalized Lawrence-Krammer representations for finite type Coxeter groups introduced by Marin (including a slightly further generalization by the author) play important roles. When W is a finite pseudo reflection group, the algebra $Br_W(\Pi)$ supports a flat, W-invariant connection $\Omega_W(Definition 2.3)$, which can be used to deform every finite dimensional representation of $Br_W(\Pi)$ to a one parameter class of representations of the associated braid group B_W . In this paper we try to introduce BMW algebras for every Coxeter group, as deformations of the Brauer type algebras of [CH]. The starting point of our construction is the following observation.

Let Γ be A_2 , the Dynkin diagram with two vertex ν_0, ν_1 and a simple lace connecting them. So the BMW algebra $B_{A_2}(\tau, l) = B_3(\tau, l)$ is generated by $\{X_0, X_1, E_0, E_1\}$ with relations as in above table. Denote the free monoid generated by $\{X_0, X_1, E_0, E_1\}$ as Λ .

Then for i = 0, 1, there is a function Φ^i on Λ such that the following relation holds in $B_{A_2}(\tau, l)$:

$$E_i x E_i = \Phi^i(x) E_i \tag{1}$$

These relations are clear from the diagram presentation of the BMW algebras. It is an interesting fact that we can obtain above Φ^i from the LK representation of the braid group B₃. Let σ_0, σ_1 be the generators of B₃ in its canonical presentation. The LK representation is a 3 dimensional representation with 2 parameters $\{\tau, l\} \rho_{LK} : B_3 \rightarrow GL(V)$. We can show (Lemma 3.1) that for i = 0, 1, the operator $e_i = \frac{1}{\nu}(\rho(\sigma_i)^2 + \nu\rho(\sigma_i) - 1)$ is a projector to some line $\mathbb{C}u_i$ in V, where $\nu = \frac{l-l^{-1}}{l-\tau}$. The LK representation can be looked as a representation of $B_{A_2}(\tau, l)$ by sending X_i, E_i to $\rho_i(\sigma_i), e_i$ respectively for i = 0, 1. We denote this representation by the same symbol ρ_{LK} . For any word $W = A_1A_2 \cdots A_k$ in Λ , where

 $A_i \in \{X_0, X_1, E_0, E_1\}$ we set $\rho_{LK}(W) = \rho_{LK}(A_1) \cdots \rho_{LK}(A_k) \in gl(V)$. Then there are functions $\bar{\Phi}^i$ (i = 0, 1) on Λ such that $e_i \rho_{LK}(W) e_i = \bar{\Phi}^i(W) e_i$, since e_i is a projector. Comparing above two identities, we have $\Phi^i = \bar{\Phi}^i$.

Denote the Dihedral group of type $I_2(m)$ as D_m , whose Dynkin diagram consists of two vertexes v_0, v_1 and an edge with weight m connecting them. Denote the associated Artin group as A_{D_m} . The group A_{D_m} has a canonical presentation with two generators σ_0, σ_1 .

The group D_m has a natural representation on \mathbb{C}^2 as a reflection group of order 2m. It has m reflection hyperplanes $H_0, H_1, \cdots, H_{m-1}$. Denote the reflection by H_i as s_i , and chose a linear function f_i on \mathbb{C}^2 such that $H_i = \ker f_i$. Set $\omega_i = df_i/f_i$, which is a holomorphic 1-form on M. Denote the complementary space $\mathbb{C}^2 \setminus \bigcup_{i=0}^{m-1} H_i$ as M_m . The Artin group A_{D_m} also has a generalize Lawrence-Krammer representation as special cases of the one constructed by Marin [Ma2] as follows. Set $V_m = \mathbb{C} < v_0, v_1, \cdots, v_m >$ be a vector space with a basis in one to one correspondence with the set of reflection hyperplanes of D_m . The action of D_m on the second set naturally induces a representation of D_m on V_m which will be denoted as ι . Then on the trivial bundle $M_m \times V_m$, we have Marin's connection: $\Omega_{LK} = \sum_{i=0}^{m-1} \kappa(k\iota(s_i) - p_i)\omega_i$, where $p_i \in End(V_m)$ is suitable projector to the line $\mathbb{C}v_i$. It is proved [Ma2] that Ω_{LK} is flat and D_m - invariant. So it induces a flat connection on the flat bundle $M \times_{D_m} V$, whose monodromy gives the generalized LK representation of A_{D_m} , which will be denoted as $T_{LK} : A_{D_m} \to Aut(V_m)$.

First we suppose the cases when m is odd, we have

Lemma 3.1 Set
$$q = \exp(\kappa k \pi \sqrt{-1})$$
 and $l = \exp(\kappa \alpha \pi \sqrt{-1})/q$. For generic κ ,

- (1) $(T_{LK}(\sigma_i) l^{-1})(T_{LK}(\sigma_i) q)(T_{LK}(\sigma_i) + q^{-1}) = 0$ for i = 0, 1.
- (2) $e_i = \frac{1}{q^{-1}-q}(T_{LK}(\sigma_i) q)(T_{LK}(\sigma_i) + q^{-1})$ is a projector, i.e, $Ranke_i = 1$ for i = 0, 1.

Still let Λ be the monoid freely generated by X_0, X_1, E_0, E_1 . Define a morphism $f : \Lambda \to \text{End}(V_m)$ by setting: $f(X_i) = T_{\text{LK}}(\sigma_i)$, $f(E_i) = e_i$ (i = 0, 1). Considering (2) of above Lemma 3.1, we define a function $\Phi^i_{m,\Pi,\kappa} : \Lambda \to \mathbb{C}$ by: $e_i f(X) e_i = \Phi^i_{m,\Pi}(X) e_i$. Then we define the BMW algebra $B_{D_m}(\Pi, \kappa)$ associated with D_m as the following Definition 3.4.

Suppose $k \in \mathbb{C} \setminus \{0\}$, $\alpha \in \mathbb{C}$, denote $\Pi = \{k, \alpha\}$. Set $q = \exp(\kappa \sqrt{-1}\pi k)$ and $l = \exp(\kappa \alpha \pi \sqrt{-1})/q$.

Definition 3.4 The algebra $B_{D_m}(\Pi, \kappa)$ is generated by X_0, X_1, E_0, E_1 with following relations.

1) $X_i X_i^- = 1$, i = 0, 1; 2) $[X_0 X_1 \cdots]_m = [X_1 X_0 \cdots]_m$; 3) $(q^{-1} - q) E_i = l(X_i - q)(X_i + q^{-1})$ for i = 0, 1; 4) $X_i E_i = E_i X_i = l^{-1} E_i$ for i = 0, 1; 5) $E_i X E_i = \Phi_{m,\Pi,\kappa}^i(X) E_i$ for any $X \in \Lambda, i = 0, 1$.

The relation 1) to 4) have clear analogues with the BMW algebra $B_3(\tau, l)$. Relation 5)

is similar with the equation (1) for $B_3(\tau, l)$. From above definition we see $B_{D_m}(\Pi, \kappa)$ is a quotient of the group algebra $\mathbb{C}A_{D_m}$, and has the Hecke algebra of type W as its quotient.

When m is even, we have a pair of projectors e_0, e_1 acting on Marin's generalized LK representation as in Lemma 3.2. By using them we define a function $\Phi^{i}_{m,\Pi,\kappa} : \Lambda \to \mathbb{C}$, which is used in Definition 3.3 of the BMW algebra $B_{D_m}(\Pi, \kappa)$.

In section 4 we study the algebras $B_{D_m}(\Pi, \kappa)$, the following results support that they are indeed the deformation of the Brauer type algebras $Br_{D_m}(\Pi)$. Our Theorem 4.1 says that for generic parameters Π , dim $Br_{D_m}(\Pi) = \dim B_{D_m}(\Pi, \kappa)$. Proposition says the representations of A_{D_m} obtained from monodromy of the KZ connection Ω_{D_m} (associated with finite dimensional representations of $Br_{D_m}(\Pi)$ factor through $B_{D_m}(\Pi, \kappa)$.

Once the definition of BMW algebras associated with Dihedral groups are given, they induce a natural definition of BMW algebra $B_{W_{\Gamma}}(\Pi, \kappa)$ for any Coxeter group W_{Γ} , as in Definition 4.1.

We hope these algebras $B_{W_{\Gamma}}(\Pi, \kappa)$ be deformation of the Brauer type algebras $Br_{W_{\Gamma}}(\Pi)$ as the dihedral cases. We can't give a proof of them in this paper. We only prove (Theorem 5.1) that $B_{W_{\Gamma}}(\Pi, \kappa)$ is finite dimensional when W is a finite group, by similar arguments used in [CGW1] in the proof of Theorem 1.1 of that paper. Proposition 5.2 says the algebra $B_{W_{\Gamma}}(\Pi, \kappa)$ is isomorphic to the simply laced BMW algebra defined in [CGW1] when W is a simply laced Coxeter group. At last, Theorem 5.2 says for finite W, for generic parameters Π , the representations of A_W resulted from monodromy of the KZ connection Ω_W all factor through $B_W(\Pi, \kappa)$ for suitable parameters.

Acknowledgements I 'd like to present my thanks to Ivan Marin for his detailed explanation of some tools involved in proof of Lemma 3.5.

2 preliminaries

Let E be a complex linear space. An hyperplane arrangement (or arrangement simply) in E means a finite set of hyperplanes contained in E. Let $\mathcal{A} = \{H_i\}_{i \in I}$ be an arrangement in E, we denote the complementary space $E - \bigcup_{i \in I} H_i$ as $M_{\mathcal{A}}$. Intersection of any subset of \mathcal{A} is called an edge. If L is an edge of \mathcal{A} , define $\mathcal{A}_L = \{H_i \in \mathcal{A} | L \subset H_i\} = \{H_i\}$ and $I_L = \{i \in I | L \subset H_i\}$.

For every $i \in I$, chose a linear form f_i with kernel H_i . Set $\omega_i = d \log f_i$, which is a holomorphic closed 1-form on M_A . Consider the formal connection $\Omega = \kappa \sum_{i \in I} X_i \omega_i$. Here X_i are linear operators to be determined. When we take X_i 's as endomorphisms of some linear space E, then Ω is realized as a connection on the bundle $M_A \times E$. We have the following theorem of Kohno.

Theorem 2.1 (Kohno [Ko1]). The formal connection Ω is flat if and only if:

 $[X_i, \sum_{j \in I_L} X_j] = 0 \text{ for any codimension 2 edge } L \text{ of } \mathcal{A}, \text{and for any } i \in I_L. Where } [A, B] \text{ means } AB - BA.$

Let $W \subset U(E)$ be a finite pseudo reflection group. Let R be the set of pseudo reflections contained in W and let $\mathcal{A} = \{H_i\}_{i \in P}$ be the set of reflection hyperplanes in E. Since the

action of W on E send reflection hyperplanes to reflection hyperplanes, we have a natural action of W on P which will denoted simply by $(w, i) \mapsto w(i)$ for $w \in W, i \in P$. For s, s' $\in R$, denote $s \sim s'$ if they lie in the same conjugacy class. For $i, i' \in P$, denote $i \sim i'$ if there exist $w \in W$ such that w(i) = i'. For $i, j \in P$, denote $R(i, j) = \{s \in R | s(H_i) = H_i\}$. For $i \in P$, let W_i be the subgroup of W consisting of elements that fixing H_i pointwise. Let $m_i = |W_i|$. It is well known that W_i is a cyclic group and we denote the element in W_i with exceptional eigenvalue $e^{\frac{2\pi\sqrt{-1}}{m_i}}$ as s_i . For $s \in R$, define $i(s) \in P$ by requesting $H_{i(s)}$ to be the reflection hyperplane of s. Set $M_W = E - \bigcup_{i \in P} H_i$. The action of W on M_W is free, by a famous theorem of Steinberg. The groups $B_W = \pi_1(M_W/W)$ and $P_W = \pi_1(M_W)$ are called the complex braid group and the complex pure braid group associated with W respectively. For $i \in P$, suppose ω_i is the 1-form on M_W associated with H_i as above. Suppose (V, ρ) is a complex linear representation of W. Suppose $\Omega = \sum_{i \in P} X_i \omega_i$ is a connection on the trivial vector bundle $M_W \times V$, where $X_i \in End(V)$ for any $i \in P$. In this case we can get a vector bundle on M_W/W as the orbit space of the free action of W on $M_W\times V$: $w \cdot (p, v) = (w \cdot p, \rho(w)(v))$, for any $w \in W$, $p \in M_W$ and $v \in V$. Denote the quotient bundle as $M_W \times_W V$, and the quotient map from $M_W \times V$ to $M_W \times_W V$ as π . We have the following Lemma, for details and backgrounds about these connections see Section 4 of [?].

Lemma 2.1. If $\rho(w)X_i\rho(w)^{-1} = X_{w(i)}$ for any $i \in P$ and $w \in W$, then Ω induce a connection $\overline{\Omega}$ on $M_W \times_W V$, whose pull back through by π is Ω .

If a connection satisfy the condition in above Lemma, we call it as W-invariant.

Suppose $E^{\mathbb{R}}$ is a n dimensional Eclidean space, denote the bilinear form on $E^{\mathbb{R}}$ as <,>. $W \subset O(E^{\mathbb{R}})$ is a finite reflection group with the set of reflections denoted as R. Suppose W is essential, which means the fixed subspace of W in $E^{\mathbb{R}}$ is 0. Denote the complexification $E^{\mathbb{R}} \oplus \sqrt{-1}E^{\mathbb{R}}$ of $E^{\mathbb{R}}$ as E. There is a natural positive definite Hermitian form on E extending the metric on $E^{\mathbb{R}}$ defined as:

$$<\nu+\sqrt{-1}w,\nu^{'}+\sqrt{-1}w^{'}>=<\nu,\nu^{'}>+\sqrt{-1}< w,\nu^{'}>-\sqrt{-1}<\nu,w^{'}>+< w,w^{'}>.$$

Denote the distance of two points $p, q \in E$ determined by this Hermitian metric as d(p, q). Because R contain only order 2 elements, the map i from R to P (index set of reflection hyperplanes) is bijective. So we simply set P = R and denote the reflection hyperplane in $E^{\mathbb{R}}$ of s as H_s . Denote the complexification of H_s as $H_s^{\mathbb{C}}$, which is a hyperplane in E. It is a well-known fact (for example see [Hu]) that each path component of $E^{\mathbb{R}} - \bigcup_{s \in R} H_s$ is a simplicial cone. Choose one from these cones, denote the closure of it as Δ . Denote the n reflection hyperplanes (of W) corresponding to n codimension one faces of Δ as H_{s_1}, \dots, H_{s_n} . The following Theorem is canonical (see [Hu]).

Theorem 2.2. The group W is generated by s_1, \dots, s_n with the following relations. 1) $s_i^2 = 1$, $1 \le i \le n$; 2) $[s_i s_j \cdots]_{m_{i,j}} = [s_j s_i \cdots]_{m_{i,j}}$, $1 \le i \ne j \le n$.

Here the expression $[xy \cdots]_m$ means a word with length m in which the letters x, y appear alternatively, and whose left most subword is xy. Similarly we define $[\cdots xy]_m$.

The symmetric matrix $\Gamma = (m_{i,j})_{n \times n}$ is called the Coxeter matrix of W, which is equivalent to a Dynkin diagram under certain simple rules.

Now consider the space M_W/W . Denote the quotient map from M_W to M_W/W as π . Choose a inner point $p \in \Delta$. Notice since $E^{\mathbb{R}}$ is a subspace of its complexification E, p belongs to M_W . Denote $\pi(p)$ as \bar{p} . Denote the segment connecting p with $s_i(p)$ as γ_i , which intersecting H_{s_i} in one point p_i . Chose $0 < \epsilon << 1$. Let p'_i be the point on $\gamma_i \cap \Delta$ satisfying: $d(p'_i, p_i) = \epsilon$. Set $p''_i = s_i(p'_i)$ (also belong to Γ_i). Denote the vector $s_i(p) - p$ as α_i , then the reflection s_i can be presented as $s_i(\nu) = \nu - \frac{2 < \nu, \alpha_i >}{<\alpha_i, \alpha_i >} \alpha_i$. denote the path from p to p'_i as γ'_i , and the path from p''_i to $s_i(p)$ as γ''_i . Define the following path η_i from p''_i to p''_i as: $\eta_i : [0, 1] \to E$, $t \mapsto p'_i - (1 - \exp \pi \sqrt{-1}t) \frac{< p'_i, \alpha_i >}{<\alpha_i, \alpha_i >} \alpha_i$. Denote the composed path $\gamma''_i \circ \eta_i \circ \gamma'_i$ as l_i . It is easy to see when ϵ is small enough, the path l_i lie in M_W , and it project to a closed path \bar{l}_i in M_W/W based at \bar{p} . Denote the element of $\pi_1(M_W/W, \bar{p})$ represented by \bar{l}_i as σ_i .

Theorem 2.3 (Brieskorn [Bri]). The group $\pi_1(M_W/W, \bar{p})$ is generated by $\sigma_1, \dots, \sigma_n$ with the following relations.

 $[\sigma_i\sigma_j\cdots]_{\mathfrak{m}_{i,j}}=[\sigma_j\sigma_i\cdots]_{\mathfrak{m}_{i,j}} \text{ for } 1\leq i\neq j\leq n.$

The group $\pi_1(M_W/W, \bar{p})$ is the Artin group of type W, which will be denoted as A_W . The following results about monodromy of flat connections will be used later. Suppose W is a finite Coxeter group, and $M_W, \Delta, s_i, H_i, p, \bar{p}$ defined as above. Suppose $\rho : W \to GL(V)$ (or Aut(V)) is a finite dimensional complex linear representation of W, and $\Omega = \kappa \sum_{s \in \mathbb{R}} X_s \omega_s$ is a W-invariant flat connection on a trivial vector bundle $M_W \times V$, where $X_s \in End(V)$ for any $s \in \mathbb{R}$. Denote the flat connection on $M_W \times_W V$ induced by Ω as $\bar{\Omega}$. Denote the monodromy representation of $\bar{\Omega}$ as $T : \pi_1(M_W/W, \bar{p}) \to GL(V)$. Then

Lemma 2.2 (Proposition 2.3 of [Ma2]). For generic κ , the operator $T(\sigma_i)$ is conjugated to $\rho(s_i) \exp(\kappa \sqrt{-1}\pi X_{s_i})$ in GL(V).

Now suppose L is an edge of the arrangement $\{H_s\}_{s\in R}$, which means $L \subset E$ is the nonempty intersection of several members of $\{H_s\}_{s\in R}$. Set $R_L = \{s \in R | L \subset H_s\}$, and $M_{W_L} = E - \bigcup_{s\in R_L} H_s$. It is a canonical result (see [Hu]) that the subgroup W_L generated by R_L is the maximal subgroup of W fixing L pointwise. As a reflection group it has its own associated Artin group A_{W_L} and pure Artin group P_{W_L} . It is clear that we have identifications: $P_{W_L} \cong \pi_1(M_{W_L})$ and $A_{W_L} \cong \pi_1(M_{W_L}/W_L)$. Now choose a point $q \in L$ such that $q \notin H_s$ for any $s \notin R_L$. We can choose $\epsilon > 0$ small enough such that the ball $D_{\epsilon}(q) = \{a \in E | d(a, q) < \epsilon\}$ has empty intersection with any $H_s(s \notin R_L)$. It is easy to see the natural morphism $\pi_1(D_{\epsilon}(q) \cap M_W) \to \pi_1(M_{W_L})$ induced by the inclusion map is an isomorphism, thus the morphism $\pi_1(D_{\epsilon}(q) \cap M_W) \to \pi_1(M_W)$ induced by inclusion map gives us an injective morphism: $\lambda_L : P_{W_L} \to P_W$. Since $D_{\epsilon}(q)$ is setwise stabilized by W_L , λ_L can be extend to an injective morphism $\overline{\lambda}_L : A_{W_L} \to A_W$.

Assume (V, ρ) , Ω as above. On the trivial bundle $M_{W_L} \times V$ we define the following connection: $\Omega' = \kappa \sum_{s \in R_L} X_s \omega_s$. it is easy to see Ω' is also a flat and W_L -invariant connection, so it induce a flat connection $\overline{\Omega}'$ on the quotient bundle $M_{W_L} \times_{W_L} V$. The connection

 $\overline{\Omega}'$ induce a monodromy representation $T_L : A_{W_L} \to GL(V)$. On the other hand, by the injection $\overline{\lambda}_L$ we have another representation $T \circ \overline{\lambda}_L$ of $A_{W_L} : A_{W_L} \to A_W \to GL(V)$. In [Ma2] the following theorem is presented.

Theorem 2.4. Assume above conventions. For generic κ , the representation (V, T_L) is isomorphic to $(V, T \circ \overline{\lambda}_L)$.

Let $W \subset U(E)$ be a finite pseudo reflection group. In [CH] the author introduced a Brauer type algebra $Br_W(\Pi)$ associated with W, which is defined as follows. Set notations $R, \mathcal{A}, P, i(s), s \sim s', i \sim i', R(i, j)$ as above. Choose constants $\{k_s\}_{s \in R} \cup \{\alpha_i\}_{i \in P}$ such that $k_s = k_{s'}$ if $s \sim s'$ and $\alpha_i = \alpha_{i'}$ if $i \sim i'$. Denote $\{k_s\}_{s \in R} \cup \{\alpha_i\}_{i \in P}$ by one symbol Π . For $i \neq j$ we denote $H_i \pitchfork H_j$ if $\{k \in P \mid H_i \cap H_j \subset H_k\} = \{i, j\}$. A codimension 2 edge L of \mathcal{A} will be called a crossing edge if there exists $i, j \in P$ such that $H_i \pitchfork H_j$ and $L = H_i \cap H_j$, otherwise L will be called a noncrossing edge.

Definition 2.1. The algebra $Br_W(\Pi)$ associated with pseudo reflection group $W \subset U(E)$ is generated by the set $\{T_w\}_{w \in G} \cup \{e_i\}_{i \in P}$ which satisfies the following relations.

0) $T_{w_1}T_{w_2} = T_{w_3}$ if $w_1w_2 = w_3$.

 $1) \ T_{s_{\mathfrak{i}}}e_{\mathfrak{i}}=e_{\mathfrak{i}}T_{s_{\mathfrak{i}}}=e_{\mathfrak{i}}, \ \textit{for} \ \mathfrak{i}\in P.$

1)' $T_w e_i = e_i T_w = e_i$, for $w \in G$ such that $w(H_i) = H_i$, and $H_i \cap E_w$ is a noncrossing edge. Where $E_w = \{v \in E | w(v) = v\}$.

2) $e_i^2 = \alpha_i e_i$.

3) $T_w e_j = e_i T_w$, if $w \in G$ satisfies $w(H_j) = H_i$.

4) $e_i e_j = e_j e_i$, if $H_i \pitchfork H_j$.

5) $e_ie_j = (\sum_{s \in R(i,j)} k_s T_s)e_j = e_i(\sum_{s \in R(i,j)} k_s T_s)$, if $H_i \cap H_j$ is a noncrossing edge, and $R(i,j) \neq \emptyset.$

6) $e_i e_j = 0$, if $H_i \cap H_j$ is a noncrossing edge, and $R(i, j) = \emptyset$.

It is proved in [CH](Theorem 8.4) that when W is a finite type Coxeter group with Coxeter matrix $(m_{i,j})_{n \times n}$, the algebra $Br_W(\Pi)$ is isomorphic to the following algebra $Br'_{W_M}(\Pi)$ with a canonical presentation.

Definition 2.2. For any Coxeter matrix $M = (m_{i,j})_{n \times n}$, the algebra $Br'_{W_M}(\Pi)$ is defined as follows. Let Π be as in Definition 2.1. Denote α_{s_i} in Υ as α_i . If we don't give range for an index then it means "for all". The generators are $S_1, \dots, S_n, E_1, \dots, E_n$.

The relations are

$$\begin{split} 1)S_{i}^{2} &= 1; \\ 2)[S_{i}S_{j}\cdots]_{m_{i,j}} &= [S_{j}S_{i}\cdots]_{m_{i,j}}; \\ 3)S_{i}E_{i} &= E_{i} &= E_{i}S_{i}; \\ 4)E_{i}^{2} &= \alpha_{i}E_{i}; \\ 5)S_{i}E_{j} &= E_{j}S_{i}if\ m_{i,j} &= 2; \\ 6)E_{i}E_{j} &= E_{j}E_{i}if\ m_{i,j} &= 2; \\ 7)[S_{j}S_{i}\cdots]_{2k-1}E_{i} \\ &= E_{i}[S_{j}S_{i}\cdots]_{2k-1}E_{i} \\ &= E_{i}[S_{j}S_{i}\cdots]_{2k-1} &= E_{i}, \\ if\ m_{i,j} &= 2k > 2; \\ \end{split}$$

An important feature of the algebra $Br_W(\Pi)$ is it supports the following W-invariant, flat formal connection

Definition 2.3. The KZ connection of $Br_W(\Pi)$ is the following formal connection on $M_W \times Br_W(\Pi)$: $\Omega_W = \kappa \sum_{i \in P} (\sum_{s:i(s)=i} k_s T_s - e_i) \omega_i$.

The following fact is from Proposition 5.1 of [CH].

Proposition 2.1. The connection Ω_W is flat and W-invariant.

This connection can deform every finite dimensional representation of $Br_W(\Pi)$ to a representation of B_W , the associated braid group. We have the following example. Let R, P, M_W, ω_i be defined as above. Set $V_W = \mathbb{C} < \nu_i >_{i \in P}$ be a complex linear space with a basis in one to one correspondence with the set of reflection hyperplanes of W. The permutating action of W on P induces a representation ι of W on V_W in a natural way: $\iota(w)(\nu_i) = \nu_{w(i)}$ for any $w \in W$ and $i \in P$. Let the data $\{k_s\}_{s \in R} \cup \{\alpha_i\}_{i \in P}$ be Π in $Br_W(\Pi)$. For $i \in P$, define an element $p_i \in End(V_W)$ as by $p_i(\nu_i) = \alpha_i \nu_i$; $p_i(\nu_j) = (\sum_{t \in R:t(j)=i} k_t)\nu_i$. So p_i is a projector to the line $\mathbb{C}\nu_i$.

Lemma 2.3. The map $T_w \mapsto \iota(w)$ for $w \in W$ and $e_i \mapsto p_i$ for $i \in P$ extends to a representation $\rho_{LK} : Br_W(\Pi) \to End(V_W)$.

We call this representation as the infinitesimal Lawrence-Krammer representation. So the connection $\Omega_W^{LK} = \kappa \sum_{i \in P} (\sum_{s \in R:i(s)=i} k_s \iota(s) - p_i) \omega_i$ on the vector bundle $M_W \times V_W$ is flat and W-invariant by Proposition 2.1. Ω_W^{LK} induces a flat connection $\overline{\Omega}_W^{LK}$ on the bundle $M_W \times_W V_W$, whose monodromy representation is defined as the generalized Lawrence-Krammer representations (of A_W). The generalized Lawrence-Krammer representations for finite type simply-laced Artin groups are invented by Cohen and Wales in [CW], and Digne in [Di] independently. In [Ma2] Marin introduced generalized Lawrence-Krammer representations for pseudo reflection groups whose pseudo reflections all have order 2. Above slightly further generalization of Marin's work can be found in [CH], which is put in the frame of $Br_W(\Pi)$ for later convenience.

Consider the permutating action of W on P, denote the space of orbits as P/W. For each $c \in P/W$, it is easy to see $V_{W,c} = \mathbb{C} < v_i >_{i \in c}$ is a sub representation of ρ_{LK} . Thus we have a decomposition $\rho_{LK} = \bigoplus_{c \in P/W} \rho_{LK,c}$ according to the decomposition $V_W = \bigoplus_{c \in P/W} V_{W,c}$. In dihedral cases (where we can set P = R), when m is odd, the set of reflections R of D_m contains only one orbit. When m is even, let $c_0 \subset R$, $c_1 \subset R$ be those reflections conjugated to s_0 , s_1 respectively. Then we have $V_{D_m} = V_{D_m,c_0} \oplus V_{D_m,c_1}$, correspondingly $\rho_{LK} = \rho_{LK,c_0} \oplus \rho_{LK,c_1}$.

The following result follows from Proposition 3.4 and Proposition 4.2 of [Ma2].

Lemma 2.4. When m is odd, for generic k, α , the infinitesimal Lawrence-Krammer representation ρ_{LK} is irreducible. When m is even, for generic $k_0, \alpha_0, k_1, \alpha_1$, the representations $\rho_{LK,c_0}, \rho_{LK,c_1}$ are irreducible.

3 Generalized Lawrence-Krammer Representations

Suppose W is a finite Coxeter group. We assume the conventions for E, R, M_W as in Section 2. We study the generalized Lawrence-Krammer representation in detail for the cases when W is a dihedral group, as they will play an important role in the next section.

Denote the dihedral group of type $I_2(m)$ as D_m . The arrangement of its reflection hyperplanes can be explained with the following Figure 1.



Figure 1: The arrangement of D_6

There are m lines(hyperplanes) passing the origin. The angle between every two neighboring lines is π/m . Suppose the x-axis is one of the reflection lines and denote it as H₀, we denote these lines by H₀, H₁, \cdots , H_{m-1} in anticlockwise order as shown in above graph. Denote the reflection by H_i as s_i. The set of reflections in D_m is $R = \{s_i\}_{0 \le i \le m-1}$. Denote the rotation of $2j\pi/m$ in anticlockwise order as r_j . It is well known that D_m is generated by s₀, s₁ with the following presentation

$$< s_0, s_1, |(s_0s_1)^m = 1, s_0^2 = s_1^2 = 1 > .$$

The associated Artin group A_{W_m} has a presentation:

$$< \sigma_0, \ \sigma_1, \ |[\sigma_0 \sigma_1 \cdots]_m = [\sigma_1 \sigma_0 \cdots]_m > .$$

As a set $D_m = \{s_i, r_i\}_{0 \le i \le m-1}$. Under this presentation, s_i can be determined inductively in the following way. $s_1 = s_1$, $s_2 = s_1 s_0 s_1$ and $s_i = s_{i-1} s_{i-2} s_{i-1}$. Recall by $[s_i s_j \cdots]_k$ we denote the length k word starting with $s_i s_j$, in which s_i and s_j appear alternatively. The word $[\cdots s_0 s_1]_k$ is defined similarly. Then $s_i = [s_1 s_0 \cdots]_{2i-1}$, where $s_m = [s_1 s_0 \cdots]_{2m-1} = s_0$.

The cases when m is odd Recall the connection Ω_W^{LK} on $M_W \times V_W$ in Lemma 2.2. We denote D_m simply as W in this paragraph. For these cases all reflections in W lie in one conjugating class. So the constants k_s all equal to some k and constants α_s all equal to some α . We simply denote the base elements v_{s_i} of V_W as v_i , the projector p_{s_i} as p_i , and ω_{s_i} as ω_i for $0 \le i \le m - 1$. The connection can be rewritten as:

$$\Omega_{\rm m} = \sum_{i=0}^{\rm m-1} \kappa(k\iota(s_i) - p_i)\omega_i, \qquad (2)$$

Where the W representation (V_W, ι) is as in Section 2, and p_i is defined by $p_i(\nu_i) = \alpha \nu_i; p_i(\nu_j) = k\nu_i$ if $j \neq i$. Denote the induced flat connection on $M_W \times_W V_W$ as $\bar{\Omega}_m$ and the monodromy representation of $\bar{\Omega}_m$ as $T_{LK}: A_W \to GL(V_W)$.

Lemma 3.1. Set $q = \exp(\kappa k \pi \sqrt{-1})$ and $l = \exp(\kappa \alpha \pi \sqrt{-1})/q$. For generic κ ,

(1)
$$(T_{LK}(\sigma_i) - l^{-1})(T_{LK}(\sigma_i) - q)(T_{LK}(\sigma_i) + q^{-1}) = 0$$
 for $i = 0, 1$.

(2)
$$e_i = \frac{l}{q^{-1}-q}(T_{LK}(\sigma_i) - q)(T_{LK}(\sigma_i) + q^{-1})$$
 is a projector, i.e., $Ranke_i = 1$ for $i = 0, 1$.

Proof. We only need to prove the case i = 0. Another case is similar. First we diagonalize $\iota(s_0)$ under certain basis $\{u_0, \cdots, u_{m-1}\}$ where $u_0 = \nu_0$. Under the same basis $\kappa(\kappa\iota(s_0) - p_0)$ is presented by a matrix whose entries other than the diagonal and the first column are zero. Now by Lemma 2.2, the operator $T_{LK}(\sigma_0)$ is conjugated to $\iota(s_0) \exp(\sqrt{-1}\pi\kappa(\kappa\iota(s_0) - p_0))$. Under the basis $\{u_i\}$, the operator $\iota(s_0) \exp(\sqrt{-1}\pi\kappa(\kappa\iota(s_0) - p_0))$ is still presented by a matrix whose entries other than the diagonal and the first column are zero, and whose (1, 1) entry is l^{-1} , other diagonal entries are q, q^{-1} . When $\kappa(k - \alpha) \notin \{\kappa k + \mathbb{Z}\} \cup \{-\kappa k + \mathbb{Z}\}\}$, so $l^{-1} \notin \{q, q^{-1}\}$, we see $\iota(s_0) \exp(\sqrt{-1}\pi\kappa(\kappa\iota(s_0) - p_0))$ is also diagonalizable with eigenvalues $q, -q^{-1}, l^{-1}$, and the eigenspace of l^{-1} is one dimensional. So complete the proof.

The cases when m = 2k is even In this paragraph denote $D_m(m \text{ being even})$ as W. In these cases the set of reflections in W consists of two conjugacy classes, represented by s_0 , s_1 respectively. So constants for the connection consists of $k_{s_0}, k_{s_1}, \alpha_{s_0}, \alpha_{s_1}$. Denote them simply by $k_0, k_1, \alpha_0, \alpha_1$. Denote the basis element v_{s_i} of V_W as v_i , projectors p_{s_i} as p_i , and ω_{s_i} as ω_i for $0 \le i \le m - 1$. The connection Ω_W^{LK} can be rewritten as

$$\Omega_{\mathfrak{m}} = \kappa (\sum_{i=0}^{k-1} (k_0 \iota(s_{2i}) - \mathfrak{p}_{2i}) \omega_{2i} + \sum_{i=0}^{k-1} (k_1 \iota(s_{2i+1}) - \mathfrak{p}_{2i+1}) \omega_{2i+1}).$$
(3)

Where $p_i \ (0 \le i \le m-1)$ is determined by: $p_i(\nu_j) = 0$ if $2 \nmid i-j$; $p_i(\nu_j) = 2\nu_i$ if $2 \mid i-j$ and $i \ne j$; $p_i(\nu_i) = k_0\nu_i$ if i is odd; $p_i(\nu_i) = k_1\nu_i$ if i is even. Denote the flat connection on $M_W \times_W V_W$ induced from Ω_m as $\overline{\Omega}_m$, and denote the monodromy representation of $\overline{\Omega}_m$ as $T_{LK} : A_W \to GL(V_W)$. We have the following Lemma. The proof is similar to Lemma 3.1.

Lemma 3.2. Set $q_i = \exp(\kappa k_i \pi \sqrt{-1})$ and $l_i = \exp(\kappa \alpha_i \pi \sqrt{-1})/q_i$ for i = 0, 1. For generic k_i and α_i ,

- $(1) \ (T_{LK}(\sigma_i) l_i^{-1})(T_{LK}(\sigma_i) q_i)(T_{LK}(\sigma_i) + q_i^{-1}) = 0 \ \text{for} \ i = 0, 1.$
- (2) $e_i = \frac{l_i}{q_i^{-1} q_i} (T_{LK}(\sigma_i) q_i) (T_{LK}(\sigma_i) + q_i^{-1})$ is a projector, *i.e.* Rank $e_i = 1$ for i = 0, 1.

The following lemma is a easy fact.

are irreducible.

Lemma 3.3. Suppose V is a Hermitian vector space whose distance is denoted as d(,). Suppose $\{v_i\}_{i=1,\cdots,m} \subset V$ span V. Then there exists $\varepsilon > 0$, such that $d(v'_i, v_i) < \varepsilon$ for $i = 1, \cdots, m$ implies $\{v'_i\}_{i=1,\cdots,m}$ span V.

Definition 3.1. Let V be a linear space. A set $\{A_i\}_{i \in I} \subset gl(V)$ is called irreducible on V if for any $v \in V$, the set $\{A_i(v)\}_{i \in I}$ span V.

Lemma 3.4. Let V be a finite dimensional Hermitian linear space. Suppose $\{A_i\}_{i=1,\cdots,m} \subset gl(V)$ is a finite set being irreducible on V. Then there exists $\varepsilon > 0$ such that $d(A'_i, A_i) < \varepsilon$ for $i = 1, \cdots, m$ implies that $\{A'_i\}_{i=1,\cdots,m}$ is irreducible on V. Where $A'_i \in gl(V)$, and d(,) is a metric on gl(V) defined by $d((a_{i,j}), (b_{i,j})) = \sum_{i,j} |a_{i,j} - b_{i,j}|^2$ (Here we suppose a basis of V has been chosen, so identify gl(V) with the matrix algebra.)

Proof. Denote the projective space of V as P(V). By Lemma 4.3, for any $[v] \in P(V)$ we can easily find a neighborhood U([v]) of [v] and a constant $\epsilon_{[v]} > 0$ such that $[v'] \in U([v])$ and $d(A'_i, A_i) < \epsilon$ for $i = 1, \cdots, m$ implies that $\{A'_iv'\}_{i=1,\cdots,m}$ span V. From this we can find a constant $\epsilon > 0$ such that if $d(A'_i, A_i) < \epsilon$ for $i = 1, \cdots, m$, then $\{A'_i\}_{i=1,\cdots,m}$ is irreducible on V, because P(V) is compact.

Lemma 3.5. When m is odd, for generic data Π and κ , the representation T_{LK} is irreducible. When m is even, for generic data Π and κ , the representations $T_{LK,0}$, $T_{LK,1}$

Proof. First consider the cases when m is odd. As in section 2, set $\Pi = \{k, \alpha\}$, the map $s_i \mapsto \iota(s_i), e_i \mapsto p_i \ (i = 0, 1)$ extends to a representation $\rho_{LK} : Br_{D_m}(\Pi) \to End(V_{D_m})$. Where $p_i, \iota(w)$ are defined as in section 3. Then the connection Ω_m (equation (1)) can be rewritten as $\Omega_m = \sum_{i=0}^{m-1} (k\rho_{LK}(s_i) - \rho_{LK}(e_i))\omega_i$. By Theorem 6.1 of [CH] we know $Br_{D_m}(\Pi)$ has dimension $2m + m^2$. Suppose x_1, \cdots, x_{2m+m^2} is a basis of $Br_{D_m}(\Pi)$, and suppose the generating set $\{s_0, s_1, e_0, e_1\} \subset x_1, \cdots, x_{2m+m^2}$. By Lemma 2.4, we know for generic Π , the set of operators $\{\rho_{LK}(x_1), \cdots, \rho_{LK}(x_{2m+m^2})\}$ is irreducible on V_{D_m} . Now by Lemma 2.2, we have

$$T_{LK}(\sigma_0) = \rho_{LK}(s_0) \exp(\sqrt{-1\pi\kappa(k\rho_{LK}(s_0) - \rho_{LK}(e_0))}) = \rho_{LK}(s_0) + O(\kappa)$$
(4)

and

$$T_{LK}(\sigma_1) = A(\kappa)\rho_{LK}(s_1) \exp(\sqrt{-1}\pi\kappa(k\rho_{LK}(s_1) - \rho_{LK}(e_1)))A(\kappa)^{-1} = \rho_{LK}(s_1) + O(\kappa)$$
(5)

for suitable matrix $A(\kappa)$ such that $A(\kappa) = I + O(\kappa)$. So we have

$$\frac{1}{2\sqrt{-1}\pi\kappa}(T_{LK}(\sigma_0)^2 - 1) = k\rho_{LK}(s_0) - \rho_{LK}(e_0) + O(\kappa).$$
(6)

Similarly we have

$$\frac{1}{2\sqrt{-1}\pi\kappa}(T_{LK}(\sigma_1)^2 - 1) = A(\kappa)(k\rho_{LK}(s_1) - \rho_{LK}(e_1))A(\kappa)^{-1} + O(\kappa) = (k\rho_{LK}(s_1) - \rho_{LK}(e_1)) + O(\kappa)$$
(7)

Apply Lemma 3.4 to the set $\{\rho_{LK}(x_1), \cdots, \rho_{LK}(x_{2m+m^2})\}\)$, we obtain some $\epsilon > 0$. Since $\{s_0, s_1, ks_0 - e_0, ks_1 - e_1\}\)$ generate $Br_{D_m}(\Pi)$, by above equations (3), (4), (5), (6), we can find elements $\{X_1, \cdots, X_{2m+m^2}\} \subset \mathbb{C}A_{D_m}\)$ and $\eta > 0$, such that if $\kappa < \eta$ then $d(T_{LK}(X_i), \rho_{LK}(x_i)) < \epsilon$ for $1 \le i \le 2m + m^2$. Where d(A, B) is the distance introduced in Lemma 3.4. So by Lemma 3.4 for those Π and $\kappa < \eta$, the representation T_{LK} is irreducible. So we see the subset of κ with which the representations are irreducible contains an open set. Since the subset of κ with which the representation T_{LK} being reducible is a subvariety, we see for generic κ the representation T_{LK} is irreducible. The cases when m is even can be proved similarly.

Denote the free monoid generated by X_0, X_1, E_0, E_1 as Λ . Let m be an odd positve integer, and $\Pi = (k, \alpha) \in \mathbb{C}^{\times} \times \mathbb{C}$, or m being an even positive integer and $\Pi = (k_0, k_1, \alpha_0, \alpha_1) \in (\mathbb{C}^{\times})^2 \times \mathbb{C}^2$. For $X \in \Lambda$, define $f(X) \in End(V_W)$ to be the morphism obtained by replace letters X_0, X_1, E_0, E_1 in X with $T_{LK}(\sigma_0), T_{LK}(\sigma_1), e_0, e_1$ respectively. For example, $f(X_0E_0E_1) = T_{LK}(\sigma_0)e_0e_1 \in End(V_W)$.

Definition 3.2. Choose generic constants system Π and κ . The map $\Phi^{i}_{m,\Pi,\kappa} : \Lambda \to \mathbb{C}$ (i = 0, 1) is defined by the following identity. $e_i f(X) e_i = \Phi^{i}_{m,\Pi,\kappa}(X) e_i$.

Remark 3.1. Above definition makes sense because for any $\phi \in \text{End}(V)$, $e_i \phi e_i \in \mathbb{C}e_i$ since e_i 's are projectors by Lemma 3.1 and Lemma 3.2.

Now we present the definition of Birman-Murakami-Wenzl algebras for dihedral groups. Let W be a finite Coxeter group, suppose the set of reflections R in W consists of n_W conjugacy classes C_i $(i = 1, \dots, n_W)$. We associate a pair of numbers $(k_i, \alpha_i) \in \mathbb{C}^{\times} \times \mathbb{C}$ with each class C_i , and denote the data of all (k_i, α_i) as Π . We denote the BMW algebra associated with W (to be defined) as $B_W(\Pi, \kappa)$, where $\kappa \in \mathbb{C}$. If W is of type Γ (Γ is a Coxeter matrix or Dynkin diagram), we also denote the associated BMW algebra as $B_{\Gamma}(\Pi, \kappa)$. Suppose m = 2l. So $\Pi = \{k_0, \alpha_0, k_1, \alpha_1\}$. Set $q_i = \exp(\kappa \sqrt{-1}\pi k_i)$, $l_i = \exp(\kappa \alpha_i \pi \sqrt{-1})/q_i$ for i = 0, 1.

Definition 3.3. The algebra $B_{D_m}(\Pi)$ is generated by $X_0^{\pm}, X_1^{\pm}, E_0, E_1$ with following relations.

1) $X_i X_i^- = 1$, i = 0, 1; 2) $[X_0 X_1 \cdots]_m = [X_1 X_0 \cdots]_m$; 3) $(q_i^{-1} - q_i) E_i = l_i (X_i - q_i) (X_i + q_i^{-1})$ for i = 0, 1; 4) $X_i E_i = E_i X_i = l_i^{-1} E_i$ for i = 0, 1; 5) $E_i X E_i = \Phi_{m,\Pi,\kappa}^i (X) E_i$ for any $X \in \Lambda$, i = 0, 1; 6) $E_0 X E_1 = 0 = E_1 X E_0$ for any $X \in \Lambda$.

When is odd, the data $\Pi = \{k, \alpha\}$. Set $q = \exp(\kappa \sqrt{-1}\pi k)$ and $l = \exp(\kappa \alpha \pi \sqrt{-1})/q$.

Definition 3.4. The algebra $B_{D_m}(\Pi, \kappa)$ is generated by X_0, X_1, E_0, E_1 with following relations.

1) $X_i X_i^- = 1$, i = 0, 1; 2) $[X_0 X_1 \cdots]_m = [X_1 X_0 \cdots]_m$; 3) $(q^{-1} - q) E_i = l(X_i - q)(X_i + q^{-1})$ for i = 0, 1; 4) $X_i E_i = E_i X_i = l^{-1} E_i$ for i = 0, 1; 5) $E_i X E_i = \Phi_{m,\Pi,\kappa}^i(X) E_i$ for any $X \in \Lambda, i = 0, 1$.

Remark 3.2. Above 5) of Definition 3.3 and 4) of Definition 3.4 looks containing infinite many relations. But it isn't hard to see we can choose finite many X in these relations to obtain the same algebra.

4 BMW type algebra associated with dihedral groups

In this section we study the algebras $B_{D_m}(\Pi, \kappa)$ to certify from several aspects that they are suitable deformations of the Brauer type algebras introduced in [CH]. The following Proposition 4.1 is easy to see.

Proposition 4.1. Denote the ideal in $B_{D_m}(\Pi, \kappa)$ generated by E_0, E_1 as I_m . Then when m is odd, the quotient algebra $B_{D_m}(\Pi, \kappa)/I_m$ is isomorphic to the Hecke algebra $H_{D_m}(q)$. When m is even, $B_{D_m}(\Pi, \kappa)/I_m$ is isomorphic to the Hecke algebra $H_{D_m}(q_0, q_1)$ of multiparameters.

Set
$$\tau = \frac{l(l^{-1}-q)(l^{-1}+q^{-1})}{q^{-1}-q}; \tau_i = \frac{l_i(l_i^{-1}-q_i)(l_i^{-1}+q_i^{-1})}{q_i^{-1}-q_i}$$
 for $i = 0, 1$.

Lemma 4.1. In $B_{D_m}(\Pi, \kappa)$ for odd m we have

(1) $E_i^2 = \tau E_i \text{ for } i = 0, 1;$ (2) $E_1[X_0X_1\cdots]_{m-1} = [X_0X_1\cdots]_{m-1}E_0.$ In $B_{D_m}(\Pi)$ for even m we have (3) $E_i^2 = \tau_i E_i \text{ for } i = 0, 1;$ (4) $E_1[X_0X_1\cdots]_{m-1} = [X_0X_1\cdots]_{m-1}E_1;$ (5) $E_0[X_1X_0\cdots]_{m-1} = [X_1X_0\cdots]_{m-1}E_0;$ (6) $[\cdots X_1X_0]_{m-1}E_1 = E_1[\cdots X_1X_0]_{m-1} = \tau_1^{-1}\Phi_{m,\Pi}^1([\cdots X_1X_0]_{m-1})E_1;$ (7) $[\cdots X_0X_1]_{m-1}E_0 = E_0[\cdots X_0X_1]_{m-1} = \tau_0^{-1}\Phi_{m,\Pi}^0([\cdots X_0X_1]_{m-1})E_0.$ *Proof.* (1), (3) are by simple computations. To show (2), we use relation (3) of Definition 3.3 and the identities $X_1^i[X_0X_1\cdots]_{m-1} = [X_0X_1\cdots]_{m-1}X_0^i$ for nonzero i's. The proof of (4), (5) are similar to (2). For (6), we start from the equation $E_1[\cdots X_1X_0]_{m-1}E_1 = \Phi_{m,\Pi}^1([\cdots X_1X_0]_{m-1})E_1$, which is a special case of relation (5) in Definition 3.2. Then by the proved equations (4), (1) of this lemma, the left side equals $\tau_1^{-1}[\cdots X_1X_0]_{m-1}E_1$ and (6) is proved. The proof of (7) is similar.

Lemma 4.2. (1) When m is odd, the algebra $B_{D_m}(\Pi, \kappa)$ is spanned by the subset

$$\{[X_0X_1\cdots]_i | 0 \leq i \leq m\} \cup \{[X_1X_0\cdots]_j | 1 \leq j \leq m-1\} \cup \{[\cdots X_0X_1]_i E_0[X_1X_0\cdots]_j | 0 \leq i,j \leq m-1\}.$$

 $(2) \ \ When \ m=2k \ is \ even, \ the \ algebra \ B_{D_m}(\Pi,\kappa) \ is \ spanned \ by \ the \ subset \ \{[X_0X_1\cdots]_i|0\leq i\leq m\}\cup\{[X_1X_0\cdots]_j|1\leq j\leq m-1\}\cup\{[\cdots X_0X_1]_iE_0[X_1X_0\cdots]_j|0\leq i,j\leq k-1\}\cup\{[\cdots X_1X_0]_iE_1[X_0X_1\cdots]_j|0\leq i,j\leq k-1\}.$

Proof. First we prove (1). It is a well known fact that the Hecke algebra $H_m(q)$ can be spanned by the subset $\{[X_0X_1\cdots]_i|0 \le i \le m\} \cup \{[X_1X_0\cdots]_j|1 \le j \le m-1\}$ for any q. So by Proposition 4.1, we only need to show the ideal I_m is spanned by the subset $\{[\cdots X_0X_1]_iE_0[X_1X_0\cdots]_j|0 \le i, j \le m-1\}$. Now by (2) of Lemma 4.1, I_m can be spanned by $\{f(X_0^{\pm}, X_1^{\pm}, E_0)\}$, which is the set of words composed by letters E_0, X_i^{\pm} . By (5) of Definition 3.4 and (1) of Lemma 4.1 we see I_m can be spanned by those words in $\{f(X_0^{\pm}, X_1^{\pm}, E_0)\}$ which contain only one E_0 . Then by (3), (4) of Definition 3.3, I_m can be spanned by the set $\{[\cdots X_0X_1]_{n_1}E_0[X_1X_0\cdots]_{n_2}\}$. Denote the space spanned by $\{[\cdots X_0X_1]_iE_0[X_1X_0\cdots]_j|0 \le i, j \le m-1\}$ as I'_m . We prove I'_m is stable under the multiplication by X_0, X_1 from left side.

Case 1. i < m - 1 and i is odd, then the left most letter in $[\cdots X_0 X_1]_i$ is X_1 , so we write the word in discussion as $[X_1 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j$. Since i < m - 1, $X_0 \cdot [X_1 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j \in I'_m$. And we have $X_1 \cdot [X_1 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j =$

 $[\cdots X_0 X_1]_{m-1} E_0 [\cdots X_0 X_1]_{m-1}^{-1} [\cdots X_0 X_1]_{i-1} E_0 [X_1 X_0 \cdots]_j = \gamma [\cdots X_0 X_1]_{m-1} E_0 [X_1 X_0 \cdots]_j \text{ for some } \gamma \in \mathbb{C} \text{ by } (5) \text{ of Definition 3.4.}$

Case 2. i < m-1 and i is even, then the word can be written as $[X_0 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j$. Since i < m - 1, $X_1 \cdot [X_0 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j \in I'_m$. It is easy to see we have also $X_0 \cdot [X_0 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j \in I'_m$.

Case 3. i = m - 1. It is easy to see $X_0[\cdots X_0X_1]_{m-1}E_0[X_1X_0\cdots]_j \in I'_m$. And we have $X_1[\cdots X_0X_1]_{m-1}E_0[X_1X_0\cdots]_j = X_1E_1[\cdots X_0X_1]_{m-1}[X_1X_0\cdots]_j =$

 $l^{-1}E_{1}[\cdots X_{0}X_{1}]_{m-1}[X_{1}X_{0}\cdots]_{j} = l^{-1}[\cdots X_{0}X_{1}]_{m-1}E_{0}[X_{1}X_{0}\cdots]_{j} \in I'_{m}.$

Similarly we can prove I'_m is stable by multiplication by X_0, X_1 from the right side. So (1) is proved.

For (2), denote the subspace in $B_{D_m}(\Pi, \kappa)$ spanned by

 $\{[\cdots X_0X_1]_i E_0[X_1X_0\cdots]_j | 0 \le i, j \le k-1\} \cup \{[\cdots X_1X_0]_i E_1[X_0X_1\cdots]_j | 0 \le i, j \le k-1\}$

as I'_m . We only need to show $I_m = I'_m$ as well. Similar discussions by using Definition 3.3 (Notice the relation (6)) and Lemma 4.1 show that I_m is spanned by the set $\{[\cdots X_0X_1]_{n_1}E_0[X_1X_0\cdots]_{n_2}\}\cup\{[\cdots X_1X_0]_{n_1}E_1[X_0X_1\cdots]_{n_2}\}$. So it is enough to show that I'_m is stable under the left and right multiplication by X_0, X_1 . It suffice to show the left case and the cases for words $\{[\cdots X_0X_1]_iE_0[X_1X_0\cdots]_j|0 \le i, j \le k-1\}$. Other cases are similar.

Case 1. i < k - 1 and i is odd. Then the word in discussion can be presented as $[X_1 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j$. Since i < k - 1 we have $X_0 \cdot [X_1 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j \in I'_m$. On the other hand, $X_1 \cdot [X_1 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j = X_1^2 [\cdots X_0 X_1]_{i-1} E_0 [X_1 X_0 \cdots]_j \in I'_m$ by (3), (6) of Definition 3.3.

Case 2. i < k-1 and i is even. The word can be written as $[X_0 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j$. Since i < k-1 we have $X_1 \cdot [X_0 \cdots X_0 X_1]_i E_0 [X_1 X_0 \cdots]_j \in I'_m$. On the other hand,

 $X_{0} \cdot [X_{0} \cdots X_{0}X_{1}]_{i} E_{0}[X_{1}X_{0} \cdots]_{j} = X_{0}^{2}[\cdots X_{0}X_{1}]_{i-1} E_{0}[X_{1}X_{0} \cdots]_{j} \in I_{m}^{'} \text{ by } (3), (5) \text{ of Definition}$ 3.3.

Case 3. i = k-1. If k is odd, then the word can be written as $[X_0 \cdots X_0 X_1]_{k-1} E_0 [X_1 X_0 \cdots]_j$. Now we have

 $\begin{array}{l} X_{0} \cdot [X_{0} \cdots X_{0} X_{1}]_{k-1} E_{0}[X_{1} X_{0} \cdots]_{j} \in I_{m}^{'} \text{ by } (3), (5) \text{ of Definition 3.3. And we have} \\ X_{1} \cdot [X_{0} \cdots X_{0} X_{1}]_{k-1} E_{0}[X_{1} X_{0} \cdots]_{j} = \tau_{0}^{-1} \Phi_{m,\Pi}^{0}([\cdots X_{0} X_{1}]_{m-1})[\cdots X_{1} X_{0}]_{k-1}^{-1} E_{0}[X_{1} X_{0} \cdots]_{j} = \\ \tau_{0}^{-1} \Phi_{m,\Pi}^{0}([\cdots X_{0} X_{1}]_{m-1})[\cdots X_{0}^{-1} X_{1}^{-1}]_{k-1} E_{0}[X_{1} X_{0} \cdots]_{j} = \gamma [\cdots X_{0} X_{1}]_{k-1} E_{0}[X_{1} X_{0} \cdots]_{j} \mod I_{m}^{'} \\ \text{for some } \gamma \in \mathbb{C}. \text{ Where the first " = " in above equation is by (7) of Lemma 4.1, the third} \\ " = " is by iterated use of Definition 3.3. The cases of k being even and i = k - 1 are easy \\ \text{to see.} \end{array}$

The dihedral group D_3 is just the symmetric group S_3 . We have

Proposition 4.2. The algebra $B_{D_3}(\Pi, \kappa)$ in Definition 3.4 is isomorphic the BMW algebra $B_3(q, l)$.

Proof. By comparing Definition 1.1 for $B_3(q, l)(B_{A_2}(q, l))$ with Definition 2.1 for $B_{D_3}(\Pi, \kappa)$, we only need to prove that the relation 5) of Definition 3.4 can be induced from relations in Definition 1.1 for $B_3(q, l)$. First, for $B_3(q, l)$, let X be any word composed from X_i, E_i (i = 0, 1), we claim there exist $\phi(X), \psi(X) \in \mathbb{C}[q^{\pm}, l^{\pm}]$, such that $E_0XE_0 = \phi(X)E_0, E_1XE_1 = \psi(X)E_1$. Denote the length of a word X as l(X). Call letters X_0, E_0 (X_1, E_1) as of type 0 (type1). By relations in Definition 1.1, we only need to consider those X in which type 0 letters and type 1 letters appear alternatively and l(x) being odd. It is proved in Proposition 2.3 of [CGW1] that $E_0E_1E_0 = E_0$ and $E_1E_0E_1 = E_1$. So the claim is true for l(X) = 0, 1. Suppose we have proved the claim for $l(X) \leq 2k + 1$. Now suppose l(X) = 2k + 3. For the case E_0XE_0 , we can assume the letter E_0 doesn't appear in X and E_1 appear in X at most once. Because otherwise we can use induction. So we only need to consider those X with $l(X) \leq 3$. Because if $l(X) \geq 5$, by assumption above $X = X_1X_0A_1Y$ or $X = YA_1X_0X_1$, where $A_1 = X_1$ or E_1 . Now $E_0X_1X_0A_1YE_0 = X_1X_0E_1A_1YE_0 = \lambda X_1X_0E_1YE_0 = \lambda E_0X_1X_0YE_0$ for some $\lambda \in \mathbb{C}[q^{\pm}, l^{\pm}]$, which implies $E_0X_1X_0A_1YE_0 \in \mathbb{C}[q^{\pm}, l^{\pm}]E_0$ because $l(X_1X_0Y) < l(X)$. So to

prove the claim for E_0XE_0 , we are left with the cases $X \in \{X_1X_0X_1, E_1X_0X_1, X_1X_0E_1\}$, which can be certified by simple computations. The cases for E_1XE_1 can be proved similarly.

Since the Lawrence-Krammer representation factor through $B_3(q,l)$, we see $\phi(X) = \Phi^0_{3,\Pi,\kappa}(X)$ and $\psi(X) = \Phi^1_{3,\Pi,\kappa}(X)$, where $\Pi = \{\alpha, k\}$ such that $q = \exp(\kappa \sqrt{-1}\pi k)$, $l = q^{\alpha-1}$.

When m is odd, suppose (V, ρ) is a finite dimensional representation of $Br_{D_m}(\Pi)$, where $\Pi = \{k, \alpha\}$. Then on the trivial bundle $M_{D_m} \times V$ we have a D_m -invariant flat connection

$$\rho(\Omega_{D_m}) = \kappa \sum_{i=0}^{m-1} (k\rho(s_i) - \rho(e_i))\omega_i.$$
(8)

It induces a flat connection $\rho(\bar{\Omega}_{D_m})$ on $M_{D_m} \times_{D_m} V$, whose monodromy representation will be denoted as $T_{\rho} : A_{D_m} \to GL(V)$. When m = 2k is even, suppose (V, ρ) is a finite dimensional representation of $Br_{D_m}(\Pi)$, where $\Pi = \{k_0, \alpha_0; k_1, \alpha_1\}$. Similarly by using the D_m -invariant flat connection

$$\rho(\Omega_{D_m}) = \kappa[\sum_{i=0}^{k-1} (k_0 \rho(s_{2i}) - \rho(e_{2i}))\omega_{2i} + \sum_{i=0}^{k-1} (k_1 \rho(s_{2i+1}) - \rho(e_{2i+1}))\omega_{2i+1}],$$
(9)

we obtain a monodromy representation $T_{\rho} \! : \ A_{D_{\mathfrak{m}}} \to GL(V).$

Proposition 4.3. (1) Suppose m is odd. Let the data Π be generic so that $Br_{D_m}(\Pi)$ is a semisimple algebra. Then for generic κ , the representation T_{ρ} of A_{D_m} factors through the algebra $B_{D_m}(\Pi, \kappa)$. (2) Suppose m is even. Let the data Π be generic so that $Br_{D_m}(\Pi)$ is a semisimple algebra. Then for generic κ , the representation T_{ρ} of A_{D_m} factors through the algebra $B_{D_m}(\Pi, \kappa)$.

Proof. We prove (2), the proof of (1) is similar and easier. So let m be even. Denote the natural quotient map from $Br_{D_m}(\Pi)$ to $\mathbb{C}D_m$ as π_m . Suppose $\rho_1, \dots, \rho_{\nu}$ are all the irreducible representations of $\mathbb{C}D_m$, denote the representation of $Br_{D_m}(\Pi)$ induced from ρ_i through π_m as $\bar{\rho}_i$. As in Lemma 2.4, we have two more representation $\rho_{LK,c_0}, \rho_{LK,c_1}$ of $B_{D_m}(\Pi)$. By Theorem 6.1 ,Theorem 6.3 of [CH], we know for generic data Π , the algebra $Br_{D_m}(\Pi)$ is semisimple and $\bar{\rho}_1, \dots, \bar{\rho}_{\nu}, \rho_{LK,c_0}, \rho_{LK,c_1}$ are all the irreducible representations of $Br_{D_m}(\Pi)$. So because of semisimplicity of $Br_{D_m}(\Pi)$, we only need to prove the proposition when ρ is one of above listed irreducible representations. Set $q_i = \exp(\sqrt{-1}\pi\kappa k_i)$, $l_i =$ $\exp(\kappa\alpha_i\pi\sqrt{-1})/q_i. \text{ Set } e_i^{\rho} = \frac{l_i}{q_i^{-1}-q_i}(T_{\rho}(\sigma_i)-q_i)(T_{\rho}(\sigma_i)+q_i^{-1}) \text{ for } i=0,1. \text{ We want to show } i=0,1. \text{ We want } i=0,1. \text{ We want to show } i=0,1. \text{ We want } i=0,1. \text{ We wa$ the map $\phi_m : E_i \mapsto e_i^{\rho}, X_i \mapsto T_{\rho}(\sigma_i)$ extends to an algebraic morphism. Recall by Lemma 2.2, $T_{\rho}(\sigma_i)$ is conjugated to $\rho(s_i) \exp(\kappa \sqrt{-1}\pi(k_i\rho(s_i) - \rho(e_i)))$ where $k_i = k_0$ for even i, and $k_i = k_1$ for odd i. Suppose $\rho = \bar{\rho}_i$, then $k_i \rho(s_i) - \rho(e_i) = k_i \rho(s_i)$ can be presented by a diagonal matrix whose diagonal elements being in $\{\pm k_i\}$. So we know $e_i^{\rho} = 0$ and ϕ_m keep all relations in Definition 3.3 and these cases are done. Now suppose $\rho = \rho_{LK,c_0}$. By (1) of lemma 3.2, we see ϕ_m satisfy (4) of Definition 3.3. The cases for (1), (2), (3) of Definition 3.3 are trivial. Since T_{ρ} is a subrepresentation of T_{LK} , by definition of $\Phi_{m,\Pi}^{i}(X)$ we know

relation (5) of Definition 3.3 is also satisfied by ϕ_m . Since $T_{\rho_{LK,c_0}}(e_1) = 0$, we see $e_1^{\rho} = 0$ so relation (6) of Definition 3.3 is also satisfied by ϕ_m . The case for $\rho = \rho_{LK,c_1}$ is similar. \Box

Remark 4.1. If (V, ρ) is the infinitesimal Lawrence-Krammer representation of $Br_m(\Pi)$ (in Lemma 2.3), then the connection $\rho(\Omega_{D_m})$ of equation (7) ((8)) coincide with the connection Ω_m of equation (1) ((2)). So above proposition shows the generalized Lawrence-Krammer representation of A_{D_m} factor through the algebra $B_{D_m}(\Pi, \kappa)$. It thus provide us a way to compute the monodromy of Marin's flat connections by using the generalized BMW type algebras, at least in cases of dihedral groups.

Theorem 4.1. (1) When m is odd, for generic data Π, κ , the algebra $B_{D_m}(\Pi, \kappa)$ is semisimple with dimension $2m + m^2$; (2) When m is even, for generic data Π , the algebra $B_{D_m}(\Pi, \kappa)$ is semisimple with dimension $2m + \frac{m^2}{2}$.

Proof. First we prove (1). According to Proposition 4.1, denote the quotient map from $B_{D_m}(\Pi, \kappa)$ to the Hecke algebra $H_{D_m}(q)$ as π . For generic q, $H_{D_m}(q)$ is semisimple and we denote its irreducible representations as ρ_1, \dots, ρ_k . Through π , each ρ_i induces an irreducible representation $\bar{\rho}_i$ of $B_{D_m}(\Pi, \kappa)$. These representations are different from each other and we have $\sum_{i=1}^{k} (\dim \bar{\rho}_i)^2 = \dim H_{D_m}(q) = 2m$. Because for any $1 \leq i \leq k$, the annihilating polynomial of $\bar{\rho}_i(X_j)$ has degree 2, and the annihilating polynomial of $\bar{T}_{LK}(X_j)$ has degree 3, so by Lemma 3.5 we know \bar{T}_{LK} is an irreducible representation different with any $\bar{\rho}_i$. So by Wedderburn-Artin theorem, we have dim $B_{D_m}(\Pi, \kappa) \geq \sum_{i=1}^{k} (\dim \bar{\rho}_i)^2 + (\dim \bar{T}_{LK})^2 = 2m + m^2$. So by (1) of Lemma 4.2 and by Wedderburn-Artin theorem again, we know for generic data $\Pi, \kappa, B_{D_m}(\Pi, \kappa)$ is a semisimple algebra with dimension $2m + m^2$. The proof of (2) is similar, by using (2) of Lemma 4.2 and the Wedderburn-Artin theorem.

Remark 4.2. By Theorem 6.1 of [CH], we have dim $\operatorname{Br}_{D_m}(\Upsilon) = m^2 + 2m$ for odd m, and dim $\operatorname{Br}_{D_m}(\Upsilon) = m^2 + \frac{m}{2}$ for even m. Above Theorem 4.1 and Proposition 4.2, 4.3 say that the algebra $\operatorname{B}_{D_m}(\Pi, \kappa)$ is a satisfactory deformation of the Brauer type algebra $\operatorname{Br}_{D_m}(\Upsilon)$.

Now let W_{Γ} be a Coxeter group with Coxeter matrix $\Gamma = (m_{i,j})_{n \times n}$. Which has a presentation as in Theorem 2.2. For each $1 \leq i \leq n$, choose a pair of constants $(k_i, \alpha_i) \in \mathbb{C}^{\times} \times \mathbb{C}$ such that $(k_i, \alpha_i) = (k_j, \alpha_j)$ if s_i is conjugated to s_j . Denote the data consisting of $\{(k_i, \alpha_i)\}_{i=1}^m$ as one symbol Π . Let $\kappa \in \mathbb{C}$. Set $q_i = \exp(\kappa \sqrt{-1}\pi k_i)$, $l_i = (q_i)^{\alpha_i - 1}$ for $1 \leq i \leq n$. The definition of BMW type algebras of Dihedral groups naturally inspire us to present the following definition of $B_W(\Pi, \kappa)$, the BMW type algebra of type W. These algebras will be generated by elements $X_i, E_i, 1 \leq i \leq n$. For $1 \leq i < j \leq n$, denote the parabolic subgroup of W generated by s_i, s_j as $W_{i,j}$. Then $W_{i,j}$ is isomorphic to $D_{m_{i,j}}$. Denote the data $\{(k_i, \alpha_i), (k_j, \alpha_j)\}$ as $\Pi_{i,j}$. Denote the free monoid generated by X_i, E_i, X_j, E_j as $\Lambda_{i,j}$. Repeat the definition of function $\Phi^i_{m,\Pi,\kappa}$ in Definition 3.1, but replace the roles of $(D_m, X_0, E_0, X_1, E_1, \Lambda, \Pi)$ by $(W_{i,j}, X_i, E_i, X_j, E_j, \Lambda_{i,j}, \Pi_{i,j})$, we obtain functions $\Phi^i_{m_{i,j}, \Pi_{i,j}}$ and $\Phi^j_{m_{i,j}, \Pi_{i,j}}$ from $\Lambda_{i,j}$ to \mathbb{C} , which are similar to $\Phi^0_{m,\Pi,\kappa}$ and $\Phi^1_{m,\Pi,\kappa}$ in Definition 3.1 respectively.

Definition 4.1. The algebra $B_{\Gamma}(\Pi, \kappa)$ is generated by elements $X_i^{\pm}, E_i, 1 \leq i \leq n$ with the following relations.

- (1). $X_i X_i^- = X_i^- X_i = 1$ for any i;
- (2). $[X_iX_j\cdots]_{m_{i,j}} = [X_jX_i\cdots]_{m_{i,j}}$ for any $i \neq j$;
- (3). $(q_i^{-1} q_i)E_i = l_i(X_i q_i)(X_i + q_i^{-1})$ for any i;
- (4). $X_i E_i = E_i X_i = l_i^{-1} E_i$ for any i;
- $(5). E_i X E_i = \Phi^i_{\mathfrak{m}_{i,j}, \Pi_{i,j}, \kappa}(X) E_i \text{ and } E_j X E_j = \Phi^j_{\mathfrak{m}_{i,j}, \Pi_{i,j}, \kappa}(X) E_j \text{ for any } i < j \text{ and } X \in \Lambda_{i,j};$
- (6). $E_i X E_j = 0$ for any $X \in \Lambda_{i,j}$ if $m_{i,j}$ is an even number greater than 2.

5 General BMW Type Algebras

Let $\Gamma = (m_{i,j})_{n \times n}$ be a Coxeter matrix. For each $1 \leq i \leq n$, choose $q_i \in \mathbb{C}^{\times}$ such that $q_i = q_j$ if s_i is conjugated to s_j . Denote the data $\{q_1, \cdots, q_n\}$ as \bar{q} . The type Γ Hecke algebra $H_{\Gamma}(\bar{q})$ is the quotient algebra of the group algebra $\mathbb{C}A_{\Gamma}$ to the ideal generated by $\{(\sigma_i - q_i)(\sigma_i + q_i^{-1})\}$. The following Proposition is evident.

Proposition 5.1. The map $X_i \mapsto \sigma_i$, $E_i \mapsto 0$ $(1 \le i \le n)$ extends to a surjection from $B_{\Gamma}(\Pi, \kappa)$ to $H_{\Gamma}(\bar{q})$ if the data Π and \bar{q} satisfies $q_i = \exp \kappa \sqrt{-1}\pi k_i$ for $1 \le i \le n$.

The following Theorem 5.1 is proved using similar method as in the proof of Proposition 2.9 of [CGW1]. Suppose $\Gamma = (m_{i,j})_{n \times n}$ is a finite type Coxeter matrix. Let w_0 be the longest element of W_{Γ} and denote the length of w_0 as L. A sequence in [1, n] is a sequence (i_1, i_2, \dots, i_k) where $1 \le i_v \le n$ for any v. The sequence (i_1, i_2, \dots, i_k) is called reducible if $i_v = i_{v+1}$ for some v. Otherwise it is called irreducible. A basic transformation of a sequence Q is, replacing a subsequence $[i, j, \dots]_{m_{i,j}}$ in Q by $[j, i, \dots]_{m_{i,j}}$. For example, transform (3, 1, 2, 1, 2, 4) to (3, 2, 1, 2, 1, 4) when $m_{1,2} = 4$. By the theory of Coxeter groups, we see the following fact: if a sequence Q has length greater than L, then Q can be transformed into a reducible sequence.

Theorem 5.1. If Γ is of finite type, then $B_{\Gamma}(\Pi, \kappa)$ is a finite dimensional algebra.

Proof. Denote the set of words composed by $\{X_i, E_i\}_{1 \le i \le n}$ as S. By (3) of Definition 4.1, we see $B_{\Gamma}(\Pi, \kappa)$ can be spanned by S. Let w_0 be the longest element of W_{Γ} , denote the length of w_0 as L. For a word W, denote the length of W as l(W). Let $S_0 = \{W \in S | l(W) \le L\}$. Denote the subspace of $B_{\Gamma}(\Pi, \kappa)$ spanned by S_0 as A. The theorem follows from the following assertion: Any word in S lie in A as an element of the algebra. Let $W \in S$. When l(W) < L, the assertion is evident. Suppose we have prove that for any $W \in S$ with $l(W) \le K$, we have $W \in A$. Here we can ask $K \ge L$. Let $W \in S$ is a word with length K + 1. Each word $W \in S$ determine a sequence of indices Q_W . For example, if $W = E_1X_2X_3E_2$ then $Q_W = (1, 2, 3, 2)$. Since the length of Q_W is greater than L, we see Q_W can be transformed into a reducible sequence by M times of basic transformations. Then we do induction on M. If M = 0, then Q_W is reducible, which mean W contains a subword with one of the following type: E_iE_i , X_iE_i , E_iX_i , X_iX_i . By relation (3), (4) of Definition 4.1 and by induction we see $W \in A$.

Now suppose the assertion of $l(W) \leq K$ and $l(W) = K + 1, M \leq J$ are proved. Suppose we have a word W such that Q_W is irreducible, l(W) = K + 1 and M = J + 1. Then there is a series of sequence Q_1, Q_2, \dots, Q_{J+2} such that $Q_1 = Q_W, Q_{J+2}$ is reducible and Q_{i+1} is obtained by doing one basic transformation to Q_i . Denote the subword of W in the position where the basic transformation to Q_2 occur as V. And suppose the basic transformation from Q_1 to Q_2 is by replacing a subsequence $[i, j \dots]_{m_{i,j}}$ with $[j, i \dots]_{m_{i,j}}$. For example when $W = X_1 E_2 X_3 X_2 E_3$, and $Q_2 = (1, 3, 2, 3, 2)$ then $V = E_2 X_3 X_2 E_3$. First we observe if there are two E_i , or two E_j in V then we can shorten V by Relation (5) of Definition 4.1 and the assertion for W can be proved. So we can suppose there is at most one E_i and one E_j in V.

Case 1. $m_{i,j} = 2k + 1$ is odd, $V = AE_iE_jC$ where A, C contain only X_i, X_j . Denote the length of A, C as a, c respectively, so by above arguments $A = [\cdots X_iX_j]_a$ and $C = [\cdots X_jX_i]_c$. We have a + c = 2k - 1. Suppose a > c, so $a \ge k$. By applying (2) of Lemma 4.1 to the subalgebra generated by $\{X_i, X_j, E_i, E_j\}$, we have $AE_iE_jC = [\cdots]_{2k-a}E_jDE_jC = \lambda[\cdots]_{2k-a}E_jC$, where $[\cdots]_{2k-a}$ denote some word with length 2k - a, and D is some word. Since $l([\cdots]_{2k-a}E_jC) \le K$, by replace X_i^{-1} in $[\cdots]_{2k-a}$ with linear sum of X_i, E_i , we complete the proof of this case by induction. The case of a < c is similar.

Case 2. $m_{i,j} = 2k + 1$ is odd, $V = AE_iBE_jC$ Where A, B, C are nonempty words contain only X_i, X_j . Denote the length of A, B, C as a, b, c respectively. We have $a \neq c$ because otherwise $V = AE_iBE_iC$ which contradicts our assumption. We can suppose a > c. The cases of a < c is similar. Again by using (2) of Lemma 4.1, $AE_iBE_jC = [\cdots]_{2k-a}E_jDE_jC =$ $\lambda[\cdots]_{2k-a}E_jC$. Now $l(\lambda[\cdots]_{2k-a}E_jC) = 2k - a + 1 + c < 2k - 1$ so by induction we complete the proof of these cases.

Case 3. $m_{i,j} = 2k + 1$ is odd, $V = AE_iB$ where A, B are nonempty words contain only X_i, X_j . Denote the length of A, B as a, b respectively. Suppose B is nonempty. Relation (3) of Definition 4.1 gives us an identity $X_i = \lambda X_i^{-1} + \mu E_i + \gamma$. Replace every X_i in B with $\lambda X_i^{-1} + \mu E_i + \gamma$ and every X_j in B with $\lambda X_j^{-1} + \mu E_j + \gamma$, we see $V = \lambda' AE_i[X_j^{-1}X_i^{-1}\cdots]_b + U$ where U is a liner sum of words containing two E_i , or containing more than one E_i and containing one E_j with length K + 1, or with smaller length. By Case 1 and 2, we see $U \in A$. And by (2) of Lemma 4.1, we have $AE_i[X_j^{-1}X_i^{-1}\cdots]_b = [\cdots X_j^{-1}X_i^{-1}]_bE_jA'$. By replacing X_i^{-1}, X_j^{-1} in the subword $[\cdots X_j^{-1}X_i^{-1}]_b$ we see W = a linear sum of words with sequence $Q_2 \mod A$. Since Q_2 can be transformed to an reducible sequence with fewer basic transformations, we proved these cases.

Case 4. $m_{i,j} = 2k + 1$ is odd, $V = [X_iX_j \cdots]_{2k+1}$. By replacing the subword V in W by $[X_jX_i \cdots]_{2k+1}$, we see the resulted word (equal W) has sequence Q_2 , then we use induction.

For the cases when $m_{i,j} = 2k$, the same argument shows we only need to consider those cases when V contain at most one E_i and one E_j . Now (6) of Definition 4.1 show we only need to consider the cases (1) that V contain one E_i or one E_j ; (2) the case V contain only X_i, X_j . We can prove the cases (1) similarly as above case 3, and prove the cases (2) as above case 4.

When Γ is a simply laced type Coxeter group, then all reflections of W_{Γ} lie in the same

conjugacy class. So the data Π of $B_{\Gamma}(\Pi, \kappa)$ consists of $\{k, \alpha\}$ essentially. As a corollary of Proposition 4.2, we have

Proposition 5.2. The algebra $B_{\Gamma}(\Pi, \kappa)$ defined in Definition 4.1 is isomorphic to the simply laced BMW algebra $B_{\Gamma}(q, l)$ of [CGW1] for $q = \exp(\kappa \sqrt{-1}\pi k)$, $l = \frac{\exp(\alpha \pi \sqrt{-1})}{q}$.

Let W be a finite type Coxeter group. Let R be the set of reflections in W. Suppose (V, ρ) is a finite dimensional representation of $Br_W(\Pi)$. Where $\Pi = \{k_s, \alpha_s\}_{s \in \mathbb{R}}$. Then by using the KZ connection (Definition 2.3), we have a flat, W-invariant connection $\rho(\Omega_W) = \kappa \sum_{s \in \mathbb{R}} (k_s \rho(s) - \rho(e_s)) \omega_s$ on $M_W \times V$, which induces a representation $T_\rho : A_W \to GL(V)$. Suppose W is of rank n and has a canonical set of generators $\{s_i\}_{1 \leq i \leq n}$ as in Theorem 2.2. Suppose the Artin group A_W has a canonical set of generators $\{\sigma_i\}_{1 < i < n}$ as in Theorem 2.3.

Theorem 5.2. For generic Π and κ , the representation T_{ρ} factor through $B_W(\Pi, \kappa)$.

Proof. We want to show if set $X_i = T_{\rho}(\sigma_i)$ and $E_i = \frac{l_i}{q_i^{-1}-q_i}(T_{\rho}(\sigma_i)-q_i)(T_{\rho}(\sigma_i)+q_i^{-1})$ for $1 \le i \le n$, where $q_i = \exp(\kappa k_{s_i}\pi\sqrt{-1})$ and $l_i = \exp(\kappa \alpha_i\pi\sqrt{-1})/q_i$, then $\{X_i, E_i\}_{1\le i\le n}$ satisfies the relations in Definition 4.1. If W is a dihedral group D_m , this is true by Proposition 4.3. The cases for general W can be reduced to the cases for dihedral groups by using Theorem 2.4 as follows. Recall the notations " Δ , H_{s_i} " above Theorem 2.2, and the notations " Π , $\Pi_{i,j}$ " before Definition 4.1. For $1 \le i < j \le n$, first we suppose $m_{i,j}$ is odd. Denote $L = H_{s_i} \cap H_{s_j}$, define R_L , W_L , A_{W_L} and M_{W_L} as in section 2 (above Theorem 2.4). We can identify A_{W_L} with the Artin group $A_{D_{m_{i,j}}}$ in suitable way, so the morphism $\overline{\lambda}_L$ from A_{W_L} to A_W maps σ_0, σ_1 to σ_i, σ_j respectively.

In this case we have a natural isomorphism $\phi_{i,j}$ from $W_{D_{m_{i,j}}}$ to W_L extending the map $s_0 \mapsto s_i, s_1 \mapsto s_j$, and we can identify $M_{W_{D_{m_{i,j}}}}$ with M_{W_L} (actually $M_{W_L} \cong M_{W_{D_{m_{i,j}}}} \times \mathbb{C}^{n-2}$, but it will not make any difference.) The morphism $\phi_{i,j}$ can be extended to a morphism from $Br_{D_{m_{i,j}}}(\Pi_{i,j})$ to $Br_W(\Pi)$ by sending $e_0, e_1 \in Br_{D_{m_{i,j}}}(\Pi_{i,j})$ to $e_i, e_j \in Br_W(\Pi)$ respectively. And we know the morphism $\overline{\lambda}_L$ maps A_{W_L} isomorphically to the parabolic subgroup of A_W generated by σ_i, σ_j . Through $\phi_{i,j}$ we have a representation $\rho \circ \phi_{i,j}$: $Br_{D_{m_{i,j}}}(\Pi_{i,j}) \to gl(V)$. As in section 2, set $\Omega' = \kappa \sum_{s \in R_L} (k_s \rho(s) - \rho(e_s)) \omega_s$. It isn't hard to see Ω' coincide with the connection $\Omega_{\rho \circ \phi_{i,j}}$ on M_{W_L} . So by Proposition 4.3, if we set $X_0 = T_L(\sigma_0)$ and $X_1 = T_L(\sigma_1)$, then X_0, X_1 satisfy the relations in Definition 3.4 for $m = m_{i,j}$ and suitable data Π . (since by (3) of Definition 3.4, E_i can be presented by X_i for generic data, all those relations can be seen as relations between X_0, X_1 .) By Theorem 2.4, there is a element $A \in GL(V)$, so $T \circ \overline{\lambda}_L(\rho_i) = AT_L(\rho_i)A^{-1}$ for i = 0, 1. Since $T_\rho \circ \overline{\lambda}_L(\rho_0) = T_\rho(\sigma_i), T_\rho \circ \overline{\lambda}_L(\rho_1) = T(\sigma_j)$, we have that $X_l = T_\rho(\sigma_l)$ and $E_l = \frac{l_l}{q_l^{-1}-q_l}(T_\rho(\sigma_l) - q_l)(T_\rho(\sigma_l) + q_l^{-1})$ (l = i, j) satisfy the relation (5) of Definition 4.1.

Similarly we can prove if $m_{i,j}$ is even, then X_i, X_j, E_i, E_j satisfy relation (5), (6) of Definition 4.1, and we can prove X_i, E_i satisfy relation (3), (4), by using Theorem 2.4.

References

- [BW] J.S.BIRMAN, H.WENZL, Braids, link polynomials and a new algebra, Tran. Amer. Math. Soc 313(1989), 249-273.
- [Bri] E.BRIESKORN, Die Fundamentalgruppe des Raumes der regularen Orbits einer endlichen komplexen Spiegelungsgruppe *Invent. Math.* **12**(1971),57-61.
- [BS] E.BRIESKORN, K.SAITO, Artin-Gruppen und Coxeter-Gruppen, Invent. Math 17 (1972),245-271.
- [Bi] S.BIGELOW, Braid groups are linear, J.Am. Math. Soc. 14 (2001), 471-486.
- [CH] Z.CHEN, Flat connections and Brauer type algebras, preprint 2010.
- [CW] A.M.COHEN, D.B.WALES, Linearity of Artin groups of finite type, Israel Journal of Mathematics131(2002), 101-123
- [CW2] A.M.COHEN, D.B.WALES, The Birman-Murakami-Wenzl algebras of type E_n , Transformation Groups 16 (2011),681-715.
- [CGW1] A.M.COHEN, D.A.H.GIJSBERS, D.B.WALES, BMW algebras of simply laced type, Journal of Algebra 286 (2005), 107-153.
- [1] A.M.COHEN, D.A.H.GIJSBERS, D.B.WALES, The BMW Algebras of Type Dn, arXiv:0704.2743(2007)
- [Di] F.DIGNE, On the linearity of Artin braid groups, J. Algebra 268 (2003), 39-57.
- [2] FM.GOODMAN, Cellularity of cyclotomic Birman-Wenzl-Murakami algebras, Journal of algebra 321 (2009), 3299-3320
- [GH] FM.GOODMAN, HM.HAUSCHILD, Affine Birman-Wenzl-Murakami algebras and tangles in the solid torus, Arxiv preprint math/0411155.
- [Kr] D.KRAMMER, Braid groups are linear, Annals of Math. 155 (2002), 131-156.
- [Ha] R.HÄRING-OLDENBERG, Cyclotomic Birman-Murakami-Wenzl algebras, Journal of Pure and applied algebras161(2001),113-144
- [Hu] J.HUMPHREYS, Reflection groups and Coxeter groups, Cambridge studies in adv. math 29, (1992)
- [K01] T.KOHNO, Monodromy representations of braid groups and Yang-Baxter equations Ann.Inst.Fourier 37 (1987), 139-160.
- [Ma1] I.MARIN , Quotients infinitesimaux du groupe de tresses , Ann. Inst. Fourier, Grenoble 53 No.5, (2003) , 1323-1364.

- [Ma2] I.MARIN, Krammer representations for complex braid groups, arXiv:0711.3096v1 (2007).
- [Ma3] I.MARIN, Sur les representations de Krammer gnriques. Ann. Inst. Fourier, Grenoble 57 no. 6, (2007),1883-1925.
- [Ma4] I.MARIN, On the representation theory of braid groups. arXiv:math/0502118v3.
- [Mu] J.MURAKAMI, The Kauffman polynomial of links and representation theory, Osaka J. Math 24 (1987), 745-758.
- [RX] H.RUI J.XU The representations of cyclotomic BMW algebras, Journal of Pure and Applied Algebra 213 (2009), 2262-2288.
- [RX] H.RUI, J.XU, On the semisimplicity of the cyclotomic brauer algebras ii Journal of algebra 312 issue 2, (2007),995-1010.
- [Wi] S.WILCOX, On the freeness of the cyclotomic BMW algebras: admissibility and an isomorphism with the cyclotomic Kauffman tangle algebras, Arxiv preprint arXiv:0911.5284.
- [Xi2] C. XI, On the Quasi-Heredity of Birman-Wenzl Algebras, Advances in Mathematics 154, Issue 2, (2000), 280-298.
- [Yu] SH. YU, The Cyclotomic BMW algebras, arXiv:0810.0069v1.
- [Z] M.ZINNO, On Krammer's representations of the braid groups, Math.Ann. 321, (2001), 197-211.