

# FREE GROUP $C^*$ -ALGEBRAS ASSOCIATED WITH $\ell_p$

RUI OKAYASU

**ABSTRACT.** For every  $p \geq 2$ , we give a characterization of positive definite functions on a free group with finitely many generators, which can be extended to the positive linear functionals on the free group  $C^*$ -algebra associated with the ideal  $\ell_p$ . This is a generalization of Haagerup's characterization for the case of the reduced free group  $C^*$ -algebra. As a consequence, the associated  $C^*$ -algebras are mutually non-isomorphic, and they have a unique tracial state.

## 1. INTRODUCTION

N. P. Brown and E. Guentner introduce new  $C^*$ -completion of the group ring of a countable discrete group  $\Gamma$  in [2]. More precisely, for a given algebraic two-sided ideal in  $\ell_\infty(\Gamma)$ , they define the associated group  $C^*$ -algebra. These recover the full group  $C^*$ -algebra for  $\ell_\infty(\Gamma)$  itself, and the reduced group  $C^*$ -algebra for  $c_c(\Gamma)$ , respectively. Hence if we take  $c_0(\Gamma)$  or  $\ell_p(\Gamma)$  with  $p \in [1, \infty)$  for example, we may obtain new group  $C^*$ -algebra. We remark that a standard characterization of amenability implies that the associated  $C^*$ -algebras of an amenable group are all isomorphic for any ideals. In [2], they also give a characterization of the Haagerup property and Property (T) in terms of ideal completions.

In this paper, we study their  $C^*$ -algebra of a free group associated with  $\ell_p$ . By [2], for any  $p \in [1, 2]$ , the group  $C^*$ -algebra of  $\Gamma$  associated with  $\ell_p$  is isomorphic to the reduced group  $C^*$ -algebra. Therefore the case where  $p \in (2, \infty)$  is essential. Our main result is a characterization of positive definite functions on a free group, which can be extended to the positive linear functionals on the free group  $C^*$ -algebra associated with  $\ell_p$ . In [5], U. Haagerup gives its characterization for the case of the reduced free group  $C^*$ -algebra. Thus our theorem is a generalization of his result. As a consequence, the free group  $C^*$ -algebras associated with  $\ell_p$  are mutually non-isomorphic for every  $p \in [2, \infty]$ . We also obtain that for each  $p \in [2, \infty)$ , the free group  $C^*$ -algebra associated with  $\ell_p$  has a unique tracial state. Moreover, we consider algebraic ideals

$$D_p^-(\Gamma) = \bigcup_{\varepsilon > 0} \ell_{p-\varepsilon}(\Gamma)$$

for  $1 < p \leq \infty$ , and

$$D_p^+(\Gamma) = \bigcap_{\varepsilon > 0} \ell_{p+\varepsilon}(\Gamma)$$

for  $1 \leq p < \infty$ . Then, by using our characterization, it is also shown that the free group  $C^*$ -algebra associated with  $D_p^\pm$  coincides with the free group  $C^*$ -algebra associated with  $\ell_p$ .

---

2000 *Mathematics Subject Classification.* Primary 46L05; Secondary 22D25.

*Key words and phrases.*  $C^*$ -algebras, free groups, positive definite functions.

The author was supported in part by JSPS.

## 2. PRELIMINARIES

In this section, we fix the notations for the convenience of the reader and recall some results in [2].

Let  $\Gamma$  be a countable discrete group and  $\pi$  be a unitary representation of  $\Gamma$  on a Hilbert space  $\mathcal{H}$ . For  $\xi, \eta \in \mathcal{H}$ , we denote the matrix coefficient of  $\pi$  by

$$\pi_{\xi, \eta}(s) = \langle \pi(s)\xi, \eta \rangle.$$

Note that  $\pi_{\xi, \eta} \in \ell_\infty(\Gamma)$ , where  $\ell_\infty(\Gamma)$  is the abelian  $C^*$ -algebra of all bounded functions on  $\Gamma$ .

Let  $D$  be an algebraic two-sided ideal of  $\ell_\infty(\Gamma)$ . If there exists a dense subspace  $\mathcal{H}_0$  of  $\mathcal{H}$  such that  $\pi_{\xi, \eta} \in D$  for all  $\xi, \eta \in \mathcal{H}_0$ , then  $\pi$  is called  $D$ -representation. If  $D$  is invariant under the left and right translation of  $\Gamma$  on  $\ell_\infty(\Gamma)$ , then it is said to be translation invariant.

Throughout this paper, we assume that  $D$  is a non-zero translation invariant ideal of  $\ell_\infty(\Gamma)$ . For each  $p \in [1, \infty)$ , we denote the norm on  $\ell_p(\Gamma)$  by

$$|f|_p = \left( \sum_{s \in \Gamma} |f(s)|^p \right)^{\frac{1}{p}} \quad \text{for } f \in \ell_p(\Gamma).$$

Note that  $\ell_p(\Gamma)$  is a translation invariant ideal of  $\ell_\infty(\Gamma)$ . We denote by  $c_0(\Gamma)$  the of functions on  $\Gamma$ , vanishing at infinity. It is the non-trivial closed translation invariant ideal of  $\ell_\infty(\Gamma)$ .

Under our assumption,  $D$  contains  $c_c(\Gamma)$ , which is the ideal of all finitely supported functions on  $\Gamma$ . Moreover, if  $\pi$  has a cyclic vector  $\xi$  such that  $\pi_{\xi, \xi} \in D$ , then  $\pi$  is a  $D$ -representation with respect to a dense subspace

$$\mathcal{H}_0 = \text{span}\{\pi(s)\xi : s \in \Gamma\}.$$

We denote by  $\lambda$  the left regular representation of  $\Gamma$ . It is easy to see that  $\lambda$  is a  $c_c$ -representation, or a  $D$ -representation for any  $D$ .

The  $C^*$ -algebra  $C_D^*(\Gamma)$  is the  $C^*$ -completion of the group ring  $\mathbb{C}\Gamma$  by  $\|\cdot\|_D$ , where

$$\|f\|_D = \sup\{\|\pi(f)\| : \pi \text{ is a } D\text{-representation}\} \quad \text{for } f \in \mathbb{C}\Gamma.$$

Note that if  $D_1$  and  $D_2$  are ideals of  $\ell_\infty(\Gamma)$  with  $D_1 \supset D_2$ , then there exists the canonical quotient map from  $C_{D_1}^*(\Gamma)$  onto  $C_{D_2}^*(\Gamma)$ . We denote by  $C^*(\Gamma)$  the full group  $C^*$ -algebra, and by  $C_\lambda^*(\Gamma)$  the reduced group  $C^*$ -algebra, respectively. In [2], the following results are obtained:

- $C^*(\Gamma) = C_{\ell_\infty}^*(\Gamma)$  and  $C_\lambda^*(\Gamma) = C_{c_c}^*(\Gamma)$ .
- $C_{\ell_p}^*(\Gamma) = C_\lambda^*(\Gamma)$  for every  $p \in [1, 2]$ .
- $C^*(\Gamma) = C_D^*(\Gamma)$  if and only if there exists a sequence  $(h_n)$  of positive definite functions in  $D$  such that  $h_n \rightarrow 1$ .
- If  $C^*(\Gamma) = C_{\ell_p}^*(\Gamma)$  for some  $p \in [1, \infty)$ , then  $\Gamma$  is amenable.
- $\Gamma$  has the Haagerup property if and only if  $C^*(\Gamma) = C_{c_0}^*(\Gamma)$ .

## 3. POSITIVE DEFINITE FUNCTIONS ON A FREE GROUP

Let  $\mathbb{F}_d$  be the free group on finitely many generators  $a_1, \dots, a_d$  with  $d \geq 2$ . We denote by  $|s|$  the word length of  $s \in \mathbb{F}_d$  with respect to the canonical generating set  $\{a_1, a_1^{-1}, \dots, a_d, a_d^{-1}\}$ . For  $k \geq 0$ , we put

$$W_k = \{s \in \mathbb{F}_d : |s| = k\}.$$

We denote by  $\chi_k$  the characteristic function for  $W_k$ .

In the following lemma, the case where  $q = 2$  is given by Haagerup in [5, Lemma 1.3]. His proof also works for  $q \in [1, 2]$  by using Hölder's inequality, instead of Cauchy-Schwarz inequality. We remark that this is also appeared in [1].

**Lemma 3.1.** *Let  $q \in [1, 2]$ . Let  $k, \ell$  and  $m$  be non negative integers. Let  $f$  and  $g$  be functions on  $\mathbb{F}_d$  such that  $\text{supp}(f) \subset W_k$  and  $\text{supp}(g) \subset W_\ell$ , respectively. If  $|k - \ell| \leq m \leq k + \ell$  and  $k + \ell - m$  is even, then*

$$|(f * g)\chi_m|_q \leq |f|_q |g|_q,$$

and if  $m$  is any other value, then

$$|(f * g)\chi_m|_q = 0.$$

*Proof.* It is shown by an argument similar as in [5, Lemma 1.3]. However for convenience, we give the complete proof.

Note that

$$(f * g)(s) = \sum_{\substack{t, u \in \mathbb{F}_d \\ s = tu}} f(t)g(u) = \sum_{\substack{|t|=k \\ |u|=\ell \\ s=tu}} f(t)g(u).$$

Since the possible values of  $|tu|$  are  $|k - \ell|, |k - \ell| + 2, \dots, k + \ell$ , we have

$$|(f * g)\chi_m|_q = 0$$

for any other values of  $m$ .

The case where  $q = 1$  is trivial. So we consider the case where  $q \neq 1$ .

First we assume that  $m = k + \ell$ . In this case, if  $|s| = m$ , then  $s$  can be uniquely written as a product  $tu$  with  $|t| = k$  and  $|u| = \ell$ . Hence

$$(f * g)(s) = f(t)g(u).$$

Therefore

$$|(f * g)\chi_m|_q^q = \sum_{\substack{|tu|=k+\ell \\ |t|=k \\ |u|=\ell}} |f(t)|^q |g(u)|^q \leq \sum_{\substack{|t|=k \\ |u|=\ell}} |f(t)|^q |g(u)|^q = |f|_q^q |g|_q^q.$$

Next we assume that  $m = |k - \ell|, |k - \ell| + 2, \dots, k + \ell - 2$ . In these cases, we have  $m = k + \ell - 2j$  for  $1 \leq j \leq \min\{k, \ell\}$ . Let  $s = tu$  with  $|s| = m$ ,  $|t| = k$  and  $|u| = \ell$ . Then  $s$  can be uniquely written as a product  $t'u'$  such that  $t = t'v$ ,  $u = v^{-1}u'$  with  $|t'| = k - j$ ,  $|u'| = \ell - j$  and  $|v| = |v^{-1}| = j$ . We define

$$f'(t) = \left( \sum_{|v|=j} |f(tv)|^q \right)^{\frac{1}{q}} \text{ if } |t| = k - j, \text{ and } f'(t) = 0 \text{ otherwise.}$$

We also define

$$g'(u) = \left( \sum_{|v|=j} |g(v^{-1}u)|^q \right)^{\frac{1}{q}} \text{ if } |u| = \ell - j, \text{ and } g'(u) = 0 \text{ otherwise.}$$

Note that  $\text{supp}(f') \subset W_{k-j}$  and  $\text{supp}(g') \subset W_{\ell-j}$ . Moreover

$$|f'|_q^q = \sum_{|t|=k-j} \left( \sum_{|v|=j} |f(tv)|^q \right) = |f|_q^q,$$

and similarly  $|g'|_q = |g|_q$ . Take a real number  $p$  with  $1/p + 1/q = 1$ . Since  $1 < q \leq 2$ , we have  $2 \leq p < \infty$ . In particular,  $q \leq p$ . Thanks to Hölder's inequality,

$$\begin{aligned} |(f * g)(s)| &= \left| \sum_{\substack{|t|=k \\ |u|=\ell \\ s=tu}} f(t)g(u) \right| \\ &= \left| \sum_{|v|=j} f(t'v)g(v^{-1}u') \right| \\ &\leq \left( \sum_{|v|=j} |f(t'v)|^q \right)^{\frac{1}{q}} \left( \sum_{|v|=j} |g(v^{-1}u')|^p \right)^{\frac{1}{p}} \\ &\leq \left( \sum_{|v|=j} |f(t'v)|^q \right)^{\frac{1}{q}} \left( \sum_{|v|=j} |g(v^{-1}u')|^q \right)^{\frac{1}{q}} \\ &= f'(t')g'(u') \\ &= (f' * g')(s). \end{aligned}$$

Hence  $|(f * g)\chi_m| \leq (f' * g')\chi_m$ . Since  $(k - j) + (\ell - j) = m$ , it follows from the first part of the proof that

$$|(f * g)\chi_m|_q \leq |(f' * g')\chi_m|_q \leq |f'|_q |g'|_q = |f|_q |g|_q.$$

□

In the following lemma, the case where  $p = q = 2$  is given in the proof of [4, Theorem 1].

**Lemma 3.2.** *Let  $1 \leq q \leq p \leq \infty$  with  $1/p + 1/q = 1$ . Let  $\pi$  be a unitary representation of  $\Gamma$  on a Hilbert space  $\mathcal{H}$  with a cyclic vector  $\iota$  such that  $\pi_{\xi, \xi} \in \ell_p(\Gamma)$ . Then*

$$\|\pi(f)\| \leq \liminf_{n \rightarrow \infty} \left| (f^* * f)^{(*2n)} \right|_q^{\frac{1}{4n}}$$

for  $f \in c_c(\Gamma)$ .

*Proof.* Let  $f \in c_c(\Gamma)$ . In the proof of [4, Theorem 1], the following formula is obtained:

$$\|\pi(f)\| = \sup_{g \in c_c(\Gamma)} \lim_{n \rightarrow \infty} \left( \sum_{s \in \Gamma} (f^* * f)^{(*2n)}(s) \langle \pi(s)\pi(g)\xi, \pi(g)\xi \rangle \right)^{\frac{1}{4n}}.$$

Fix  $g \in c_c(\Gamma)$  and we put  $\varphi(s) = \langle \pi(s)\pi(g)\xi, \pi(g)\xi \rangle$ . Note that

$$\varphi(s) = \langle \pi(s)\pi(g)\xi, \pi(g)\xi \rangle = \sum_{t, u \in \mathbb{F}_d} \overline{g(u)} \pi_{\xi, \xi}(u^{-1}st)g(t) = (\overline{g} * \pi_{\xi, \xi} * g^\vee)(s),$$

where  $g^\vee(s) = g(s^{-1})$ . Consequently,  $\pi_{\xi, \xi} \in \ell_p(\Gamma)$  implies  $\varphi \in \ell_p(\Gamma)$ . Then by Hölder's inequality,

$$\left| \sum_{s \in \Gamma} (f^* * f)^{(*2n)}(s) \varphi(s) \right| \leq \left| (f^* * f)^{(*2n)} \right|_q |\varphi|_p.$$

Therefore it follows that

$$\|\pi(f)\| \leq \liminf_{n \rightarrow \infty} \left| (f^* * f)^{(*2n)} \right|_q^{\frac{1}{4n}}.$$

□

By combining Lemma 3.1 and Lemma 3.2, we can prove the following.

**Lemma 3.3.** *Let  $k$  be a non negative integer. Let  $1 \leq q \leq p \leq \infty$  with  $1/p + 1/q = 1$ . If a unitary representation  $\pi$  of  $\mathbb{F}_d$  on a Hilbert space  $\mathcal{H}$  has a cyclic vector  $\xi$  such that  $\pi_{\xi, \xi} \in \ell_p(\mathbb{F}_d)$ , then*

$$\|\pi(f)\| \leq (k+1)|f|_q.$$

for  $f \in c_c(\mathbb{F}_d)$  with  $\text{supp}(f) \subset W_k$ .

*Proof.* The case where  $q = 1$  and  $p = \infty$  is trivial. So we may assume that  $1 < q \leq 2$  and  $2 \leq p < \infty$  with  $1/p + 1/q = 1$ . It is also shown by an argument similar as in [5, Lemma 1.4].

We consider the norm  $\left| (f^* * f)^{(*2n)} \right|_q$ . Write  $f_{2j-1} = f^*$  and  $f_{2j} = f$  for  $j = 1, 2, \dots, 2n$ . Then

$$(f^* * f)^{(*2n)} = f_1 * f_2 * \dots * f_{4n}.$$

We also denote  $g = f_2 * \dots * f_{4n}$ . So we have

$$(f^* * f)^{(*2n)} = f_1 * g.$$

Note that  $\text{supp}(f_j) \subset W_k$  for  $j = 1, 2, \dots, 4n$  and  $g \in c_c(\mathbb{F}_d)$ . Put  $g_\ell = g\chi_\ell$ . Then  $\text{supp}(g_\ell) \subset W_\ell$  and

$$|g|_q^q = \sum_{\ell=0}^{\infty} |g_\ell|_q^q.$$

Here, remark that  $|g_\ell|_q = 0$  for all but finitely many  $\ell$ . Moreover set

$$h = f_1 * g = \sum_{\ell=0}^{\infty} f_1 * g_\ell$$

and  $h_m = h\chi_m$ . Then  $h \in c_c(\mathbb{F}_d)$  and

$$|h|_q^q = \sum_{m=0}^{\infty} |h_m|_q^q.$$

Here, notice that  $|h_m|_q = 0$  for all but finitely many  $m$ . By Lemma 3.1,

$$|(f_1 * g_\ell)\chi_m|_q \leq |f_1|_q |g_\ell|_q$$

in the case where  $|k - \ell| \leq m \leq k + \ell$  and  $k + \ell - m$  is even. We also have

$$|(f_1 * g_\ell)\chi_m|_q = 0$$

for any other values of  $m$ . Hence

$$\begin{aligned} |h_m|_q &= \left| \sum_{\ell=0}^{\infty} (f_1 * g_\ell)\chi_m \right|_q \\ &\leq \sum_{\ell=0}^{\infty} |(f_1 * g_\ell)\chi_m|_q \\ &= |f_1|_q \sum_{\substack{\ell=|m-k| \\ m+k-\ell \text{ even}}}^{m+k} |g_\ell|_q. \end{aligned}$$

By writing  $\ell = m + k - 2j$ ,

$$\begin{aligned} |h_m|_q &\leq |f_1|_q \sum_{j=0}^{\min\{m,k\}} |g_{m+k-2j}|_q \\ &\leq |f_1|_q \left( \sum_{j=0}^{\min\{m,k\}} |g_{m+k-2j}|_q^q \right)^{\frac{1}{q}} \left( \sum_{j=0}^{\min\{m,k\}} 1^p \right)^{\frac{1}{p}} \\ &\leq (k+1)^{\frac{1}{p}} |f_1|_q \left( \sum_{j=0}^{\min\{m,k\}} |g_{m+k-2j}|_q^q \right)^{\frac{1}{q}}. \end{aligned}$$

Then

$$\begin{aligned} |h|_q^q &= \sum_{m=0}^{\infty} |h_m|_q^q \\ &\leq (k+1)^{\frac{q}{p}} |f_1|_q^q \sum_{m=0}^{\infty} \sum_{j=0}^{\min\{m,k\}} |g_{m+k-2j}|_q^q \\ &= (k+1)^{\frac{q}{p}} |f_1|_q^q \sum_{j=0}^k \sum_{m=j}^{\infty} |g_{m+k-2j}|_q^q \\ &= (k+1)^{\frac{q}{p}} |f_1|_q^q \sum_{j=0}^k \sum_{\ell=k-j}^{\infty} |g_\ell|_q^q \\ &\leq (k+1)^{\frac{q}{p}} |f_1|_q^q \sum_{j=0}^k |g|_q^q \\ &= (k+1)^{\frac{q}{p}+1} |f_1|_q^q |g|_q^q. \end{aligned}$$

Hence  $|f_1 * g|_q \leq (k+1)|f_1|_q |g|_q$ , i.e.,

$$|f_1 * (f_2 * \cdots * f_{4n})|_q \leq (k+1)|f_1|_q |f_2 * \cdots * f_{4n}|_q.$$

Moreover, we inductively get

$$|(f^* * f)^{(*2n)}|_q \leq (k+1)^{4n-1} |f|_q^{4n}.$$

Therefore it follows from Lemma 3.2 that

$$\|\pi(f)\| \leq \liminf_{n \rightarrow \infty} |(f^* * f)^{(*2n)}|_q^{\frac{1}{4n}} \leq (k+1) |f|_q.$$

□

For a function  $\varphi$  on  $\Gamma$ , we denote the corresponding linear functional on  $c_c(\Gamma)$  by

$$\omega_\varphi(f) = \sum_{s \in \Gamma} f(s) \varphi(s) \quad \text{for } f \in c_c(\Gamma).$$

Note that  $\varphi$  is positive definite if and only if the functional  $\omega_\varphi$  extends to a positive linear functional on  $C^*(\Gamma)$ , (see, e.g., [3, Theorem 2.5.11]).

Now we can give a characterization of positive definite functions on  $\mathbb{F}_d$ , which can be extended to the positive linear functionals on  $C_{\ell_p}^*(\mathbb{F}_d)$  for any  $p \in [2, \infty)$ . The case of  $C_\lambda^*(\mathbb{F}_d)$  is given in [5, Theorem 3.1]. We remind the reader that  $C_\lambda^*(\mathbb{F}_d) = C_{\ell_2}^*(\mathbb{F}_d)$ . Hence the following theorem is a generalization to the case of  $C_{\ell_p}^*(\mathbb{F}_d)$  for any  $p \in [2, \infty)$ .

For  $0 < \alpha < 1$ , we set  $\varphi_\alpha(s) = \alpha^{|s|}$ , and it is positive definite on  $\mathbb{F}_d$  by [5, Lemma 1.2].

**Theorem 3.4.** *Let  $2 \leq p < \infty$ . Let  $\varphi$  be a positive definite function on  $\mathbb{F}_d$ . Then the following conditions are equivalent:*

- (1)  $\varphi$  can be extended to the positive linear functional on  $C_{\ell_p}^*(\mathbb{F}_d)$ .
- (2)  $\sup_k |\varphi \chi_k|_p (k+1)^{-1} < \infty$ .
- (3) The function  $s \mapsto \varphi(s)(1+|s|)^{-1-\frac{2}{p}}$  belongs to  $\ell_p(\mathbb{F}_d)$ .
- (4) For any  $\alpha \in (0, 1)$ , the function  $s \mapsto \varphi(s)\alpha^{|s|}$  belongs to  $\ell_p(\mathbb{F}_d)$ .

*Proof.* Without loss of generality, we may assume that  $\varphi(e) = 1$ . The proof is based on the one in [5, Theorem 3.1].

(1) $\Rightarrow$ (2): It follows from (1) that  $\omega_\varphi$  extends to the state on  $C_{\ell_p}^*(\mathbb{F}_d)$ . Hence for  $f \in c_c(\mathbb{F}_d)$ , we have

$$|\omega_\varphi(f)| \leq \|f\|_{\ell_p}$$

Put

$$f = |\varphi|^{p-2} \overline{\varphi} \chi_k.$$

Then

$$|\omega_\varphi(f)| = |\varphi \chi_k|_p^p.$$

Notice that

$$\|f\|_{\ell_p} = \sup\{\|\pi(f)\| : \pi \text{ is an } \ell_p\text{-representation}\}.$$

Let  $\pi$  be an  $\ell_p$ -representation of  $\mathbb{F}_d$  on a Hilbert space  $\mathcal{H}$  with a dense subspace  $\mathcal{H}_0$ . Then

$$\|\pi(f)\|^2 = \sup_{\substack{\xi \in \mathcal{H}_0 \\ \|\xi\|=1}} \langle \pi(f^* * f) \xi, \xi \rangle_{\mathcal{H}}.$$

Fix  $\xi \in \mathcal{H}_0$  with  $\|\xi\| = 1$ . We denote by  $\sigma$  the restriction of  $\pi$  onto the subspace

$$\mathcal{H}_\sigma = \overline{\text{span}}\{\pi(s)\xi : s \in \mathbb{F}_d\} \subset \mathcal{H}.$$

Then

$$\langle \pi(f^* * f)\xi, \xi \rangle_{\mathcal{H}} = \langle \sigma(f^* * f)\xi, \xi \rangle_{\mathcal{H}_\sigma}.$$

Note that  $\xi$  is cyclic for  $\sigma$  such that  $\sigma_{\xi, \xi} \in \ell_p(\mathbb{F}_d)$ . Take a real number  $q$  with  $1/p + 1/q = 1$ . Since  $2 \leq p < \infty$ , we have  $1 < q \leq 2$ . By Lemma 3.3,

$$\|\sigma(f)\| \leq (k+1)|f|_q.$$

Hence

$$\|\sigma(f^* * f)\| = \|\sigma(f)\|^2 \leq (k+1)^2 |f|_q^2.$$

Therefore we obtain

$$\|f\|_{\ell_p}^2 \leq (k+1)^2 |f|_q^2 = (k+1)^2 |\varphi \chi_k|_p^{2(p-1)},$$

namely,

$$\|f\|_{\ell_p} \leq (k+1) |\varphi \chi_k|_p^{p-1}.$$

Consequently,

$$|\varphi \chi_n|_p \leq k+1.$$

(2) $\Rightarrow$ (3):

$$\begin{aligned} \sum_{s \in \mathbb{F}_d} |\varphi(s)|^p (1+|s|)^{-p-2} &= \sum_{k=0}^{\infty} \sum_{|s|=k} |\varphi(s)|^p (1+k)^{-p-2} \\ &= \sum_{k=0}^{\infty} |\varphi \chi_k|_p^p (1+k)^{-p} (1+k)^{-2} \\ &\leq \left\{ \sup_k |\varphi \chi_k|_p (k+1)^{-1} \right\}^p \sum_{k=0}^{\infty} \frac{1}{(k+1)^2} < \infty. \end{aligned}$$

(3) $\Rightarrow$ (4): Easy.

(4) $\Rightarrow$ (1): Note that  $\psi_\alpha(s) = \varphi(s)\alpha^{|s|}$  is also positive definite. By the GNS construction, we obtain the unitary representation  $\sigma_\alpha$  of  $\mathbb{F}_d$  with the cyclic vector  $\xi_\alpha$  such that

$$\psi_\alpha(s) = \langle \sigma_\alpha(s)\xi_\alpha, \xi_\alpha \rangle.$$

Since  $\sigma_\alpha$  is an  $\ell_p$ -representation,  $\psi_\alpha$  can be seen as a state on  $C_{\ell_p}^*(\mathbb{F}_d)$ . By taking the weak-\* limit of  $\psi_\alpha$  as  $\alpha \nearrow 1$ , we conclude that  $\varphi$  can be extended to the state on  $C_{\ell_p}^*(\mathbb{F}_d)$ .  $\square$

**Corollary 3.5.** *Let  $p \in [2, \infty)$  and  $\alpha \in (0, 1)$ . The positive definite function  $\varphi_\alpha$  can be extended to the state on  $C_{\ell_p}^*(\mathbb{F}_d)$  if and only if*

$$\alpha \leq (2d-1)^{-\frac{1}{p}}.$$

*Proof.* Note that

$$\begin{aligned} \varphi_\alpha \in \ell_p(\mathbb{F}_d) &\iff \sum_{k=1}^{\infty} (2d-1)^{k-1} \alpha^{pk} < \infty \\ &\iff (2d-1)\alpha^p < 1 \\ &\iff \alpha < (2d-1)^{-\frac{1}{p}}. \end{aligned}$$

Hence the corollary follows from Theorem 3.4.  $\square$



*Remark 3.6.* Let  $\pi_\alpha$  be the GNS representation of  $\varphi_\alpha$ . Then  $\pi_\alpha$  is weakly contained in  $\lambda$  if and only if  $\alpha \leq (2d-1)^{-\frac{1}{2}}$ . For  $\alpha > (2d-1)^{-\frac{1}{2}}$ , we refer the reader to [7] and [8].

As a consequence, we can obtain the following result. See also [2, Proposition 4.4].

**Corollary 3.7.** *For  $2 \leq q < p \leq \infty$ , the canonical quotient map from  $C_{\ell_p}^*(\mathbb{F}_d)$  onto  $C_{\ell_q}^*(\mathbb{F}_d)$  is not injective.*

*Proof.* It suffices to consider the case where  $p \neq \infty$ , because  $\mathbb{F}_d$  is not amenable.

Suppose that the canonical quotient map from  $C_{\ell_p}^*(\mathbb{F}_d)$  onto  $C_{\ell_q}^*(\mathbb{F}_d)$  is injective for some  $q < p$ . Take a real number  $\alpha$  with

$$(2d-1)^{-\frac{1}{q}} < \alpha \leq (2d-1)^{-\frac{1}{p}}.$$

By using Corollary 3.5,

$$|\omega_{\varphi_\alpha}(f)| \leq \|f\|_{\ell_p} = \|f\|_{\ell_q} \quad \text{for } f \in c_c(\mathbb{F}_d).$$

Therefore it follows that  $\varphi_\alpha$  can be also extended to the state on  $C_{\ell_q}^*(\mathbb{F}_d)$ , but it contradicts to the choice of  $\alpha$ .  $\square$

*Remark 3.8.* The previous result has also shown by N. Higson and N. Ozawa, independently. See also [2, Remark 4.5].

In [6], Powers proves that  $C_\lambda^*(\mathbb{F}_2)$  has a unique tracial state. In [5], Haagerup gives another proof of the uniqueness. Thanks to Theorem 3.4, Haagerup's argument also works for the case of  $C_{\ell_p}^*(\mathbb{F}_d)$ .

**Corollary 3.9.** *For each  $p \in [2, \infty)$ , the  $C^*$ -algebra  $C_{\ell_p}^*(\mathbb{F}_d)$  has a unique tracial state  $\tau \circ \lambda_p$ , where  $\lambda_p$  is the canonical quotient map  $C_{\ell_p}^*(\mathbb{F}_d)$  onto  $C_\lambda^*(\mathbb{F}_d)$  and  $\tau$  is the unique tracial state on  $C_\lambda^*(\mathbb{F}_d)$ .*

*Proof.* Any tracial state on  $C_{\ell_p}^*(\mathbb{F}_d)$  corresponds a positive definite function on  $\mathbb{F}_d$ . Take such a positive definite function  $\varphi$ . Then  $\varphi|_K$  is constant for any conjugacy class  $K$  in  $\mathbb{F}_d$ . Take a conjugacy class  $K$  in  $\mathbb{F}_d$  such that  $K \neq \{e\}$ . Put  $k = \min\{|s| : s \in K\}$ . Then  $|W_{k+2n} \cap K| \geq (2d-1)^{n-1}$  for  $n \geq 1$ . Hence if  $\varphi|_K = C$  for some non-zero constant  $C$ , then

$$\sum_{s \in C} |\varphi(s)|^p \alpha^{p|s|} = \sum_{n=0}^{\infty} \sum_{s \in W_{k+2n} \cap K} C^p \alpha^{pn} \geq C^p \sum_{n=0}^{\infty} (2d-1)^{n-1} \alpha^{pn} = \infty$$

for  $\alpha \geq (2d-1)^{-\frac{1}{p}}$ . This contradicts (4) in Theorem 3.4.  $\square$

We define two algebraic ideals by

$$D_p^-(\Gamma) = \bigcup_{\varepsilon > 0} \ell_{p-\varepsilon}(\Gamma)$$

for  $1 < p \leq \infty$ , and

$$D_p^+(\Gamma) = \bigcap_{\varepsilon > 0} \ell_{p+\varepsilon}(\Gamma)$$

for  $1 \leq p < \infty$ . Note that  $\ell_q(\Gamma) \subsetneq D_p^-(\Gamma) \subsetneq \ell_p(\Gamma)$  for  $1 \leq q < p \leq \infty$ , and  $\ell_p(\Gamma) \subsetneq D_p^+(\Gamma) \subsetneq \ell_q(\Gamma)$  for  $1 \leq p < q \leq \infty$ . Then we also obtain the following.

**Corollary 3.10.** (1) For  $2 \leq p < \infty$ , the  $C^*$ -algebra  $C_{\ell_p}^*(\mathbb{F}_d)$  is canonically isomorphic to  $C_{D_p^+}^*(\mathbb{F}_d)$ . In particular,  $C_\lambda^*(\mathbb{F}_d) = C_{D_2^+}^*(\mathbb{F}_d)$ .  
 (2) For  $2 < p \leq \infty$ , the  $C^*$ -algebra  $C_{\ell_p}^*(\mathbb{F}_d)$  is canonically isomorphic to  $C_{D_p^-}^*(\mathbb{F}_d)$ . In particular,  $C^*(\mathbb{F}_d) = C_{D_\infty^-}^*(\mathbb{F}_d)$ .

*Proof.* (1) It suffices to show that if  $\varphi$  is a positive definite function on  $\mathbb{F}_d$  which can be extended the positive linear functional on  $C_{D_p^+}^*(\mathbb{F}_d)$ , then  $\varphi$  can be also extended to the one on  $C_{\ell_p}^*(\mathbb{F}_d)$ .

Now assume that  $\varphi$  is a positive definite function on  $\mathbb{F}_d$ , which can be extend to the positive linear functional on  $C_{D_p^+}^*(\mathbb{F}_d)$ . Then for any  $q \in (p, \infty)$ ,  $\varphi$  can be also extended to the positive linear functional on  $C_{\ell_q}^*(\mathbb{F}_d)$ . By Theorem 3.4,  $\varphi\varphi_\alpha \in \ell_q(\mathbb{F}_d)$  for any  $\alpha \in (0, 1)$ . We set

$$r = \frac{pq}{q-p}.$$

Then  $1/p = 1/q + 1/r$ . If we take  $(2d-1)^{-2/r} < \beta < (2d-1)^{-1/r}$ , then  $\varphi_\beta \in \ell_r(\mathbb{F}_d)$ . Since

$$|\varphi\varphi_\alpha\varphi_\beta|_p \leq |\varphi\varphi_\alpha|_q |\varphi_\beta|_r,$$

we have  $\varphi\varphi_\alpha\varphi_\beta \in \ell_p(\mathbb{F}_d)$ . Namely,  $(\varphi\varphi_\beta)\varphi_\alpha \in \ell_p(\mathbb{F}_d)$  for any  $\alpha \in (0, 1)$ . Thus  $\varphi\varphi_\beta$  can be extended to the positive linear functional on  $C_{\ell_p}^*(\mathbb{F}_d)$ . If  $q \searrow p$ , then  $r \nearrow \infty$  and  $\beta \nearrow 1$ . Hence  $\varphi\varphi_\beta \rightarrow \varphi$  in the weak-\* topology. Therefore  $\varphi$  can be also extended to the positive linear functional on  $C_{\ell_p}^*(\mathbb{F}_d)$ .

(2) The proof is quite similar as in (1). Assume that  $\varphi$  is a positive definite function on  $\mathbb{F}_d$  which can be extended to the positive linear functional on  $C_{\ell_p}^*(\mathbb{F}_d)$ . By Theorem 3.4, we have  $\varphi\varphi_\alpha \in \ell_p(\mathbb{F}_d)$  for any  $\alpha \in (0, 1)$ . For any  $q \in [2, p)$ , we set

$$r = \frac{pq}{p-q}.$$

Then  $1/q = 1/p + 1/r$ . If we take  $(2d-1)^{-2/r} < \beta < (2d-1)^{-1/r}$ , then  $\varphi_\beta \in \ell_r(\mathbb{F}_d)$ . Since

$$|\varphi\varphi_\alpha\varphi_\beta|_q \leq |\varphi\varphi_\alpha|_p |\varphi_\beta|_r,$$

we have  $\varphi\varphi_\alpha\varphi_\beta \in \ell_q(\mathbb{F}_d)$ . Namely,  $(\varphi\varphi_\beta)\varphi_\alpha \in \ell_q(\mathbb{F}_d)$  for any  $\alpha \in (0, 1)$ . Thus  $\varphi\varphi_\beta$  can be extended to the positive linear functional on  $C_{\ell_q}^*(\mathbb{F}_d)$ , and so  $\varphi\varphi_\beta$  can be extended to the one on  $C_{D_p^-}^*(\mathbb{F}_d)$ . If  $q \nearrow p$ , then  $r \nearrow \infty$  and  $\beta \nearrow 1$ . Hence  $\varphi\varphi_\beta \rightarrow \varphi$  in the weak-\* topology. Therefore  $\varphi$  can be also extended to the positive linear functional on  $C_{D_p^-}^*(\mathbb{F}_d)$ .  $\square$

*Remark 3.11.* The isomorphism  $C_\lambda^*(\Gamma) = C_{D_2^+}^*(\Gamma)$  has been already obtained in [2] for any  $\Gamma$ .

## REFERENCES

1. M. Bożejko; *Remark on Herz-Schur multipliers on free groups*. Math. Ann. **258** (1981/82), no. 1, 11–15.
2. N. P. Brown and E. Guentner; *New  $C^*$ -completions of discrete groups and related spaces*. Preprint. <http://arxiv.org/abs/1205.4649>.
3. N. P. Brown and N. Ozawa;  *$C^*$ -algebras and finite-dimensional approximations*. Graduate Studies in Mathematics, **88**. American Mathematical Society, Providence, RI, 2008. xvi+509
4. M. Cowling, U. Haagerup and R. Howe; *Almost  $L^2$  matrix coefficients*. J. Reine Angew. Math. **387** (1988), 97–110.

5. U. Haagerup; *An example of a nonnuclear  $C^*$ -algebra, which has the metric approximation property.* Invent. Math. **50** (1978/79), no. 3, 279–293.
6. R. T. Powers; *Simplicity of the  $C^*$ -algebra associated with the free group on two generators.* Duke Math. J. **42** (1975), 151–156.
7. T. Pytlik and R. Szwarc; *An analytic family of uniformly bounded representations of free groups.* Acta Math. **157** (1986), no. 3-4, 287–309.
8. R. Szwarc; *An analytic series of irreducible representations of the free group.* Ann. Inst. Fourier (Grenoble) **38** (1988), no. 1, 87–110.

DEPARTMENT OF MATHEMATICS EDUCATION, OSAKA KYOIKU UNIVERSITY, KASHIWARA, OSAKA  
582-8582, JAPAN

*E-mail address:* `rui@cc.osaka-kyoiku.ac.jp`