*Bernoulli* **20**(1), 2014, 231–244 DOI: 10.3150/12-BEJ483

# Chernoff's density is log-concave

FADOUA BALABDAOUI<sup>1</sup> and JON A. WELLNER<sup>2</sup>

 <sup>1</sup>Centre de Recherche en Mathématiques de la Décision, Université Paris-Dauphine, Paris, France. E-mail: fadoua@ceremade.dauphine.fr
 <sup>2</sup>Department of Statistics, University of Washington, Seattle, WA 98195-4322, USA. E-mail: jaw@stat.washington.edu

We show that the density of  $Z = \operatorname{argmax}\{W(t) - t^2\}$ , sometimes known as Chernoff's density, is log-concave. We conjecture that Chernoff's density is strongly log-concave or "super-Gaussian", and provide evidence in support of the conjecture.

*Keywords:* airy function; Brownian motion; correlation inequalities; hyperbolically monotone; log-concave; monotone function estimation; Polya frequency function; Prekopa–Leindler theorem; Schoenberg's theorem; slope process; strongly log-concave

## 1. Introduction: Two limit theorems

We begin by comparing two limit theorems.

First the usual central limit theorem: Suppose that  $X_1, \ldots, X_n$  are i.i.d.  $EX_1 = \mu$ ,  $E(X^2) < \infty$ ,  $\sigma^2 = \operatorname{Var}(X)$ . Then, the classical Central Limit theorem says that

$$\sqrt{n}(\overline{X}_n - \mu) \to_d N(0, \sigma^2).$$

The Gaussian limit has density

$$\phi_{\sigma}(x) = \frac{1}{\sqrt{2\pi\sigma}} \exp\left(-\frac{x^2}{2\sigma^2}\right) = e^{-V(x)},$$
$$V(x) = -\log\phi_{\sigma}(x) = \frac{x^2}{2\sigma^2} + \log(\sqrt{2\pi\sigma}),$$
$$V''(x) = (-\log\phi_{\sigma})''(x) = \frac{1}{\sigma^2} > 0.$$

Thus,  $\log \phi_{\sigma}$  is concave, and hence  $\phi_{\sigma}$  is a *log-concave density*. As is well known, the normal distribution arises as a natural limit in a wide range of settings connected with sums of independent and weakly dependent random variables; see, for example, Le Cam [30] and Dehling and Philipp [11].

This is an electronic reprint of the original article published by the ISI/BS in *Bernoulli*, 2014, Vol. 20, No. 1, 231–244. This reprint differs from the original in pagination and typographic detail.

1350-7265 © 2014 ISI/BS

Now for a much less well-known limit theorem in the setting of monotone regression. Suppose that the real-valued function r(x) is monotone increasing for  $x \in [0, 1]$ . For  $i \in \{1, ..., n\}$ , suppose that  $x_i = i/(n+1)$ ,  $\varepsilon_i$  are i.i.d. with  $E(\varepsilon_i) = 0$ ,  $\sigma^2 = E(\varepsilon_i^2) < \infty$ , and suppose that we observe  $(x_i, Y_i)$ , i = 1, ..., n, where

$$Y_i = r(x_i) + \varepsilon_i \equiv \mu_i + \varepsilon_i, \qquad i \in \{1, \dots, n\}.$$

The isotonic estimator  $\underline{\widehat{\mu}}$  of  $\underline{\mu} = (\mu_1, \dots, \mu_n)$  is given by

$$\begin{aligned} \widehat{\mu}_{j} &= \max_{i \leq j} \min_{k \geq j} \left\{ \frac{\sum_{l=i}^{k} Y_{l}}{k - i + 1} \right\}, \\ \underline{\widehat{\mu}} &= (\widehat{\mu}_{1}, \dots, \widehat{\mu}_{n}) \equiv T \underline{Y} \\ &= \text{least squares projection of } \underline{Y} \text{ onto } K_{n}, \end{aligned}$$

$$K_n = \{ y \in \mathbb{R}^n \colon y_1 \le \dots \le y_n \}.$$

For fixed  $x_0 \in (0,1)$  with  $x_j \leq x_0 < x_{j+1}$  we set  $\hat{r}_n(x_0) \equiv \hat{r}_n(x_j) = \hat{\mu}_j$ .

Brunk [7] showed that if  $r'(x_0) > 0$  and if r' is continuous in a neighborhood of  $x_0$ , then

$$n^{1/3}(\widehat{r}_n(x_0) - r(x_0)) \to_d (\sigma^2 r'(x_0)/2)^{1/3}(2Z_1),$$

where, with  $\{W(t): t \in \mathbb{R}\}$  denoting a two-sided standard Brownian motion process started at 0,

 $2Z_1$  = slope at zero of the greatest convex minorant of  $W(t) + t^2$ 

 $\stackrel{d}{=}$  slope at zero of the least concave majorant of  $W(t) - t^2$  (1.1)

 $\stackrel{d}{=} 2 \operatorname{argmin}\{W(t) + t^2\}.$ 

The density f of  $Z_1$  is called Chernoff's density. Chernoff's density appears in a number of nonparametric problems involving estimation of a monotone function:

- Estimation of a monotone regression function r: see, for example, Ayer et al. [1], van Eeden [44], Brunk [7], and Leurgans [31].
- Estimation of a monotone decreasing density: see Grenander [12], Prakasa Rao [37], and Groeneboom [14].
- Estimation of a monotone hazard function: Grenander [13], Prakasa Rao [38], Huang and Zhang [26], Huang and Wellner [25].
- Estimation of a distribution function with interval censoring: Groeneboom and Wellner [21], Groeneboom [16].

In each case:

- There is a monotone function m to be estimated.
- There is a natural nonparametric estimator  $\widehat{m}_n$ .

• If  $m'(x_0) \neq 0$  and m' continuous at  $x_0$ , then

$$n^{1/3}(\widehat{m}_n(x_0) - m(x_0)) \to_d C(m, x_0) 2Z_1,$$

where  $2Z_1$  is as in (1.1).

See Kim and Pollard [29] for a unified approach to these types of problems.

The first appearance of  $Z_1$  was in Chernoff [9]. Chernoff [9] considered estimation of the mode of a (symmetric unimodal) density f via the following simple estimator: if  $X_1, \ldots, X_n$  are i.i.d. with density h and distribution function H, then for each fixed a > 0let

 $\hat{x}_a \equiv \text{center of the interval of length } 2a \text{ containing the most observations.}$ 

Let  $x_a$  be the center of the interval of length 2a maximizing  $H(x + a) - H(x - a) = P(X \in (x - a, x + a])$ . (Note that this  $x_a$  is not the mode if f is not symmetric.) Then Chernoff shows:

$$n^{1/3}(\hat{x}_a - x_a) \to_d \left(\frac{h(x_a + a)}{c}\right)^{1/3} 2Z_1,$$

where  $c \equiv h'(x_a - a) - h'(x_a + a)$ . Chernoff also showed that the density  $f_{Z_1} = f$  of  $Z_1$  has the form

$$f(z) \equiv f_{Z_1}(z) = \frac{1}{2}g(z)g(-z), \tag{1.2}$$

where

$$g(t) \equiv \lim_{x \nearrow t^2} \frac{\partial}{\partial x} u(t,x),$$

where, with W standard Brownian motion,

$$u(t,x) \equiv P^{(t,x)}(W(z) > z^2, \text{for some } z \ge t)$$

is a solution to the backward heat equation

$$\frac{\partial}{\partial t}u(t,x)=-\frac{1}{2}\frac{\partial^2}{\partial x^2}u(t,x)$$

under the boundary conditions

$$u(t,t^2) = \lim_{x \not \to t^2} u(t,x) = 1, \qquad \lim_{x \to -\infty} u(t,x) = 0.$$

Again let W(t) be standard two-sided Brownian motion starting from zero, and let c > 0. We now define

$$Z_c \equiv \sup\{t \in \mathbb{R}: \ W(t) - ct^2 \text{ is maximal}\}.$$
(1.3)

As noted above,  $Z_c$  with c = 1 arises naturally in the limit theory for nonparametric estimation of monotone (decreasing) functions. Groeneboom [15] (see also Daniels and Skyrme [10]) showed that for all c > 0 the random variable  $Z_c$  has density

$$f_{Z_c}(t) = \frac{1}{2}g_c(t)g_c(-t),$$

where  $g_c$  has Fourier transform given by

$$\hat{g}_c(\lambda) = \int_{-\infty}^{\infty} e^{i\lambda s} g_c(s) \, \mathrm{d}s = \frac{2^{1/3} c^{-1/3}}{\mathrm{Ai}(\mathrm{i}(2c^2)^{-1/3}\lambda)}.$$
(1.4)

Groeneboom and Wellner [22] gave numerical computations of the density  $f_{Z_1}$ , distribution function, quantiles, and moments.

Recent work on the distribution of the supremum  $M_c \equiv \sup_{t \in \mathbb{R}} (W(t) - ct^2)$  is given in Janson, Louchard and Martin-Löf [27] and Groeneboom [17]. Groeneboom [18] studies the number of vertices of the greatest convex minorant of  $W(t) + t^2$  in intervals [a, b] with  $b - a \to \infty$ ; the function  $g_c$  with c = 1 also plays a key role there.

Our goal in this paper is to show that the density  $f_{Z_c}$  is log-concave. We also present evidence in support of the conjecture that  $f_{Z_c}$  is strongly log-concave: that is,  $(-\log f_{Z_c})''(t) \ge \text{some } c > 0$  for all  $t \in \mathbb{R}$ .

The organization of the rest of the paper is as follows: log-concavity of  $f_{Z_c}$  is proved in Section 2 where we also give graphical support for this property and present several corollaries and related results. In Section 3, we give some partial results and further graphical evidence for strong log-concavity of  $f \equiv f_{Z_1}$ : that is,

$$(-\log f)''(t) \ge (-\log f)''(0) = 3.4052... = 1/(0.541912...)^2 \equiv 1/\sigma_0^2$$

for all  $t \in \mathbb{R}$ . As will be shown in Section 3, this is equivalent to  $f(t) = \rho(t)\phi_{\sigma_0}(t)$  with  $\rho$  log-concave. In Section 4, we briefly discuss some of the consequences and corollaries of log-concavity and strong log-concavity, sketch connections to some results of Bondesson [5, 6], and list a few of the many further problems.

## 2. Chernoff's density is log-concave

Recall that a function h is a Pólya frequency function of order m (and we write  $h \in PF_m$ ) if  $K(x,y) \equiv h(x-y)$  is totally positive of order m: that is,  $\det(H_m(\underline{x},\underline{y}) \geq 0$  for all choices of  $x_1 \leq \cdots \leq x_m$  and  $y_1 \leq \cdots \leq y_m$  where  $H_m \equiv H_m(\underline{x},\underline{y}) = (h(x_i - y_j))_{i,j=1}^m$ . It is well known and easily proved that a density f is PF<sub>2</sub> if and only if it is log-concave. Furthermore, h is a Pólya frequency function (and we write  $h \in PF_\infty$ ) if  $K(x,y) \equiv h(x - y)$  is totally positive of all orders m; see, for example, Schoenberg [42], Karlin [28], and Marshall, Olkin and Arnold [33]. Following Karlin [28], we say that h is strictly PF<sub>∞</sub> if all the determinants  $\det(H_m)$  are strictly positive.

**Theorem 2.1.** For each c > 0 the density  $f_{Z_c}(x) = (1/2)g_c(x)g_c(-x)$  is PF<sub>2</sub>; that is, log-concave.

The Fourier transform in (1.4) implies that  $g_c$  has bilateral Laplace transform (with a slight abuse of notation)

$$\hat{g}_c(z) = \int e^{zs} g_c(s) \, \mathrm{d}s = \frac{2^{1/3} c^{-1/3}}{\operatorname{Ai}((2c^2)^{-1/3} z)}$$
(2.1)

for all z such that  $\operatorname{Re}(z) > -a_1/(2c^2)^{-1/3}$  where  $-a_1$  is the largest zero of  $\operatorname{Ai}(z)$  in  $(-\infty, 0)$ .

To prove Theorem 2.1, we first show that  $g_c$  is  $PF_{\infty}$  by application of the following two results.

**Theorem 2.2 (Schoenberg, 1951).** A necessary and sufficient condition for a (density) function g(x),  $-\infty < x < \infty$ , to be a  $PF_{\infty}$  (density) function is that the reciprocal of its bilateral Laplace transform (i.e., Fourier) be an entire function of the form

$$\psi(s) \equiv \frac{1}{\hat{g}(s)} = C e^{-\gamma s^2 + \delta s} s^k \prod_{j=1}^{\infty} (1 + b_j s) \exp(-b_j s), \qquad (2.2)$$

where C > 0,  $\gamma \ge 0$ ,  $\delta \in \mathbb{R}$ ,  $k \in \{0, 1, 2, ...\}$ ,  $b_j \in \mathbb{R}$ ,  $\sum_{j=1}^{\infty} |b_j|^2 < \infty$ . (For the subclass of densities, the if and only if statement holds for  $1/\hat{g}$  of this form with  $\psi(0) = C = 1$  and k = 0.)


**Proposition 2.1 (Merkes and Salmassi).** Let  $\{-a_k\}$  be the zeros of the Airy function Ai (so that  $a_k > 0$  for each k). The Hadamard representation of Ai is given by

$$\operatorname{Ai}(z) = \operatorname{Ai}(0)e^{-\nu z} \prod_{k=1}^{\infty} (1 + z/a_k) \exp(-z/a_k),$$

where

$$\begin{aligned} \operatorname{Ai}(0) &= \frac{1}{3^{2/3}\Gamma(2/3)} = \frac{\Gamma(1/3)}{3^{1/6}2\pi} \approx 0.35503, \\ \operatorname{Ai}'(0) &= -\frac{1}{3^{1/3}\Gamma(1/3)} = -\frac{3^{1/6}\Gamma(2/3)}{2\pi} \approx -0.25882 \quad and \\ \nu &= -\operatorname{Ai}'(0)/\operatorname{Ai}(0) = \frac{3^{1/3}\Gamma(2/3)}{\Gamma(1/3)} = \frac{2\pi}{3^{1/6}\Gamma(1/3)^2} \approx 0.729011\dots \end{aligned}$$

Proposition 2.1 is given by Merkes and Salmassi [34]; see their Lemma 1, page 211. This is also Lemma 1 of Salmassi [41]. Our statement of Proposition 2.1 corrects the constants  $c_1$  and  $c_2$  given by Merkes and Salmassi [34]. Figure 1 shows Ai(z) (black) and m term approximations to Ai(z) based on Proposition 2.1 with m = 25 (green), 125 (magenta), and 500 (blue).



**Figure 1.** Product approximations of Ai(x).

**Proposition 2.2.** The functions  $t \mapsto g_c(t)$  are in  $\operatorname{PF}_{\infty} \subset \operatorname{PF}_2$  for every c > 0. Thus, they are log-concave. In fact,  $t \mapsto g_c(t)$  is strictly  $\operatorname{PF}_{\infty}$  for every c > 0.

**Proof.** By Proposition 2.1,

$$\operatorname{Ai}((2c^2)^{-1/3}z) = \operatorname{Ai}(0)e^{-\nu(2c^2)^{-1/3}z} \prod_{j=1}^{\infty} \left(1 + \frac{z}{(2c^2)^{1/3}a_j}\right) \exp\left(-\frac{z}{(2c^2)^{1/3}a_j}\right)$$
$$= \operatorname{Ai}(0)e^{\delta z} \prod_{j=1}^{\infty} (1 + b_j z) \exp(-b_j z),$$

which is of the form (2.2) required in Schoenberg's theorem with k = 0,

$$\delta = -(2c^2)^{-1/3}\nu = -\frac{(3/2)^{1/3}\Gamma(2/3)}{c^{2/3}\Gamma(1/3)},$$
(2.3)

$$C = \operatorname{Ai}(0) = 1/(3^{2/3}\Gamma(2/3))$$
 and (2.4)

$$b_j = \frac{1}{(2c^2)^{1/3}a_j}, \qquad j \ge 1,$$
(2.5)

where  $\{-a_j\}$  are the zeros of the Airy function Ai. Thus, we conclude from Schoenberg's theorem that  $g_c$  is  $\mathrm{PF}_{\infty}$  for each c > 0.

The strict  $PF_{\infty}$  property follows from Karlin [28], Theorem 6.1(a), page 357: note that in the notation of Karlin [28],  $\gamma = 0$  and Karlin's  $a_i$  is our  $1/a_k$  with  $\sum_k (1/a_k) = \infty$ in view of the fact that  $a_k \sim ((3/8)\pi(4k-1))^{2/3}$  via 9.9.6 and 9.9.18, page 18, Olver *et al.* [36].

Now we are in position to prove Theorem 2.1.

Proof of Theorem 2.1. This follows from Proposition 2.2: note that

$$-\log f_{Z_c}(x) = -\log g_c(x) - \log g_c(-x),$$

SO

$$w(x) \equiv (-\log f_{Z_c})''(x) = (-\log g_c)''(x) + (-\log g_c)''(-x) \equiv v(x) + v(-x) \ge 0$$
  
$$g_c \in \operatorname{PF}_{\infty} \subset \operatorname{PF}_2.$$

since  $g_c \in \mathrm{PF}_{\infty} \subset \mathrm{PF}_2$ .

Some scaling relations: From the Fourier transform of  $g_c$  given above, it follows that

$$g_c(x) = \frac{(2/c)^{1/3}}{2\pi} \int_{-\infty}^{\infty} \frac{e^{-iux}}{\operatorname{Ai}(\mathrm{i}(2c^2)^{-1/3}u)} \,\mathrm{d}u$$
$$= \frac{(2/c)^{1/3}(2c^2)^{1/3}}{2\pi} \int_{-\infty}^{\infty} \frac{e^{-\mathrm{i}v(2c^2)^{1/3}x}}{\operatorname{Ai}(\mathrm{i}v)} \,\mathrm{d}v$$
$$\equiv 2^{1/6}c^{1/3}g_{2^{-1/2}}((2c^2)^{1/3}x).$$

Thus it follows that

$$(\log g_c)''(x) = (2c^2)^{2/3} \cdot (\log g_{2^{-1/2}})''((2c^2)^{1/3}x),$$

and, in particular,

$$(\log g_c(x))''|_{x=0} = (2c^2)^{2/3} \cdot (\log g_{2^{-1/2}})''(x)|_{x=0}$$

When c = 1, the conversion factor is  $2^{2/3}$ . Furthermore we compute

$$f_{Z_c}(t) = \frac{1}{2}g_c(t)g_c(-t) = \frac{1}{2}2^{1/3}c^{2/3}g_{2^{-1/2}}((2c^2)^{1/3}t)g_{2^{-1/2}}(-(2c^2)^{1/3}t)$$
$$\equiv c^{2/3}f_1(c^{2/3}t),$$

where

$$f_1(t) \equiv f_{Z_1}(t) = \frac{1}{2}g_1(t)g_1(-t)$$
  
=  $\frac{1}{2}2^{1/3}g_{2^{-1/2}}(2^{1/3}t)g_{2^{-1/2}}(-2^{1/3}t)$ 

Thus we see that

 $Z_c \stackrel{d}{=} c^{-2/3} Z_1$ 

for all c > 0.

Figure 2 gives a plot of  $f_Z$ ; Figure 3 gives a plot of  $-\log f_Z$ ; and Figure 4 gives a plot of  $(-\log f_Z)''$ .




Figure 2. The density  $f_Z$ .

If we use the inverse Fourier transform to represent g via (1.4), and then calculate directly, some interesting correlation type inequalities involving the Airy kernel emerge. Here is one of them.

Let  $h(u) \equiv 1/|\operatorname{Ai}(iu)| \sim 2\sqrt{\pi}u^{1/4} \exp(-(\sqrt{2}/3)u^{3/2})$  as  $u \to \infty$  by Groeneboom [15], page 95. We also define  $\varphi(u, x/2) = \operatorname{Re}(e^{iux/2}\operatorname{Ai}(iu))h(u)$  and  $\psi(u, x/2) = \operatorname{Im}(e^{iux/2} \times \operatorname{Ai}(iu))h(u)$ .





Figure 3.  $-\log f_Z$ .

Chernoff's density is log-concave



**Figure 4.**  $(-\log f_Z)''$ .

Corollary 2.1. With the above notation,

$$\int_0^\infty \sin^2(uy)\varphi(u,x)h(u)\,\mathrm{d} u \cdot \int_0^\infty \cos^2(uy)\varphi(u,x)h(u)\,\mathrm{d} u$$
$$+\int_0^\infty \sin(uy)\cos(uy)\psi(u,x)h(u)\,\mathrm{d} u \ge 0 \qquad \text{for all } x,y \in \mathbb{R}.$$

## 3. Is Chernoff's density strongly log-concave?

From Rockafellar and Wets [40] page 565,  $h: \mathbb{R}^d \to \overline{\mathbb{R}}$  is strongly convex if there exists a constant c > 0 such that

$$h(\theta x + (1-\theta)y) \le \theta h(x) + (1-\theta)h(y) - \frac{1}{2}c\theta(1-\theta)\|x-y\|^2$$

for all  $x, y \in \mathbb{R}^d$ ,  $\theta \in (0, 1)$ . It is not hard to show that this is equivalent to convexity of

$$h(x) - \frac{1}{2}c||x||^2$$

for some c > 0. This leads (by replacing h by  $-\log f$ ) to the following definition of *strong* log-concavity of a (density) function:  $f : \mathbb{R}^d \to \overline{\mathbb{R}}$  is strongly log-concave if and only if

$$-\log f(x) - \frac{1}{2}c||x||^2$$

is convex for some c > 0. Defining  $-\log g(x) \equiv -\log f(x) - (1/2)c||x||^2$ , it is easily seen that f is strongly log-concave if and only if

$$f(x) = g(x) \exp(-(1/2)c||x||^2)$$

for some c > 0 and log-concave function g. Thus if  $f \in C^2(\mathbb{R}^d)$ , a sufficient condition for strong log-concavity is:  $\operatorname{Hess}(-\log f)(x) \ge cI_d$  for all  $x \in \mathbb{R}^d$  and some c > 0 where  $I_d$  is the  $d \times d$  identity matrix.

Figure 4 provides compelling evidence for the following conjecture concerning strong log-concavity of Chernoff's density.

**Conjecture 3.1.** Let  $Z_1$  again be a "standard" Chernoff random variable. Then for  $\sigma \geq \sigma_0 \approx 0.541912... = (-(\log f_{Z_1}(z))''|_{z=0})^{-1/2}$  the density  $f_{Z_1}$  can be written as

$$f_{Z_1}(x) = \rho(x) \frac{1}{\sigma} \varphi(x/\sigma),$$

where  $\varphi(x) = (2\pi)^{-1/2} \exp(-x^2/2)$  is the standard normal density and  $\rho$  is log-concave. Equivalently, if  $c \ge \sigma_0^{3/2} \approx 0.398927...$ , then

$$f_{Z_c}(x) = \tilde{\rho}(x)\varphi(x),$$

where  $\tilde{\rho}$  is log-concave.

**Proof.** (Partial) Let  $w \equiv (-\log f_{Z_c})''$  and  $v \equiv (-\log g_c)''$ . Then

$$w(t) = v(t) + v(-t) \ge 2v(0) = w(0) > 0$$

is implied by convexity of v and strict positivity of w(0). Thus, we want to show that  $v^{(2)} = (-\log g_c)^{(4)} \ge 0.$ 

To prove this, we investigate the normalized version of  $g_c$  given by  $\tilde{g}_c(x) = g_c(x) \operatorname{Ai}(0)/(2/c)^{1/3} = g_c(x)/\int g_c(y) \, dy$  so that  $\int \tilde{g}_c(x) \, dx = 1$ . Suppose that  $b_i$  is given in (2.5), and let  $X_i \sim \operatorname{Exp}(1/b_i)$  be independent exponential random variables for  $i = 1, 2, \ldots$  Since  $\sum_{i=1}^{\infty} b_i^2 < \infty$ , the random variable  $Y_0 = \sum_{i=1}^{\infty} (X_i - b_i)$  is finite almost surely (see, e.g., Shorack [43], Theorem 9.2, page 241) and the Laplace transform of  $-(\delta + Y_0)$  is given by

$$\varphi(s) \equiv e^{-\delta s} E e^{-sY_0} = \exp(-\delta s) \cdot \frac{1}{\prod_{i=1}^{\infty} (1+b_i s) e^{-b_i s}}$$
$$= \frac{1}{e^{\delta s} \cdot \prod_{i=1}^{\infty} (1+b_i s) e^{-b_i s}},$$

exactly the form of the Laplace transform of g implicit in the proof of Proposition 2.2, but without the Gaussian term. Thus, we conclude that  $\tilde{g}_c$  is the density of  $Y \equiv -\delta - Y_0 = -\delta - \sum_{i=1}^{\infty} (X_j - b_j)$ .

 $-\delta - \sum_{j=1}^{\infty} (X_j - b_j).$ Now let  $\lambda_i = 1/b_i$  for  $i \ge 1$ . Thus,  $X_i \sim \operatorname{Exp}(\lambda_i)$ . A closed form expression for the density of  $Y_m \equiv \sum_{i=1}^m X_i$  has been given by Harrison [24]. From Harrison's theorem 1,  $Y_m$  has density

$$f_m(t) = \sum_{j=1}^m \lambda_j \exp(-\lambda_j t) \prod_{i \neq j} \frac{\lambda_i}{\lambda_i - \lambda_j}.$$
(3.1)

10

If we could show that  $v_m(t) \equiv (-\log f_m)''(t)$  is convex, then we would be done! Direct calculation shows that this holds for m = 2, but our attempts at a proof for general m have not (yet) been successful. On the other hand, we know that for  $t \ge 0$ ,

$$w(t) = v(t) + v(-t) \ge v(t) \ge v(0) > 0$$

if v satisfies  $v(t) \ge v(0)$  for all  $t \ge 0$ , so we would have strong log-concavity with the constant v(0).

## 4. Discussion and open problems

Log-concavity of Chernoff's density implies that the peakedness results of Proschan [39] and Olkin and Tong [35] apply. See also Marshall and Olkin [32], page 373, and Marshall, Olkin and Arnold [33].

Note that the conclusion of Conjecture 3.1 is exactly the form of the hypothesis of the inequality of Hargé [23] and of Theorem 11, page 559, of Caffarelli [8]; see also Barthe [4], Theorem 2.4, page 1532. Wellner [47] shows that the class of strongly log-concave densities is closed under convolution, so in particular if Conjecture 3.1 holds, then the sum of two independent Chernoff random variables is again strongly log-concave.

Another implication is that a theorem of Caffarelli [8] applies: the transportation map  $T = \nabla \varphi$  is a contraction. In our particular one-dimensional special case, the transportation map T satisfying  $T(X) \stackrel{d}{=} Z$  for  $X \sim N(0,1)$  is just the solution of  $\Phi(z) = F_Z(T(z))$ , or equivalently  $T(z) = F_Z^{-1}(\Phi(z))$ . This function is apparently connected to another question concerning convex ordering of  $F_Z$  and  $\Phi(\cdot)$  in the sense of van Zwet [46]; see also van Zwet [45]: is  $T^{-1}(w) = \Phi^{-1}(F_Z(w))$  convex for w > 0?

As we have seen above, Chernoff's density has the symmetric product form (1.2) where g has Fourier transform given in (1.4). In this case, we know from Section 2 that  $g \in PF_{\infty}$ .

As is shown in the longer technical report version of this paper Balabdaoui and Wellner [3], it follows from the results of Bondesson [5, 6] that the standard normal density  $\phi$  can be written in the same structural form as that of Chernoff's density (1.2); that is:

$$\phi(z) = \frac{1}{2}g(z)g(-z), \tag{4.1}$$

where now

$$g(z) \equiv (2/\pi)^{1/4} \exp(z) \exp\left(\int_0^\infty \log\left(\frac{e^s + 1}{e^s + e^z}\right) ds\right)$$
  
=  $(2/\pi)^{1/4} \exp\left(\frac{\pi^2}{12} + z - \int_0^{e^z} \frac{\log(1+t)}{t} dt\right)$  (4.2)

is log-concave, integrable, and  $g \in \log(HM_{\infty})$ , the log-transform (in terms of random variables) of the Hyperbolically Completely Monotone class of Bondesson [5, 6]. Two natural questions are: (a) Does the function g in (1.2) satisfy  $g \in \log(HM_{\infty})$ ? (b) Does the function g in (4.2) satisfy  $g \in PF_{\infty}$ ?

A further question remaining from Section 3: Is Chernoff's density strongly logconcave?

A whole class of further problems involves replacing the (ordered) convex cone  $K_n$  in Section 1 by the convex cone  $\tilde{\mathcal{K}}_n$  corresponding to a convexity restriction as in Section 2 of Groeneboom, Jongbloed and Wellner [20]. In this latter case, the limiting distribution depends on an "invelope" of the integral of a two-sided Brownian motion plus a polynomial drift as follows: it is the density of the second derivative at zero of the "invelope". See Groeneboom, Jongbloed and Wellner [19, 20] for further details and Balabdaoui, Rufibach and Wellner [2] for another convexity related shape constraint where this limiting distribution occurs. However, virtually nothing is known concerning the analytical properties of this distribution.

## Acknowledgements

We owe thanks to Guenther Walther for pointing us to Karlin [28] and Schoenberg's theorem. We also thank Tilmann Gneiting for several helpful discussions. Supported in part by NSF Grants DMS-08-04587 and DMS-11-04832, by NI-AID Grant 2R01 AI291968-04, and by the Alexander von Humboldt Foundation.

## References

- AYER, M., BRUNK, H.D., EWING, G.M., REID, W.T. and SILVERMAN, E. (1955). An empirical distribution function for sampling with incomplete information. *Ann. Math. Statist.* 26 641–647. MR0073895
- BALABDAOUI, F., RUFIBACH, K. and WELLNER, J.A. (2009). Limit distribution theory for maximum likelihood estimation of a log-concave density. Ann. Statist. 37 1299–1331. MR2509075
- [3] BALABDAOUI, F. and WELLNER, J.A. (2012). Chernoff's distribution is log-concave. Technical Report No. tr595.pdf. Dept. Statistics, Univ. Washington. Available at arXiv:1203.0828v1.
- [4] BARTHE, F. (2006). The Brunn-Minkowski theorem and related geometric and functional inequalities. In International Congress of Mathematicians. Vol. II 1529–1546. Zürich: Eur. Math. Soc. MR2275657
- BONDESSON, L. (1992). Generalized Gamma Convolutions and Related Classes of Distributions and Densities. Lecture Notes in Statistics 76. New York: Springer. MR1224674
- BONDESSON, L. (1997). On hyperbolically monotone densities. In Advances in the Theory and Practice of Statistics. Wiley Ser. Probab. Statist. Appl. Probab. Statist. 299–313. New York: Wiley. MR1481175
- BRUNK, H.D. (1970). Estimation of isotonic regression. In Nonparametric Techniques in Statistical Inference (Proc. Sympos., Indiana Univ., Bloomington, Ind., 1969) 177–197. London: Cambridge Univ. Press. MR0277070
- [8] CAFFARELLI, L.A. (2000). Monotonicity properties of optimal transportation and the FKG and related inequalities. Comm. Math. Phys. 214 547–563. MR1800860
- [9] CHERNOFF, H. (1964). Estimation of the mode. Ann. Inst. Statist. Math. 16 31–41. MR0172382

12

- [10] DANIELS, H.E. and SKYRME, T.H.R. (1985). The maximum of a random walk whose mean path has a maximum. Adv. in Appl. Probab. 17 85–99. MR0778595
- [11] DEHLING, H. and PHILIPP, W. (2002). Empirical Process Techniques for Dependent Data. Boston, MA: Birkhäuser. MR1958776
- [12] GRENANDER, U. (1956). On the theory of mortality measurement. I. Skand. Aktuarietidskr. 39 70–96. MR0086459
- [13] GRENANDER, U. (1956). On the theory of mortality measurement. II. Skand. Aktuarietidskr. 39 125–153. MR0093415
- [14] GROENEBOOM, P. (1985). Estimating a monotone density. In Proceedings of the Berkeley Conference in Honor of Jerzy Neyman and Jack Kiefer, Vol. II (Berkeley, CA, 1983). Wadsworth Statist./Probab. Ser. 539–555. Belmont, CA: Wadsworth. MR0822052
- [15] GROENEBOOM, P. (1989). Brownian motion with a parabolic drift and Airy functions. Probab. Theory Related Fields 81 79–109. MR0981568
- [16] GROENEBOOM, P. (1996). Lectures on inverse problems. In Lectures on Probability Theory and Statistics (Saint-Flour, 1994). Lecture Notes in Math. 1648 67–164. Berlin: Springer. MR1600884
- [17] GROENEBOOM, P. (2010). The maximum of Brownian motion minus a parabola. *Electron.* J. Probab. 15 1930–1937. MR2738343
- [18] GROENEBOOM, P. (2011). Vertices of the least concave majorant of Brownian motion with parabolic drift. *Electron. J. Probab.* 16 2234–2258. MR2861676
- [19] GROENEBOOM, P., JONGBLOED, G. and WELLNER, J.A. (2001). A canonical process for estimation of convex functions: The "invelope" of integrated Brownian motion +t<sub>4</sub>. Ann. Statist. **29** 1620–1652. MR1891741
- [20] GROENEBOOM, P., JONGBLOED, G. and WELLNER, J.A. (2001). Estimation of a convex function: Characterizations and asymptotic theory. Ann. Statist. 29 1653–1698. MR1891742
- [21] GROENEBOOM, P. and WELLNER, J.A. (1992). Information Bounds and Nonparametric Maximum Likelihood Estimation. DMV Seminar 19. Basel: Birkhäuser. MR1180321
- [22] GROENEBOOM, P. and WELLNER, J.A. (2001). Computing Chernoff's distribution. J. Comput. Graph. Statist. 10 388–400. MR1939706
- [23] HARGÉ, G. (2004). A convex/log-concave correlation inequality for Gaussian measure and an application to abstract Wiener spaces. Probab. Theory Related Fields 130 415–440. MR2095937
- [24] HARRISON, P.G. (1990). Laplace transform inversion and passage-time distributions in Markov processes. J. Appl. Probab. 27 74–87. MR1039185
- [25] HUANG, J. and WELLNER, J.A. (1995). Estimation of a monotone density or monotone hazard under random censoring. Scand. J. Stat. 22 3–33. MR1334065
- [26] HUANG, Y. and ZHANG, C.H. (1994). Estimating a monotone density from censored observations. Ann. Statist. 22 1256–1274. MR1311975
- [27] JANSON, S., LOUCHARD, G. and MARTIN-LÖF, A. (2010). The maximum of Brownian motion with parabolic drift. *Electron. J. Probab.* 15 1893–1929. MR2738342
- [28] KARLIN, S. (1968). Total Positivity. Vol. I. Stanford, CA: Stanford Univ. Press. MR0230102
- [29] KIM, J. and POLLARD, D. (1990). Cube root asymptotics. Ann. Statist. 18 191–219. MR1041391
- [30] LE CAM, L. (1986). The central limit theorem around 1935. Statist. Sci. 1 78–96. MR0833276
- [31] LEURGANS, S. (1982). Asymptotic distributions of slope-of-greatest-convex-minorant estimators. Ann. Statist. 10 287–296. MR0642740

- [32] MARSHALL, A.W. and OLKIN, I. (1979). Inequalities: Theory of Majorization and Its Applications. Mathematics in Science and Engineering 143. New York: Academic Press [Harcourt Brace Jovanovich Publishers]. MR0552278
- [33] MARSHALL, A.W., OLKIN, I. and ARNOLD, B.C. (2011). Inequalities: Theory of Majorization and Its Applications, 2nd ed. Springer Series in Statistics. New York: Springer. MR2759813
- [34] MERKES, E.P. and SALMASSI, M. (1997). On univalence of certain infinite products. Complex Variables Theory Appl. 33 207–215. MR1624939
- [35] OLKIN, I. and TONG, Y.L. (1988). Peakedness in multivariate distributions. In Statistical Decision Theory and Related Topics, IV, Vol. 2 (West Lafayette, IN, 1986) 373–383. New York: Springer. MR0927147
- [36] OLVER, F.W.J., LOZIER, D.W., BOISVERT, R.F. and CLARK, C.W. (2010). NIST Handbook of Mathematical Functions. Washington, DC: U.S. Department of Commerce National Institute of Standards and Technology. MR2723248
- [37] PRAKASA RAO, B.L.S. (1969). Estkmation of a unimodal density. Sankhyā Ser. A 31 23–36. MR0267677
- [38] PRAKASA RAO, B.L.S. (1970). Estimation for distributions with monotone failure rate. Ann. Math. Statist. 41 507–519. MR0260133
- [39] PROSCHAN, F. (1965). Peakedness of distributions of convex combinations. Ann. Math. Statist. 36 1703–1706. MR0187269
- [40] ROCKAFELLAR, R.T. and WETS, R.J.B. (1998). Variational Analysis. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences] 317. Berlin: Springer. MR1491362
- [41] SALMASSI, M. (1999). Inequalities satisfied by the Airy functions. J. Math. Anal. Appl. 240 574–582. MR1731663
- [42] SCHOENBERG, I.J. (1951). On Pólya frequency functions. I. The totally positive functions and their Laplace transforms. J. Anal. Math. 1 331–374. MR0047732
- [43] SHORACK, G.R. (2000). Probability for Statisticians. Springer Texts in Statistics. New York: Springer. MR1762415
- [44] VAN EEDEN, C. (1957). Maximum likelihood estimation of partially or completely ordered parameters. I. Nederl. Akad. Wetensch. Proc. Ser. A. 60 = Indag. Math. 19 128–136. MR0083869
- [45] VAN ZWET, W.R. (1964). Convex transformations: A new approach to skewness and kurtosis. Stat. Neerl. 18 433–441. MR0175217
- [46] VAN ZWET, W.R. (1964). Convex Transformations of Random Variables. Mathematical Centre Tracts 7. Amsterdam: Mathematisch Centrum. MR0176511
- [47] WELLNER, J.A. (2013). Strong log-concavity is preserved by convolution. In *High Dimensional Probability VI: The Banff Volume (Progress in Probability)* (C. Houdré, D. M. Mason, J. Rosiński and J. A. Wellner, eds.) 95–103. Basel: Birkhauser.

Received March 2012 and revised October 2012