

ELEMENTARY APPROACH TO HOMOGENEOUS C^* -ALGEBRAS

PIOTR NIEMIEC

ABSTRACT. A C^* -algebra is n -homogeneous (where n is finite) if every its nonzero irreducible representation acts on an n -dimensional Hilbert space. An elementary proof of Fell's characterization of n -homogeneous C^* -algebras (by means of their spectra) is presented. A spectral theorem and a functional calculus for finite systems of elements which generate n -homogeneous C^* -algebras are proposed.

1. INTRODUCTION

In 1961 Fell [9] gave a characterization of n -homogeneous C^* -algebras in terms of fibre bundles. It is a natural generalization of the commutative Gelfand-Naimark theorem, which gives models for commutative C^* -algebras. However, Fell's proof involves the machinery of (general) operator fields and as such is more advanced than Gelfand's theory of commutative Banach algebras. In this paper we propose a new proof of his theorem (starting from the very beginning), which is elementary and resembles the standard proof of the commutative Gelfand-Naimark theorem. We avoid the abstract language of fibre bundles—instead of them we introduce n -spaces, which are counterparts of locally compact Hausdorff spaces in the commutative case. These are locally compact Hausdorff spaces endowed with a (continuous) free action of the group $\mathfrak{U}_n = \mathcal{U}_n/Z(\mathcal{U}_n)$ where \mathcal{U}_n is the unitary group of $n \times n$ -matrices and $Z(\mathcal{U}_n)$ is its center.

Our approach to the subject mentioned above enables us to generalize the spectral theorem (for a normal Hilbert space operator) to the context of finite systems generating homogeneous C^* -algebras. It also allows building so-called n -functional calculus for such systems. These and related topics are discussed in the present paper.

The paper is organized as follows. Section 2 is devoted to operator-valued version of the Stone-Weierstrass theorem, which plays an important role in our proof of Fell's theorem on homogeneous C^* -algebras (presented in Section 5). In Section 3 we define and establish basic properties of so-called n -spaces (X, \cdot) (which, in fact, are the same as Fell's fibre bundles) and corresponding to them C^* -algebras $C^*(X, \cdot)$. These investigations are continued in the next part where we define spectral n -measures and characterize by means of them *all* representations of $C^*(X, \cdot)$ for any n -space (X, \cdot) . In Section 5 we give a new proof of Fell's characterization of homogeneous C^* -algebras. In the last, sixth, part we formulate the spectral theorem for finite systems of elements which generate n -homogeneous C^* -algebras and build the n -functional calculus for them.

2010 *Mathematics Subject Classification.* Primary 46L05; Secondary 46L35.

Key words and phrases. Homogeneous C^* -algebra; spectrum of a C^* -algebra; Stone-Weierstrass theorem; commutative Gelfand-Naimark theorem; spectral theorem; spectral measure; functional calculus.

The author gratefully acknowledges the assistance of the Polish Ministry of Sciences and Higher Education grant NN201 546438 for the years 2010–2013.

Notation and terminology. If a C^* -algebra \mathcal{A} has a unit e , the spectrum of x is denoted by $\sigma(x)$ and it is the set of all $\lambda \in \mathbb{C}$ for which $x - \lambda e$ is noninvertible in \mathcal{A} . For two selfadjoint elements a and b of \mathcal{A} we write $a \leq b$ provided $b - a$ is nonnegative. If $a \leq b$ and $b - a$ is invertible in \mathcal{A} , we shall express this by writing $a < b$ or $b > a$. The C^* -algebra of all bounded operators on a (complex) Hilbert space \mathcal{H} is denoted by $\mathcal{B}(\mathcal{H})$. Representations of unital C^* -algebra need not preserve unities and they are understood as $*$ -homomorphisms into $\mathcal{B}(\mathcal{H})$ for some Hilbert space \mathcal{H} . A representation of a C^* -algebra is n -dimensional if it acts on an n -dimensional Hilbert space. A *map* is a continuous function.

2. OPERATOR-VALUED STONE-WEIERSTRASS THEOREM

The classical Stone-Weierstrass theorem finds many applications in functional analysis and approximation theory. It reached many generalizations as well — see e.g. [10], [2], [17], [21], [11], [14] or [27] and the references there (consult also Theorem 1.4 in [9], §4.7 in [23] and Corollary 11.5.3 in [5]). A first significant counterpart of it for general C^* -algebras was established by Glimm [10]. Much later Longo [17] and Popa [21] proved independently a stronger version of Glimm's result, solving a long-standing problem in theory of C^* -algebras. In comparison to the classical Stone-Weierstrass theorem or, for example, to its generalization by Timofte [27], Glimm's and Longo's-Popa's theorems are not settled in function spaces. In this section we propose another version of the theorem under discussion which takes place in spaces of functions taking values in C^* -algebras. As such, it may be considered as its very natural generalization. Although the results of Glimm and Longo and Popa are stronger and more general than ours, they involve advanced machinery of C^* -algebras and advanced language of this theory, while our approach is very elementary and its proof is similar to Stone's [24, 25]. To formulate our result, we need to introduce the following notion.

2.1. Definition. Let X be a set, x and y be distinct points of X and let \mathcal{A} be a unital C^* -algebra. A collection \mathcal{F} of functions from X to \mathcal{A} *spectrally separates* points x and y if there is $f \in \mathcal{F}$ such that $f(x)$ and $f(y)$ are normal elements of \mathcal{A} and their spectra are disjoint. If \mathcal{F} spectrally separates any two distinct points of X , we say that \mathcal{F} *spectrally separates points* of X .

The reader should notice that a collection of complex-valued functions spectrally separates two points iff it separates them.

Whenever \mathcal{A} is a unital C^* -algebra and a is a selfadjoint element of \mathcal{A} , let us denote by $M(a)$ the real number $\max \sigma(a)$. Further, if X is a locally compact Hausdorff space and $f: X \rightarrow \mathcal{A}$ is a map, we say that f *vanishes at infinity* iff for every $\varepsilon > 0$ there is a compact set $K \subset X$ such that $\|f(x)\| < \varepsilon$ for any $x \in X \setminus K$. The set of all \mathcal{A} -valued maps on X vanishing at infinity is denoted by $\mathcal{C}_0(X, \mathcal{A})$. Notice that $\mathcal{C}_0(X, \mathcal{A})$ is a C^* -algebra when it is equipped with pointwise actions and the supremum norm induced by the norm of \mathcal{A} . Moreover, $\mathcal{C}_0(X, \mathcal{A})$ is unital iff X is compact (recall that we assume here that \mathcal{A} is unital).

A full version of our Stone-Weierstrass type theorem has the following form.

2.2. Theorem. *Let X be a locally compact Hausdorff space and let \mathcal{A} be a unital C^* -algebra. Let \mathcal{E} be a $*$ -subalgebra of $\mathcal{C}_0(X, \mathcal{A})$ such that:*

(AX0) *if X is noncompact, then for each $z \in X$ either $f_0(z)$ is invertible in \mathcal{A} for some $f_0 \in \mathcal{E}$ or $f(z) = 0$ for any $f \in \mathcal{E}$;*

and for any two points x and y of X one of the following two conditions is fulfilled:

(AX1) *either x and y are spectrally separated by \mathcal{E} , or*

(AX2) *$M(f(x)) = M(f(y))$ for any $f \in \mathcal{E}$.*

Then the (uniform) closure of \mathcal{E} in $\mathcal{C}_0(X, \mathcal{A})$ coincides with the $*$ -algebra $\Delta_2(\mathcal{E})$ of all maps $u \in \mathcal{C}_0(X, \mathcal{A})$ such that for any $x, y \in X$ and each $\varepsilon > 0$ there exists $v \in \mathcal{E}$ with $\|v(z) - u(z)\| < \varepsilon$ for $z \in \{x, y\}$.

As a consequence of the above result we obtain the following result, which is a special case of Corollary 11.5.3 in [5].

2.3. Proposition. *Let X be a locally compact Hausdorff space and let \mathcal{A} be a unital C^* -algebra. A $*$ -subalgebra \mathcal{E} of $\mathcal{C}_0(X, \mathcal{A})$ is dense in $\mathcal{C}_0(X, \mathcal{A})$ iff \mathcal{E} spectrally separates points of X and for every $x \in X$ the set $\mathcal{E}(x) := \{f(x) : f \in \mathcal{E}\}$ is dense in \mathcal{A} .*

It is worth noting that we know no characterization of dense $*$ -subalgebras of $\mathcal{C}_0(X, \mathcal{A})$ in case \mathcal{A} does not have a unit.

The proof of Theorem 2.2 is partially based on the original proof of the Stone-Weierstrass theorem given by Stone [24, 25]. However, the key tool in our proof is the so-called Loewner-Heinz inequality (for the discussion on this inequality, see page 150 in [1]), first proved by Loewner [18]:

2.4. Theorem. *Let a and b be two selfadjoint nonnegative elements in a C^* -algebra such that $a \leq b$. Then for every $s \in (0, 1)$, $a^s \leq b^s$.*

The proof of Theorem 2.2 is preceded by several auxiliary results. For simplicity, the unit of \mathcal{A} will be denoted by 1 and the function from X to \mathcal{A} constantly equal to 1 will be denoted by 1_X . We also preserve the notation of Theorem 2.2. Additionally, $\bar{\mathcal{E}}$ stands for the (uniform) closure of \mathcal{E} in $\mathcal{C}_0(X, \mathcal{A})$.

2.5. Lemma. *Suppose X is compact. Then $1_X \in \bar{\mathcal{E}}$ if and only if for every $x \in X$ there is $f \in \mathcal{E}$ such that $f(x)$ is invertible in \mathcal{A} .*

Proof. The necessity is clear (since the set of all invertible elements is open in \mathcal{A}). To prove the sufficiency, for each $x \in X$ take $f_x \in \mathcal{E}$ such that $f_x(x)$ is invertible. Put $u_x = f_x^* f_x \in \mathcal{E}$ and let $V_x \subset X$ consist of all $y \in X$ such that $u_x(y) > 0$. It follows from the continuity of u_x that V_x is open. By the compactness of X , $X = \bigcup_{j=1}^p V_{x_j}$ for some finite system x_1, \dots, x_p . Put $u = \sum_{j=1}^p u_{x_j} \in \mathcal{E}$ and note that $u(x) > 0$ for each $x \in X$. This implies that u is invertible in $\mathcal{C}_0(X, \mathcal{A})$. Let $f : [0, \|u\|] \rightarrow \mathbb{R}$ be a map with $f(0) = 0$ and $f|_{\sigma(u)} \equiv 1$. There is a sequence of real polynomials p_1, p_2, \dots which converge uniformly to f on $[0, \|u\|]$. Then $p_n(u) \rightarrow f(u) = 1_X$ (in the norm topology) and hence $1_X \in \bar{\mathcal{E}}$. \square

2.6. Lemma. *Suppose X is compact and $1_X \in \bar{\mathcal{E}}$. Let $x \in X$ and $\delta > 0$ be arbitrary. For any selfadjoint $f \in \Delta_2(\mathcal{E})$ there are selfadjoint $g, h \in \bar{\mathcal{E}}$ such that $g(x) = f(x) = h(x)$ and $g - \delta \cdot 1_X \leq f \leq h + \delta \cdot 1_X$.*

Proof. It follows from the definition of $\Delta_2(\mathcal{E})$ (and the fact that $*$ -homomorphisms between C^* -algebras have closed ranges) that for every $y \in X$ there is $f_y \in \bar{\mathcal{E}}$ with $f_y(z) = f(z)$ for $z \in \{x, y\}$. Replacing, if needed, f_y by $\frac{1}{2}(f_y + f_y^*)$, we may assume that f_y is selfadjoint. Let $U_y \subset X$ consist of all $z \in X$ such that $\|f_y(z) - f(z)\| < \delta$. Take a finite number of points x_1, \dots, x_p for which $X = \bigcup_{j=1}^p U_{x_j}$. For simplicity, put $V_j = U_{x_j}$ and $g_j = f_{x_j}$ ($j = 1, \dots, p$). Observe that $f - \delta \cdot 1_X \leq g_j$ on V_j and $g_j(x) = f(x)$. We define by induction functions $h_1, \dots, h_p \in \bar{\mathcal{E}}$: $h_1 = g_1$ and $h_k = \frac{1}{2}(h_{k-1} + g_k + |h_{k-1} - g_k|)$ for $k = 2, \dots, p$ where $|u| = \sqrt{u^* u}$ for each $u \in \bar{\mathcal{E}}$. Since $\bar{\mathcal{E}}$ is a C^* -algebra, we clearly have $h_k \in \bar{\mathcal{E}}$. Use induction to show that $h_j(x) = f(x)$ and $g_j \leq h_p$ for $j = 1, \dots, p$. Then $h = h_p$ is the function we searched for. Indeed, $h(x) = f(x)$ and for any $y \in X$ there is $j \in \{1, \dots, p\}$ such that $y \in V_j$, which implies that $f(y) - \delta \cdot 1 \leq g_j(y) \leq h(y)$.

Now if we apply the above argument to the function $-f$, we shall obtain a selfadjoint function $h' \in \bar{\mathcal{E}}$ such that $-f(x) = h'(x)$ and $-f \leq h' + \delta \cdot 1_X$. Then put $g := -h'$ to complete the proof. \square

2.7. Lemma. *Let $\varepsilon > 0$, $r > 0$ and $k \geq 1$ be given. There is a natural number $N = N(\varepsilon, r, k)$ with the following property. If a_1, \dots, a_k, b are selfadjoint elements of \mathcal{A} such that $0 \leq a_j \leq b$, $ba_j = a_jb$ ($j = 1, \dots, k$) and $\|b\| \leq r$, then $a_s \leq (\sum_{j=1}^k a_j^n)^{\frac{1}{n}} \leq b + \varepsilon \cdot 1$ for any $s \in \{1, \dots, k\}$ and $n \geq N$.*

Proof. Let $N \geq 2$ be such that $\sqrt[n]{k} \leq 1 + \frac{\varepsilon}{r}$ for each $n \geq N$ and let a_1, \dots, a_k, b be as in the statement of the lemma. Since then $a_s^n \leq \sum_{j=1}^k a_j^n$, Theorem 2.4 yields $a_s \leq (\sum_{j=1}^k a_j^n)^{\frac{1}{n}}$. Further, since b commutes with a_j , we get $a_j^n \leq b^n$ and consequently $\sum_{j=1}^k a_j^n \leq kb^n$. So, another application of Theorem 2.4 gives us $(\sum_{j=1}^k a_j^n)^{\frac{1}{n}} \leq \sqrt[n]{k}b$. So, it suffices to have $\sqrt[n]{k}b \leq b + \varepsilon \cdot 1$ which is fulfilled for $n \geq N$ because $\|(\sqrt[n]{k} - 1)b\| \leq (\sqrt[n]{k} - 1)r \leq \varepsilon$. \square

2.8. Lemma. *Suppose X is compact and $1_X \in \bar{\mathcal{E}}$. If $f \in \Delta_2(\mathcal{E})$ commutes with every member of \mathcal{E} , then $f \in \bar{\mathcal{E}}$.*

Proof. Since $\Delta_2(\mathcal{E})$ is a $*$ -algebra, we may assume that f is selfadjoint. Fix $\delta > 0$. By Lemma 2.6, for every $x \in X$ there is $f_x \in \bar{\mathcal{E}}$ with $f_x(x) = f(x)$ and $f_x \leq f + \delta \cdot 1_X$. Let $U_x \subset X$ consist of all $y \in X$ such that $f_x(y) > f(y) - \delta \cdot 1$. We infer from the compactness of X that $X = \bigcup_{j=1}^k U_{x_j}$ for some points $x_1, \dots, x_k \in X$. For simplicity, we put $V_j = U_{x_j}$ and $g_j = f_{x_j}$. We then have

$$(2-1) \quad g_j(x) \geq f(x) - \delta \cdot 1 \quad \text{for any } x \in V_j$$

and

$$(2-2) \quad g_j(x) \leq f(x) + \delta \cdot 1 \quad \text{for any } x \in X.$$

It follows from the compactness of X that there is a constant $c > 0$ such that $g_j + c \cdot 1_X \geq 0$ ($j = 1, \dots, k$) and $f + (c - \delta) \cdot 1_X \geq 0$. Further, there is $r > 0$ such that $f + (c + \delta) \cdot 1_X \leq r \cdot 1_X$. Now let $N = N(\delta, r, k)$ be as in Lemma 2.7. Since f commutes with each member of $\bar{\mathcal{E}}$, we conclude from that lemma and from (2-2) that $g_s(x) + c \cdot 1 \leq [\sum_{j=1}^k (g_j(x) + c \cdot 1)^n]^{\frac{1}{n}} \leq f(x) + (c + 2\delta) \cdot 1$ for any $x \in X$. Finally, since $1_X \in \bar{\mathcal{E}}$, the function $g := [\sum_{j=1}^k (g_j + c \cdot 1_X)^n]^{\frac{1}{n}} - c \cdot 1_X$ belongs to $\bar{\mathcal{E}}$. What is more, $g \leq f + 2\delta \cdot 1_X$ and $g(x) \geq g_j(x) \geq f(x) - \delta \cdot 1$ for $x \in V_j$ (cf. (2-1)). This gives $f - \delta \cdot 1_X \leq g$ on the whole space X and therefore $-\delta \cdot 1_X \leq g - f \leq 2\delta \cdot 1_X$, which is equivalent to $\|g - f\| \leq 2\delta$ and finishes the proof. \square

2.9. Lemma. *Suppose X is compact, $1_X \in \bar{\mathcal{E}}$ and there exists an equivalence relation \mathcal{R} on X such that two points x and y are spectrally separated by \mathcal{E} whenever $(x, y) \notin \mathcal{R}$. Then every map $g: X \rightarrow \mathbb{C} \cdot 1 \subset \mathcal{A}$ which is constant on each equivalence class with respect to \mathcal{R} belongs to $\bar{\mathcal{E}}$.*

Proof. By Lemma 2.8, we only need to check that $g \in \Delta_2(\mathcal{E})$. We may assume that $g: X \rightarrow \mathbb{R} \cdot 1$. Let x and y be arbitrary. Write $g(x) = \alpha \cdot 1$ and $g(y) = \beta \cdot 1$. If $(x, y) \in \mathcal{R}$, then both x and y belong to the same equivalence class and hence $\alpha = \beta$. Then $g(z) = (\alpha \cdot 1_X)(z)$ for $z \in \{x, y\}$ (and $\alpha \cdot 1_X \in \bar{\mathcal{E}}$). Now assume that $(x, y) \notin \mathcal{R}$. Then—by assumption—there is $f \in \mathcal{E}$ such that both $f(x)$ and $f(y)$ are normal and $\sigma(f(x)) \cap \sigma(f(y)) = \emptyset$. Let $\varphi: \mathbb{C} \rightarrow \mathbb{R}$ be a map such that $\varphi|_{\sigma(f(x))} \equiv \alpha$ and $\varphi|_{\sigma(f(y))} \equiv \beta$. There is a sequence of polynomials $p_1(z, \bar{z}), p_2(z, \bar{z}), \dots$ which converge uniformly to φ on $K := \sigma(f(x)) \cup \sigma(f(y))$. Then $p_n(f, f^*) \in \mathcal{E}$ and, for $w \in \{x, y\}$, $[p_n(f, f^*)](w) = p_n(f(w), [f(w)]^*)$. Since $f(w)$ is normal and its

spectrum is contained in K , we see that $\lim_{n \rightarrow \infty} [p_n(f, f^*)](w) = \varphi(f(w))$. Now the notice that $\varphi(f(x)) = \alpha \cdot 1 = g(x)$ and $\varphi(f(y)) = \beta \cdot 1 = g(y)$ finishes the proof. \square

We recall that if X is a compact Hausdorff space and \mathcal{R} is a closed equivalence relation on X , then the quotient topological space X/\mathcal{R} is Hausdorff as well.

2.10. Lemma. *Suppose X is compact and there is a closed equivalence relation \mathcal{R} on X such that $M(f(x)) = M(f(y))$ for each selfadjoint $f \in \mathcal{E}$ whenever $(x, y) \in \mathcal{R}$. Let $\pi: X \rightarrow X/\mathcal{R}$ denote the canonical projection, $f \in \bar{\mathcal{E}}$ be selfadjoint, a and b be two real numbers and let $U = \{x \in X: a \cdot 1 < f(x) < b \cdot 1\}$. Then $\pi^{-1}(\pi(U)) = U$ and $\pi(U)$ is open in X/\mathcal{R} .*

Proof. Recall that $\pi(U)$ is open in X/\mathcal{R} iff $\pi^{-1}(\pi(U))$ is open in X . Therefore it suffices to show that $\pi^{-1}(\pi(U)) = U$. Of course, the inclusion ' \supset ' is immediate. And if $y \in \pi^{-1}(\pi(U))$, then there is $x \in U$ such that $(x, y) \in \mathcal{R}$. We then have $a \cdot 1 < f(x) < b \cdot 1$, $M(f(x)) = M(f(y))$ and $M(-f(x)) = M(-f(y))$ (the last two relations follows from the fact that $f \in \bar{\mathcal{E}}$). The first of these relations says that $[-M(-f(x)), M(f(x))] \subset (a, b)$ from which we infer that $[-M(-f(y)), M(f(y))] \subset (a, b)$ and consequently $y \in U$. \square

The following is a special case of Theorem 2.2.

2.11. Lemma. *Suppose X is compact, $1_X \in \bar{\mathcal{E}}$ and for any $x, y \in X$ one of conditions (AX1)–(AX2) is fulfilled. Then $\Delta_2(\mathcal{E}) = \bar{\mathcal{E}}$.*

Proof. We only need to show that $\Delta_2(\mathcal{E})$ is contained in $\bar{\mathcal{E}}$. Let $f \in \Delta_2(\mathcal{E})$ be selfadjoint and let $\delta > 0$. We shall construct $w \in \bar{\mathcal{E}}$ such that $\|w - f\| \leq 3\delta$. By Lemma 2.6, for each $x \in X$ there are functions $u_x, v_x \in \bar{\mathcal{E}}$ such that $u_x(x) = f(x) = v_x(x)$ and $u_x - \delta \cdot 1_X < f < v_x + \delta \cdot 1_X$. Let $G_x \subset X$ consist of all $y \in X$ such that $v_x(y) - \delta \cdot 1 < f(y) < u_x(y) + \delta \cdot 1$. Since $x \in G_x$ and X is compact, there is a finite system $x_1, \dots, x_k \in X$ for which $X = \bigcup_{j=1}^k G_{x_j}$. For simplicity, we put $W_j = G_{x_j}$, $p_j = u_{x_j}$ and $q_j = v_{x_j}$. Observe that then

$$(2-3) \quad p_j(x) - \delta \cdot 1 < f(x) < q_j(x) + \delta \cdot 1 \quad \text{for any } x \in X$$

and

$$(2-4) \quad q_j(x) - \delta \cdot 1 < f(x) < p_j(x) + \delta \cdot 1 \quad \text{for any } x \in W_j.$$

Let D_j consist of all $x \in X$ such that $-2\delta \cdot 1 < p_j(x) - q_j(x) < 2\delta \cdot 1$. We infer from (2-3) and (2-4) that $W_j \subset D_j$ and thus $X = \bigcup_{j=1}^k D_j$. Further, let \mathcal{R} be an equivalence relation on X given by the rule: $(x, y) \in \mathcal{R} \iff M(u(x)) = M(u(y))$ for each selfadjoint $u \in \mathcal{E}$. It follows from the definition of \mathcal{R} that \mathcal{R} is closed in $X \times X$. Denote by $\pi: X \rightarrow X/\mathcal{R}$ the canonical projection. We deduce from Lemma 2.10 that the sets $\pi(D_1), \dots, \pi(D_k)$ form an open cover of the space X/\mathcal{R} (which is compact and Hausdorff). Now let $\beta_1, \dots, \beta_k: X/\mathcal{R} \rightarrow [0, 1]$ be a partition of unity such that $\beta_j^{-1}((0, 1]) \subset \pi(D_j)$ for $j = 1, \dots, k$. Put $\alpha_j = (\beta_j \circ \pi) \cdot 1: X \rightarrow \mathbb{C} \cdot 1 \subset \mathcal{A}$. Lemma 2.9 combined with conditions (AX1)–(AX2) yields that $\alpha_1, \dots, \alpha_k \in \bar{\mathcal{E}}$. Define $w \in \bar{\mathcal{E}}$ by $w = \sum_{j=1}^k \alpha_j p_j$. Since $\sum_{j=1}^k \alpha_j = 1_X$, we conclude from (2-3) that $w \leq f + \delta \cdot 1_X$. So, to end the proof, it is enough to check that $f(x) \leq w(x) + 3\delta \cdot 1$ for each $x \in X$. This inequality will be satisfied provided

$$(2-5) \quad \alpha_j(x)(f(x) - 3\delta \cdot 1) \leq \alpha_j(x)p_j(x)$$

for any j . We consider two cases. If $x \in D_j$, then $p_j(x) > q_j(x) - 2\delta \cdot 1 > f(x) - 3\delta \cdot 1$ (by (2-3)) and consequently (2-5) holds. Finally, if $x \notin D_j$, then $\pi(x) \notin \pi(D_j)$ (see Lemma 2.10) and therefore $\alpha_j(x) = 0$, which easily gives (2-5). \square

Proof of Theorem 2.2. We only need to check that $\Delta_2(\mathcal{E}) \subset \bar{\mathcal{E}}$. We consider two cases.

First assume X is compact. Let $\mathcal{E}' = \mathcal{E} + \mathbb{C} \cdot 1_X$. Observe that \mathcal{E}' is a $*$ -algebra and for any two points x and y one of conditions (AX1)–(AX2) is fulfilled with \mathcal{E} replaced by \mathcal{E}' . Consequently, it follows from Lemma 2.11 that $\overline{\mathcal{E}'} = \Delta_2(\mathcal{E}')$. But $\overline{\mathcal{E}'} = \bar{\mathcal{E}} + \mathbb{C} \cdot 1_X$. So, for any $g \in \Delta_2(\mathcal{E})$ we clearly have $g \in \Delta_2(\mathcal{E}')$ and hence $g = f + \lambda \cdot 1_X$ for some $f \in \bar{\mathcal{E}}$ and $\lambda \in \mathbb{C}$. If $\lambda = 0$, then $g = f \in \bar{\mathcal{E}}$ and we are done. Otherwise, $1_X = \frac{1}{\lambda}(g - f) \in \Delta_2(\mathcal{E})$ which implies that the assumptions of Lemma 2.5 are satisfied. We infer from that lemma that $1_X \in \bar{\mathcal{E}}$ and therefore $g \in \bar{\mathcal{E}}$ as well.

Now assume X is noncompact. Let $\hat{X} = X \cup \{\infty\}$ be the one-point compactification of X . Every function $f \in \mathcal{C}_0(X, \mathcal{A})$ admits a unique continuous extension $\hat{f}: \hat{X} \rightarrow \mathcal{A}$, given by $\hat{f}(\infty) = 0$. Denote by $\hat{\mathcal{E}}$ the $*$ -subalgebra of $\mathcal{C}(\hat{X}, \mathcal{A})$ consisting of all extensions of (all) functions from \mathcal{E} . We claim that for any $x, y \in \hat{X}$ one of conditions (AX1)–(AX2) is fulfilled with \mathcal{E} replaced by $\hat{\mathcal{E}}$. Indeed, if both x and y differ from ∞ , this follows from our assumptions about \mathcal{E} . And if, for example, $y = \infty \neq x$, condition (AX0) implies that either $M(\hat{f}(x)) = M(\hat{f}(y))$ for each $f \in \mathcal{E}$ or $\hat{u}(x)$ is invertible in \mathcal{A} for some $u \in \mathcal{E}$. But then $f = u^*u \in \mathcal{E}$ is normal and $0 \notin \sigma(\hat{f}(x))$, while $\sigma(\hat{f}(y)) = \{0\}$, which shows that x and y are spectrally separated by $\hat{\mathcal{E}}$. So, it follows from the first part of the proof that the closure of $\hat{\mathcal{E}}$ in $\mathcal{C}(\hat{X}, \mathcal{A})$ coincides with $\Delta_2(\hat{\mathcal{E}})$. But the closure of $\hat{\mathcal{E}}$ coincides with $\{\hat{f}: f \in \bar{\mathcal{E}}\}$ and $\Delta_2(\hat{\mathcal{E}}) = \{\hat{f}: f \in \Delta_2(\mathcal{E})\}$. We infer from these that $\Delta_2(\mathcal{E}) = \bar{\mathcal{E}}$ and the proof is complete. \square

Proof of Proposition 2.3. The necessity of the condition is clear (since for any two distinct points x and y in X and any elements a and b of \mathcal{A} there is a function $f \in \mathcal{C}_0(X, \mathcal{A})$ such that $f(x) = a$ and $f(y) = b$). To prove the sufficiency, assume \mathcal{E} spectrally separates points of X and for each $x \in X$ the set $\mathcal{E}(x)$ is dense in \mathcal{A} . First notice that then for each $x \in X$ there is $f \in \mathcal{E}$ such that $f(x)$ is invertible in \mathcal{A} . This shows that all assumptions of Theorem 2.2 are satisfied. According to that result, we only need to show that for any two distinct points x and y of X the set $L := \{(f(x), f(y)): f \in \mathcal{E}\}$ is dense in $\mathcal{A} \times \mathcal{A}$. Since x and y are spectrally separated by \mathcal{E} , the proof of Lemma 2.9 shows that $(1, 0), (0, 1) \in \bar{L}$. Further, since both $\mathcal{E}(x)$ and $\mathcal{E}(y)$ are dense in \mathcal{A} , we conclude that $\{f(x): f \in \bar{\mathcal{E}}\} = \{f(y): f \in \bar{\mathcal{E}}\} = \mathcal{A}$ and therefore for arbitrary two elements a and b of \mathcal{A} there are $u, v \in \bar{\mathcal{E}}$ for which $u(x) = a$ and $v(y) = b$. Then $(a, b) = (u(x), u(y)) \cdot (1, 0) + (v(x), v(y)) \cdot (0, 1) \in \bar{L}$ (we use here the coordinatewise multiplication) and we are done. \square

Taking into account Proposition 2.3, the following question may be interesting.

2.12. Problem. Let \mathcal{A} be a C^* -algebra without unit and let X be a locally compact Hausdorff space. Is it true that a $*$ -subalgebra \mathcal{E} of $\mathcal{C}_0(X, \mathcal{A})$ is dense in $\mathcal{C}_0(X, \mathcal{A})$ iff for any two points x and y of X the set $\{(f(x), f(y)): f \in \mathcal{E}\}$ is dense in $\mathcal{A} \times \mathcal{A}$?

3. TOPOLOGICAL n -SPACES

In Fell's characterization of homogeneous C^* -algebras [9] (consult also Theorem IV.1.7.23 in [3]) special fibre bundles appear. To make our lecture as simple and elementary as possible, we avoid this language and instead of using fibre bundles we shall introduce so-called n -spaces (see Definition 3.1 below). To this end, let M_n be the C^* -algebra of all complex $n \times n$ -matrices. Let \mathcal{U}_n be the unitary group of M_n and I be its neutral element. Let $\mathbb{T} = \{z \in \mathbb{C}: |z| = 1\}$. Let \mathfrak{U}_n denote the compact topological group $\mathcal{U}_n/(\mathbb{T} \cdot I)$ and let $\pi_n: \mathcal{U}_n \rightarrow \mathfrak{U}_n$ be the canonical homomorphism. Members of \mathfrak{U}_n will be denoted by \mathfrak{u} and \mathfrak{v} , and j is reserved for

the neutral element of \mathfrak{U}_n . The (probabilistic) Haar measure on \mathfrak{U}_n will be denoted by du . For any $A \in M_n$ and $u \in \mathfrak{U}_n$ let $u.A$ denote the matrix UAU^{-1} where $U \in \mathcal{U}_n$ is such that $\pi_n(U) = u$. It is easily seen that the function

$$\mathfrak{U}_n \times M_n \ni (u, A) \mapsto u.A \in M_n$$

is a well defined continuous action of \mathfrak{U}_n on M_n (which means that $j.A = A$ and $u.(v.A) = (uv).A$ for any $u, v \in \mathfrak{U}_n$ and $A \in M_n$). More generally, for any C^* -algebra \mathcal{A} let $M_n(\mathcal{A})$ be the algebra of all $n \times n$ -matrices with entries in \mathcal{A} . ($M_n(\mathcal{A})$ may naturally be identified with $\mathcal{A} \otimes M_n$.) For any matrix $A \in M_n(\mathcal{A})$ and each $u \in \mathfrak{U}_n$, $u.A$ is defined as UAU^{-1} where $U \in \mathcal{U}_n$ is such that $\pi_n(U) = u$, and UAU^{-1} is computed in a standard manner.

3.1. Definition. A pair (X, \cdot) is said to be an n -space if X is a locally compact Hausdorff space and $\mathfrak{U}_n \times X \ni (u, x) \mapsto u.x \in X$ is a continuous **free** action of \mathfrak{U}_n on X . Recall that the action is free iff the equality $u.x = x$ (for some $x \in X$) implies $u = j$.

Let (X, \cdot) be an n -space. Let $C^*(X, \cdot)$ be the $*$ -algebra of all maps $f \in \mathcal{C}_0(X, M_n)$ such that $f(u.x) = u.f(x)$ for any $u \in \mathfrak{U}_n$ and $x \in X$. $C^*(X, \cdot)$ is a C^* -subalgebra of $\mathcal{C}_0(X, M_n)$.

By a *morphism* between two n -spaces (X, \cdot) and $(Y, *)$ we mean any proper map $\psi: X \rightarrow Y$ such that $\psi(u.x) = u * \psi(x)$ for any $u \in \mathfrak{U}_n$ and $x \in X$. (A map is proper if the inverse images of compact sets under this map are compact.) A morphism which is a homeomorphism is said to be an *isomorphism*. Two n -spaces are *isomorphic* if there exists an isomorphism between them.

The reader should notice that the (natural) action of \mathfrak{U}_n on M_n is not free. However, one may check that the set \mathfrak{M}_n of all irreducible matrices $A \in M_n$ (that is, $A \in \mathfrak{M}_n$ iff every matrix $X \in M_n$ which commutes with both A and A^* is of the form λI where $\lambda \in \mathbb{C}$) is open in M_n (and thus \mathfrak{M}_n is locally compact) and the action $\mathfrak{U}_n \times \mathfrak{M}_n \ni (u, A) \mapsto u.A \in \mathfrak{M}_n$ is free, which means that (\mathfrak{M}_n, \cdot) is an n -space.

In this section we establish basic properties of C^* -algebras of the form $C^*(X, \cdot)$ where (X, \cdot) is an n -space. To this end, recall that whenever $(\Omega, \mathfrak{M}, \mu)$ is a finite measure space and $f: \Omega \ni \omega \mapsto (f_1(\omega), \dots, f_k(\omega)) \in \mathbb{C}^k$ is an \mathfrak{M} -measurable (which means that $f^{-1}(U) \in \mathfrak{M}$ for every open set $U \subset \mathbb{C}^k$) bounded function, then $\int_{\Omega} f(\omega) d\mu(\omega)$ is (well) defined as $(\int_{\Omega} f_1(\omega) d\mu(\omega), \dots, \int_{\Omega} f_k(\omega) d\mu(\omega))$. If $\|\cdot\|$ is any norm on \mathbb{C}^k , then $\|\int_{\Omega} f(\omega) d\mu(\omega)\| \leq \int_{\Omega} \|f(\omega)\| d\mu(\omega)$. In particular, the above rules apply to matrix-valued measurable functions.

From now on, $n \geq 1$ and an n -space (X, \cdot) are fixed. A set $A \subset X$ is said to be *invariant* provided $u.a \in A$ for any $u \in \mathfrak{U}_n$ and $a \in A$. Observe that if A is closed or open and A is invariant, then A is locally compact and consequently (A, \cdot) is an n -space (when the action of \mathfrak{U}_n is restricted to A). We begin with

3.2. Lemma. For each $f \in \mathcal{C}_0(X, M_n)$ let $f^{\mathfrak{U}}: X \rightarrow M_n$ be given by:

$$f^{\mathfrak{U}}(x) = \int_{\mathfrak{U}_n} u^{-1}.f(u.x) du \quad (x \in X).$$

- (a) For any $f \in \mathcal{C}_0(X, M_n)$, $f^{\mathfrak{U}} \in C^*(X, \cdot)$.
- (b) If $f \in \mathcal{C}_0(X, M_n)$ and $x \in X$ are such that $f(u.x) = u.f(x)$ for any $u \in \mathfrak{U}_n$, then $f^{\mathfrak{U}}(x) = f(x)$.
- (c) Let $A \subset X$ be a closed invariant nonempty set. Every map $g \in C^*(A, \cdot)$ extends to a map $\tilde{g} \in C^*(X, \cdot)$ such that $\sup_{a \in A} \|g(a)\| = \sup_{x \in X} \|\tilde{g}(x)\|$.
- (d) For any $x \in X$ and $A \in M_n$ there is $f \in C^*(X, \cdot)$ with $f(x) = A$.

- (e) Let x and y be two points of X such that there is no $u \in \mathfrak{U}_n$ for which $u.x = y$. Then for any $A, B \in M_n$ there is $f \in C^*(X, \cdot)$ such that $f(x) = A$ and $f(y) = B$.
- (f) $C^*(X, \cdot)$ has a unit iff X is compact.

Proof. It is clear that $f^{\mathfrak{U}}$ is continuous for every $f \in \mathcal{C}_0(X, M_n)$. Further, if $K \subset X$ is a compact set such that $\|f(x)\| \leq \varepsilon$ for each $x \in X \setminus K$, then $\|f^{\mathfrak{U}}(z)\| \leq \varepsilon$ for any $z \in X \setminus \mathfrak{U}_n.K$ where $\mathfrak{U}_n.K = \{u.x : u \in \mathfrak{U}_n, x \in K\}$. The note that $\mathfrak{U}_n.K$ is compact leads to the conclusion that $f^{\mathfrak{U}} \in \mathcal{C}_0(X, M_n)$. Finally, for any $v \in \mathfrak{U}_n$, any representant $V \in \mathcal{U}_n$ of v and each $x \in X$, we have

$$\begin{aligned} f^{\mathfrak{U}}(v.x) &= \int_{\mathfrak{U}_n} u^{-1}.f(uv.x) du = \int_{\mathfrak{U}_n} (uv^{-1})^{-1}.f(u.x) du \\ &= \int_{\mathfrak{U}_n} v.[u^{-1}.f(u.x)] du = \int_{\mathfrak{U}_n} V[u^{-1}.f(u.x)]V^{-1} du \\ &= V \cdot \left(\int_{\mathfrak{U}_n} u^{-1}.f(u.x) du \right) \cdot V^{-1} = v.f^{\mathfrak{U}}(x), \end{aligned}$$

which proves (a). Point (b) is a simple consequence of the definition of $f^{\mathfrak{U}}$. Further, if g is as in (c), it follows from Tietze's type theorem that there is $G \in \mathcal{C}_0(X, M_n)$ which extends g and satisfies $\sup_{a \in A} \|g(a)\| = \sup_{x \in X} \|G(x)\|$ (if X is noncompact, consider the one-point compactification $\hat{X} = X \cup \{\infty\}$ of X and note that then the set $\hat{A} = A \cup \{\infty\}$ is closed in \hat{X} and g extends continuously to \hat{A}). Then $\tilde{g} = G^{\mathfrak{U}}$ is a member of $C^*(X, \cdot)$ (by (a)) which we searched for (see (b)).

We turn to (d) and (e). Let $K = \mathfrak{U}_n.\{x\}$ and $f_0: K \rightarrow M_n$ be given by $f_0(u.x) = u.A$ ($u \in \mathfrak{U}_n$). Since the action of \mathfrak{U}_n on X is free, f_0 is a well defined map. Since K is compact, (c) yields the existence of $f \in C^*(X, \cdot)$ which extends f_0 . To prove (e), we argue similarly: put $L = \mathfrak{U}_n.\{x, y\}$ and let $g_0: L \rightarrow M_n$ be given by $g_0(u.x) = u.A$ and $g_0(u.y) = u.B$ ($u \in \mathfrak{U}_n$). We infer from the assumption of (e) that g_0 is a well defined map. Consequently, since L is compact, there exists, by (c), a map $g \in C^*(X, \cdot)$ which extends g_0 . This finishes the proof of (e), while point (f) immediately follows from (d). \square

3.3. Proposition. (a) For every closed two-sided ideal \mathcal{I} in $C^*(X, \cdot)$ there exists a (unique) closed invariant set $A \subset X$ such that \mathcal{I} coincides with the ideal \mathcal{I}_A of all functions $f \in C^*(X, \cdot)$ which vanish on A . Moreover, $C^*(X, \cdot)/\mathcal{I}$ is “naturally” isomorphic to $C^*(A, \cdot)$.

- (b) Let $k \leq n$ and let $\pi: C^*(X, \cdot) \rightarrow M_k$ be a nonzero representation. Then $k = n$ and there is a unique point $x \in X$ such that $\pi(f) = f(x)$ for $f \in C^*(X, \cdot)$.
- (c) Let $(Y, *)$ be an n -space. For every $*$ -homomorphism $\Phi: (X, \cdot) \rightarrow (Y, *)$ there is a unique pair (U, φ) where U is an open invariant subset of Y , $\varphi: (U, *) \rightarrow (X, \cdot)$ is a morphism of n -spaces and

$$(3-1) \quad [\Phi(f)](y) = \begin{cases} f(\varphi(y)) & \text{if } y \in U \\ 0 & \text{if } y \notin U. \end{cases}$$

In particular, $C^*(X, \cdot)$ and $C^*(Y, *)$ are isomorphic iff so are (X, \cdot) and $(Y, *)$.

Proof. The uniqueness of A in (a) follows from point (e) of Lemma 3.2. To show its existence, let A consist of all $x \in X$ such that $f(x) = 0$ for any $f \in \mathcal{I}$. It is clear that A is closed and invariant, and that $\mathcal{I} \subset \mathcal{I}_A$. To prove the converse inclusion, we shall involve Theorem 2.2 for $\mathcal{E} = \mathcal{I}$. First of all, it follows from point (d) of Lemma 3.2 that for each $x \in X$ the set $\mathcal{I}(x) := \{f(x) : f \in \mathcal{I}\}$ is a two-sided ideal in M_n . Since $\{0\}$ is the only proper ideal of M_n , we conclude that $\mathcal{I}(x) = \{0\}$

for $x \in A$ and $\mathcal{J}(x) = M_n$ for $x \in X \setminus A$. This shows that condition (AX0) of Theorem 2.2 is satisfied. Further, if x and y are arbitrary points of X , then either

- $x, y \in A$; in that case (AX2) is fulfilled; or
- $x \in A$ and $y \notin A$ (or conversely); in that case there is $f \in \mathcal{J}$ such that $f(y) = I$, and $f(x) = 0$ (since $x \in A$)—this implies that x and y are spectrally separated by \mathcal{J} ; or
- $x, y \notin A$ and $y = u.x$ for some $u \in \mathfrak{U}_n$; in that case (AX2) is fulfilled since for any selfadjoint $f \in \mathcal{J}$, $f(y) = u.f(x)$ and consequently $\sigma(f(x)) = \sigma(f(y))$; or
- $x, y \notin A$ and $y \notin \mathfrak{U}_n.\{x\}$; in that case there are $f_1 \in \mathcal{J}$ and $f_2 \in C^*(X, \cdot)$ such that $f_1(x) = I = f_2(x)$ and $f_2(y) = 0$ (cf. point (e) of Lemma 3.2), then $f = f_1 f_2 \in \mathcal{J}$ is such that $f(x) = I$ and $f(y) = 0$ and hence x and y are spectrally separated by \mathcal{J} .

Now according to Theorem 2.2, it suffices to check that $\mathcal{J}_A \subset \Delta_2(\mathcal{J})$ (since \mathcal{J} is closed). To this end, we fix $f \in \mathcal{J}_A$ and two arbitrary points x and y of X . We consider similar cases as above:

- (1°) If $x, y \in A$, we have nothing to do because then $f(x) = f(y) = 0$.
- (2°) If $x \in A$ and $y \notin A$ (or conversely), then there is $g \in \mathcal{J}$ such that $g(y) = f(y)$. But also $g(x) = 0 = f(x)$ and we are done.
- (3°) If $x, y \notin A$ and $y = u.x$ for some $u \in \mathfrak{U}_n$, then there is $g \in \mathcal{J}$ with $g(x) = f(x)$. Then also $g(y) = g(u.x) = u.g(x) = u.f(x) = f(y)$ and we are done.
- (4°) Finally, if $x, y \notin A$ and $y \notin \mathfrak{U}_n.\{x\}$, there are functions $g_1, g_2 \in \mathcal{J}$ and $h_1, h_2 \in C^*(X, \cdot)$ such that $g_1(x) = f(x)$, $g_2(y) = f(y)$, $h_1(x) = I = h_2(y)$ and $h_1(y) = 0 = h_2(x)$. Then $g = g_1 h_1 + g_2 h_2 \in \mathcal{J}$ satisfies $g(z) = f(z)$ for $z \in \{x, y\}$.

The arguments (1°)–(4°) show that $f \in \Delta_2(\mathcal{J})$ and thus $\mathcal{J} = \mathcal{J}_A$. It follows from point (c) of Lemma 3.2 that the $*$ -homomorphism $C^*(X, \cdot) \ni f \mapsto f|_A \in C^*(A, \cdot)$ is surjective. What is more, its kernel coincides with $\mathcal{J}_A = \mathcal{J}$ and therefore $C^*(X, \cdot)$ and $C^*(A, \cdot)$ are isomorphic.

We now turn to (b). We infer from (a) that there is a closed invariant set $A \subset X$ such that $\ker(\pi) = \mathcal{J}_A$. Since π is nonzero, A is nonempty. Further, $k^2 \geq \dim \pi(C^*(X, \cdot)) = \dim(C^*(X, \cdot) / \ker(\pi)) = \dim C^*(A, \cdot) \geq n^2$ (by point (d) of Lemma 3.2 and by (a)) and thus $k = n$, $\dim C^*(A, \cdot) = n^2$ and π is surjective. Fix $a \in A$ and observe that $A = \mathfrak{U}_n.\{a\}$, because otherwise $\dim C^*(A, \cdot) > n^2$ (thanks to point (e) of Lemma 3.2). Now define $\Phi: M_n \rightarrow M_n$ by the rule $\Phi(X) = f(a)$ where $\pi(f) = X$. It may easily be checked (using the fact that $\ker(\pi) = \mathcal{J}_{\mathfrak{U}_n.\{a\}}$) that Φ is a well defined one-to-one $*$ -homomorphism of M_n . We conclude that there is $u \in \mathfrak{U}_n$ for which $\Phi(X) = u.X$ (in the algebra of matrices this is quite an elementary fact; however, this follows also from [23, Corollary 2.9.32]). Put $x = u^{-1}.a$ and note that then $f(a) = \Phi(\pi(f)) = u.\pi(f)$ and consequently $\pi(f) = u^{-1}.f(a) = f(x)$ for each $f \in C^*(X, \cdot)$. The uniqueness of x follows from points (d) and (e) of Lemma 3.2.

We turn to (c). Let $\Phi: C^*(X, \cdot) \rightarrow C^*(Y, *)$ be a $*$ -homomorphism of C^* -algebras. Put $U = Y \setminus \{y \in Y : [\Phi(f)](y) = 0 \text{ for each } f \in C^*(X, \cdot)\}$. It is clear that U is invariant and open in Y . For any $y \in U$ the function $C^*(X, \cdot) \ni f \mapsto [\Phi(f)](y) \in M_n$ is a nonzero representation and therefore, thanks to (b), there is a unique point $\varphi(y) \in X$ such that $[\Phi(f)](y) = f(\varphi(y))$ for each $f \in C^*(X, \cdot)$. In this way we have obtained a function $\varphi: U \rightarrow X$ for which (3-1) holds. By the uniqueness in (b), we see that $\varphi(u.y) = u.\varphi(y)$ for any $u \in \mathfrak{U}_n$ and $y \in U$. So, to prove that φ is a morphism of n -spaces, it remains to check that φ is a proper map. First we shall show that φ is continuous. Suppose, for the contrary, that there is a set $D \subset U$ and a point $b \in U \cap \bar{D}$ (\bar{D} is the closure of D in Y) such that $a := \varphi(b) \notin \overline{\varphi(D)}$ (the closure taken in X). Let V be an open neighbourhood

of a whose closure is compact and disjoint from $F := \overline{\varphi(D)}$. Let $\langle \cdot, - \rangle$ be the standard inner product on M_n , that is, $\langle X, Y \rangle = \text{tr}(Y^*X)$ ('tr' is the trace) and let $\|X\|_2 := \sqrt{\text{tr}(X^*X)}$. Take an irreducible matrix $Q \in M_n$ with $\|Q\|_2 = 1$. For simplicity, put $\mathcal{B} = \{X \in M_n : \|X\|_2 \leq 1\}$. Our aim is to construct $f \in C^*(X, \cdot)$ such that $f(a) = Q$ and $f^{-1}(\{Q\}) \subset V$. Observe that there is a compact convex nonempty set \mathcal{K} such that

$$(3-2) \quad Q \notin \mathcal{K} \subset \mathcal{B} \quad \text{and} \quad \{u.a : u \in \mathfrak{U}_n, u.Q \notin \mathcal{K}\} \subset V.$$

(Indeed, it suffices to define \mathcal{K} as the convex hull of the set $\{X \in \mathcal{B} : \|X - Q\|_2 \geq r\}$ where $r > 0$ is such that $u.a \in V$ whenever $u \in \mathfrak{U}_n$ satisfies $\|u.Q - Q\|_2 < r$. Such r exists because Q is irreducible and hence the maps $\mathfrak{U}_n \ni u \mapsto u.b \in X$ and $\mathfrak{U}_n \ni u \mapsto u.Q \in M_n$ are embeddings.) Let $W = \mathfrak{U}_n.a$ and let $g_0 : W \rightarrow M_n$ be given by $g_0(u.a) = u.Q$. Since $g_0(W \setminus V) \subset \mathcal{K}$ (by (3-2)) and the set \mathcal{K} (being compact, convex and nonempty) is a retract of M_n , there is a map $g_1 \in \mathcal{C}_0(X \setminus V, M_n)$ such that $g_1(X \setminus V) \subset \mathcal{K}$ and $g_1(x) = g_0(x)$ for $x \in W \setminus V$. Finally, there is $g \in \mathcal{C}_0(X, M_n)$ which extends both g_0 and g_1 , and $g(X) \subset \mathcal{B}$. Now put $f = g^{\mathfrak{U}} \in C^*(X, \cdot)$ and notice that $f(a) = Q$ (by point (b) of Lemma 3.2). We claim that

$$(3-3) \quad f^{-1}(\{Q\}) \subset V.$$

Let us prove the above relation. Let $x \in X \setminus V$. Then $g(x) = g_1(x) \in \mathcal{K}$ and hence $g(x) \neq Q$ (see (3-2)). The set $\mathfrak{G} := \{u \in \mathfrak{U}_n : u^{-1}.g(u.x) \neq Q\}$ is open in \mathfrak{U}_n and nonempty, which implies that its Haar measure is positive. Further, $|\langle u^{-1}.g(u.x), Q \rangle| \leq 1$ for any $u \in \mathfrak{U}_n$ and $\langle u^{-1}.g(u.x), Q \rangle \neq 1$ for $u \in \mathfrak{G}$ (since $g(X) \subset \mathcal{B}$). We infer from these remarks that $\int_{\mathfrak{U}_n} \langle u^{-1}.g(u.x), Q \rangle du \neq 1$. Equivalently, $\langle f(x), Q \rangle \neq 1$, which implies that $f(x) \neq Q$ and finishes the proof of (3-3). For $m \geq 1$ let $C_m = \{y \in Y : \|[\Phi(f)](y) - Q\|_2 \leq 2^{-m}\}$ and $F_m = \{x \in X : \|f(x) - Q\|_2 \leq 2^{-m}\}$. Since $f \in \mathcal{C}_0(X, M_n)$ and $\Phi(f) \in \mathcal{C}_0(Y, M_n)$, F_m is compact and C_m is a compact neighbourhood of b . Consequently, $C_m \cap D \neq \emptyset$. We infer from (3-1) that $\varphi(C_m \cap D) \subset F_m \cap F$. Now the compactness argument gives $F \cap \bigcap_{m=1}^{\infty} F_m \neq \emptyset$. Let c belong to this intersection. Then $f(c) = Q$ and $c \notin V$, which contradicts (3-3) and finishes the proof of the continuity of φ .

To see that φ is proper, take a compact set $K \subset X$ and note that $L = \mathfrak{U}_n.K$ is compact as well. Let $G \subset X$ be an open neighbourhood of L with compact closure. Take a map $\beta \in \mathcal{C}_0(X, M_n)$ such that $\beta(x) = I$ for $x \in L$ and β vanishes off G . Let $f = \beta^{\mathfrak{U}} \in C^*(X, \cdot)$ and observe that $f(x) = I$ for $x \in L$. Since $\Phi(f) \in \mathcal{C}_0(Y, M_n)$, the set $Z := \{y \in Y : [\Phi(f)](y) = I\}$ is a compact subset of Y . But (3-1) implies that $Z \subset U$ and $\varphi^{-1}(K) \subset Z$. This finishes the proof of the fact that φ is a morphism. The uniqueness of the pair (U, φ) follows from Lemma 3.2 and is left to the reader.

Now if Φ is a $*$ -isomorphism of C^* -algebras, then $U = Y$ (by point (d) of Lemma 3.2) and thus $\Phi(f) = f \circ \varphi$. Similarly, Φ^{-1} is of the form $\Phi^{-1}(g) = g \circ \psi$ for some morphism $\psi : (X, \cdot) \rightarrow (Y, *)$. Then $f = f \circ (\varphi \circ \psi)$ for each $f \in C^*(X, \cdot)$ and the uniqueness in (c) gives $(\varphi \circ \psi)(x) = x$ for each $x \in X$. Similarly, $(\psi \circ \varphi)(y) = y$ for any $y \in Y$ and consequently φ is an isomorphism of n -spaces. The proof is complete. \square

4. REPRESENTATIONS OF $C^*(X, \cdot)$

In this section we will characterize all representations of $C^*(X, \cdot)$ for an arbitrary n -space (X, \cdot) . But first we shall give a 'canonical' description of all continuous linear functionals on $C^*(X, \cdot)$. We underline here that we are not interested in the formula for the norm of a functional. The results of the section will be applied in the next two parts where we formulate our version of Fell's characterization of

homogeneous C^* -algebras (Section 5) and a counterpart of the spectral theorem for finite systems of operators which generate n -homogeneous C^* -algebras (Section 6).

4.1. Definition. Let (X, \cdot) be an n -space. Let $\mathfrak{B}(X)$ denote the σ -algebra of all Borel subsets of X ; that is, $\mathfrak{B}(X)$ is the smallest σ -algebra of subsets of X which contains all open sets. For any $\mathbf{u} \in \mathfrak{U}_n$ and $A \in \mathfrak{B}(X)$ the set $\mathbf{u}.A := \{\mathbf{u}.a : a \in A\}$ is Borel as well. We shall denote by $\chi_A : X \rightarrow \{0, 1\}$ the characteristic function of A . Further, $\mathfrak{B}C^*(X, \cdot)$ stands for the C^* -algebra of all bounded Borel (i.e. $\mathfrak{B}(X)$ -measurable) functions $f : X \rightarrow M_n$ such that $f(\mathbf{u}.x) = \mathbf{u}.f(x)$ for any $\mathbf{u} \in \mathfrak{U}_n$ and $x \in X$.

An n -measure on (X, \cdot) is an $n \times n$ -matrix $\mu = [\mu_{jk}]$ where $\mu_{jk} : \mathfrak{B}(X) \rightarrow \mathbb{C}$ is a regular (complex-valued) measure and $\mu(\mathbf{u}.A) = \mathbf{u}.\mu(A)$ for any $\mathbf{u} \in \mathfrak{U}_n$ and $A \in \mathfrak{B}(X)$ (here, of course, $\mu(A) = [\mu_{jk}(A)] \in M_n$). The set of all n -measures on (X, \cdot) is denoted by $\mathcal{M}(X, \cdot)$.

For any bounded Borel function $f : X \rightarrow M_n$ and an $n \times n$ -matrix $\mu = [\mu_{jk}]$ of complex-valued regular Borel measures we define the integral $\int f d\mu$ as the complex number $\sum_{j,k} \int_X f_{jk} d\mu_{kj}$ where $f(x) = [f_{jk}(x)]$ for $x \in X$. We underline that in the formula for $\int f d\mu$, f_{jk} meets μ_{kj} (not μ_{jk} [!]).

The first purpose of this section is to prove the following

4.2. Theorem. For every continuous linear functional $\varphi : C^*(X, \cdot) \rightarrow \mathbb{C}$ there exists a unique $\mu \in \mathcal{M}(X, \cdot)$ such that $\varphi(f) = \int f d\mu$ for any $f \in C^*(X, \cdot)$.

The above result is a simple consequence of the next one.

4.3. Proposition. Let $\mu = [\mu_{jk}]$ be an $n \times n$ -matrix of complex-valued regular Borel measures on X . Then $\mu \in \mathcal{M}(X, \cdot)$ iff for every map $f \in \mathcal{C}_0(X, M_n)$,

$$(4-1) \quad \int f d\mu = \int f^\mu d\mu.$$

Proof. For any $n \times n$ -matrix A we shall write A_{jk} to denote the suitable entry of A . We adapt the same rule for functions $f \in \mathcal{C}_0(X, M_n)$ and matrix-valued measures. Further, for arbitrarily fixed two indices (j, k) and (p, q) , the function $\mathfrak{U}_n \ni \mathbf{u} \mapsto \mathbf{u}_{jk}\bar{\mathbf{u}}_{pq} \in \mathbb{C}$ is well defined and continuous (although ' \mathbf{u}_{jk} ' is not well defined). Observe that for any $A \in M_n$, $\mathbf{u} \in \mathfrak{U}_n$ and an index (p, q) one has:

$$(\mathbf{u}.A)_{p,q} = \sum_{j,k} \mathbf{u}_{pj}\bar{\mathbf{u}}_{qk} \cdot A_{jk} \quad \text{and} \quad (\mathbf{u}^{-1}.A)_{p,q} = \sum_{j,k} \mathbf{u}_{kq}\bar{\mathbf{u}}_{jp} \cdot A_{jk}.$$

Further, for $\mathbf{u} \in \mathfrak{U}_n$ and a complex-valued regular Borel measure ν on X let $\nu^\mathbf{u}$ be the (complex-valued regular Borel) measure on X given by $\nu^\mathbf{u}(A) = \nu(\mathbf{u}.A)$ ($A \in \mathfrak{B}(X)$). It follows from the transport measure theorem that for any $g \in \mathcal{C}_0(X, \mathbb{C})$:

$$\int_X g(\mathbf{u}.x) d\nu^\mathbf{u}(x) = \int_X g(x) d\nu(x).$$

We adapt the above notation also for $n \times n$ -matrix μ of measures: $\mu^\mathbf{u}(A) = \mu(\mathbf{u}.A)$. Notice that $(\mu^\mathbf{u})_{jk} = (\mu_{jk})^\mathbf{u}$.

Now assume that $\mu \in \mathcal{M}(X, \cdot)$. This means that for any $\mathbf{u} \in \mathfrak{U}_n$, $\mathbf{u}.\mu = \mu^\mathbf{u}$. For $f \in \mathcal{C}_0(X, M_n)$ and $x \in X$ we have

$$(f^\mu)_{pq}(x) = \sum_{j,k} \int_{\mathfrak{U}_n} \mathbf{u}_{kq}\bar{\mathbf{u}}_{jp} \cdot f_{jk}(\mathbf{u}.x) d\mathbf{u}$$

and therefore, by Fubini's theorem,

$$\begin{aligned}
\int f^{\mathfrak{U}} d\mu &= \sum_{p,q} \int_X (f^{\mathfrak{U}})_{p,q} d\mu_{qp} = \sum_{p,q} \sum_{j,k} \int_X \int_{\mathfrak{U}_n} \mathfrak{u}_{kq} \bar{\mathfrak{u}}_{jp} \cdot f_{jk}(\mathfrak{u}.x) d\mathfrak{u} d\mu_{qp}(x) \\
&= \sum_{j,k} \int_{\mathfrak{U}_n} \int_X f_{jk}(\mathfrak{u}.x) d\left(\sum_{p,q} \mathfrak{u}_{kq} \bar{\mathfrak{u}}_{jp} \cdot \mu_{qp}\right)(x) d\mathfrak{u} \\
&= \sum_{j,k} \int_{\mathfrak{U}_n} \int_X f_{jk}(\mathfrak{u}.x) d(\mathfrak{u}.\mu)_{kj}(x) d\mathfrak{u} = \sum_{j,k} \int_{\mathfrak{U}_n} \int_X f_{jk}(\mathfrak{u}.x) d(\mu_{kj})^{\mathfrak{u}}(x) d\mathfrak{u} \\
&= \sum_{j,k} \int_{\mathfrak{U}_n} \int_X f_{jk}(x) d\mu_{kj}(x) d\mathfrak{u} = \sum_{j,k} \int_X f_{jk}(x) d\mu_{kj}(x) = \int f d\mu,
\end{aligned}$$

which gives (4-1). Conversely, assume (4-1) if fulfilled for any $f \in \mathcal{C}_0(X, M_n)$ and fix a compact \mathcal{G}_δ subset K of X and an index (p, q) . Let $g \in \mathcal{C}_0(X, \mathbb{C})$ be arbitrary and let $f \in \mathcal{C}_0(X, M_n)$ be such that $f_{pq} = g$ and $f_{jk} = 0$ for $(j, k) \neq (p, q)$. Applying (4-1) for such f , we obtain

$$(4-2) \quad \int_X g d\mu_{qp} = \sum_{j,k} \int_X \int_{\mathfrak{U}_n} \mathfrak{u}_{qk} \bar{\mathfrak{u}}_{pj} \cdot g(\mathfrak{u}.x) d\mathfrak{u} d\mu_{kj}(x).$$

Further, since K is compact and \mathcal{G}_δ , there is a sequence $(g_k)_{k=1}^\infty \subset \mathcal{C}_0(X, \mathbb{C})$ such that $g_k(X) \subset [0, 1]$ and $\lim_{k \rightarrow \infty} g_k(x) = \chi_K(x)$ for any $x \in X$. Substituting $g = g_k$ in (4-2) and letting $k \rightarrow \infty$, we obtain (by Lebesgue's dominated convergence theorem and Fubini's one):

$$\begin{aligned}
\mu_{q,p}(K) &= \sum_{j,k} \int_{\mathfrak{U}_n} \int_X \mathfrak{u}_{qk} \bar{\mathfrak{u}}_{pj} \cdot \chi_K(\mathfrak{u}.x) d\mu_{kj}(x) d\mathfrak{u} \\
&= \int_{\mathfrak{U}_n} \left(\sum_{j,k} \mathfrak{u}_{qk} \bar{\mathfrak{u}}_{pj} \cdot \mu_{kj}(\mathfrak{u}^{-1}.K) \right) d\mathfrak{u} = \int_{\mathfrak{U}_n} (\mathfrak{u}.\mu)_{q,p}(\mathfrak{u}^{-1}.K) d\mathfrak{u}.
\end{aligned}$$

We infer from the arbitrariness of (p, q) in the above formula that

$$\mu(K) = \int_{\mathfrak{U}_n} \mathfrak{u}.\mu(\mathfrak{u}^{-1}.K) d\mathfrak{u}.$$

Now if $\mathfrak{v} \in \mathfrak{U}_n$, the set $\mathfrak{v}.K$ is also compact and \mathcal{G}_δ and therefore

$$\begin{aligned}
\mu(\mathfrak{v}.K) &= \int_{\mathfrak{U}_n} \mathfrak{u}.\mu(\mathfrak{u}^{-1}\mathfrak{v}.K) d\mathfrak{u} = \int_{\mathfrak{U}_n} \mathfrak{v}[\mathfrak{u}.\mu(\mathfrak{u}^{-1}.K)] d\mathfrak{u} \\
&= \mathfrak{v} \cdot \left(\int_{\mathfrak{U}_n} \mathfrak{u}.\mu(\mathfrak{u}^{-1}.K) d\mathfrak{u} \right) = \mathfrak{v}.\mu(K).
\end{aligned}$$

Finally, since μ is regular, the relation $\mu(\mathfrak{v}.A) = \mathfrak{v}.\mu(A)$ holds for any $A \in \mathfrak{B}(X)$ and we are done. \square

Proof of Theorem 4.2. Note that the function $P: \mathcal{C}_0(X, M_n) \ni f \mapsto f^{\mathfrak{U}} \in C^*(X, \cdot)$ is a continuous linear projection (that is, $P(f) = f$ for $f \in C^*(X, \cdot)$). So, if $\varphi: C^*(X, \cdot) \rightarrow \mathbb{C}$ is a continuous linear functional, so is $\psi := \varphi \circ P: \mathcal{C}_0(X, M_n) \rightarrow \mathbb{C}$. Since $\mathcal{C}_0(X, M_n)$ is isomorphic, as a Banach space, to $[\mathcal{C}_0(X, \mathbb{C})]^n$, the Riesz-type representation theorem yields that there is a unique $n \times n$ -matrix μ of complex-valued regular Borel measures such that $\psi(f) = \int f d\mu$. Observe that $\psi(f^{\mathfrak{U}}) = \psi(f)$ for any $f \in \mathcal{C}_0(X, M_n)$ and hence $\mu \in \mathcal{M}(X, \cdot)$, thanks to Proposition 4.3. The uniqueness of μ follows from the above construction, Proposition 4.3 and the uniqueness in the Riesz-type representation theorem. \square

Now we turn to representations of $C^*(X, \cdot)$. To this end, we introduce

4.4. Definition. An operator-valued n -measure on the n -space (X, \cdot) is any function of the form $E: \mathfrak{B}(X) \ni A \mapsto [E_{jk}(A)] \in M_n(\mathcal{B}(\mathcal{H}))$ (where $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ is a Hilbert space) such that:

- (M1) for any $h, w \in \mathcal{H}$ and $j, k \in \{1, \dots, n\}$, the function $E_{jk}^{(h, w)}: \mathfrak{B}(X) \ni A \mapsto \langle E_{jk}(A)h, w \rangle \in \mathbb{C}$ is a (complex-valued) measure,
- (M2) for any $u \in \mathfrak{U}_n$ and $A \in \mathfrak{B}(X)$, $E(u.A) = u.E(A)$.

In other words, an operator-valued n -measure is an $n \times n$ -matrix of operator-valued measures which satisfies axiom (M2). The operator-valued n -measure E is *regular* iff $E_{jk}^{(h, w)}$ is regular for any h, w and j, k .

Recall that if $\mu: \mathfrak{B}(X) \rightarrow \mathcal{B}(\mathcal{H})$ is an operator-valued measure and $f: X \rightarrow \mathbb{C}$ is a bounded Borel function, $\int_X f d\mu$ is a bounded linear operator on \mathcal{H} , defined by an implicit formula:

$$\left\langle \left(\int_X f d\mu \right) h, w \right\rangle = \int_X f d\mu^{(h, w)} \quad (h, w \in \mathcal{H}),$$

where $\mu^{(h, w)}(A) = \langle \mu(A)h, w \rangle$ ($A \in \mathfrak{B}(X)$). Now assume that $E = [E_{jk}]: \mathfrak{B}(X) \rightarrow M_n(\mathcal{B}(\mathcal{H}))$ is an n -measure and $f = [f_{jk}]: X \rightarrow M_n$ is a bounded Borel function. We define $\int f dE$ as a bounded linear operator on \mathcal{H} given by $\int f dE = \sum_{j, k} \int_X f_{jk} dE_{kj}$. We are now ready to introduce

4.5. Definition. A *spectral* n -measure is any operator-valued regular n -measure $E: \mathfrak{B}(X) \rightarrow M_n(\mathcal{B}(\mathcal{H}))$ such that

$$(4-3) \quad \left(\int f dE \right)^* = \int f^* dE,$$

$$(4-4) \quad \int f \cdot g dE = \int f dE \cdot \int g dE$$

for any $f, g \in \mathfrak{B}C^*(X, \cdot)$. (The product $f \cdot g$ is computed pointwise as the product of matrices.) In other words, a spectral n -measure is an operator-valued regular n -measure $E: \mathfrak{B}(X) \rightarrow M_n(\mathcal{B}(\mathcal{H}))$ such that the operator

$$\mathfrak{B}C^*(X, \cdot) \ni f \mapsto \int f dE \in \mathcal{B}(\mathcal{H})$$

is a representation of a C^* -algebra $\mathfrak{B}C^*(X, \cdot)$.

The main result of this section is the following

4.6. Theorem. Let (X, \cdot) be an n -space and $\pi: C^*(X, \cdot) \rightarrow \mathcal{B}(\mathcal{H})$ be a representation. There is a unique spectral n -measure $E: \mathfrak{B}(X) \rightarrow M_n(\mathcal{B}(\mathcal{H}))$ such that

$$(4-5) \quad \pi(f) = \int f dE \quad (f \in C^*(X, \cdot)).$$

In particular, every representation of $C^*(X, \cdot)$ admits an extension to a representation of $\mathfrak{B}C^*(X, \cdot)$.

In the proof of the above result we shall involve the following

4.7. Lemma. Let $\mu: \mathfrak{B}(X) \rightarrow \mathbb{R}_+$ be a regular measure. For any $f \in \mathfrak{B}C^*(X, \cdot)$ and $\varepsilon > 0$ there exists $g \in C^*(X, \cdot)$ such that $\sup_{x \in X} \|g(x)\| \leq \sup_{x \in X} \|f(x)\|$ and $\int_X \|f(x) - g(x)\| d\mu(x) < \varepsilon$.

Proof. Let $f = [f_{jk}] \in \mathfrak{B}C^*(X, \cdot)$ and let $M > 0$ be such that

$$\sup_{x \in X} \|f(x)\| \leq M.$$

It follows from the regularity of μ that for each (j, k) there is a compact set L_{jk} such that $\mu(X \setminus L_{jk}) \leq \frac{\varepsilon}{2Mn^2}$ and $f_{jk}|_{L_{jk}}$ is continuous. Put $L = \bigcap_{j, k} L_{jk}$ and

$K = \mathfrak{U}_n.L$. Then K is compact and invariant, and $\mu(X \setminus K) \leq \frac{\varepsilon}{2M}$. What is more, $f|_K$ is continuous (this follows from the facts that $f|_L$ is continuous and $f(\mathbf{u}.x) = \mathbf{u}.f(x)$). Now point (c) of Lemma 3.2 yields the existence of $g \in C^*(X, \cdot)$ such that $\sup_{x \in X} \|g(x)\| \leq \sup_{x \in X} \|f(x)\|$ and $g|_K = f|_K$. Then:

$$\int_X \|f(x) - g(x)\| d\mu(x) = \int_{X \setminus K} \|f(x) - g(x)\| d\mu \leq 2M \cdot \mu(X \setminus K) = \varepsilon$$

and we are done. \square

4.8. Proposition. *Let $E = [E_{jk}]: \mathfrak{B}(X) \rightarrow M_n(\mathcal{B}(\mathcal{H}))$ be a regular n -measure.*

- (a) *E satisfies (4-3) for any $f \in \mathfrak{B}C^*(X, \cdot)$ iff (4-3) is fulfilled for any $f \in C^*(X, \cdot)$, iff $(E_{jk}(A))^* = E_{kj}(A)$ for each $A \in \mathfrak{B}(X)$,*
- (b) *E is spectral iff (4-3) and (4-4) are satisfied for any $f, g \in C^*(X, \cdot)$.*

Proof. For any complex-valued regular Borel measure ν on X we shall denote by $|\nu|$ the variation of ν . Recall that $|\nu|$ is a nonnegative finite regular Borel measure on X . Further, for any $h, w \in \mathcal{H}$ and $j, k \in \{1, \dots, n\}$ let $E_{jk}^{(h,w)}$ be as in Definition 4.4. Finally, $\langle \cdot, - \rangle$ stands for the scalar product of \mathcal{H} .

We begin with (a). Fix $h, w \in \mathcal{H}$ and $j, k \in \{1, \dots, n\}$. First assume that (4-3) is fulfilled for any $f \in C^*(X, \cdot)$. Let $E^{(h,w)} := [E_{jk}^{(h,w)}]$ and note that $E^{(h,w)} \in \mathcal{M}(X, \cdot)$, since $E_{pq}(\mathbf{u}.A) = \sum_{j,k} \mathbf{u}_{pj} \bar{\mathbf{u}}_{qk} \cdot E_{jk}(A)$ and thus $E_{pq}^{(h,w)}(\mathbf{u}.A) = \sum_{j,k} \mathbf{u}_{pj} \bar{\mathbf{u}}_{qk} \cdot E_{jk}^{(h,w)}(A) = (\mathbf{u}.E^{(h,w)}(A))_{pq}$. Observe that $(E^{(h,w)})^* \in \mathcal{M}(X, \cdot)$ as well where $(E^{(h,w)})^*(A) = (E^{(h,w)}(A))^*$ (because $(\mathbf{u}.P)^* = \mathbf{u}.P^*$ for any $P \in M_n$). Further, for each $f \in C^*(X, \cdot)$ we have

$$\begin{aligned} \overline{\int f^* dE^{(h,w)}} &= \sum_{j,k} \overline{\int_X (f^*)_{jk} dE_{kj}^{(h,w)}} = \sum_{j,k} \int_X f_{kj} d\overline{E_{kj}^{(h,w)}} \\ &= \sum_{j,k} \int_X f_{kj} d(E^{(h,w)})_{jk}^* = \int f d(E^{(h,w)})^* \end{aligned}$$

and, on the other hand,

$$\begin{aligned} \overline{\int f^* dE^{(h,w)}} &= \overline{\left\langle \left(\int f^* dE \right) h, w \right\rangle} = \overline{\left\langle \left(\int f dE \right)^* h, w \right\rangle} \\ &= \left\langle \left(\int f dE \right) w, h \right\rangle = \int f dE^{(w,h)}. \end{aligned}$$

The uniqueness in Theorem 4.2 implies that $(E^{(h,w)})^* = E^{(w,h)}$, which means that for each $A \in \mathfrak{B}(X)$, $\langle (E_{jk}(A))w, h \rangle = \overline{\langle (E_{kj}(A))h, w \rangle} = \langle (E_{kj}(A))^*w, h \rangle$. We conclude that $(E_{jk}(A))^* = E_{kj}(A)$. Finally, if the last relation holds for any $j, k \in \{1, \dots, n\}$, then for every $f \in \mathfrak{B}C^*(X, \cdot)$ we get

$$\begin{aligned} \left(\int f dE \right)^* &= \sum_{j,k} \left(\int_X f_{jk} dE_{kj} \right)^* = \sum_{j,k} \int_X \bar{f}_{jk} d(E_{kj})^* \\ &= \sum_{j,k} \int_X (f^*)_{kj} dE_{jk} = \int f^* dE. \end{aligned}$$

This completes the proof of (a). We now turn to (b). We assume that (4-3) and (4-4) are fulfilled for any $f, g \in C^*(X, \cdot)$. We know from (a) that actually (4-3) is satisfied for any $f \in \mathfrak{B}C^*(X, \cdot)$. The proof of (4-4) is divided into three steps stated below.

Step 1. If $\xi \in \mathfrak{B}C^*(X, \cdot)$ is such that

$$(4-6) \quad \int g \cdot \xi \, dE = \int g \, dE \cdot \int \xi \, dE$$

for any $g \in C^*(X, \cdot)$, then $\int f \cdot \xi \, dE = \int f \, dE \cdot \int \xi \, dE$ for any $f \in \mathfrak{B}C^*(X, \cdot)$.

Proof of step 1. Fix $f \in \mathfrak{B}C^*(X, \cdot)$, $h, w \in \mathcal{H}$ and $\varepsilon > 0$. Let $M \geq 1$ be such that $\sup_{x \in X} \|\xi(x)\| \leq M$. Put $v = (\int \xi \, dE)h$ and $\mu = \sum_{j,k} (|E_{jk}^{(h,w)}| + |E_{jk}^{(v,w)}|)$. Since μ is finite and regular, Lemma 4.7 gives us a map $g \in C^*(X, \cdot)$ such that $\int_X \|f(x) - g(x)\| \, d\mu(x) \leq \frac{\varepsilon}{M}$. Then (4-6) holds and therefore (remember that $M \geq 1$):

$$\begin{aligned} & \left| \left\langle \left(\int f \cdot \xi \, dE - \int f \, dE \cdot \int \xi \, dE \right) h, w \right\rangle \right| \leq \left| \left\langle \left(\int f \cdot \xi \, dE - \int g \cdot \xi \, dE \right) h, w \right\rangle \right| \\ & \quad + \left| \left\langle \left(\int g \, dE \cdot \int \xi \, dE - \int f \, dE \cdot \int \xi \, dE \right) h, w \right\rangle \right| \\ & = \left| \sum_{j,k} \int_X ((f - g)\xi)_{jk} \, dE_{kj}^{(h,w)} \right| + \left| \sum_{j,k} \int_X (g_{jk} - f_{jk}) \, dE_{kj}^{(v,w)} \right| \\ & \leq \sum_{j,k} \int_X \|(f(x) - g(x))\xi(x)\| \, d|E_{kj}^{(h,w)}|(x) + \sum_{j,k} \int_X \|g(x) - f(x)\| \, d|E_{kj}^{(v,w)}|(x) \\ & \leq M \int_X \|f(x) - g(x)\| \, d\mu(x) \leq \varepsilon. \end{aligned}$$

Step 2. For any $f \in \mathfrak{B}C^*(X, \cdot)$ and $g \in \mathcal{C}(X, \cdot)$, (4-4) holds.

Proof of step 2. It follows from step 1 and our assumptions in (b) that $\int g^* \cdot f^* \, dE = \int g^* \, dE \cdot \int f^* \, dE$. Now it suffices to apply (4-3):

$$\int f \cdot g \, dE = \left(\int g^* \cdot f^* \, dE \right)^* = \left(\int g^* \, dE \cdot \int f^* \, dE \right)^* = \int f \, dE \cdot \int g \, dE.$$

Step 3. The condition (4-4) is satisfied for any $f, g \in \mathfrak{B}C^*(X, \cdot)$.

Proof of step 3. Just apply step 2 and then step 1. \square

Proof of Theorem 4.6. According to point (b) of Proposition 4.8, it suffices to show that there exists a regular n -measure $E: \mathfrak{B}(X) \rightarrow M_n(\mathcal{B}(\mathcal{H}))$ such that (4-5) holds and that such E is unique. According to Theorem 4.2, for any $h, w \in \mathcal{H}$ there is a unique $\mu^{(h,w)} = [\mu_{jk}^{(h,w)}] \in \mathcal{M}(X, \cdot)$ such that

$$(4-7) \quad \langle \pi(f)h, w \rangle = \int f \, d\mu^{(h,w)}$$

for each $f \in C^*(X, \cdot)$ ($\langle \cdot, \cdot \rangle$ is the scalar product of \mathcal{H}). Now for any $j, k \in \{1, \dots, n\}$ and each $A \in \mathfrak{B}(X)$ there is a unique bounded operator on \mathcal{H} , denoted by $E_{jk}(A)$, for which $\mu_{jk}^{(h,w)}(A) = \langle (E_{jk}(A))h, w \rangle$ ($h, w \in \mathcal{H}$). We put $E(A) = [E_{jk}(A)] \in M_n(\mathcal{B}(\mathcal{H}))$. We want to show that $E(u.A) = u.E(A)$. Since $\mu^{(h,w)} \in \mathcal{M}(X, \cdot)$, we obtain

$$\begin{aligned} \langle (E_{pq}(u.A))h, w \rangle &= (\mu^{(h,w)}(u.A))_{pq} = (u.\mu^{(h,w)}(A))_{pq} = \sum_{j,k} u_{pj} \bar{u}_{qk} \cdot \mu_{jk}^{(h,w)}(A) \\ &= \sum_{j,k} u_{pj} \bar{u}_{qk} \cdot \langle (E_{jk}(A))h, w \rangle = \langle (u.E(A))_{pq}h, w \rangle, \end{aligned}$$

which shows that indeed $E(u.A) = u.E(A)$. Further, observe that $E_{jk}^{(h,w)} = \mu_{jk}^{(h,w)}$ and thus E is an operator-valued regular n -measure and $\langle (\int f \, dE)h, w \rangle = \langle \pi(f)h, w \rangle$ (thanks to (4-7)). Consequently, $\int f \, dE = \pi(f)$ and we are done.

The uniqueness of E follows from the above construction and its proof is left to the reader. \square

4.9. Example. Let (X, \cdot) be an n -space and let $E = [E_{jk}]: \mathfrak{B}(X) \rightarrow M_n(\mathcal{B}(\mathcal{H}))$ be a spectral n -measure. We denote by $\mathfrak{B}_{inv}(X)$ the σ -algebra of all invariant Borel subsets of X (that is, $A \in \mathfrak{B}(X)$ belongs to $\mathfrak{B}_{inv}(X)$ iff $\mathbf{u}.A = A$ for any $\mathbf{u} \in \mathfrak{U}_n$). Let $F: \mathfrak{B}_{inv}(X) \ni A \mapsto \sum_j E_{jj}(A) \in \mathcal{B}(\mathcal{H})$. Then for every $A \in \mathfrak{B}_{inv}(X)$ one has:

$$(E1) \quad E_{jk}(A) = 0 \text{ whenever } j \neq k,$$

$$(E2) \quad E_{11}(A) = \dots = E_{nn}(A) = \frac{1}{n}F(A)$$

and F is a spectral measure (possibly with $F(X) \neq I_{\mathcal{H}}$ where $I_{\mathcal{H}}$ is the identity operator on \mathcal{H}). Let us briefly prove these claims. Since $E(A) = E(\mathbf{u}.A) = \mathbf{u}.E(A)$ for any $\mathbf{u} \in \mathfrak{U}_n$, conditions (E1)–(E2) are fulfilled. Further, if $j_A: X \rightarrow M_n$ is given by $j_A(x) = \chi_A(x) \cdot I$ where $I \in M_n$ is the unit matrix, then $j_A \in \mathfrak{B}C^*(X, \cdot)$ and for $B \in \mathfrak{B}_{inv}(X)$, $F(A \cap B) = \int j_{A \cap B} dE = \int j_A \cdot j_B dE = \int j_A dE \cdot \int j_B dE = F(A)F(B)$. What is more, point (a) of Proposition 4.8 implies that $F(A)$ is selfadjoint and hence F is indeed a spectral measure. One may also easily check that $F(X) = I_{\mathcal{H}}$ iff the representation $\pi_E: C^*(X, \cdot) \ni f \mapsto \int f dE \in \mathcal{B}(\mathcal{H})$ is nondegenerate.

The spectral measure F defined above corresponds to the representation of the center Z of $C^*(X, \cdot)$. It is a simple exercise that Z consists precisely of all $f \in C_0(X, \mathbb{C} \cdot I)$ which are constant on the sets of the form $\mathfrak{U}_n \cdot \{x\}$ ($x \in X$). Thus, $\mathfrak{B}_{inv}(X)$ may naturally be identified with the Borel σ -algebra of the spectrum of Z and consequently F is the spectral measure induced by the representation $\pi_E|_Z$ of Z .

Conditions (E1)–(E2) show that a nonzero spectral n -measure E for $n > 1$ *never* satisfies the condition of a spectral measure — that $E(A \cap B) = E(A)E(B)$. Indeed, $E(X) \neq (E(X))^2$.

The next result is well known. For reader convenience, we give its short proof.

4.10. Lemma. *Let \mathcal{A} be a C^* -algebra and let $\pi: \mathcal{A} \rightarrow M_n$ (where $n \geq 1$ is finite) be a nonzero irreducible representation of \mathcal{A} . Then π is surjective.*

Proof. Let $J = \pi(\mathcal{A})$. Since π is irreducible, $J' = \mathbb{C} \cdot I$ and consequently $J'' = M_n$. But it follows from von Neumann's double commutant theorem that $J'' = J + \mathbb{C} \cdot I$ (here we use the fact that n is finite). So, the facts that J is a $*$ -algebra and $M_n = J + \mathbb{C} \cdot I$ imply that J is a two-sided ideal in M_n . Consequently, $J = \{0\}$ or $J = M_n$. But $\pi \neq 0$ and hence $J = M_n$. \square

With the aid of the above lemma and Theorem 4.6 we shall now characterize all irreducible representations of $C^*(X, \cdot)$.

4.11. Proposition. *Every nonzero irreducible representation π of $C^*(X, \cdot)$ (where (X, \cdot) is an n -space) is n -dimensional and has the form $\pi(f) = f(x)$ (for some $x \in X$).*

Proof. Let $\pi: C^*(X, \cdot) \rightarrow \mathcal{B}(\mathcal{H})$ be a nonzero irreducible representation. It follows from Theorem 4.6 that there is a spectral n -measure $E: \mathfrak{B}(X) \rightarrow \mathcal{B}(\mathcal{H})$ such that (4-5) holds. Let $\mathfrak{B}_{inv}(X)$ and $F: \mathfrak{B}_{inv}(X) \rightarrow \mathcal{B}(\mathcal{H})$ be as in Example 4.9. Then F is a (regular) spectral measure (with $F(X) = I_{\mathcal{H}}$ because π is nondegenerate) and for any $A \in \mathfrak{B}_{inv}(X)$ and $f \in C^*(X, \cdot)$ we have $\int f dE \cdot \int \chi_A I dE = \int \chi_A I dE \cdot \int f dE$ (where I is the unit $n \times n$ -matrix). Since π is irreducible, we deduce that for every $A \in \mathfrak{B}_{inv}(X)$, $\int \chi_A I dE$ is a scalar multiple of the identity operator on \mathcal{H} . This implies that we may think of F as a complex-valued (spectral) measure. But $\mathfrak{B}_{inv}(X)$ is naturally 'isomorphic' to the σ -algebra of all Borel sets of X/\mathfrak{U}_n (which is locally compact) and thus F is supported on a set $S := \mathfrak{U}_n \cdot a$ for some $a \in X$. But then $\int \chi_X I dE = \int \chi_S I dE$ and consequently $\pi(f) = \int f|_S dE_S$ where E_S is

the restriction of E to $\mathfrak{B}(S)$. Since the vector space $\{f|_S: f \in C^*(X, \cdot)\}$ is finite dimensional (and its dimension is equal to n^2), we infer that $\mathcal{A} := \pi(C^*(X, \cdot))$ is finite dimensional as well and $\dim \mathcal{A} \leq n^2$. So, the irreducibility of π implies that \mathcal{H} is finite dimensional, while Lemma 4.10 shows that $\dim \mathcal{H} \leq n$. Finally, point (b) of Proposition 3.3 completes the proof. \square

5. HOMOGENEOUS C^* -ALGEBRAS

5.1. Definition. A C^* -algebra is said to be n -homogeneous (where n is finite) iff every its nonzero irreducible representation is n -dimensional.

Our version of Fell's characterization of n -homogeneous C^* -algebras [9] reads as follows.

5.2. Theorem. For a C^* -algebra \mathcal{A} and finite $n \geq 1$ the following conditions are equivalent:

- (i) \mathcal{A} is an n -homogeneous C^* -algebra;
- (ii) there is an n -space (X, \cdot) such that \mathcal{A} is isomorphic (as a C^* -algebra) to $C^*(X, \cdot)$.

What is more, if \mathcal{A} is n -homogeneous, the n -space (X, \cdot) appearing in (ii) is unique up to isomorphism.

Proof of Theorem 5.2. We infer from point (c) of Proposition 3.3 that the n -space (X, \cdot) appearing in (ii) is unique up to isomorphism. What is more, it easily follows from Proposition 4.11 that $C^*(X, \cdot)$ is n -homogeneous for any n -space (X, \cdot) . So, it remains to show that (i) implies (ii). To this end, assume \mathcal{A} is n -homogeneous and let \mathfrak{X} be the set of all representations (including the zero one) $\pi: \mathcal{A} \rightarrow M_n$, equipped with the topology of pointwise convergence. Since each representation is a bounded linear operator of norm not greater than 1, \mathfrak{X} is compact. Consequently, $X := \mathfrak{X} \setminus \{0\}$ is locally compact. We define an action of \mathfrak{U}_n on X by the formula: $(u.\pi)(a) = u.\pi(a)$ ($a \in \mathcal{A}$, $\pi \in X$, $u \in \mathfrak{U}_n$). It is easily seen that the action is continuous. What is more, Lemma 4.10 ensures us that it is free as well. So, (X, \cdot) is an n -space. The next step of construction is very common. For any $a \in \mathcal{A}$ let $\hat{a}: X \rightarrow M_n$ be given by $\hat{a}(\pi) = \pi(a)$. It is clear that $\hat{a} \in C_0(X, M_n)$ (indeed, if X is noncompact, then $\mathfrak{X} = X \cup \{0\}$ is a one-point compactification of X and \hat{a} extends to a map on \mathfrak{X} which vanishes at 0). We also readily have $\hat{a}(u.\pi) = u.\hat{a}(\pi)$ for any $u \in \mathfrak{U}_n$. So, we have obtained a $*$ -homomorphism $\Phi: \mathcal{A} \ni a \mapsto \hat{a} \in C^*(X, \cdot)$. It follows from (i) (and the fact that all irreducible representations separate points of a C^* -algebra) that Φ is one-to-one and, consequently, Φ is isometric. So, to end the proof, it suffices to show that $\mathcal{E} = \Phi(\mathcal{A})$ is dense in $C^*(X, \cdot)$. To this end, we involve Theorem 2.2. It follows from Lemma 4.10 that condition (AX0) is fulfilled. Further, let π_1 and π_2 be arbitrary members of X . We consider two cases. First assume that $\pi_2 = u.\pi_1$ for some $u \in \mathfrak{U}_n$. Then $\hat{a}(\pi_2) = u.\hat{a}(\pi_1)$ and consequently $\sigma(\hat{a}(\pi_1)) = \sigma(\hat{a}(\pi_2))$ ($a \in \mathcal{A}$). So, in that case (AX2) holds. Now assume that there is no $u \in \mathfrak{U}_n$ for which $\pi_2 = u.\pi_1$. We shall show that in that case:

$$(5-1) \quad \pi_1(a) = 0 \quad \text{and} \quad \pi_2(a) = I \quad \text{for some } a \in \mathcal{A}.$$

Let $\mathcal{M} \subset M_{2n}$ consist of all matrices of the form $\begin{pmatrix} \pi_1(x) & 0 \\ 0 & \pi_2(x) \end{pmatrix}$ with $x \in \mathcal{A}$. Since \mathcal{M} is a finite-dimensional C^* -algebra, it is singly generated (see e.g. [22]) and unital (cf. [26, §1.11]). Thanks to Lemma 4.10, \mathcal{M} contains matrices of the form $\begin{pmatrix} I & 0 \\ 0 & A \end{pmatrix}$ and $\begin{pmatrix} B & 0 \\ 0 & I \end{pmatrix}$ (for some $A, B \in M_n$). We conclude that the unit of

\mathcal{M} coincides with the unit of M_{2n} . This, combined with the fact that \mathcal{M} is singly generated, yields that there is $z \in \mathcal{A}$ such that for $A_j = \pi_j(z)$ ($j = 1, 2$) we have

$$\mathcal{M} = \left\{ \begin{pmatrix} p(A_1, A_1^*) & 0 \\ 0 & p(A_2, A_2^*) \end{pmatrix} : p \in \mathcal{P} \right\}$$

where \mathcal{P} is the free algebra of all polynomials in two noncommuting variables. Observe that then $M_n = \pi_j(\mathcal{A}) = \{p(A_j, A_j^*) : p \in \mathcal{P}\}$ ($j = 1, 2$), which means that A_1 and A_2 are irreducible matrices. What is more, A_1 and A_2 are not unitarily equivalent, that is, there is no $u \in \mathfrak{U}_n$ for which $A_2 = u.A_1$ (indeed, if $A_2 = u.A_1$, then $\pi_2 = u.\pi_1$, since for every $x \in \mathcal{A}$ there is $p \in \mathcal{P}$ such that $\pi_j(x) = p(A_j, A_j^*)$).

These two remarks imply that $\begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix} \in \mathcal{M}$, because the $*$ -commutant in M_{2n} of

the matrix $\begin{pmatrix} A_1 & 0 \\ 0 & A_2 \end{pmatrix}$ consists of matrices of the form $\begin{pmatrix} X & 0 \\ 0 & Y \end{pmatrix}$ (this follows from the so-called Schur's lemma on intertwining transformations—see Theorem 1.5 in [6] and Corollary 1.8 there; cf. also Proposition 5.2.1 in [19]) and consequently $\begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix} \in \mathcal{M}'' = \mathcal{M}$. So, there is $a \in \mathcal{A}$ such that $\begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix} = \begin{pmatrix} \pi_1(a) & 0 \\ 0 & \pi_2(a) \end{pmatrix}$,

which gives (5-1). Replacing a by $\frac{a+a^*}{2}$, we may assume that a is selfadjoint. Then $f = \hat{a} \in \mathcal{E}$ is selfadjoint (and hence normal) and $\sigma(f(\pi_1)) \cap \sigma(f(\pi_2)) = \emptyset$, which shows that π_1 and π_2 are spectrally separated by \mathcal{E} . According to Theorem 2.2, it therefore suffices to check that each $g \in C^*(X, \cdot)$ belongs to $\Delta_2(\mathcal{E})$. To this end, we fix $\pi_1, \pi_2 \in X$ and consider the same two cases as before. If $\pi_2 = u.\pi_1$, it follows from Lemma 4.10 that there is $x \in \mathcal{A}$ for which $\pi_1(x) = g(\pi_1)$. Then $\hat{x}(\pi_1) = g(\pi_1)$ and $\hat{x}(\pi_2) = u.\hat{x}(\pi_1) = u.g(\pi_1) = g(\pi_2)$ and we are done.

Finally, if $\pi_2 \neq u.\pi_1$ for any $u \in \mathfrak{U}_n$, (5-1) implies that there are points a_1 and a_2 in \mathcal{A} such that $\pi_1(a_1) = I = \pi_2(a_2)$ and $\pi_1(a_2) = 0 = \pi_2(a_1)$. Moreover, there are points $x, y \in \mathcal{A}$ such that $\pi_1(x) = g(\pi_1)$ and $\pi_2(y) = g(\pi_2)$ (by Lemma 4.10). Put $z = xa_1 + ya_2 \in \mathcal{A}$ and note that $\hat{z}(\pi_j) = g(\pi_j)$ for $j = 1, 2$, which means that $g \in \Delta_2(\mathcal{E})$. The whole proof is complete. \square

5.3. Definition. Let \mathcal{A} be an n -homogeneous C^* -algebra. By an n -spectrum of \mathcal{A} we mean any n -space (X, \cdot) such that \mathcal{A} is isomorphic to $C^*(X, \cdot)$. It follows from Theorem 5.2 that an n -spectrum of \mathcal{A} is unique up to isomorphism of n -spaces. By *concrete n -spectrum* of \mathcal{A} we mean the n -space of all nonzero representations $\pi : \mathcal{A} \rightarrow M_n$ endowed with the pointwise convergence topology and the natural action of \mathfrak{U}_n .

The trivial algebra $\{0\}$ is n -homogeneous and its n -spectrum is the empty n -space.

The reader interested in general ideas of operator spectra should consult [7, 8, 9]; [4] as well as [12, 13]; [20]; §2.5 in [6]; [15] or [16].

Our approach to n -homogeneous C^* -algebras allows us to prove briefly the following

5.4. Proposition. *Let \mathcal{A}_1 and \mathcal{A}_2 be two n -homogeneous C^* -algebras such that $\mathcal{A}_1 \subset \mathcal{A}_2$.*

- (a) *Every representation $\pi_1 : \mathcal{A}_1 \rightarrow M_n$ is extendable to a representation $\pi_2 : \mathcal{A}_2 \rightarrow M_n$.*
- (b) *If every n -dimensional representation (including the zero one) of \mathcal{A}_1 has a unique extension to an n -dimensional representation of \mathcal{A}_2 , then $\mathcal{A}_1 = \mathcal{A}_2$.*

Proof. We begin with (a). We may and do assume that π_1 is nonzero. For $j = 1, 2$ let (X_j, \cdot) denote an n -spectrum of \mathcal{A}_j and let $\Psi_j : \mathcal{A}_j \rightarrow C^*(X_j, \cdot)$ be a $*$ -isomorphism of C^* -algebras. Let $j : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ be the inclusion map. Then $\Phi :=$

$\Psi_2 \circ j \circ \Psi_1^{-1}: C^*(X_1, \cdot) \rightarrow C^*(X_2, \cdot)$ is a one-to-one $*$ -homomorphism. We infer from Proposition 3.3 that there are an invariant open (in X_2) set U and a morphism $\varphi: (U, \cdot) \rightarrow (X_1, \cdot)$ such that (3-1) holds. We claim that

$$(5-2) \quad \varphi(U) = X_1.$$

Since φ is proper, the set $F := \varphi(U)$ is closed in X_1 . It is also invariant. So, if $F \neq X_1$, we may take $b \in X_1 \setminus F$ and apply point (c) of Lemma 3.2 to get a function $f \in C^*(X_1, \cdot)$ such that $f|_F \equiv 0$ and $f(b) = I$. Then $\Phi(f) = 0$, by (3-1), which contradicts the fact that Φ is one-to-one. So, (5-2) is fulfilled.

Further, Proposition 3.3 yields that there is $x \in X_1$ such that $\pi_1(\Psi_1^{-1}(f)) = f(x)$ for any $f \in C^*(X_1, \cdot)$. It follows from (5-2) that we may find $z \in U$ for which $\varphi(z) = x$. Now define $\pi_2: \mathcal{A}_2 \rightarrow M_n$ by $\pi_2(a) = [\Psi_2(a)](z)$ ($a \in \mathcal{A}_2$). It remains to check that π_2 extends π_1 . To see this, for $a \in \mathcal{A}_1$ put $f = \Psi_1(a)$ and note that $\pi_2(a) = [\Psi_2(a)](z) = [\Psi_2(\Psi_1^{-1}(f))](z) = [\Phi(f)](z) = f(\varphi(z)) = f(x) = \pi_1(a)$ (cf. (3-1)).

Now if the assumption of (b) is satisfied, the above argument shows that φ is one-to-one (since different points of X_2 correspond to different n -dimensional representations of \mathcal{A}_2). It may also easily be checked that for every $z \in X_2 \setminus U$ the representation $\mathcal{A}_2 \ni a \mapsto [\Psi_2(a)](z) \in M_n$ vanishes on \mathcal{A}_1 (use (3-1) and the definition of Φ). So, we conclude from the uniqueness of the extension of the zero representation of \mathcal{A}_1 that $U = X_2$ and hence both φ and Φ are isomorphisms. Consequently, $\mathcal{A}_1 = \mathcal{A}_2$ and we are done. \square

6. SPECTRAL THEOREM AND n -FUNCTIONAL CALCULUS

Whenever \mathcal{A} is a unital C^* -algebra and x_1, \dots, x_k are arbitrary elements of \mathcal{A} , let $C^*(x_1, \dots, x_k)$ denote the C^* -subalgebra of \mathcal{A} generated by x_1, \dots, x_k and let $C_1^*(x_1, \dots, x_k)$ be the smallest C^* -subalgebra of \mathcal{A} which contains x_1, \dots, x_k as well as the unit of \mathcal{A} (so, $C_1^*(x_1, \dots, x_k) = C^*(x_1, \dots, x_k) + \mathbb{C} \cdot 1$ where 1 is the unit of \mathcal{A}). We would like to distinguish those systems (x_1, \dots, x_k) for which one of these two C^* -algebras defined above is n -homogeneous. However, the property of being n -homogeneous is not hereditary for $n > 1$. That is, when $n > 1$, *every* nonzero n -homogeneous C^* -algebra contains a C^* -subalgebra which is not n -homogeneous (namely, a nonzero commutative one). This causes that the class of distinguished systems may depend on the choice of C^* -algebras related to them. Fortunately, this does not happen, which is explained in the following

6.1. Lemma. *Let \mathcal{A} be a unital C^* -algebra and $x_1, \dots, x_k \in \mathcal{A}$. If $C_1^*(x_1, \dots, x_k)$ is n -homogeneous for some $n > 1$, then $C_1^*(x_1, \dots, x_k) = C^*(x_1, \dots, x_k)$.*

Proof. Suppose, for the contrary, that the assertion is false. Observe that $\mathcal{I} := C^*(x_1, \dots, x_k)$ is a two-sided ideal in $\mathcal{B} := C_1^*(x_1, \dots, x_k)$, since $\mathcal{B} = \mathcal{I} + \mathbb{C} \cdot 1$ (1 = the unit of \mathcal{A}). Moreover, \mathcal{B}/\mathcal{I} is isomorphic (as a C^* -algebra) to \mathbb{C} , which means that the canonical projection $\pi: \mathcal{B} \rightarrow \mathcal{B}/\mathcal{I}$ may be considered as a 1-dimensional (nonzero) representation. It is obviously irreducible, which contradicts the fact that \mathcal{B} is n -homogeneous (since $n > 1$). \square

Taking into account the above result, we may now introduce

6.2. Definition. A system (x_1, \dots, x_k) of elements of an (unnecessarily unital) C^* -algebra \mathcal{A} is said to be *n -homogeneous* (where $n \geq 1$ is finite) if the C^* -subalgebra $C^*(x_1, \dots, x_k)$ of \mathcal{A} generated by x_1, \dots, x_k is n -homogeneous.

This part of the paper is devoted to studies of (finite) n -homogeneous systems. We begin with

6.3. Proposition. Let (x_1, \dots, x_k) be an n -homogeneous system in a C^* -algebra A . Let (\mathfrak{X}, \cdot) be the concrete n -spectrum of $C^*(x_1, \dots, x_k)$ and let

$$(6-1) \quad \sigma_n(x_1, \dots, x_k) := \{(\pi(x_1), \dots, \pi(x_k)) : \pi \in \mathfrak{X}\}$$

be equipped with the topology inherited from $(M_n)^k$ and with the action

$$\mathbf{u} \cdot (A_1, \dots, A_k) := (\mathbf{u} \cdot A_1, \dots, \mathbf{u} \cdot A_k) \quad (\mathbf{u} \in \mathfrak{U}_n, (A_1, \dots, A_k) \in \sigma_n(x_1, \dots, x_k)).$$

(Sp1) The pair $(\sigma_n(x_1, \dots, x_k), \cdot)$ is an n -space.

(Sp2) The function $H : (\mathfrak{X}, \cdot) \ni \pi \mapsto (\pi(x_1), \dots, \pi(x_k)) \in (\sigma_n(x_1, \dots, x_k), \cdot)$ is an isomorphism of n -spaces.

(Sp3) Every member of $\sigma_n(x_1, \dots, x_k)$ is irreducible; that is, if $(A_1, \dots, A_k) \in \sigma_n(x_1, \dots, x_k)$ and $T \in M_n$ commutes with each of $A_1, A_1^*, \dots, A_k, A_k^*$, then T is a scalar multiple of the unit matrix.

(Sp4) The set $\sigma_n(x_1, \dots, x_k)$ is either compact or its closure in $(M_n)^k$ coincides with $\sigma_n(x_1, \dots, x_k) \cup \{0\}$.

Proof. Let $\pi_0 : A \rightarrow M_n$ be the zero representation and let $\Omega = \mathfrak{X} \cup \{\pi_0\}$ be equipped with the pointwise convergence topology. Then Ω is compact (cf. the proof of Theorem 5.2). If $\pi_1, \pi_2 \in \Omega$, then the set $\{x \in C^*(x_1, \dots, x_k) : \pi_1(x) = \pi_2(x)\}$ is a C^* -subalgebra of $C^*(x_1, \dots, x_k)$. This implies that the function $\tilde{H} : \Omega \ni \pi \mapsto (\pi(x_1), \dots, \pi(x_k)) \in \sigma_n(x_1, \dots, x_k) \cup \{0\}$ is one-to-one. It is obviously seen that \tilde{H} is surjective and continuous. Consequently, \tilde{H} is a homeomorphism (since Ω is compact). This proves (Sp4) and shows that $\sigma_n(x_1, \dots, x_k)$ is locally compact. It is also clear that $H(\mathbf{u} \cdot \pi) = \mathbf{u} \cdot H(\pi)$, which is followed by (Sp1) and (Sp2). Finally, for any $\pi \in \mathfrak{X}$, $C^*(\pi(x_1), \dots, \pi(x_k)) = \pi(C^*(x_1, \dots, x_k)) = M_n$ (see Lemma 4.10), which yields (Sp3) and completes the proof. \square

6.4. Definition. Let (x_1, \dots, x_k) be an n -homogeneous system in a C^* -algebra. The n -space $(\sigma_n(x_1, \dots, x_k), \cdot)$ defined by (6-1) is said to be the n -spectrum of (x_1, \dots, x_k) . According to Proposition 6.3, the n -spectrum of (x_1, \dots, x_k) is an n -spectrum of $C^*(x_1, \dots, x_k)$.

6.5. Proposition. Let $\mathbf{x} = (x_1, \dots, x_k)$ be an n -homogeneous system in a C^* -algebra. There exists a unique $*$ -homomorphism

$$\Phi_{\mathbf{x}} : C^*(\sigma_n(\mathbf{x}), \cdot) \rightarrow C^*(\mathbf{x})$$

such that $\Phi_{\mathbf{x}}(p_j) = x_j$ where $p_j : \sigma_n(\mathbf{x}) \ni (A_1, \dots, A_k) \mapsto A_j \in M_n$ ($j = 1, \dots, k$). Moreover, $\Phi_{\mathbf{x}}$ is a $*$ -isomorphism of C^* -algebras.

Proof. Let (\mathfrak{X}, \cdot) be the concrete n -spectrum of $C^*(\mathbf{x})$ and let $H : (\mathfrak{X}, \cdot) \rightarrow (\sigma_n(\mathbf{x}), \cdot)$ be the isomorphism as in point (Sp2) of Proposition 6.3. For $x \in C^*(\mathbf{x})$ let $\hat{x} \in \mathcal{C}(\mathfrak{X}, \cdot)$ be given by $\hat{x}(\pi) = \pi(x)$. The proof of Theorem 5.2 shows that the function $C^*(\mathbf{x}) \ni x \mapsto \hat{x} \in C^*(\mathfrak{X}, \cdot)$ is a $*$ -isomorphism of C^* -algebras. Consequently, $\Psi : C^*(\mathbf{x}) \ni x \mapsto \hat{x} \circ H^{-1} \in C^*(\sigma_n(\mathbf{x}), \cdot)$ is a $*$ -isomorphism as well. A direct calculation shows that $\Psi(x_j) = p_j$ ($j = 1, \dots, k$). This implies that $C^*(p_1, \dots, p_k) = C^*(\sigma_n(\mathbf{x}), \cdot)$, from which we infer the uniqueness of $\Phi_{\mathbf{x}}$. To convince about its existence, just put $\Phi_{\mathbf{x}} = \Psi^{-1}$. \square

We are now ready to introduce

6.6. Definition. Let $\mathbf{x} = (x_1, \dots, x_k)$ be an n -homogeneous system and let $\Phi_{\mathbf{x}}$ be as in Proposition 6.5. For every $f \in C^*(\sigma_n(x_1, \dots, x_k), \cdot)$ we denote by $f(x_1, \dots, x_k)$ the element $\Phi_{\mathbf{x}}(f)$. The assignment $f \mapsto f(x_1, \dots, x_k)$ is called the n -functional calculus.

The reader familiar with functional calculus on normal operators (or normal elements in C^* -algebras) has to be careful with the n -functional calculus, because its main disadvantage is that its values are not n -homogeneous elements in general. Therefore we cannot speak of the n -spectrum of $f(x_1, \dots, x_k)$ in general. What is more, it may happen that $\sigma_n(x_1, \dots, x_k)$ is compact, but $j(x_1, \dots, x_k)$, where j is constantly equal to the unit matrix, differs from the unit of the underlying C^* -algebra \mathcal{A} from which x_1, \dots, x_k were taken. This happens precisely when $C^*(x_1, \dots, x_k)$ has unit, but this unit is not the unit of \mathcal{A} .

As a consequence of Proposition 6.5 and Theorem 4.6 we obtain the *spectral theorem* (for n -homogeneous systems) announced before.

6.7. Theorem. *Let $\mathbf{T} = (T_1, \dots, T_k)$ be an n -homogeneous system of bounded linear operators acting on a Hilbert space \mathcal{H} . There exists a unique spectral n -measure $E_{\mathbf{T}}: \mathfrak{B}(\sigma_n(\mathbf{T})) \rightarrow M_n(\mathcal{B}(\mathcal{H}))$ such that*

$$\int p_j dE_{\mathbf{T}} = T_j \quad (j = 1, \dots, k)$$

where $p_j: \sigma_n(\mathbf{T}) \ni (A_1, \dots, A_k) \mapsto A_j \in M_n$.

6.8. Definition. Let $\mathbf{T} = (T_1, \dots, T_k)$ be an n -homogeneous system of bounded Hilbert space operators and let $E_{\mathbf{T}}$ be the spectral n -measure as in Theorem 6.7. $E_{\mathbf{T}}$ is called the *spectral n -measure of \mathbf{T}* and the assignment $\mathfrak{B}C^*(\sigma(\mathbf{T}), \cdot) \ni f \mapsto f(T_1, \dots, T_k) := \int f dE_{\mathbf{T}} \in \mathcal{B}(\mathcal{H})$ is called the *extended n -functional calculus*.

There is nothing surprising in the following

6.9. Proposition. *Let \mathcal{M} be a von Neumann algebra acting on a Hilbert space \mathcal{H} and let $\mathbf{T} = (T_1, \dots, T_k)$ be an n -homogeneous system of operators belonging to \mathcal{M} . Let $X = \sigma_n(\mathbf{T})$.*

- (a) *For any $f \in \mathfrak{B}C^*(X, \cdot)$, $f(\mathbf{T}) \in \mathcal{M}$.*
- (b) *If $f^{(1)}, f^{(2)}, \dots \in \mathfrak{B}C^*(X, \cdot)$ converge pointwise to $f: X \rightarrow M_n$ and*

$$\sup_{\substack{m \geq 1 \\ x \in X}} \|f^{(m)}(x)\| < \infty,$$

then $f \in \mathfrak{B}C^(X, \cdot)$ and $\lim_{m \rightarrow \infty} (f^{(m)}(\mathbf{T}))h = (f(\mathbf{T}))h$ for each $h \in \mathcal{H}$.*

Proof. We begin with (a). It is clear that $g(\mathbf{T}) \in \mathcal{M}$ for $g \in C^*(X, \cdot)$. Let $E_{\mathbf{T}} = [E_{pq}]$. Denote by $\langle \cdot, - \rangle$ the scalar product of \mathcal{H} and fix $f = [f_{pq}] \in \mathfrak{B}C^*(X, \cdot)$. We shall show that $f(\mathbf{T})$ belongs to the closure of $\{g(\mathbf{T}): g \in C^*(X, \cdot)\}$ in the weak operator topology of $\mathcal{B}(\mathcal{H})$, which will give (a). To this end, we fix $h_1, w_1, \dots, h_r, w_r \in \mathcal{H}$ and $\varepsilon > 0$. Put $\mu = \sum_{s=1}^r \sum_{p,q} |E_{pq}^{(h_s, w_s)}|$. By Lemma 4.7, there is $g = [g_{pq}] \in C^*(X, \cdot)$ such that $\int_X \|f(x) - g(x)\| d\mu(x) \leq \varepsilon$. But then, for each $s \in \{1, \dots, r\}$,

$$\begin{aligned} \left| \left\langle \left(\int f dE_{\mathbf{T}} - \int g dE_{\mathbf{T}} \right) h_s, w_s \right\rangle \right| &= \left| \sum_{p,q} \int_X (f_{pq} - g_{pq}) dE_{qp}^{(h_s, w_s)} \right| \\ &\leq \sum_{p,q} \int_X |f_{pq} - g_{pq}| d|E_{qp}^{(h_s, w_s)}| \leq \int_X \|f(x) - g(x)\| d\mu(x) \leq \varepsilon \end{aligned}$$

and we are done (since $f(\mathbf{T}) = \int f dE_{\mathbf{T}}$ and $g(\mathbf{T}) = \int g dE_{\mathbf{T}}$).

We turn to (b). It is clear that $f \in \mathfrak{B}C^*(X, \cdot)$. Replacing $f^{(m)}$ by $f^{(m)} - f$, we may assume $f = 0$. Observe that then $\lim_{m \rightarrow \infty} ((f^{(m)})^* f^{(m)})_{pq}(x) = 0$ for any $x \in X$ and $p, q \in \{1, \dots, n\}$ and the functions $((f^{(1)})^* f^{(1)})_{pq}, ((f^{(2)})^* f^{(2)})_{pq}, \dots$ are

uniformly bounded. Therefore (by Lebesgue's dominated convergence theorem) for any $h \in H$,

$$\begin{aligned} \|(f^{(m)}(\mathbf{T}))h\|^2 &= \langle (f^{(m)}(\mathbf{T}))^* (f^{(m)}(\mathbf{T}))h, h \rangle = \left\langle \left(\int (f^{(m)})^* f^{(m)} dE_{\mathbf{T}} \right) h, h \right\rangle \\ &= \sum_{p,q} \int_X ((f^{(m)})^* f^{(m)})_{pq} dE_{qp}^{(h,h)} \rightarrow 0 \quad (m \rightarrow \infty), \end{aligned}$$

which finishes the proof. \square

We end the paper with the note that the above result enables defining the extended n -functional calculus for n -homogeneous systems in W^* -algebras.

REFERENCES

- [1] R. Bhatia, *Matrix Analysis*, Springer, New York, 1997.
- [2] E. Bishop, *A generalization of the Stone-Weierstrass theorem*, Pacific J. Math. **11** (1961), 777–783.
- [3] B. Blackadar, *Operator Algebras. Theory of C^* -algebras and von Neumann algebras* (Encyclopaedia of Mathematical Sciences, vol. 122: Operator Algebras and Non-Commutative Geometry III), Springer-Verlag, Berlin-Heidelberg, 2006.
- [4] W.-M. Ching, *Topologies on the quasi-spectrum of a C^* -algebra*, Proc. Amer. Math. Soc. **46** (1974), 273–276.
- [5] J. Dixmier, *C^* -algebras*, North-Holland Publ. Co., Amsterdam, 1977.
- [6] J. Ernest, *Charting the operator terrain*, Mem. Amer. Math. Soc. **171** (1976), 207 pp.
- [7] J.M.G. Fell, *C^* -algebras with smooth dual*, Illinois J. Math. **4** (1960), 221–230.
- [8] J.M.G. Fell, *The dual spaces of C^* -algebras*, Trans. Amer. Math. Soc. **94** (1960), 365–403.
- [9] J.M.G. Fell, *The structure of algebras of operator fields*, Acta Math. **106** (1961), 233–280.
- [10] J. Glimm, *A Stone-Weierstrass theorem for C^* -algebras*, Ann. Math. **72** (1960), 216–244.
- [11] M.I. Garrido and F. Montalvo, *On some generalizations of the Kakutani-Stone and Stone-Weierstrass theorems*, Extracta Math. **6** (1991), 156–159.
- [12] D.W. Hadwin, *An operator-valued spectrum*, Notices Amer. Math. Soc. **23** (1976), A-163.
- [13] D.W. Hadwin, *An operator-valued spectrum*, Indiana Univ. Math. J. **26** (1977), 329–340.
- [14] D. Hofmann, *On a generalization of the Stone-Weierstrass theorem*, Appl. Categ. Structures **10** (2002), 569–592.
- [15] J.S. Kim, Ch.R. Kim, S.G. Lee, *Reducing operator valued spectra of a Hilbert space operator*, J. Korean Math. Soc. **17** (1980), 123–129.
- [16] S.G. Lee, *Remarks on reducing operator valued spectrum*, J. Korean Math. Soc. **16** (1980), 131–136.
- [17] R. Longo, *Solution of the factorial Stone-Weierstrass conjecture. An application of the theory of standard split W^* -inclusions*, Invent. Math. **76** (1984), 145–155.
- [18] K. Löwner, *Über monotone Matrixfunctionen*, Math. Z. **38** (1934), 177–216.
- [19] P. Niemiec, *Unitary equivalence and decompositions of finite systems of closed densely defined operators in Hilbert spaces*, Dissertationes Math. (Rozprawy Mat.) **482** (2012), 1–106.
- [20] C. Pearcy and N. Salinas, *Finite-dimensional representations of separable C^* -algebras*, Notices Amer. Math. Soc. **21** (1974), A-376.
- [21] S. Popa, *Semiregular maximal abelian $*$ -subalgebras and the solution to the factor state Stone-Weierstrass problem*, Invent. Math. **76** (1984), 157–161.
- [22] T. Saitô, *Generations of von Neumann algebras*, Lecture Notes in Math. vol. 247 (*Lecture on Operator Algebras*), Springer, Berlin-Heidelberg-New York, 1972, 435–531.
- [23] S. Sakai, *C^* -Algebras and W^* -Algebras*, Springer-Verlag, Berlin-Heidelberg-New York, 1971.
- [24] M.H. Stone, *Application of the theory of Boolean rings to general topology*, Trans. Amer. Math. Soc. **41** (1937), 375–481.
- [25] M.H. Stone, *The generalized Weierstrass approximation theorem*, Math. Mag. **21** (1948), 167–184.
- [26] M. Takesaki, *Theory of Operator Algebras I* (Encyclopaedia of Mathematical Sciences, Volume 124), Springer-Verlag, Berlin-Heidelberg-New York, 2002.
- [27] V. Timofte, *Stone-Weierstrass theorems revisited*, J. Approx. Theory **136** (2005), 45–59.

PIOTR NIEMIEC, INSTYTUT MATEMATYKI, WYDZIAŁ MATEMATYKI I INFORMATYKI, UNIWERSYTET JAGIELLONSKI, UL. ŁOJASIEWICZA 6, 30-348 KRAKÓW, POLAND
E-mail address: piotr.niemiec@uj.edu.pl