

Sharp constants in weighted trace inequalities on Riemannian manifolds

Tianling Jin and Jingang Xiong

February 19, 2022

Abstract

We establish some sharp weighted trace inequalities $W^{1,2}(\rho^{1-2\sigma}, M) \hookrightarrow L^{\frac{2n}{n-2\sigma}}(\partial M)$ on $n + 1$ dimensional compact smooth manifolds with smooth boundaries, where ρ is a defining function of M and $\sigma \in (0, 1)$. This is stimulated by some recent work on fractional (conformal) Laplacians and related problems in conformal geometry, and also motivated by a conjecture of Aubin.

1 Introduction

Let Ω be an open set in \mathbb{R}^n , $n \geq 1$, and $\rho(x) = \text{dist}(x, \partial\Omega)$ for $x \in \Omega$. There have been much work devoted to the structures of weighted Sobolev spaces of the type $W^{k,p}(\rho^\alpha, \Omega)$ where $\alpha \in \mathbb{R}$, $k \in \mathbb{N}$ and $1 \leq p \leq \infty$, as well as to their applications in different areas such as (stochastic) partial differential equations and Riemannian manifolds with fractal boundaries or boundary singularities. We refer to the book [36] of Maz'ya and references therein for these topics.

In this paper, we would like to study sharp constants in weighted trace type inequalities $W^{1,2}(\rho^{1-2\sigma}) \hookrightarrow L^{\frac{2n}{n-2\sigma}}(\partial M)$ on Riemannian manifolds M with boundaries ∂M . Let us start from Euclidean spaces. Denote $\dot{H}^\sigma(\mathbb{R}^n)$ as the σ -order homogeneous Sobolev space on \mathbb{R}^n , $n \geq 2$, which is the closure of $C_c^\infty(\mathbb{R}^n)$ under the norm

$$\|f\|_{\dot{H}^\sigma(\mathbb{R}^n)} = \left(\int_{\mathbb{R}^n} |(-\Delta)^{\sigma/2} f(x)|^2 dx \right)^{1/2}.$$

The sharp σ -order Sobolev inequality asserts that

$$\|f\|_{L^{\frac{2n}{n-2\sigma}}(\mathbb{R}^n)}^2 \leq c(n, \sigma) \|f\|_{\dot{H}^\sigma(\mathbb{R}^n)}^2$$

for all $f \in \dot{H}^\sigma(\mathbb{R}^n)$, where

$$c(n, \sigma) = 2^{-2\sigma} \pi^{-\sigma} \left(\frac{\Gamma((n-2\sigma)/2)}{\Gamma((n+2\sigma)/2)} \right) \left(\frac{\Gamma(n)}{\Gamma(n/2)} \right)^{\frac{2\sigma}{n}},$$

and the equality holds if and only if $f(x)$ takes the form

$$c \left(\frac{\lambda}{1 + \lambda^2 |x - x_0|^2} \right)^{\frac{n-2\sigma}{2}}$$

for some $c \in \mathbb{R}$, $\lambda > 0$ and $x_0 \in \mathbb{R}^n$. These have been proved by Lieb in [34]. Set $x = (x', x_{n+1}) \in \mathbb{R}_+^{n+1} := \mathbb{R}^n \times (0, \infty)$ and

$$F(x', x_{n+1}) = \int_{\mathbb{R}^n} \mathcal{P}_\sigma(x' - \xi, x_{n+1}) f(\xi) d\xi,$$

where

$$\mathcal{P}_\sigma(x', x_{n+1}) = \beta(n, \sigma) \frac{x_{n+1}^{2\sigma}}{(|x'|^2 + x_{n+1}^2)^{\frac{n+2\sigma}{2}}} \quad (1)$$

with the normalization constant $\beta(n, \sigma) > 0$ such that $\int_{\mathbb{R}^n} \mathcal{P}_\sigma(x', 1) dx' = 1$. Then one has (see, e.g., [9])

$$N_\sigma \int_{\mathbb{R}_+^{n+1}} x_{n+1}^{1-2\sigma} |\nabla F(x', x_{n+1})|^2 dx = \|f\|_{\dot{H}^\sigma(\mathbb{R}^n)}^2,$$

where $N_\sigma = 2^{2\sigma-1} \Gamma(\sigma) / \Gamma(1-\sigma)$. Hence, we have

$$\|f\|_{L^{\frac{2n}{n-2\sigma}}(\mathbb{R}^n)}^2 \leq S(n, \sigma) \int_{\mathbb{R}_+^{n+1}} x_{n+1}^{1-2\sigma} |\nabla F(x', x_{n+1})|^2 dx \quad (2)$$

for all $f \in \dot{H}^\sigma(\mathbb{R}^n)$, where $S(n, \sigma) = N_\sigma \cdot c(n, \sigma)$. Consequently, one can show (see, e.g., Proposition 2.1 below together with a density argument) that

$$\|U(\cdot, 0)\|_{L^{\frac{2n}{n-2\sigma}}(\mathbb{R}^n)}^2 \leq S(n, \sigma) \int_{\mathbb{R}_+^{n+1}} x_{n+1}^{1-2\sigma} |\nabla U(x', x_{n+1})|^2 dx \quad (3)$$

for all $U \in W^{1,2}(x_{n+1}^{1-2\sigma}, \mathbb{R}_+^{n+1})$, which is the closure of $C_c^\infty(\overline{\mathbb{R}_+^{n+1}})$ under the norm

$$\|U\|_{W^{1,2}(x_{n+1}^{1-2\sigma}, \mathbb{R}_+^{n+1})} = \sqrt{\int_{\mathbb{R}_+^{n+1}} x_{n+1}^{1-2\sigma} (|U|^2 + |\nabla U|^2) dx}.$$

Stimulated by several recent work on fractional (conformal) Laplacians and related problems in conformal geometry (see, e.g., [22, 10, 21, 26]) and a conjecture of Aubin [2], we study

weighted Sobolev trace inequalities of type (3) on Riemannian manifolds with boundaries. For $n \geq 2$, let (M, g) be an $n + 1$ dimensional, compact, smooth Riemannian manifold with smooth boundary ∂M . We say a function $\rho \in C^\infty(\overline{M})$ is a *defining function* of M if

$$\rho > 0 \quad \text{in } M, \quad \rho = 0 \text{ and } \nabla_g \rho \neq 0 \quad \text{on } \partial M.$$

Since $\rho^{1-2\sigma}$, where $\sigma \in (0, 1)$ is a constant, belongs to the Muckenhoupt A_2 class, we define the weighted Sobolev space $H^1(\rho^{1-2\sigma}, M)$ as the closure of $C^\infty(\overline{M})$ under the norm

$$\|u\|_{H^1(\rho^{1-2\sigma}, M)} = \left(\int_M \rho^{1-2\sigma} (|u|^2 + |\nabla u|^2) dv_g \right)^{\frac{1}{2}},$$

where dv_g denote the volume form of (M, g) . $H^1(\rho^{1-2\sigma}, M)$ is a Hilbert space and it has a well-defined *trace operator* T (see, e.g., [36] or [39]) which continuously maps $H^1(\rho^{1-2\sigma}, M)$ to $H^\sigma(\partial M)$, where $H^\sigma(\partial M)$ is the σ -order Sobolev space on ∂M .

Theorem 1.1. *For $n \geq 2$, let (M, g) be an $n + 1$ dimensional, compact, smooth Riemannian manifold with smooth boundary ∂M . Let $\sigma \in (0, \frac{1}{2}]$, and ρ be a defining function of M satisfying $|\nabla_g \rho| = 1$ on ∂M . Then there exists a positive constant $A = A(M, g, n, \rho, \sigma)$ such that*

$$\left(\int_{\partial M} |u|^{\frac{2n}{n-2\sigma}} ds_g \right)^{\frac{n-2\sigma}{n}} \leq S(n, \sigma) \int_M \rho^{1-2\sigma} |\nabla_g u|^2 dv_g + A \int_{\partial M} u^2 ds_g, \quad (4)$$

for all $u \in H^1(\rho^{1-2\sigma}, M)$, where ds_g denotes the induced volume form on ∂M .

For $\sigma \in (\frac{1}{2}, 1)$, we have

Theorem 1.2. *Let $\sigma \in (\frac{1}{2}, 1)$, $n \geq 4$ and (M, g) be an $n + 1$ dimensional, compact, smooth Riemannian manifold with smooth boundary ∂M . Suppose in addition that ∂M is totally geodesic. Let ρ be a defining function of M satisfying $\rho(x) = d(x) + O(d(x)^3)$ as $d(x) \rightarrow 0$, where $d(x)$ denotes the distance between x and ∂M with respect to the metric g . Then there exists a positive constant $A = A(M, g, n, \rho, \sigma)$ such that (4) holds for all $u \in H^1(\rho^{1-2\sigma}, M)$.*

Remark 1.1. *The constant $S(n, \sigma)$ in (4) is optimal for all $\sigma \in (0, 1)$, see Proposition 2.2.*

Remark 1.2. *Theorem 1.2 may fail without any geometric assumption on ∂M . For example, it is the case when the mean curvature of ∂M is positive somewhere. In particular, (4) is false on any bounded smooth domain in \mathbb{R}^{n+1} when $\sigma \in (1/2, 1)$. However, Theorem 1.1 holds for all $\sigma \in (0, 1)$ if $S(n, \sigma)$ is replaced by any $S > S(n, \sigma)$, see Proposition 2.5.*

Remark 1.3. *It is clear that we only need to consider the case when M is connected. Throughout the paper, we assume this.*

When $\sigma = \frac{1}{2}$, (4) is a standard Sobolev trace inequality which has been extensively studied, see, e.g., Lions [35], Escobar [14], Beckner [5], Adimurthi-Yadava [1], Li-Zhu [32, 33] and many others. In particular, Li-Zhu [32] established Theorem 1.1 for $\sigma = \frac{1}{2}$. The sharp inequality (4) is in the same spirit of a conjecture posed by Aubin [2] which concerns the best constants in Sobolev embedding theorems on Riemannian manifolds. Aubin's conjecture had been confirmed through the work of Hebey-Vaugon [25], Aubin-Li [4] and Druet [11, 12]. Besides, various refinements of Aubin's conjecture were obtained in Druet-Hebey [13], Li-Ricciardi [31] and etc. These sharp Sobolev type inequalities play important roles in the study of nonlinear partial differential equations, see Aubin [3], Hebey [24], Schoen-Yau [42] and references therein.

For the defining function in the above theorems, $(M, g/\rho^2)$ is *asymptotically hyperbolic* in the sense that $(M, g/\rho^2)$ is a complete manifold and along any smooth curve in $M \setminus \partial M$ tending to a point $\xi \in \partial M$ all sectional curvatures of g/ρ^2 approach to -1 (see Mazzeo [37] or Mazzeo-Melrose [38]). On the conformal infinity $(\partial M, [g|_{\partial M}])$ of $(M, g/\rho^2)$, one can define fractional order conformally invariant operators P_σ^g for $\sigma \in (0, \frac{n}{2})$ except at most finite values, via normalized scattering operators (see Graham-Zworski [22] and Chang-González [10]), which leads to σ -scalar curvature $R_\sigma^g := P_\sigma^g(1)$ on ∂M . A fractional Yamabe problem, which is to find a metric in $[g|_{\partial M}]$ of constant σ -curvature and related ones, have been studied by Qing-Raske [41], González-Mazzeo-Sire [20] and González-Qing [21]. When $\sigma \in (0, 1)$, it can be formulated (see [21]) as seeking minimizers of the energy functional

$$I^\sigma[u] = \frac{N_\sigma \int_M \rho^{1-2\sigma} |\nabla u|^2 dv_g + \int_{\partial M} R_\sigma^g u^2 ds_g}{\left(\int_{\partial M} |u|^{\frac{2n}{n-2\sigma}} ds_g \right)^{\frac{n-2\sigma}{n}}}, \quad u \in H^1(\rho^{1-2\sigma}, M), \quad u \not\equiv 0 \text{ on } \partial M, \quad (5)$$

for some proper ρ . For $\sigma = 1/2$, it is the energy functional of a Yamabe problem with boundary initially studied by Escobar [15]. A fractional Nirenberg problem about prescribing σ -scalar curvature on \mathbb{S}^n has been studied by Jin-Li-Xiong [26, 27] and a fractional Yamabe flow has been studied by Jin-Xiong [28]. Variational problems related to energy functional (5) on bounded domains in Euclidean spaces have been studied by González [19], Palatucci-Sire [40].

Finally, we provide a brief sketch of the proofs of the two main theorems. Since the right hand side of (4) does not contain terms like $\int_M \rho^{1-2\sigma} u^2 dv_g$, we adapt a global argument from Li-Zhu [32, 33]. By contradiction, we assume that for any $\alpha > 0$,

$$I_\alpha := \frac{\int_M \rho^{1-2\sigma} |\nabla_g u|^2 dv_g + \alpha \int_{\partial M} |u|^2 ds_g}{\left(\int_{\partial M} |u|^{\frac{2n}{n-2\sigma}} ds_g \right)^{\frac{n-2\sigma}{n}}} < \frac{1}{S(n, \sigma)},$$

for some $u \in H^1(\rho^{1-2\sigma}, M)$ with that $u \not\equiv 0$ on ∂M . It follows that there exists a minimizer u_α of I_α , and u_α blows up at exactly one point as $\alpha \rightarrow \infty$. One key step is the asymptotical analysis of u_α near its blow up point. Here we have to overcome difficulties from the degeneracy and the lack of conformal invariance of the Euler-Lagrange equation of I_α satisfied by u_α . Another difference from [32] (the case $\sigma = 1/2$) is that some Sobolev embedding theorems for $H^1(\rho^{1-2\sigma}, M)$,

which play important roles in establishing the blow-up profile of u_α in the interior of M in [32] in the case $\sigma = \frac{1}{2}$, fail when $\sigma > \frac{1}{2}$ (see, e.g., Theorem 1 in page 135 or Corollary 2 in page 193 of [36]). However, we succeeded in establishing the optimal asymptotical behavior of u_α on the boundary ∂M (Proposition 3.3). In this step, a Liouville type theorem in Jin-Li-Xiong [26] and *Neumann functions* for degenerate equations in Theorem 1.3 are used. The last step is to derive a contradiction by checking balance via a Pohozaev type inequality in some proper region, where a Harnack inequality established by Cabre-Sire [8] or Tan-Xiong [43] is used to obtain the asymptotical behavior of u_α near its blowup point in M from that on ∂M . Some extra arguments on ∂M are needed for $\sigma > \frac{1}{2}$.

Theorem 1.3. *Let $f \in L^1(\partial M)$ with mean value zero, i.e., $\int_{\partial M} f = 0$. Then there exists a weak solution $u \in W^{1,1+\varepsilon_0}(\rho^{1-2\sigma}, M)$ of (59) where $\varepsilon_0 > 0$ depending only on n and σ . Consequently, if $f = \delta_{x_0} - \frac{1}{|\partial M|_g}$ for some $x_0 \in \partial M$, where δ_{x_0} is the delta function at x_0 and $|\partial M|_g$ is the area of ∂M with respect to the induced metric g , then there exists a weak solution $u \in W^{1,1+\varepsilon_0}(\rho^{1-2\sigma}, M) \cap H_{loc}^1(\rho^{1-2\sigma}, \overline{M} \setminus \{x_0\})$ of (59) with mean value zero. Moreover, for all $x \in \overline{M} \setminus \{x_0\}$,*

$$A_1 \text{dist}_g(x, x_0)^{2\sigma-n} - A_0 \leq u(x) \leq A_2 \text{dist}_g(x, x_0)^{2\sigma-n}$$

where A_0, A_1, A_2 are positive constants depending only on M, g, n, σ, ρ .

The proof of Theorem 1.3 follows from Lemma A.5, Theorem A.5 and some approximation arguments. When $\sigma = 1/2$, Theorem 1.3 follows directly from Brezis-Strauss [7] and Kenig-Pipher [29].

Notations. We collect below a list of the main notations used throughout the paper.

- We always assume that $n \geq 2, \sigma \in (0, 1)$, and ρ is a smooth defining function as in Theorem 1.1 without otherwise stated. Denote $q = \frac{2n}{n-2\sigma}$.
- For a domain $D \subset \mathbb{R}^{n+1}$ with boundary ∂D , we denote $\partial' D$ as the interior of $\overline{D} \cap \partial \mathbb{R}_+^{n+1}$ in $\mathbb{R}^n = \partial \mathbb{R}_+^{n+1}$ and $\partial'' D = \partial D \setminus \partial' D$.
- For $\bar{x} \in \mathbb{R}^{n+1}$, $\mathcal{B}_r(\bar{x}) := \{x \in \mathbb{R}^{n+1} : |x - \bar{x}| = \sqrt{(x_1 - \bar{x}_1)^2 + \cdots + (x_{n+1} - \bar{x}_{n+1})^2} < r\}$, $\mathcal{B}_r^+(\bar{x}) := \mathcal{B}_r(\bar{x}) \cap \mathbb{R}_+^{n+1}$. If $\bar{x} \in \partial \mathbb{R}_+^{n+1}$, $B_r(\bar{x}) := \{x = (x', 0) : |x' - \bar{x}'| < r\}$. Hence $\partial' \mathcal{B}_r^+(\bar{x}) = B_r(\bar{x})$ if $\bar{x} \in \partial \mathbb{R}_+^{n+1}$. We will not keep writing the center \bar{x} if $\bar{x} = 0$.

Acknowledgements: Both authors thank Prof. Y.Y. Li for encouragements and useful discussions. Tianling Jin was partially supported by a University and Louis Bevier Dissertation Fellowship at Rutgers University and Rutgers University School of Art and Science Excellence Fellowship. Jingang Xiong was partially supported by CSC project for visiting Rutgers University and NSFC No. 11071020. He is very grateful to the Department of Mathematics at Rutgers University for the kind hospitality.

2 Preliminaries

Proposition 2.1. *For any $u \in C_c^\infty(\overline{\mathbb{R}_+^{n+1}})$, we have*

$$\left(\int_{\mathbb{R}^n} |u(x', 0)|^q dx' \right)^{\frac{2}{q}} \leq S(n, \sigma) \int_{\mathbb{R}_+^{n+1}} x_{n+1}^{1-2\sigma} |\nabla u(x)|^2 dx.$$

Moreover, the above inequality fails if $S(n, \sigma)$ is replaced by any smaller constant.

Proof. It follows from (3) and Lemma A.3 of [26]. See also Corollary 5.3 of [21]. \square

Proposition 2.2. *Let M be as in Theorem 1.1. Let $\sigma \in (0, 1)$, and ρ be a defining function of ∂M with $|\nabla_g \rho| = 1$ on ∂M . Suppose there exist some positive constants \tilde{S} and \tilde{A} such that, for all $u \in H^1(\rho^{1-2\sigma}, M)$,*

$$\left(\int_{\partial M} |u|^q ds_g \right)^{\frac{2}{q}} \leq \tilde{S} \int_M \rho^{1-2\sigma} |\nabla_g u|^2 dv_g + \tilde{A} \int_{\partial M} |u|^2 ds_g.$$

Then $\tilde{S} \geq S(n, \sigma)$.

Proof. Given Proposition 2.1, the proof is standard (see, e.g., Proposition 4.2 of [24]). We include it here for completeness and to illustrate the role of $|\nabla \rho| = 1$. We argue by contradiction. Suppose that there exists a Riemannian manifold (M, g) , a defining function ρ of ∂M with $|\nabla_g \rho| = 1$ on ∂M , $\sigma \in (0, 1)$, $\tilde{S} < S(n, \sigma)$ and $\tilde{A} > 0$ such that for all $u \in H^1(\rho^{1-2\sigma}, M)$,

$$\left(\int_{\partial M} |u|^q ds_g \right)^{\frac{2}{q}} \leq \tilde{S} \int_M \rho^{1-2\sigma} |\nabla_g u|^2 dv_g + \tilde{A} \int_{\partial M} |u|^2 ds_g. \quad (6)$$

Let $x \in \partial M$. For any $\varepsilon > 0$, which will be chosen sufficiently small, there exists a chart (Ω, φ) of M at x and $\delta > 0$ such that $\varphi(\Omega) = \mathcal{B}_\delta^+(0)$ the upper half Euclidean ball of center 0 and radius δ in \mathbb{R}_+^{n+1} , and

$$(1 - \varepsilon)\delta_{ij} \leq g_{ij} \leq (1 + \varepsilon)\delta_{ij}. \quad (7)$$

By assumption, (6) holds for any $u \in C_c^\infty(\Omega \cup (\partial\Omega \cap \partial M))$, i.e.,

$$\begin{aligned} \left(\int_{B_\delta(0)} |u|^q \sqrt{\det(g_{ij})} dx' \right)^{\frac{2}{q}} &\leq \tilde{S} \int_{\mathcal{B}_\delta^+(0)} \rho^{1-2\sigma} g^{ij} u_i u_j \sqrt{\det(g_{ij})} dx \\ &\quad + \tilde{A} \int_{B_\delta(0)} |u|^2 \sqrt{\det(g_{ij})} dx'. \end{aligned}$$

It follows from (7), $|\nabla_g \rho| = 1$ and $\rho = 0$ on ∂M that there exists $\delta_0 > 0$, $\tilde{S}' < S(n, \sigma)$, $\tilde{A}' > 0$ such that for all $\delta \in (0, \delta_0)$ and $u \in C_c^\infty(\mathcal{B}_\delta(0) \cup B_\delta(0))$, i.e.,

$$\left(\int_{B_\delta(0)} |u|^q dx' \right)^{\frac{2}{q}} \leq \tilde{S}' \int_{\mathcal{B}_\delta^+(0)} x_{n+1}^{1-2\sigma} |\nabla u|^2 dx + \tilde{A}' \int_{B_\delta(0)} |u|^2 dx'.$$

By Hölder's inequality, $\int_{B_\delta(x)} |u|^2 dx' \leq |B_\delta(0)|^{\frac{q-2}{q}} \left(\int_{B_\delta(0)} |u|^q dx' \right)^{\frac{2}{q}}$. By choosing δ sufficiently small, we have that there exists $\tilde{S}'' < S(n, \sigma)$ such that for all $u \in C_c^\infty(\mathcal{B}_\delta(0) \cup B_\delta(0))$

$$\left(\int_{B_\delta(0)} |u|^q dx' \right)^{\frac{2}{q}} \leq \tilde{S}'' \int_{\mathcal{B}_\delta^+(0)} x_{n+1}^{1-2\sigma} |\nabla u|^2 dx.$$

Consequently, by a scaling argument, we have

$$\left(\int_{\mathbb{R}^n} |u(x', 0)|^q dx' \right)^{\frac{2}{q}} \leq \tilde{S}'' \int_{\mathbb{R}_+^{n+1}} x_{n+1}^{1-2\sigma} |\nabla u(x)|^2 dx.$$

for any $u \in C_c^\infty(\overline{\mathbb{R}_+^{n+1}})$, which contradicts Proposition 2.1. \square

Proposition 2.3. *Assume the assumptions in Proposition 2.2. Then for any $\varepsilon > 0$ there exists a positive constant B_ε such that*

$$\left(\int_{\partial M} |u|^q ds_g \right)^{\frac{2}{q}} \leq (S(n, \sigma) + \varepsilon) \int_M \rho^{1-2\sigma} |\nabla_g u|^2 dv_g + B_\varepsilon \int_M \rho^{1-2\sigma} |u|^2 dv_g.$$

Proof. It also follows from Proposition 2.1 and a standard partition of unity argument, see, e.g., Theorem 4.5 of [24] on page 95. \square

For every $\alpha > 0$, consider the functional

$$I_\alpha[u] = \frac{\int_M \rho^{1-2\sigma} |\nabla_g u|^2 dv_g + \alpha \int_{\partial M} |u|^2 ds_g}{\left(\int_{\partial M} |u|^q ds_g \right)^{2/q}}, \quad u \in H^1(\rho^{1-2\sigma}, M), \quad u \not\equiv 0 \text{ on } \partial M.$$

Proposition 2.4. *Suppose that for some $\alpha > 0$,*

$$\xi_\alpha := \inf_{u \in H^1(\rho^{1-2\sigma}, M), u|_{\partial M} \not\equiv 0} I_\alpha[u] < \frac{1}{S(n, \sigma)}, \quad (8)$$

then ξ_α is achieved by a nonnegative function $u_\alpha \in H^1(\rho^{1-2\sigma}, M)$ with

$$\int_{\partial M} u_\alpha^q ds_g = 1. \quad (9)$$

Proof. Given Proposition 2.3, the Proposition follows from standard calculus of variations, see page 452 of [32]. \square

Proposition 2.5. *Assume the assumptions in Proposition 2.2. For any $\varepsilon > 0$, there exists a positive constant A_ε such that*

$$\left(\int_{\partial M} |u|^q ds_g \right)^{\frac{2}{q}} \leq (S(n, \sigma) + \varepsilon) \int_M \rho^{1-2\sigma} |\nabla_g u|^2 dv_g + A_\varepsilon \int_{\partial M} |u|^2 ds_g.$$

Proof. Given Propositions 2.3 and 2.4, and Corollary A.1, the proof of Proposition 2.5 is similar to Proposition 1.2 of [32] and we omit it here. \square

3 Asymptotic analysis

For brevity, from now on we write S instead of $S(n, \sigma)$. We prove Theorem 1.1 by contradiction. Namely, assume that for any $\alpha \geq 1$,

$$\xi_\alpha < \frac{1}{S}, \quad (10)$$

where ξ_α is defined as in Proposition 2.4. Let u_α be some nonnegative minimizer of I_α obtained in Proposition 2.4 which satisfies

$$\xi_\alpha = \int_M \rho^{1-2\sigma} |\nabla_g u_\alpha|^2 dv_g + \alpha \int_{\partial M} u_\alpha^2 ds_g, \quad \int_{\partial M} u_\alpha^q ds_g = 1, \quad (11)$$

and for any $\varphi \in H^1(\rho^{1-2\sigma}, M)$,

$$\int_M \rho^{1-2\sigma} \langle \nabla_g u_\alpha, \nabla_g \varphi \rangle_g dv_g + \alpha \int_{\partial M} u_\alpha \varphi ds_g = \xi_\alpha \int_{\partial M} u_\alpha^{q-1} \varphi ds_g. \quad (12)$$

The geodesic distance function $d(x) := \text{dist}(x, \partial M)$ determines for some $\varepsilon_0 > 0$ an identification of $\partial M \times [0, \varepsilon_0)$ with a neighborhood of ∂M in M : $(x', d) \in \partial M \times [0, \varepsilon_0)$ corresponds to the point obtained by following the integral curve of $\nabla_g d$ emanating from x' for d units of time. Furthermore, $\nabla_g d$ is orthogonal to the slices $\partial M \times \{d\}$. Define $\nu := -\nabla_g d$ for $d < \varepsilon_0$. It follows from Theorem A.2, Theorem A.3 and Proposition A.1 that $u_\alpha \in C^\gamma(\overline{M}) \cap C^\infty(M) \cap C^\infty(\partial M)$ for some $\gamma \in (0, 1)$ and $\rho^{1-2\sigma} \frac{\partial_g u_\alpha}{\partial \nu} \in C(\partial M \times [0, \varepsilon_0/2])$. Hence, u_α satisfies the Euler-Lagrange equation

$$\begin{cases} \text{div}_g \left(\rho^{1-2\sigma} \nabla_g u_\alpha \right) = 0, & \text{in } M, \\ \lim_{d \rightarrow 0} \rho^{1-2\sigma} (x', d) \frac{\partial_g u_\alpha}{\partial \nu} (x', \rho) = \xi_\alpha u_\alpha^{q-1}(x') - \alpha u_\alpha(x'), & \text{on } \partial M. \end{cases} \quad (13)$$

in the pointwise sense.

It follows from the maximum principle that $\max_{\overline{M}} u_\alpha = \max_{\partial M} u_\alpha$. Let $u_\alpha(x_\alpha) = \max_{\overline{M}} u_\alpha$, where $x_\alpha \in \partial M$, and $\mu_\alpha = u_\alpha(x_\alpha)^{-\frac{2}{n-2\sigma}}$. By a Hopf Lemma (see, e.g., Proposition 4.11 in [8]), we have $\xi_\alpha u_\alpha(x_\alpha)^{q-1} - \alpha u_\alpha(x_\alpha) > 0$, that is

$$\alpha \mu_\alpha^{2\sigma} < \xi_\alpha. \quad (14)$$

Hence, $\lim_{\alpha \rightarrow \infty} \mu_\alpha^{2\sigma} = 0$.

Lemma 3.1. *As $\alpha \rightarrow \infty$, we have*

$$\xi_\alpha \rightarrow \frac{1}{S}, \quad (15a)$$

$$\alpha \|u_\alpha\|_{L^2(\partial M)}^2 \rightarrow 0. \quad (15b)$$

Proof. For all small $\varepsilon > 0$, it follows from Proposition 2.5 that

$$\begin{aligned} 1 &\leq (S + \varepsilon) \int_M \rho^{1-2\sigma} |\nabla_g u_\alpha|^2 dv_g + A_\varepsilon \int_{\partial M} u_\alpha^2 ds_g \\ &= (S + \varepsilon) \xi_\alpha + (A_\varepsilon - (S + \varepsilon) \alpha) \int_{\partial M} u_\alpha^2 ds_g. \end{aligned}$$

Hence, for every $\alpha \geq \frac{2A_\varepsilon}{S+\varepsilon}$ we have

$$\frac{1}{S + \varepsilon} \leq \xi_\alpha < \frac{1}{S}, \quad \frac{S}{2} \alpha \int_{\partial M} u_\alpha^2 ds_g < \frac{\varepsilon}{S}.$$

(15a) and (15b) follow immediately. \square

Let $x = (x_1, \dots, x_n, x_{n+1}) = (x', x_{n+1})$ be *Fermi coordinates* (see, e.g., [15]) at x_α , where (x_1, \dots, x_n) are normal coordinates on ∂M at x_α and $\gamma(x_{n+1})$ is the geodesic leaving from (x_1, \dots, x_n) in the orthogonal direction to ∂M and parametrized by arc length. In this coordinate system,

$$\sum_{1 \leq i, j \leq n+1} g_{ij}(x) dx_i dx_j = dx_{n+1}^2 + \sum_{1 \leq i, j \leq n} g_{ij}(x) dx_i dx_j.$$

Moreover, g^{ij} has the following Taylor expansion near ∂M :

Lemma 3.2 (Lemma 3.2 in [15]). *For $\{x_k\}_{k=1, \dots, n+1}$ are small,*

$$g^{ij}(x) = \delta^{ij} + 2h^{ij}(x', 0)x_{n+1} + O(|x|^2), \quad (16)$$

where $i, j = 1, \dots, n$ and h_{ij} is the second fundamental form of ∂M .

For suitably small $\delta_0 > 0$ (independent of α), we define v_α in a neighborhood of $x_\alpha = 0$ by

$$v_\alpha(x) = \mu_\alpha^{(n-2\sigma)/2} u_\alpha(\mu_\alpha x), \quad x \in \mathcal{B}_{\delta_0/\mu_\alpha}^+.$$

It follows that

$$\begin{cases} \operatorname{div}_{g_\alpha} \left(\rho_\alpha^{1-2\sigma} \nabla_{g_\alpha} v_\alpha \right) = 0, & \text{in } \mathcal{B}_{\delta_0/\mu_\alpha}^+ \\ \lim_{x_{n+1} \rightarrow 0^+} \rho_\alpha^{1-2\sigma} \frac{\partial_{g_\alpha} v_\alpha}{\partial \nu} = \xi_\alpha v_\alpha^{q-1} - \alpha \mu_\alpha^{2\sigma} v_\alpha, & \text{on } \partial' \mathcal{B}_{\delta_0/\mu_\alpha}^+ = B_{\delta_0/\mu_\alpha} \\ v_\alpha(0) = 1, \quad 0 \leq v_\alpha \leq 1, \end{cases} \quad (17)$$

where $g_\alpha(x) = g_{ij}(\mu_\alpha x) dx_i dx_j$, $\rho_\alpha(x) = \rho(\mu_\alpha x)/\mu_\alpha$. It follows from (14) and Theorem A.2 in the Appendix that for all $R > 1$,

$$\|v_\alpha\|_{C^\gamma(\mathcal{B}_R^+)} + \|v_\alpha\|_{H^1(\rho_\alpha^{1-2\sigma}, \mathcal{B}_R^+)} \leq C(R), \quad \text{for all sufficiently large } \alpha, \quad (18)$$

where $\gamma \in (0, 1)$ is independent of R and α . It follows that there exists $v \in C_{loc}^\gamma(\overline{\mathbb{R}_+^{n+1}}) \cap H_{loc}^1(x_{n+1}^{1-2\sigma}, \overline{\mathbb{R}_+^{n+1}})$ such that along some subsequence,

$$\begin{cases} v_\alpha \rightarrow v \text{ in } C^{\gamma/2}(\mathcal{B}_R^+), \\ v_\alpha \rightharpoonup v \text{ weakly in } H^1(x_{n+1}^{1-2\sigma}, \mathcal{B}_R^+) \end{cases} \quad (19)$$

for any $R > 0$ as $\alpha \rightarrow \infty$. Since $v_\alpha(0) = 1$, we have

$$\begin{aligned} \int_{B_1} v_\alpha^q ds_{g_\alpha} &\geq 1/C > 0, \\ \int_{B_1} v_\alpha^2 ds_{g_\alpha} &\geq 1/C > 0. \end{aligned} \quad (20)$$

On the other hand,

$$\alpha \|u_\alpha\|_{L^2(\partial M)}^2 \geq \alpha \int_{B_{\mu_\alpha}(x_\alpha)} u_\alpha^2 = \alpha \mu_\alpha^{2\sigma} \int_{B_1} v_\alpha^2,$$

where we abused notation by denoting $B_r(x_\alpha)$ as the geodesic ball on ∂M centered at x_α with radius r . It follows from (15b) and (20) that

$$\lim_{\alpha \rightarrow \infty} \alpha \mu_\alpha^{2\sigma} = 0. \quad (21)$$

From (17), (21) and (15a), we conclude that v is a weak solution (see Section A.2 for the definition of weak solutions) of

$$\begin{cases} \operatorname{div}(x_{n+1}^{1-2\sigma} \nabla v) = 0, & \text{in } \mathbb{R}_+^{n+1}, \\ - \lim_{x_{n+1} \rightarrow 0^+} x_{n+1}^{1-2\sigma} \partial_{x_{n+1}} v = \frac{1}{S} v^{q-1}, & \text{on } \partial \mathbb{R}_+^{n+1}, \\ v(0) = 1, \quad 0 \leq v \leq 1. \end{cases} \quad (22)$$

By a Liouville type theorem, Theorem 1.5 in [26],

$$v(x', 0) = \left(\frac{1}{1 + \tilde{c}(n, \sigma)|x'|^2} \right)^{\frac{n-2\sigma}{2}}, v(x', x_{n+1}) = \int_{\mathbb{R}^n} \mathcal{P}_\sigma(x' - y', x_{n+1}) v(y', 0) dy',$$

where $\tilde{c}(n, \sigma)$ is a positive constant such that $\int_{\mathbb{R}^n} v^q(z) dz = 1$, and $\mathcal{P}_\sigma(x)$ is given in (1). Due to the uniqueness of the limit function v , we know that (19) holds for all $\alpha \rightarrow \infty$.

Proposition 3.1. *For $\delta_0 = \delta_0(M, g) > 0$ small enough,*

$$\lim_{\alpha \rightarrow \infty} \int_{B_{\delta_0/\mu_\alpha}} |v_\alpha - v|^q = 0.$$

Proof. Note that $v_\alpha \geq 0$ and

$$\int_{B_{\delta_0/\mu_\alpha}} v_\alpha^q \leq \int_{\partial M} u_\alpha^q = 1. \quad (23)$$

For any $\varepsilon > 0$, choose $R > 0$ such that $\int_{\mathbb{R}^n \setminus B_R} v^q(x', 0) dx' \leq \varepsilon$. It follows from (19) that $\int_{B_R} |v_\alpha - v|^q \leq \varepsilon$ and $1 - \int_{B_R} v_\alpha^q < 2\varepsilon$ for all α sufficiently large. Then

$$\begin{aligned} & \int_{B_{\delta_0/\mu_\alpha}} |v_\alpha - v|^q \\ &= \int_{B_{\delta_0/\mu_\alpha} \cap B_R} |v_\alpha - v|^q + \int_{B_{\delta_0/\mu_\alpha} \cap B_R^c} |v_\alpha - v|^q \\ &\leq \int_{B_{\delta_0/\mu_\alpha} \cap B_R} |v_\alpha - v|^q + 2^q \int_{B_{\delta_0/\mu_\alpha} \cap B_R^c} v_\alpha^q + 2^q \int_{B_{\delta_0/\mu_\alpha} \cap B_R^c} v^q \\ &\leq \varepsilon + 2^q(1 - \int_{B_R} v_\alpha^q) + 2^q(1 - \int_{B_R} v^q) \leq \varepsilon(1 + 3 \cdot 2^q), \end{aligned}$$

which finishes the proof. \square

Corollary 3.1. *For all $\delta_1 > 0$ we have*

$$\lim_{\alpha \rightarrow \infty} \int_{B_{\delta_1}(x_\alpha) \cap \partial M} u_\alpha^q = 1.$$

Proof. It follows immediately from Proposition 3.1. \square

Let \tilde{G}_α be the weak solution of

$$\begin{cases} -\operatorname{div}_g(\rho^{1-2\sigma} \nabla_g \tilde{G}_\alpha) = 0, & \text{in } M, \\ \lim_{y \rightarrow x \in \partial M} \rho^{1-2\sigma}(y) \frac{\partial}{\partial \nu} \tilde{G}_\alpha(y) = \delta_{x_\alpha} - \frac{1}{|\partial M|_g}, & \text{on } \partial M, \end{cases}$$

constructed in Theorem A.5. We can find a positive constant $C > 0$ sufficiently large depending only on M, g, n, σ, ρ such that $G_\alpha := \tilde{G}_\alpha + C \geq 1$ on \overline{M} .

Proposition 3.2. Let $\varphi_\alpha(x) = \mu_\alpha^{\frac{n-2\sigma}{2}} G_\alpha(x)$, $\tilde{g}_{ij} = \varphi_\alpha^{\frac{4}{n-2\sigma}} g_{ij}$ and $a = 2 - \frac{2(n-1)}{n-2\sigma}$. Then $w_\alpha := \frac{u_\alpha}{\varphi_\alpha}$ satisfies

$$\begin{cases} \operatorname{div}_{\tilde{g}} \left(\varphi_\alpha^a \rho^{1-2\sigma} \nabla_{\tilde{g}} w_\alpha \right) = 0, & \text{in } M, \\ \lim_{y \rightarrow \bar{x} \in \partial M} \varphi_\alpha^a \rho^{1-2\sigma} \frac{\partial_{\tilde{g}} w_\alpha(y)}{\partial \tilde{\nu}} \leq \xi_\alpha w_\alpha^{q-1}(\bar{x}), & \bar{x} \in \partial M \setminus \{x_\alpha\}, \end{cases} \quad (24)$$

for $\alpha \geq \frac{1}{|\partial M|_g}$.

Proof. The proof follows from some direct computations. For brevity, we drop the subscript α of φ_α and u_α . First of all,

$$\begin{aligned} & \operatorname{div}_{\tilde{g}} \left(\varphi^a \rho^{1-2\sigma} \nabla_{\tilde{g}} \frac{u}{\varphi} \right) \\ &= \varphi^{a-1-\frac{4}{n-2\sigma}} \operatorname{div}_g \left(\rho^{1-2\sigma} \nabla_g u \right) - u \varphi^{a-2-\frac{4}{n-2\sigma}} \operatorname{div}_g \left(\rho^{1-2\sigma} \nabla_g \varphi \right) \\ & \quad + \left(a - 2 + \frac{2(n-1)}{n-2\sigma} \right) \rho^{1-2\sigma} \varphi^{a-2-\frac{4}{n-2\sigma}} \left(\langle \nabla_g u, \nabla_g \varphi \rangle_g - u \varphi |\nabla_g \varphi|_g^2 \right) \\ &= 0. \end{aligned}$$

On the other hand, in Fermi coordinate system centered at \bar{x} ,

$$\begin{aligned} & \lim_{x_{n+1} \rightarrow 0} \varphi^a \rho^{1-2\sigma} \frac{\partial_{\tilde{g}}}{\partial \tilde{\nu}} \left(\frac{u}{\varphi} \right) \\ &= \lim_{x_{n+1} \rightarrow 0} \varphi^a \rho^{1-2\sigma} \left(\frac{1}{\varphi} \frac{\partial u}{\partial x_{n+1}} - \frac{u}{\varphi^2} \frac{\partial \varphi}{\partial x_{n+1}} \right) \tilde{g}^{n+1, n+1} \left\langle \frac{\partial}{\partial x_{n+1}}, \tilde{\nu} \right\rangle_{\tilde{g}} \\ &= \varphi^{a-1-\frac{2}{n-2\sigma}} \left(\xi_\alpha u^{\frac{n+2\sigma}{n-2\sigma}} - \alpha u \right) + \varphi^{a-2-\frac{2}{n-2\sigma}} u \mu_\alpha^{\frac{n-2\sigma}{2}} \frac{1}{|\partial M|} \\ &\leq \xi_\alpha \left(\frac{u}{\varphi} \right)^{\frac{n+2\sigma}{n-2\sigma}} + \varphi^{a-2-\frac{2}{n-2\sigma}} u \mu_\alpha^{\frac{n-2\sigma}{2}} \left(\frac{1}{|\partial M|_g} - \alpha \right) \\ &\leq \xi_\alpha \left(\frac{u}{\varphi} \right)^{\frac{n+2\sigma}{n-2\sigma}}, \end{aligned}$$

provided $\alpha \geq \frac{1}{|\partial M|_g}$. □

Proposition 3.3. Suppose the assumptions in Proposition 3.2. Then there exists some constant C depending only on M, g, n, ρ, σ such that for all $\alpha \geq 1$,

$$w_\alpha \leq C, \quad \text{on } \partial M.$$

Proof. In the following, C denotes some constant which may depend on M, g, n, ρ, σ but not on α and may vary from line to line.

It suffices to prove the proposition for large α , in particular, say, $\alpha \geq \max\{\frac{1}{|\partial M|_g}, 1\}$. Let $\tilde{\rho} := \varphi_\alpha^{\frac{2}{n-2\sigma}} \rho$. Then (24) can be rewritten as

$$\begin{cases} \operatorname{div}_{\tilde{g}} \left(\tilde{\rho}^{1-2\sigma} \nabla_{\tilde{g}} w_\alpha \right) = 0, & \text{in } M, \\ \lim_{y \rightarrow \bar{x}} \tilde{\rho}^{1-2\sigma} \frac{\partial_{\tilde{g}} w_\alpha(y)}{\partial \tilde{\nu}} \leq \xi_\alpha w_\alpha^{q-1}(\bar{x}), & \text{for } \bar{x} \in \partial M \setminus \{x_\alpha\}, \end{cases} \quad (25)$$

where the limit is taken in the sense explained in the paragraph above (13). In the following, we shall abuse notation a little by writing $\psi^{-1}(\mathcal{B}_\delta^+(0))$ as $\mathcal{B}_\delta^+(0)$ where $(\psi^{-1}(\mathcal{B}_\delta^+(0)), \psi)$ is a Fermi coordinate of M at x_α , and denoting $B_\delta(x_\alpha)$ as the geodesic ball on ∂M centered at x_α with radius δ as before. Note that the interior of $\overline{\mathcal{B}_\delta^+(0)} \cap \partial M$ is $B_\delta(x_\alpha)$.

Step 1. We claim that there exist some constants $0 < \delta_2 \ll 1$, $s_0 > q$ independent of α such that

$$\int_{\partial M \setminus B_{\mu_\alpha/\delta_2}(x_\alpha)} w_\alpha^{s_0} ds_{\tilde{g}} \leq C. \quad (26)$$

For any $\varepsilon > 0$, it follows from Proposition 3.1 that there exists a small δ_2 such that

$$\begin{aligned} \int_{\partial M \setminus B_{\mu_\alpha/\delta_2}(x_\alpha)} w_\alpha^q ds_{\tilde{g}} &= \int_{\partial M \setminus B_{\mu_\alpha/\delta_2}(x_\alpha)} u_\alpha^q ds_g \\ &= 1 - \int_{\partial' \mathcal{B}_{1/\delta_2}^+} v_\alpha^q \\ &\leq \varepsilon. \end{aligned} \quad (27)$$

Without loss of generality, we may assume $10\mu_\alpha/\delta_2 < \delta_0$ where δ_0 is the constant such that the Fermi coordinate system centered at x_α exists in $\mathcal{B}_{\delta_0}^+(x_\alpha)$.

We choose η to be some cutoff function satisfying

$$\begin{aligned} \eta(x) &= 1 \text{ if } |x| \geq \mu_\alpha/\delta_2, \quad \eta(x) = 0 \text{ if } |x| \leq \mu_\alpha/(2\delta_2), \\ \text{and } \eta &= \eta(|x|) \text{ in the Fermi coordinate system centered at } x_\alpha. \end{aligned}$$

Multiplying (25) by $w_\alpha^k \eta^2$ for $k > 1$ and integrating by parts, we obtain

$$\int_M \tilde{\rho}^{1-2\sigma} \nabla_{\tilde{g}} w_\alpha \nabla_{\tilde{g}} (w_\alpha^k \eta^2) dv_{\tilde{g}} \leq \xi_\alpha \int_{\partial M} w_\alpha^{q-1+k} \eta^2 ds_{\tilde{g}}.$$

By a direct computation, we see that

$$\begin{aligned}
& \int_M \tilde{\rho}^{1-2\sigma} \nabla_{\tilde{g}} w_\alpha \nabla_{\tilde{g}} (w_\alpha^k \eta^2) \, dv_{\tilde{g}} \\
&= \frac{4k}{(k+1)^2} \int_M \tilde{\rho}^{1-2\sigma} |\nabla_{\tilde{g}} (w_\alpha^{(k+1)/2} \eta)|^2 \, dv_{\tilde{g}} + \frac{k-1}{(k+1)^2} \int_M w_\alpha^{k+1} \operatorname{div}_{\tilde{g}} \left(\tilde{\rho}^{1-2\sigma} \nabla_{\tilde{g}} \eta^2 \right) \, dv_{\tilde{g}} \\
&\quad - \frac{4k}{(k+1)^2} \int_M \tilde{\rho}^{1-2\sigma} w_\alpha^{k+1} |\nabla_{\tilde{g}} \eta|^2 \, dv_{\tilde{g}},
\end{aligned}$$

where we have used that $\lim_{\rho \rightarrow 0} \tilde{\rho}^{1-2\sigma} \frac{\partial_{\tilde{g}} \eta^2}{\partial \nu} = 0$ since η is radial. In conclusion, we obtain

$$\begin{aligned}
& \int_M \tilde{\rho}^{1-2\sigma} |\nabla_{\tilde{g}} (w_\alpha^{(k+1)/2} \eta)|^2 \, dv_{\tilde{g}} \\
&\leq -\frac{k-1}{4k} \int_M w_\alpha^{k+1} \operatorname{div}_{\tilde{g}} \left(\tilde{\rho}^{1-2\sigma} \nabla_{\tilde{g}} \eta^2 \right) \, dv_{\tilde{g}} + \int_M \tilde{\rho}^{1-2\sigma} w_\alpha^{k+1} |\nabla_{\tilde{g}} \eta|^2 \, dv_{\tilde{g}} \\
&\quad + \frac{\xi_\alpha (k+1)^2}{4k} \int_{\partial M} w_\alpha^{q-1+k} \eta^2 \, ds_{\tilde{g}}.
\end{aligned} \tag{28}$$

Since $\tilde{g}^{ij} \sim \mu_\alpha^2 \delta^{ij}$ in $\mathcal{B}_{2\mu_\alpha/\delta_2}^+(x_\alpha) \setminus \mathcal{B}_{\mu_\alpha/(4\delta_2)}^+(x_\alpha)$, we have

$$|\nabla_{\tilde{g}} \eta| + |\nabla_{\tilde{g}}^2 \eta| \leq C.$$

Since η is radial in the Fermi coordinate system, using (65a), (65b) and (65c), we have

$$|\operatorname{div}_{\tilde{g}} (\tilde{\rho}^{1-2\sigma} \nabla_{\tilde{g}} \eta^2)| \leq C \tilde{\rho}^{1-2\sigma}.$$

Taking $1 < k \leq q-1$ in (28) and using Theorem A.1 and Theorem A.5, it follows that

$$\begin{aligned}
& \int_M \tilde{\rho}^{1-2\sigma} |\nabla_{\tilde{g}} (w_\alpha^{(k+1)/2} \eta)|^2 \, dv_{\tilde{g}} \\
&\leq C(k, \delta_2) + \frac{\xi_\alpha (k+1)^2}{4k} \int_{\partial M} w_\alpha^{q-1+k} \eta^2 \, ds_{\tilde{g}} \\
&\leq C(k, \delta_2) + \frac{\xi_\alpha (k+1)^2}{4k} \varepsilon^{(q-2)/q} \left(\int_{\partial M} (w_\alpha^{(1+k)/2} \eta)^q \, ds_{\tilde{g}} \right)^{2/q} \\
&\leq C(k, \delta_2) + C \varepsilon^{(q-2)/q} \int_M \tilde{\rho}^{1-2\sigma} |\nabla_{\tilde{g}} (w_\alpha^{(k+1)/2} \eta)|^2 \, dv_{\tilde{g}},
\end{aligned}$$

where we used

$$\begin{aligned}
& \int_{M \cap (\mathcal{B}_{\mu_\alpha/\delta_2}^+ \setminus \mathcal{B}_{\mu_\alpha/(2\delta_2)}^+)} \tilde{\rho}^{1-2\sigma} w_\alpha^{k+1} dv_{\tilde{g}} \\
& \leq C(\delta_2) \int_{M \cap (\mathcal{B}_{\mu_\alpha/\delta_2}^+ \setminus \mathcal{B}_{\mu_\alpha/(2\delta_2)}^+)} \left(\frac{\rho}{\mu_\alpha}\right)^{1-2\sigma} (\mu_\alpha^{(n-2\sigma)/2} u_\alpha)^{k+1} \mu_\alpha^{-(n+1)} dv_g \\
& \leq C(\delta_2) \int_{1/(2\delta_2) \leq |z| \leq 1/\delta_2} \rho_\alpha(z)^{1-2\sigma} v_\alpha(z)^{k+1} dv_{g_\alpha} \quad \text{by changing variables} \\
& \leq C(k, \delta_2),
\end{aligned} \tag{29}$$

and $\rho_\alpha(z)$, $v_\alpha(z)$ are those in (17).

Taking $\varepsilon > 0$ sufficiently small, we have

$$\int_M \tilde{\rho}^{1-2\sigma} |\nabla_{\tilde{g}}(w_\alpha^{(k+1)/2} \eta)|^2 dv_{\tilde{g}} \leq C.$$

The claim follows immediately from Theorem A.1 in the Appendix.

Step 2. We shall complete the proof by Moser's iterations. Set, for $\delta = \delta_2/10$,

$$R_l = \mu_\alpha \frac{(2 - 2^{-(l-1)})}{\delta}, \quad l = 1, 2, 3, \dots$$

We choose η_l to be some cutoff function satisfying

$$\begin{aligned}
& \eta_l(x) = 1 \text{ if } |x| \geq R_{l+1}, \quad \eta_l(x) = 0 \text{ if } |x| \leq R_l, \\
& \text{and } \eta_l = \eta_l(|x|) \text{ in the Fermi coordinate system centered at } x_\alpha.
\end{aligned}$$

Since $\tilde{g}^{ij} \sim \mu_\alpha^2 \delta^{ij}$ in $\mathcal{B}_{2\mu_\alpha/\delta_2}^+(x_\alpha) \setminus \mathcal{B}_{\mu_\alpha/(4\delta_2)}^+(x_\alpha)$ and η_l is radial in the Fermi coordinate system, we have

$$|\nabla_{\tilde{g}} \eta_l| \leq C2^l, \quad |\operatorname{div}_{\tilde{g}}(\tilde{\rho}^{1-2\sigma} \nabla_{\tilde{g}} \eta_l^2)| \leq C4^l \tilde{\rho}^{1-2\sigma}, \quad \text{and } \lim_{\rho \rightarrow 0} \tilde{\rho}^{1-2\sigma} \frac{\partial_{\tilde{g}} \eta_l^2}{\partial \tilde{\nu}} = 0.$$

In view of (28), we have

$$\begin{aligned}
& \int_M \tilde{\rho}^{1-2\sigma} |\nabla_{\tilde{g}}(w_\alpha^{(k+1)/2} \eta_l)|^2 dv_{\tilde{g}} \\
& \leq C4^l \int_{M \cap (\mathcal{B}_{R_{l+1}}^+(x_\alpha) \setminus \mathcal{B}_{R_l}^+(x_\alpha))} \tilde{\rho}^{1-2\sigma} w_\alpha^{k+1} dv_{\tilde{g}} + \frac{C(k+1)^2}{k} \int_{\partial M \setminus B_{R_l}(x_\alpha)} w_\alpha^{q-1+k} ds_{\tilde{g}}.
\end{aligned} \tag{30}$$

Set $r_0 = s_0/(q-2)$, where s_0 is given in the step 1. It follows Hölder inequality and (26) that

$$\begin{aligned} \int_{\partial M \setminus B_{R_l}(x_\alpha)} w_\alpha^{q-1+k} ds_{\tilde{g}} &= \int_{\partial M \setminus B_{R_l}(x_\alpha)} w_\alpha^{q-2} w_\alpha^{k+1} ds_{\tilde{g}} \\ &\leq C \left(\int_{\partial M \setminus B_{R_l}(x_\alpha)} w_\alpha^{(k+1)r_0/(r_0-1)} ds_{\tilde{g}} \right)^{(r_0-1)/r_0}. \end{aligned} \quad (31)$$

Computing as (29), we see that

$$\begin{aligned} &\int_{M \cap (\mathcal{B}_{R_{l+1}}^+(x_\alpha) \setminus \mathcal{B}_{R_l}^+(x_\alpha))} \tilde{\rho}^{1-2\sigma} w_\alpha^{k+1} dv_{\tilde{g}} \\ &\leq C^{k+1} \int_{2^{-2-(l-1)} \leq \delta |z| \leq 2^{-2-l}} \rho_\alpha(z)^{1-2\sigma} v_\alpha(z)^{k+1} dv_{g_\alpha} \\ &\leq C^{k+1} \delta^{-1} 2^{-l} \max_{\mathcal{B}_{2/\delta}^+} v_\alpha^{k+1}, \end{aligned}$$

and

$$\begin{aligned} &\left(\int_{\partial M \setminus B_{R_l}(x_\alpha)} w_\alpha^{(k+1)r_0/(r_0-1)} ds_{\tilde{g}} \right)^{(r_0-1)/r_0} \\ &\geq C^{-(k+1)} \left(\int_{1 \leq \delta |z'| \leq 2} \rho_\alpha(z', 0)^{1-2\sigma} v_\alpha(z)^{(k+1)r_0/(r_0-1)} ds_{g_\alpha} \right)^{(r_0-1)/r_0} \\ &\geq C^{-(k+1)} \min_{\partial' \mathcal{B}_{2/\delta}^+} v_\alpha^{k+1}. \end{aligned}$$

Hence, it follows from (19) that

$$\begin{aligned} &\left(\int_{M \cap (\mathcal{B}_{R_{l+1}}^+(x_\alpha) \setminus \mathcal{B}_{R_l}^+(x_\alpha))} \tilde{\rho}^{1-2\sigma} w_\alpha^{k+1} dv_{\tilde{g}} \right)^{1/(k+1)} \\ &\leq C \left(\int_{\partial M \setminus B_{R_l}(x_\alpha)} w_\alpha^{(k+1)r_0/(r_0-1)} ds_{\tilde{g}} \right)^{(r_0-1)/r_0(k+1)} \end{aligned} \quad (32)$$

It follows from Theorem A.1, (30), (31) and (32) that

$$\begin{aligned} &\left(\int_{\partial M \setminus B_{R_{l+1}}(x_\alpha)} w_\alpha^{(k+1)q/2} ds_{\tilde{g}} \right)^{2/(k+1)q} \\ &\leq \left(C 4^l + \frac{C(k+1)^2}{k} \right)^{1/(k+1)} \left(\int_{\partial M \setminus B_{R_l}(x_\alpha)} w_\alpha^{(k+1)r_0/(r_0-1)} ds_{\tilde{g}} \right)^{(r_0-1)/r_0(k+1)}. \end{aligned} \quad (33)$$

Set $\chi := \frac{r_0-1}{r_0} \cdot \frac{q}{2} = 1 + \frac{(s_0-q)(q-2)}{2s_0} > 1$, $q_0 = \frac{2r_0}{r_0-1}$, $q_l = q_{l-1} \cdot \chi = \chi^{l-1}q$ and $p_l = q_l(r_0 - 1)/r_0 = 2\chi^l$ where $l \geq 1$. Taking $k = p_l - 1$ in (33), we obtain

$$\|w_\alpha\|_{L^{q_{l+1}}(\partial M \setminus B_{R_{l+1}})} \leq \left(C4^l + \frac{Cp_l^2}{p_l - 1}\right)^{1/p_l} \|w_\alpha\|_{L^{q_l}(\partial M \setminus B_{R_l})}.$$

Therefore,

$$\begin{aligned} \|w_\alpha\|_{L^{q_{l+1}}(\partial M \setminus B_{R_{l+1}})} &\leq \|w_\alpha\|_{L^{q_1}(\partial M \setminus B_{R_1})} \prod_{l=1}^{\infty} \left(C4^l + \frac{Cp_l^2}{p_l - 1}\right)^{1/p_l} \\ &\leq \|w_\alpha\|_{L^{p_1}(\partial M \setminus B_{R_1})} \prod_{l=1}^{\infty} C^{1/(2\chi^l)} (4 + \chi)^{l/(2\chi^l)} \\ &\leq C \|w_\alpha\|_{L^{p_1}(\partial M \setminus B_{R_1})}. \end{aligned}$$

Sending l to ∞ , we have

$$\|w_\alpha\|_{L^\infty(\partial M \setminus B_{2\mu_\alpha/\delta}(x_\alpha))} \leq C. \quad (34)$$

By the choice of G_α , $\varphi_\alpha(x) \geq C^{-1}\mu_\alpha^{-(n-2\sigma)/2}$ for $x \in B_{2\mu_\alpha/\delta}(x_\alpha)$. Hence, for $x \in B_{2\mu_\alpha/\delta}(x_\alpha)$,

$$w_\alpha(x) = \frac{u_\alpha(x)}{\varphi_\alpha(x)} \leq C\mu_\alpha^{(n-2\sigma)/2}u_\alpha(x) \leq C. \quad (35)$$

In view of (34) and (35), we completed the proof of the proposition. \square

Corollary 3.2. *There exists a positive constant C depending only on M, g, n, ρ, σ such that*

$$u_\alpha(x) \leq Cu_\alpha(x_\alpha)^{-1} \text{dist}_{\partial M, g}(x, x_\alpha)^{2\sigma-n}, \quad \text{for all } x \in \partial M.$$

Proof. It follows immediately from Proposition 3.3. \square

4 Proofs of the main theorems

Let u_α and x_α be as in Section 3. We will still use Fermi coordinates $x = (x_1, \dots, x_{n+1})$ centered at x_α . In this coordinate system,

$$\sum_{1 \leq i, j \leq n+1} g_{ij}(x) dx_i dx_j = dx_{n+1}^2 + \sum_{1 \leq i, j \leq n} g_{ij}(x) dx_i dx_j, \quad \text{for } |x| \leq \delta_0,$$

where $\delta_0 > 0$ is independent of α . Then we have

$$\begin{cases} \text{div}_g \left(\rho(x)^{1-2\sigma} \nabla_g u_\alpha(x) \right) = 0, & \text{in } \mathcal{B}_{\delta_0}^+, \\ - \lim_{x_{n+1} \rightarrow 0^+} \rho(x)^{1-2\sigma} \frac{\partial u_\alpha}{\partial x_{n+1}} = \xi_\alpha u_\alpha^{q-1}(x', 0) - \alpha u_\alpha(x', 0), & \text{on } \partial' \mathcal{B}_{\delta_0}^+. \end{cases} \quad (36)$$

Proposition 4.1. *There exists a positive constant C independent of α such that*

$$u_\alpha(x) \leq C u_\alpha(0)^{-1} |x|^{2\sigma-n}, \quad \mathcal{B}_{10\alpha^{-1/2\sigma}}^+(0).$$

Proof. By Corollary 3.2,

$$u_\alpha(x', 0) \leq C u_\alpha(0)^{-1} |x'|^{2\sigma-n}, \quad |x'| \leq \delta_0. \quad (37)$$

Let $r := |\bar{x}| < 10\alpha^{-1/2\sigma}$, $\phi_\alpha(x) = r^{\frac{n-2\sigma}{2}} u_\alpha(rx)$. Then ϕ_α satisfies

$$\begin{cases} \operatorname{div}_{\hat{g}} \left(\hat{\rho}(x)^{1-2\sigma} \nabla_{\hat{g}} \phi_\alpha(x) \right) = 0, & \text{in } \mathcal{B}_{\delta_0/r}^+, \\ - \lim_{x_{n+1} \rightarrow 0^+} \hat{\rho}(x)^{1-2\sigma} \frac{\partial \phi_\alpha}{\partial x_{n+1}} = \xi_\alpha \phi_\alpha^{q-1}(x', 0) - \alpha r^{2\sigma} \phi_\alpha(x', 0), & \text{on } \partial' \mathcal{B}_{\delta_0/r}^+, \end{cases} \quad (38)$$

where $\hat{\rho}(x) = \rho(rx)/r$, $\hat{g}(x) = g_{ij}(rx) dx_i dx_j$. Since $x_\alpha = 0$ is a maximum point of u_α , it follows from (37) that

$$\phi_\alpha(x', 0) = r^{\frac{n-2\sigma}{2}} u_\alpha(rx', 0) \leq C r^{\frac{n-2\sigma}{2}} (r|x'|)^{-\frac{n-2\sigma}{2}} \leq C, \quad \frac{1}{2} < |x'| < 2. \quad (39)$$

Applying the Harnack inequality in [8] or [43] and standard Harnack inequality for uniformly elliptic equations to ϕ_α in $\{x : \frac{1}{2} < |x| < 2, x_{n+1} > 0\}$, we conclude that

$$\max_{\mathcal{B}_{3/2}^+ \setminus \mathcal{B}_{3/4}^+} \phi_\alpha \leq C \min_{\mathcal{B}_{3/2}^+ \setminus \mathcal{B}_{3/4}^+} \phi_\alpha.$$

Hence, by (37)

$$u_\alpha(\bar{x}) \leq C u(\tilde{x}', 0) \leq C u_\alpha(0)^{-1} |\bar{x}|^{2\sigma-n},$$

where $|\tilde{x}'| = |\bar{x}|$. By the arbitrary choice of \bar{x} , the proposition follows immediately. \square

Let $\mu_\alpha = u_\alpha(0)^{-\frac{2}{n-2\sigma}}$, $R_\alpha = (\alpha^{1/2\sigma} \mu_\alpha)^{-1}$, $g_\alpha = g_{ij}(\mu_\alpha x) dx_i dx_j$ and $\rho_\alpha(x) = \frac{\rho(\mu_\alpha x)}{\mu_\alpha}$ in $\mathcal{B}_{10R_\alpha}^+$. Set $v_\alpha(x) = \mu_\alpha^{\frac{n-2\sigma}{2}} u_\alpha(\mu_\alpha x)$ for $x \in \mathcal{B}_{10R_\alpha}^+$. It follows that

$$\begin{cases} \operatorname{div}_{g_\alpha} \left(\rho_\alpha^{1-2\sigma} \nabla_{g_\alpha} v_\alpha \right) = 0, & \text{in } \mathcal{B}_{10R_\alpha}^+, \\ \lim_{x_{n+1} \rightarrow 0} \rho_\alpha^{1-2\sigma} \frac{\partial_{g_\alpha} v_\alpha}{\partial \nu} = \xi_\alpha v_\alpha^{q-1} - \alpha \mu_\alpha^{2\sigma} v_\alpha, & \text{on } \partial' \mathcal{B}_{10R_\alpha}^+ = B_{10R_\alpha}, \\ v_\alpha(0) = 1, \quad 0 < v_\alpha \leq 1. \end{cases} \quad (40)$$

By Proposition 4.1,

$$v_\alpha(x) \leq \frac{C}{1 + |x|^{n-2\sigma}}, \quad x \in \overline{\mathcal{B}}_{10R_\alpha}^+. \quad (41)$$

Proposition 4.2. For all $\alpha \geq 1$, $x \in \mathcal{B}_{R_\alpha}^+(0)$, we have

$$\begin{aligned} |\nabla_{x'} v_\alpha(x', x_{n+1})| &\leq \frac{C}{1 + |x|^{n+1-2\sigma}}, \\ |\nabla_{x'}^2 v_\alpha(x', x_{n+1})| &\leq \frac{C}{1 + |x|^{n+2-2\sigma}}, \\ |\partial_{n+1} v_\alpha(x', x_{n+1})| &\leq \frac{C x_{n+1}^{2\sigma-1}}{1 + |x|^n}. \end{aligned}$$

Proof. Given Theorem A.3 and Proposition A.1, the proofs follow from (41) and standard rescaling arguments (see, e.g., Proposition 3.1 of [32]). \square

Proof of Theorem 1.1. We complete the proof of Theorem 1.1 by checking balance via a Pohozaev type inequality.

It follows from direct computations that

$$\begin{aligned} &2\operatorname{div}(x_{n+1}^{1-2\sigma} \nabla v_\alpha)(\nabla v_\alpha \cdot x) \\ &= \operatorname{div}(2x_{n+1}^{1-2\sigma} (\nabla v_\alpha \cdot x) \nabla v_\alpha - x_{n+1}^{1-2\sigma} |\nabla v_\alpha|^2 x) + (n - 2\sigma) x_{n+1}^{1-2\sigma} |\nabla v_\alpha|^2. \end{aligned} \quad (42)$$

Integrating both sides of (42) over $\mathcal{B}_{R_\alpha}^+$, we have

$$\begin{aligned} &\int_{\mathcal{B}_{R_\alpha}^+} \operatorname{div}(x_{n+1}^{1-2\sigma} \nabla v_\alpha)(\nabla v_\alpha \cdot x) \, dx - \frac{n - 2\sigma}{2} \int_{\mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} |\nabla v_\alpha|^2 \, dx \\ &= \frac{1}{2} \int_{\mathcal{B}_{R_\alpha}^+} \operatorname{div}(2x_{n+1}^{1-2\sigma} (\nabla v_\alpha \cdot x) \nabla v_\alpha - x_{n+1}^{1-2\sigma} |\nabla v_\alpha|^2 x) \, dx. \end{aligned} \quad (43)$$

Integrating by parts, we obtain

$$\begin{aligned} &\frac{1}{2} \int_{\mathcal{B}_{R_\alpha}^+} \operatorname{div}(2x_{n+1}^{1-2\sigma} (\nabla v_\alpha \cdot x) \nabla v_\alpha - x_{n+1}^{1-2\sigma} |\nabla v_\alpha|^2 x) \, dx \\ &= - \int_{\partial' \mathcal{B}_{R_\alpha}^+} \left(\sum_{i=1}^n x_i \frac{\partial v_\alpha}{\partial x_i} \right) \frac{\partial v_\alpha}{\partial x_{n+1}^\sigma} \, dx' + \int_{\partial'' \mathcal{B}_{R_\alpha}^+} |x| x_{n+1}^{1-2\sigma} \left(\left(\frac{\partial v_\alpha}{\partial \nu} \right)^2 - \frac{1}{2} |\nabla v_\alpha|^2 \right) \, dS \\ &= - \int_{\partial' \mathcal{B}_{R_\alpha}^+} \left(\sum_{i=1}^n x_i \frac{\partial v_\alpha}{\partial x_i} \right) \frac{\partial v_\alpha}{\partial x_{n+1}^\sigma} \, dx' + \int_{\partial'' \mathcal{B}_{R_\alpha}^+} \frac{|x|}{2} x_{n+1}^{1-2\sigma} \left(\left(\frac{\partial v_\alpha}{\partial \nu} \right)^2 - |\partial_{\tan} v_\alpha|^2 \right) \, dS, \end{aligned}$$

where $\frac{\partial v_\alpha}{\partial x_{n+1}^\sigma} := \lim_{x_{n+1} \rightarrow 0^+} x_{n+1}^{1-2\sigma} \frac{\partial v_\alpha}{\partial x_{n+1}}$ and ∂_{\tan} denotes the tangential differentiation on $\partial'' \mathcal{B}_{R_\alpha}^+$.

On the other hand,

$$\begin{aligned} \int_{\mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} |\nabla v_\alpha|^2 \, dx &= - \int_{\mathcal{B}_{R_\alpha}^+} \operatorname{div}(x_{n+1}^{1-2\sigma} \nabla v_\alpha) v_\alpha \, dx \\ &\quad - \int_{\partial' \mathcal{B}_{R_\alpha}^+} v_\alpha \frac{\partial v_\alpha}{\partial x_{n+1}^\sigma} \, dx' + \int_{\partial'' \mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} v_\alpha \frac{\partial v_\alpha}{\partial \nu} \, dS. \end{aligned}$$

In summary, we obtain

$$\begin{aligned} & \int_{\mathcal{B}_{R_\alpha}^+} \operatorname{div}(x_{n+1}^{1-2\sigma} \nabla v_\alpha) (\nabla v_\alpha \cdot x) \, dx + \frac{n-2\sigma}{2} \int_{\mathcal{B}_{R_\alpha}^+} \operatorname{div}(x_{n+1}^{1-2\sigma} \nabla v_\alpha) v_\alpha \, dx \\ &= B'(R_\alpha, v_\alpha) + B''(R_\alpha, v_\alpha), \end{aligned} \quad (44)$$

where

$$\begin{aligned} B'(R_\alpha, v_\alpha) &= -\frac{1}{2} \int_{\partial' \mathcal{B}_{R_\alpha}^+} 2 \left(\sum_{i=1}^n x_i \frac{\partial v_\alpha}{\partial x_i} \right) \frac{\partial v_\alpha}{\partial x_{n+1}^\sigma} + (n-2\sigma) v_\alpha \frac{\partial v_\alpha}{\partial x_{n+1}^\sigma} \, dx', \\ B''(R_\alpha, v_\alpha) &= \frac{1}{2} \int_{\partial'' \mathcal{B}_{R_\alpha}^+} |x| x_{n+1}^{1-2\sigma} \left(\left(\frac{\partial v_\alpha}{\partial \nu} \right)^2 - |\partial_{\tan} v_\alpha|^2 \right) + (n-2\sigma) x_{n+1}^{1-2\sigma} v_\alpha \frac{\partial v_\alpha}{\partial \nu} \, dS. \end{aligned}$$

Note that

$$\begin{aligned} & \operatorname{div}_{g_\alpha}(\rho_\alpha^{1-2\sigma} \nabla_{g_\alpha} v_\alpha) \\ &= g_\alpha^{ij} \frac{\partial v_\alpha}{\partial x_i} \frac{\partial \rho_\alpha^{1-2\sigma}}{\partial x_j} + \rho_\alpha^{1-2\sigma} g_\alpha^{ij} \left(\frac{\partial^2 v_\alpha}{\partial x_i \partial x_j} - \Gamma_{ij}^k \frac{\partial v_\alpha}{\partial x_k} \right) \\ &= \operatorname{div}(x_{n+1}^{1-2\sigma} \nabla v_\alpha) + \sum_{1 \leq i, j \leq n} g_\alpha^{ij} \frac{\partial v_\alpha}{\partial x_i} \frac{\partial \rho_\alpha^{1-2\sigma}}{\partial x_j} + \left(\frac{\partial \rho_\alpha^{1-2\sigma}}{\partial x_{n+1}} - \frac{\partial x_{n+1}^{1-2\sigma}}{\partial x_{n+1}} \right) \frac{\partial v_\alpha}{\partial x_{n+1}} \\ &\quad + \rho_\alpha^{1-2\sigma} (g_\alpha^{ij} - \delta^{ij}) \frac{\partial^2 v_\alpha}{\partial x_i \partial x_j} + (\rho_\alpha^{1-2\sigma} - x_{n+1}^{1-2\sigma}) \Delta v_\alpha - \rho_\alpha^{1-2\sigma} g_\alpha^{ij} \Gamma_{ij}^k \frac{\partial v_\alpha}{\partial x_k}, \end{aligned} \quad (45)$$

where Γ_{ij}^k is the Christoffel symbol of g_α . It is easy to see that

$$|h_\alpha^{ij}(x) - \delta^{ij}| \leq C \mu_\alpha |x|, \quad (46a)$$

$$|\Gamma_{ij}^k| \leq C \mu_\alpha, \quad (46b)$$

$$|\rho_\alpha(x)^{1-2\sigma} - x_{n+1}^{1-2\sigma}| \leq C \mu_\alpha x_{n+1}^{2-2\sigma}, \quad (46c)$$

$$\left| \frac{\partial \rho_\alpha(x)^{1-2\sigma}}{\partial x_i} \right| \leq C \mu_\alpha x_{n+1}^{1-2\sigma} \quad \text{for } i < n+1, \quad (46d)$$

$$\left| \frac{\partial \rho_\alpha(x)^{1-2\sigma}}{\partial x_{n+1}} - \frac{\partial x_{n+1}^{1-2\sigma}}{\partial x_{n+1}} \right| \leq C \mu_\alpha x_{n+1}^{1-2\sigma}. \quad (46e)$$

Indeed,

$$\begin{aligned} |\rho_\alpha(x)^{1-2\sigma} - x_{n+1}^{1-2\sigma}| &= x_{n+1}^{1-2\sigma} \left| \left(\frac{\rho(\mu_\alpha x)}{\mu_\alpha x_{n+1}} \right)^{1-2\sigma} - 1 \right| \\ &= x_{n+1}^{1-2\sigma} \left| \left(\frac{\mu_\alpha x_{n+1} + O(\mu_\alpha x_{n+1})^2}{\mu_\alpha x_{n+1}} \right)^{1-2\sigma} - 1 \right| \\ &\leq C \mu_\alpha x_{n+1}^{2-2\sigma}, \end{aligned}$$

and

$$\begin{aligned}
\frac{\partial \rho_\alpha(x)^{1-2\sigma}}{\partial x_i} &= (1-2\sigma)\rho_\alpha(x)^{-2\sigma} \left(\frac{\partial \rho_\alpha(x)}{\partial x_i} - \frac{\partial \rho_\alpha(x', 0)}{\partial x_i} \right) \\
&= O(1)\mu_\alpha \rho_\alpha^{1-2\sigma} \\
&\leq C\mu_\alpha x_{n+1}^{1-2\sigma}.
\end{aligned}$$

It follows from (40), (44), (45) and (46a)-(46e) that

$$\begin{aligned}
&B'(R_\alpha, v_\alpha) + B''(R_\alpha, v_\alpha) \\
&\leq C\mu_\alpha \int_{\mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} (v_\alpha + |\nabla v_\alpha \cdot x|)(|\nabla v_\alpha| + |x||\nabla_{x'}^2 v_\alpha| + x_{n+1}|\Delta v_\alpha|) dx.
\end{aligned} \tag{47}$$

Since $\lim_{x_{n+1} \rightarrow 0} \rho_\alpha^{1-2\sigma} \frac{\partial g_\alpha v_\alpha}{\partial \nu} = -\frac{\partial v_\alpha}{\partial x_{n+1}^\sigma}$ on $\partial' \mathcal{B}_{R_\alpha}^+$,

$$\begin{aligned}
B'(R_\alpha, v_\alpha) &= \int_{\partial' \mathcal{B}_{R_\alpha}^+} \left(\sum_{i=1}^n x_i \frac{\partial v_\alpha}{\partial x_i} \right) (\xi_\alpha v_\alpha^{q-1} - \alpha \mu_\alpha^{2\sigma} v_\alpha) + \frac{(n-2\sigma)}{2} (\xi_\alpha v_\alpha^q - \alpha \mu_\alpha^{2\sigma} v_\alpha^2) dx' \\
&= \sigma \alpha \mu_\alpha^{2\sigma} \int_{\partial' \mathcal{B}_{R_\alpha}^+} v_\alpha^2 dx' + \int_{\partial B_{R_\alpha}} \left(\frac{\xi_\alpha}{q} v_\alpha^q - \frac{\alpha \mu_\alpha^{2\sigma}}{2} v_\alpha^2 \right) R_\alpha dS,
\end{aligned}$$

where integrations by parts were used in the second equality. Clearly,

$$B''(R_\alpha, v_\alpha) = O \left(\int_{\partial'' \mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} (|x||\nabla v_\alpha|^2 + v_\alpha |\nabla v_\alpha|) dS \right).$$

Therefore, we obtain

$$\begin{aligned}
&\alpha \mu_\alpha^{2\sigma} \int_{\partial' \mathcal{B}_{R_\alpha}^+} v_\alpha^2 dx' \\
&\leq C\mu_\alpha \int_{\mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} (v_\alpha + |\nabla v_\alpha \cdot x|)(|\nabla v_\alpha| + |x||\nabla_{x'}^2 v_\alpha| + x_{n+1}|\Delta v_\alpha|) dx \\
&\quad + C \int_{\partial'' \mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} (|x||\nabla v_\alpha|^2 + v_\alpha |\nabla v_\alpha|) dS + C \int_{\partial B_{R_\alpha}} \alpha \mu_\alpha^{2\sigma} v_\alpha^2 R_\alpha dS.
\end{aligned} \tag{48}$$

Since $\operatorname{div}_{g_\alpha}(\rho_\alpha^{1-2\sigma} \nabla_{g_\alpha} v_\alpha) = 0$ and $g_\alpha^{i, n+1} = 0$ for $i < n+1$,

$$|\partial_{n+1}^2 v_\alpha(x', x_{n+1})| \leq C(\mu_\alpha |\nabla v_\alpha| + |\partial_{x+1} v_\alpha| x_{n+1}^{-1} + |\nabla_{x'}^2 v_\alpha|). \tag{49}$$

It follows from (48), (49) and Proposition 4.2 that

$$\begin{aligned}
& \alpha \mu_\alpha^{2\sigma} \int_{\partial' \mathcal{B}_{R_\alpha}^+} v_\alpha^2 \, dx' \\
& \leq C \mu_\alpha \int_{\mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} (v_\alpha + |\nabla v_\alpha \cdot x|)(|\nabla v_\alpha| + |x| |\nabla_{x'}^2 v_\alpha|) \, dx \\
& \quad + C \int_{\partial'' \mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} \left(\frac{1}{R_\alpha^{2n+1-4\sigma}} + \frac{x_{n+1}^{2\sigma-1}}{R_\alpha^{2n-2\sigma}} + \frac{x_{n+1}^{4\sigma-2}}{R_\alpha^{2n-1}} \right) \, dS + C \frac{\alpha \mu_\alpha^{2\sigma}}{R_\alpha^{n-4\sigma}} \\
& \leq C \mu_\alpha \int_{\mathcal{B}_{R_\alpha}^+} \left(\frac{x_{n+1}^{1-2\sigma}}{(1+|x|)^{2n+1-4\sigma}} + \frac{1}{(1+|x|)^{2n-2\sigma}} \right) \, dx \\
& \quad + C R_\alpha^{2\sigma-n} \int_{\partial'' \mathcal{B}_1} (y_{n+1}^{1-2\sigma} + 1 + y_{n+1}^{2\sigma-1}) \, dS + C \frac{\alpha \mu_\alpha^{2\sigma}}{R_\alpha^{n-4\sigma}} \\
& \leq \begin{cases} C \mu_\alpha \ln R_\alpha + C(\alpha \mu_\alpha^{2\sigma})^{\frac{n-2\sigma}{2\sigma}} + C \alpha \mu_\alpha^{2\sigma} R_\alpha^{4\sigma-n}, & n = 2\sigma + 1 \\ C \mu_\alpha + C(\alpha \mu_\alpha^{2\sigma})^{\frac{n-2\sigma}{2\sigma}} + C \alpha \mu_\alpha^{2\sigma} R_\alpha^{4\sigma-n}, & n > 2\sigma + 1. \end{cases}
\end{aligned}$$

For $\sigma = 1/2$ and $n = 2$, Theorem 1.1 was proved in [32]. Hence, we may assume that $n > 2\sigma + 1$. Since $\sigma \in (0, 1/2]$, $n > 2\sigma + 1 \geq 4\sigma$. Therefore,

$$0 < \frac{1}{C} \leq \int_{\partial' \mathcal{B}_{R_\alpha}^+} v_\alpha^2 \, dx' \rightarrow 0, \quad \text{as } \alpha \rightarrow \infty$$

which is a contradiction. \square

Proof of Theorem 1.2. Since ∂M is totally geodesic, Lemma 3.2 implies that

$$|h_\alpha^{ij}(x) - \delta^{ij}| \leq C \mu_\alpha^2 |x|^2, \quad (50a)$$

$$|\Gamma_{ij}^k| \leq C \mu_\alpha^2 |x|. \quad (50b)$$

Since $\rho = d(x) + O(d(x)^3)$, it follows that

$$|\rho_\alpha(x)^{1-2\sigma} - x_{n+1}^{1-2\sigma}| \leq C \mu_\alpha^2 x_{n+1}^{3-2\sigma}, \quad (51a)$$

$$\left| \frac{\partial \rho_\alpha(x)^{1-2\sigma}}{\partial x_i} \right| \leq C \mu_\alpha^2 x_{n+1}^{2-2\sigma}, \quad i < n+1, \quad (51b)$$

$$\left| \frac{\partial \rho_\alpha(x)^{1-2\sigma}}{\partial x_{n+1}} - \frac{\partial x_{n+1}^{1-2\sigma}}{\partial x_{n+1}} \right| \leq C \mu_\alpha^2 x_{n+1}^{2-2\sigma}. \quad (51c)$$

Similar to (48), we have

$$\begin{aligned}
& \alpha \mu_\alpha^{2\sigma} \int_{\partial' \mathcal{B}_{R_\alpha}^+} v_\alpha^2 dx' \\
& \leq C \mu_\alpha^2 \int_{\mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} (v_\alpha + |\nabla v_\alpha \cdot x|) (|x| |\nabla v_\alpha| + |x|^2 |\nabla_{x'}^2 v_\alpha| + x_{n+1}^2 |\Delta v_\alpha|) dx \\
& \quad + C \int_{\partial'' \mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} (|x| |\nabla v_\alpha|^2 + v_\alpha |\nabla v_\alpha|) dS + C \int_{\partial B_{R_\alpha}} \alpha \mu_\alpha^{2\sigma} v_\alpha^2 R_\alpha dS.
\end{aligned} \tag{52}$$

It follows from (49), (52) and Proposition 4.2 that

$$\begin{aligned}
& \alpha \mu_\alpha^{2\sigma} \int_{\partial' \mathcal{B}_{R_\alpha}^+} v_\alpha^2 dx' \\
& \leq C \mu_\alpha^2 \int_{\mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} (v_\alpha + |\nabla v_\alpha \cdot x|) (|x| |\nabla v_\alpha| + |x|^2 |\nabla_{x'}^2 v_\alpha|) dx \\
& \quad + C \int_{\partial'' \mathcal{B}_{R_\alpha}^+} x_{n+1}^{1-2\sigma} (|x| |\nabla v_\alpha|^2 + v_\alpha |\nabla v_\alpha|) dS + C \int_{\partial B_{R_\alpha}} \alpha \mu_\alpha^{2\sigma} v_\alpha^2 R_\alpha dS \\
& \leq C \mu_\alpha^2 \int_{\mathcal{B}_{R_\alpha}^+} \frac{x_{n+1}^{1-2\sigma}}{(1+|x|)^{2n-4\sigma}} dx + C (\alpha \mu_\alpha^{2\sigma})^{\frac{n-2\sigma}{2\sigma}} + C \alpha \mu_\alpha^{2\sigma} R_\alpha^{4\sigma-n} \\
& \leq C \mu_\alpha^2 + C (\alpha \mu_\alpha^{2\sigma})^{\frac{n-2\sigma}{2\sigma}} + C \alpha \mu_\alpha^{2\sigma} R_\alpha^{4\sigma-n},
\end{aligned}$$

provided $n > 2 + 2\sigma$ (i.e., $n \geq 4$). Therefore,

$$0 < \frac{1}{C} \leq \int_{\partial' \mathcal{B}_{R_\alpha}^+} v_\alpha^2 dx' \rightarrow 0 \quad \text{as } \alpha \rightarrow \infty,$$

which is a contradiction. \square

A Appendix

A.1 A trace inequality

Let (M, g) be a smooth, compact Riemannian manifold of dimension $n+1$ ($n \geq 2$) with boundary.

Lemma A.1. *For $n \geq 2$, there exists some positive constant $C = C(n, \sigma)$ such that for all $u \in H^1(x_{n+1}^{1-2\sigma}, \mathcal{B}_1^+)$, $u \equiv 0$ in an open neighborhood of $x = 0$, we have*

$$\left(\int_{\partial' \mathcal{B}_1^+} \frac{|u(x', 0)|^q}{|x'|^{2n}} dx' \right)^{2/q} \leq C \int_{\mathcal{B}_1^+} \frac{x_{n+1}^{1-2\sigma} |\nabla u|^2}{|x|^{2n-4\sigma}} dx.$$

Proof. By the assumption of u , there exists a positive constant $\mu = \mu(u) > 0$ such that $u \equiv 0$ for $|x| < \mu$ with $x_{n+1} > 0$. Consider

$$v(y) = u\left(\frac{y}{|y|^2}\right), \quad |y| > 1, y_{n+1} > 0.$$

It is easy to see that

$$v(y) \equiv 0, \quad \text{for all } |y| > 1/\mu, y_{n+1} > 0,$$

and for some $C(n) > 0$,

$$\int_{\partial' \mathcal{B}_1^+} \frac{|u(x', 0)|^q}{|x'|^{2n}} dx' = C(n) \int_{|y'| \geq 1} |v(y', 0)|^q dy',$$

and

$$\int_{\mathcal{B}_1^+} \frac{x_{n+1}^{1-2\sigma} |\nabla u|^2}{|x|^{2n-4\sigma}} dx = C(n) \int_{|y| \geq 1, y_{n+1} > 0} y_{n+1}^{1-2\sigma} |\nabla v(y)|^2 dy.$$

By some appropriate extension of v to $|y| < 1$, it follows from (3) that

$$\int_{|y'| \geq 1} |v(y', 0)|^q dy' \leq C(n, \sigma) \int_{|y| \geq 1, y_{n+1} > 0} y_{n+1}^{1-2\sigma} |\nabla v(y)|^2 dy.$$

The proof is completed. \square

Lemma A.2. For $\delta > 0$, there exists $C = C(M, g, n, \sigma, \delta, \rho) > 0$ such that for all $x_0 \in \partial M$, $u \in H^1(\rho^{1-2\sigma}, M \setminus \mathcal{B}_{\delta/2}(x_0))$, we have

$$\begin{aligned} & \left(\int_{\partial M \setminus B_\delta(x_0)} |u(x)|^q \right)^{2/q} + \int_{M \setminus \mathcal{B}_\delta^+(x_0)} \rho^{1-2\sigma} |u(x)|^2 \\ & \leq C \left\{ \int_{M \setminus \mathcal{B}_{\delta/2}^+(x_0)} \rho^{1-2\sigma} |\nabla_g u|^2 + \int_{\partial M \cap (B_\delta(x_0) \setminus \overline{B}_{\delta/2}(x_0))} |u(x)|^2 \right\}. \end{aligned} \quad (53)$$

Proof. We prove (53) by contradiction. Suppose the contrary of (53) that for some $\delta > 0$, there exists a sequence of points $\{x_i\} \in \partial M$, $\{u_i\} \in H^1(\rho^{1-2\sigma}, M \setminus \mathcal{B}_{\delta/2}^+(x_i))$ satisfying

$$\left(\int_{\partial M \setminus B_\delta(x_i)} |u_i(x)|^q \right)^{2/q} + \int_{M \setminus \mathcal{B}_\delta^+(x_i)} \rho^{1-2\sigma} |u_i(x)|^2 = 1, \quad (54)$$

but

$$\int_{M \setminus \mathcal{B}_{\delta/2}^+(x_i)} \rho^{1-2\sigma} |\nabla_g u_i|^2 + \int_{\partial M \cap (B_\delta(x_i) \setminus \overline{B}_{\delta/2}(x_i))} |u_i(x)|^2 \leq \frac{1}{i}. \quad (55)$$

After passing to some subsequence, $\{u_i\}$ converges weakly to u in $H^1(\rho^{1-2\sigma}, M \setminus \mathcal{B}_\delta^+(x_i))$. By (55), $u \equiv 0$. It follows from a compact Sobolev embedding in Proposition A.2 that

$$\int_{M \setminus \mathcal{B}_\delta^+(x_i)} \rho^{1-2\sigma} |u_i(x)|^2 \rightarrow 0.$$

By a trace embedding in Proposition 2.3, we also conclude that

$$\left(\int_{\partial M \setminus B_\delta(x_i)} |u(x)|^q \right)^{2/q} \rightarrow 0.$$

Therefore, we reach a contradiction to (54). \square

Theorem A.1. *There exists some constant $C = C(M, g, \rho, n, \sigma)$ such that for all $x_0 \in \partial M$, $\mu > 0$, $u \in H^1(\rho^{1-2\sigma}, M)$, $u \equiv 0$ in $\{x \in M : \text{dist}(x, x_0) < \mu\}$, we have*

$$\left(\int_{\partial M} \frac{|u(x)|^q}{\text{dist}(x, x_0)^{2n}} ds_g \right)^{2/q} \leq C \int_M \frac{\rho^{1-2\sigma} |\nabla_g u|^2}{\text{dist}(x, x_0)^{2n-4\sigma}} dv_g.$$

Proof. The theorem follows clearly from Lemma A.1 and Lemma A.2. \square

A.2 Regularity results for degenerate elliptic equations

Suppose that $a^{ij}(x)$, $1 \leq i, j \leq n+1$, is a smooth positive definite matrix-valued in \mathcal{B}_2^+ and there exists a positive constant $\Lambda \geq 1$ such that

$$\frac{1}{\Lambda} |\xi|^2 \leq a^{ij} \xi_i \xi_j \leq \Lambda |\xi|^2, \quad \forall \xi \in \mathbb{R}^{n+1}$$

Suppose also that

$$a^{i, n+1} = a^{n+1, i} = 0 \text{ for } i < n+1.$$

Consider

$$\begin{cases} \frac{\partial}{\partial x_i} (x_{n+1}^{1-2\sigma} a^{ij}(x) \frac{\partial}{\partial x_j} u(x)) = 0, & \text{in } \mathcal{B}_2^+, \\ - \lim_{x_{n+1} \rightarrow 0^+} x_{n+1}^{1-2\sigma} a^{n+1, n+1} \frac{\partial u(x)}{\partial x_{n+1}} = b(x')u + f(x'), & \text{on } \partial' \mathcal{B}_2^+. \end{cases} \quad (56)$$

We say $u \in H^1(x_{n+1}^{1-2\sigma}, \mathcal{B}_2^+)$ is a weak solution of (56) if

$$\int_{\mathcal{B}_2^+} x_{n+1}^{1-2\sigma} a^{ij}(x) \frac{\partial u}{\partial x_j} \frac{\partial \varphi}{\partial x_i} = \int_{\partial' \mathcal{B}_2^+} b(x')u(x', 0) \varphi(x', 0) + f(x') \varphi(x', 0)$$

for all $\varphi \in C_c^\infty(\mathcal{B}_2^+ \cup \partial' \mathcal{B}_2^+)$.

Theorem A.2. Suppose that $b, f \in L^p(B_2)$ for some $p > \frac{n}{2\sigma}$. Let $u \in H^1(x_{n+1}^{1-2\sigma}, \mathcal{B}_2^+)$ be a weak solution of (56). Then there exist constants $\gamma \in (0, 1)$, $C > 0$ depending only on $n, \sigma, \Lambda, p, \|b\|_{L^p(B_2)}$ such that $u \in C^\gamma(\mathcal{B}_1^+)$ and

$$\|u\|_{C^\gamma(\mathcal{B}_1^+)} \leq C(\|u\|_{L^1(x_{n+1}^{1-2\sigma}, \mathcal{B}_2^+)} + \|f\|_{L^p(B_2)}).$$

Proof. It follows from a modification of the proof of Proposition 2.4 in [26], which uses standard Moser iteration techniques. \square

Theorem A.3. Suppose that $b, f \in C^\beta(B_2)$ for some $0 < \beta \notin \mathbb{N}$. Let $u \in H^1(x_{n+1}^{1-2\sigma}, \mathcal{B}_2^+)$ be a weak solution of (56). Suppose that $2\sigma + \beta$ is not an integer. Then $x_{n+1}^{1-2\sigma} \frac{\partial u(x)}{\partial x_{n+1}} \in C(\overline{\mathcal{B}_1^+})$, and $u(\cdot, 0) \in C^{2\sigma+\beta}(B_1)$. Moreover,

$$\left| x_{n+1}^{1-2\sigma} \frac{\partial u(x)}{\partial x_{n+1}} \right|_{C(\overline{\mathcal{B}_1^+})} + \|u(\cdot, 0)\|_{C^{2\sigma+\beta}(B_1)} \leq C(\|u\|_{L^2(x_{n+1}^{1-2\sigma}, \mathcal{B}_2^+)} + \|f\|_{C^\beta(B_2)}),$$

where $C > 0$ depending only on $n, \sigma, \Lambda, \beta, \|b\|_{C^\beta(B_2)}$.

Proof. It follows from modifications of the proofs of Theorem 2.3 and Lemma 2.3 in [26]. \square

Proposition A.1. Let $b, f \in C^k(B_2)$, $u \in H^1(x_{n+1}^{1-2\sigma}, \mathcal{B}_2^+)$ be a weak solution of (56), where k is a positive integer. Then we have

$$\sum_{j=1}^k \|\nabla_{x'}^j u\|_{L^\infty(\mathcal{B}_1^+)} \leq C(\|u\|_{L^2(x_{n+1}^{1-2\sigma}, \mathcal{B}_2^+)} + \|f\|_{C^k(B_2)}),$$

where $C > 0$ depending only on $n, \sigma, \Lambda, \beta, \|b\|_{C^k(B_2)}$.

Proof. It follows from a modification of the proof of Proposition 2.5 in [26]. \square

A.3 Degenerate elliptic equations with conormal boundary conditions involving measures

We start with some Sobolev embeddings. For every $p \in [1, +\infty)$, we define $W^{1,p}(\rho^{1-2\sigma}, M)$ as the closure of $C^\infty(\overline{M})$ under the norm

$$\|u\|_{W^{1,p}(\rho^{1-2\sigma}, M)} = \left(\int_M \rho^{1-2\sigma} (|u|^p + |\nabla u|^p) dv_g \right)^{\frac{1}{p}},$$

where dv_g denote the volume form of (M, g) . $W^{1,p}(\rho^{1-2\sigma}, M)$ is a Banach space for all $p \in [1, +\infty)$ (see [30]). The following Proposition follows directly from Theorem 8.8 and Theorem 8.12 in [23].

Proposition A.2. Let Ω be a bounded domain in \mathbb{R}^{n+1} with Lipschitz boundary $\partial\Omega$. Let $\sigma \in (0, 1)$, $1 \leq p \leq q < \infty$ with $\frac{1}{n+1} > \frac{1}{p} - \frac{1}{q}$ and $d(x)$ be the distance from x to $\partial\Omega$.

(i) Suppose that $2 - 2\sigma \leq p$. Then $W^{1,p}(d^{1-2\sigma}, \Omega)$ is compactly embedded in $L^q(d^{1-2\sigma}, \Omega)$ if

$$\frac{2 - 2\sigma}{p(n + 2 - 2\sigma)} > \frac{1}{p} - \frac{1}{q}.$$

(ii) Suppose that $2 - 2\sigma > p$. Then $W^{1,p}(d^{1-2\sigma}, \Omega)$ is compactly embedded in $L^q(d^{1-2\sigma}, \Omega)$ if and only if

$$\frac{1}{n + 2 - 2\sigma} > \frac{1}{p} - \frac{1}{q}.$$

Corollary A.1. For $n \geq 2$, let (M, g) be an $n + 1$ dimensional, compact, smooth Riemannian manifold with smooth boundary ∂M . Let $\sigma \in (0, 1)$, and ρ be a defining function of M with $|\nabla_g \rho| = 1$ on ∂M . Let $1 \leq p \leq q < \infty$ with $\frac{1}{n+1} > \frac{1}{p} - \frac{1}{q}$.

(i) Suppose that $2 - 2\sigma \leq p$. Then $W^{1,p}(\rho^{1-2\sigma}, M)$ is compactly embedded in $L^q(d^{1-2\sigma}, M)$ if

$$\frac{2 - 2\sigma}{p(n + 2 - 2\sigma)} > \frac{1}{p} - \frac{1}{q}.$$

(ii) Suppose that $2 - 2\sigma > p$. Then $W^{1,p}(\rho^{1-2\sigma}, M)$ is compactly embedded in $L^q(d^{1-2\sigma}, M)$ if and only if

$$\frac{1}{n + 2 - 2\sigma} > \frac{1}{p} - \frac{1}{q}.$$

Proof. It follows from Proposition A.2 and partition of unity. \square

Proposition A.3. For $n \geq 2$, let (M, g) be an $n + 1$ dimensional, compact, smooth Riemannian manifold with smooth boundary ∂M . Let $\sigma \in (0, 1)$, ρ be a defining function of M with $|\nabla_g \rho| = 1$ on ∂M , and $(u)_{M,\rho} = \int_M \rho^{1-2\sigma} u \, dV_g / \int_M \rho^{1-2\sigma} \, dV_g$. Let $1 < p < \infty$. Then there exists a constant C , depending only on M, g, p, n, σ and ρ , such that

$$\|u - (u)_{M,\rho}\|_{L^p(\rho^{1-2\sigma}, M)} \leq C \|\nabla_g u\|_{L^p(\rho^{1-2\sigma}, M)} \quad (57)$$

for every function $u \in W^{1,p}(\rho^{1-2\sigma}, M)$.

Proof. We argue by contradiction. Were the stated estimate false, there would exist for each integer $k = 1, 2, \dots$ a function $u_k \in W^{1,p}(\rho^{1-2\sigma}, M)$ satisfying

$$\|u_k - (u_k)_{M,\rho}\|_{L^p(\rho^{1-2\sigma}, M)} > k \|\nabla_g u_k\|_{L^p(\rho^{1-2\sigma}, M)}.$$

For each k , define

$$v_k := \frac{u - (u)_{M,\rho}}{\|u - (u)_{M,\rho}\|_{L^p(\rho^{1-2\sigma}, M)}}.$$

Then

$$(v_k)_{M,\rho} = 0, \quad \|v_k\|_{L^p(\rho^{1-2\sigma}, M)} = 1, \quad \|\nabla_g v_k\|_{L^p(\rho^{1-2\sigma}, M)} < 1/k.$$

By Corollary A.1, there exists a subsequence of $\{v_k\}$, which is still denoted as $\{v_k\}$, and a function $v \in L^p(\rho^{1-2\sigma}, M)$ such that

$$v_k \rightarrow v \text{ in } L^p(\rho^{1-2\sigma}, M), \quad v_k \rightharpoonup v \text{ in } W^{1,p}(\rho^{1-2\sigma}, M).$$

Consequently,

$$(v)_{M,\rho} = 0, \quad \|v\|_{L^p(\rho^{1-2\sigma}, M)} = 1, \quad \|\nabla_g v\|_{L^p(\rho^{1-2\sigma}, M)} \leq \liminf_{k \rightarrow \infty} \|\nabla_g v_k\|_{L^p(\rho^{1-2\sigma}, M)} = 0.$$

We reach a contradiction. \square

Corollary A.2. *For $n \geq 2$, let (M, g) be an $n + 1$ dimensional, compact, smooth Riemannian manifold with smooth boundary ∂M . Let $\sigma \in (0, 1)$, ρ be a defining function of M with $|\nabla_g \rho| = 1$ on ∂M , and $(u)_{M,\rho} = \int_M \rho^{1-2\sigma} u \, dV_g / \int_M \rho^{1-2\sigma} \, dV_g$. Let $1 < p < \infty$. Then there exists a constant δ_0 depending only on n, σ, p such that for any $1 \leq k \leq 1 + \delta_0$,*

$$\|u - (u)_{M,\rho}\|_{L^{kp}(\rho^{1-2\sigma}, M)} \leq C \|\nabla_g u\|_{L^p(\rho^{1-2\sigma}, M)} \quad (58)$$

for every function $u \in W^{1,p}(\rho^{1-2\sigma}, M)$, where C is a positive constant depending only on M, g, p, n, σ and ρ ,

Proof. By Corollary A.1, there exists a constant δ_0 depending only on n, σ, p such that for any $1 \leq k \leq 1 + \delta_0$,

$$\begin{aligned} \|u - (u)_{M,\rho}\|_{L^{kp}(\rho^{1-2\sigma}, M)} &\leq C \|\nabla_g u\|_{L^p(\rho^{1-2\sigma}, M)} + C \|u - (u)_{M,\rho}\|_{L^p(\rho^{1-2\sigma}, M)} \\ &\leq C \|\nabla_g u\|_{L^p(\rho^{1-2\sigma}, M)} \end{aligned}$$

where in the last inequality we have used Proposition A.3. \square

Let (M, g) , ρ be as in Theorem 1.1. For $\sigma \in (0, 1)$, we consider

$$\begin{cases} \operatorname{div}_g(\rho^{1-2\sigma} \nabla_g u) = 0, & \text{in } M \\ \lim_{y \rightarrow x \in \partial M} \rho(y)^{1-2\sigma} \frac{\partial_g u}{\partial \nu} = f(x) & \text{on } \partial M. \end{cases} \quad (59)$$

We say $u \in W^{1,1}(\rho^{1-2\sigma}, M)$ is a weak solution of (59) if

$$\int_M \rho^{1-2\sigma} \langle \nabla_g u, \nabla_g \varphi \rangle \, dv_g = \int_{\partial' M} f \varphi \, ds_g \quad (60)$$

for all $\varphi \in C^\infty(\overline{M})$. Define $\tilde{H}^1 := \{u \in H^1(\rho^{1-2\sigma}, M) : \int_M \rho^{1-2\sigma} u \, dv_g = 0\}$.

Lemma A.3. Let $f \in H^{-\sigma}(\partial M) := (H^\sigma(\partial M))^*$, the dual of $H^{-\sigma}(\partial M)$, such that $\langle f, 1 \rangle = 0$. Then (59) admits a unique weak solution $u \in \tilde{H}^1$.

Proof. The lemma follows immediately from Proposition A.3 and the Lax-Milgram theorem. \square

Lemma A.4. Let $f \in L^2(\partial M)$ with zero mean value, $u \in \tilde{H}^1$ be the weak solution of (59). Then for any $\theta > 1$,

$$\int_M \rho^{1-2\sigma} \frac{|\nabla_g u|^2}{(1+|u|)^\theta} dv_g \leq \frac{1}{\theta-1} \|f\|_{L^1(\partial M)}.$$

Proof. In our proofs of this and the next lemma, we adapt some arguments from [6] and [18]. For $\theta > 0$, let $\phi_\theta(r) = \int_0^r \frac{dt}{(1+t)^\theta}$ if $r \geq 0$ and $\phi_\theta(r) = -\phi_\theta(-r)$ if $r < 0$. It is easy to see that $\varphi_\theta := \phi_\theta(u) \in H^1(\rho^{1-2\sigma}, M)$ and $|\varphi_\theta| \leq 1/(\theta-1)$ on \overline{M} if $\theta > 1$. Hence, the Lemma follows from multiplying (60) by letting $\varphi = \varphi_\theta$. \square

Lemma A.5. Let $f \in L^2(\partial M)$ with zero mean value, $u \in \tilde{H}^1$ be the weak solution of (59). Then there exists $\varepsilon_0 > 0$ depending only on n and σ such that for any $1 \leq \tau \leq 1 + \varepsilon_0$, we have

$$\|u\|_{W^{1,\tau}(\rho^{1-2\sigma}, M)} \leq C,$$

where $C > 0$ depends only on $M, g, \sigma, \rho, \|f\|_{L^1(\partial M)}$.

Proof. By the Hölder inequality,

$$\begin{aligned} & \int_M \rho^{1-2\sigma} |\nabla_g u|^\tau dv_g \\ & \leq \left(\int_M \rho^{1-2\sigma} \frac{|\nabla_g u|^2}{(1+|u|)^\theta} dv_g \right)^{\tau/2} \left(\int_M \rho^{1-2\sigma} (1+|u|)^{\frac{\tau\theta}{2-\tau}} dv_g \right)^{(2-\tau)/2} \\ & \leq C(\theta) \left(\int_M \rho^{1-2\sigma} (1+|u|)^{\frac{\tau\theta}{2-\tau}} dv_g \right)^{(2-\tau)/2}, \end{aligned} \quad (61)$$

where we used Lemma A.4 in the last inequality and $\theta \in (1, 2)$ will be chosen later. Applying Corollary A.2 (see also [17]) to $\varphi_{\theta/2}$ yields that for any $1 \leq k \leq 1 + \delta_0$

$$\left(\int_M \rho^{1-2\sigma} |\varphi_{\theta/2} - \fint_M \rho^{1-2\sigma} \varphi_{\theta/2} dv_g|^{2k} dv_g \right)^{1/k} \leq C \int_M \rho^{1-2\sigma} \frac{|\nabla_g u|^2}{(1+|u|)^\theta} dv_g, \quad (62)$$

where $\delta_0 > 0$ depends only on n, σ , and C depends only on M, g, σ, ρ, k . Since $\phi_{\theta/2}(r) \approx |r|^{1-\frac{\theta}{2}}$ for $|r|$ large, it follows from (62) and Lemma A.4 that

$$\left(\int_M \rho^{1-2\sigma} |u|^{k(2-\theta)} \right)^{1/2k} dv_g \leq C + C \int_M \rho^{1-2\sigma} |u|^{1-\frac{\theta}{2}} dv_g. \quad (63)$$

Choosing θ close to 1 such that $k(2 - \theta) = \frac{\tau\theta}{2-\tau}$ (this can be achieved as long as τ is closed to 1) and inserting (63) to (61), we obtain

$$\begin{aligned} \left(\int_M \rho^{1-2\sigma} |\nabla_g u|^\tau dv_g \right)^{1/\tau} &\leq C \left(1 + \int_M \rho^{1-2\sigma} |u|^{1-\frac{\theta}{2}} dv_g \right)^{\frac{\theta}{2-\theta}} \\ &\leq C + C \left(\int_M \rho^{1-2\sigma} |u| dv_g \right)^{\frac{\theta}{2}} \end{aligned} \quad (64)$$

Since $\int_M \rho^{1-2\sigma} u dv_g = 0$, by the Poincaré-Sobolev inequality, Hölder inequality and (64), we have

$$\|u\|_{L^1(\rho^{1-2\sigma}, M)} \leq C \int_M \rho^{1-2\sigma} |\nabla_g u| dv_g \leq C(1 + \|u\|_{L^1(\rho^{1-2\sigma}, M)}^{\frac{\theta}{2}}).$$

Thus, $\|u\|_{L^1(\rho^{1-2\sigma}, M)} \leq C$ because $\frac{\theta}{2} < 1$. Therefore, the lemma follows immediately from (64) and the Poincaré-Sobolev inequality. \square

Theorem A.4. *For any bounded radon measure f defined on ∂M with $\langle f, 1 \rangle = 0$, there exists a weak solution $u \in W^{1,1+\varepsilon_0}(\rho^{1-2\sigma}, M)$ of (59).*

Proof. The proof follows from Lemma A.3 and A.5 and some standard approximating procedure, see, e.g., [18]. We omit the details here. \square

Theorem A.5. *For $x_0 \in \partial M$, let $f = \delta_{x_0} - \frac{1}{|\partial M|_g}$, where $|\partial M|_g$ is the area of ∂M with respect to the induced metric g . Then there exists a weak solution $u \in W^{1,1+\varepsilon_0}(\rho^{1-2\sigma}, M)$ of (59) with mean value zero and for all $x \in \overline{M} \setminus \{x_0\}$,*

$$A_1 \text{dist}_g(x, x_0)^{2\sigma-n} - A_0 \leq u(x) \leq A_2 \text{dist}_g(x, x_0)^{2\sigma-n}, \quad (65a)$$

$$|\nabla_{\tan} u| \leq A_3 \text{dist}_g(x, x_0)^{2\sigma-n-1}, \quad (65b)$$

$$\left| \frac{\partial u}{\partial \nu} \right| \leq A_4 \rho^{2\sigma-1} \text{dist}_g(x, x_0)^{-n}, \quad (65c)$$

where A_0, A_1, A_2, A_3, A_4 are positive constants depending only on M, g, n, σ, ρ .

Proof. Let $f_k \in C^1(\partial M)$ with $\int_{\partial M} f_k ds_g = 0$, $\|f_k\|_{L^1(\partial M)} \leq C$ independent of k , such that $f_k \rightarrow f$ in distribution sense as $k \rightarrow \infty$. We can also assume that $f_k \rightarrow f$ in $C_{loc}^1(\partial M \setminus \{x_0\})$. By Lemma A.3 and Lemma A.5, there exists a unique solution $u_k \in \tilde{H}^1$ of (59) with f replaced by f_k , and

$$\|u_k\|_{W^{1,1+\varepsilon_0}(\rho^{1-2\sigma}, M)} \leq C(\|f_k\|_{L^1(\partial M)}) \leq C.$$

Moreover, it follows from Moser's iterations (see, e.g., the proof of Theorem A.2) that there exists some $\alpha > 0$ such that

$$\|u_k\|_{C^\alpha(M \setminus \mathcal{B}_r(x_0))} \leq C(r) \quad (66)$$

for any $r > 0$. By standard compactness arguments, $u_k \rightharpoonup u$ in $W^{1,1+\varepsilon_0}(\rho^{1-2\sigma}, M)$ for some u , which is a weak solution of (59) and satisfies

$$\|u\|_{C^{\alpha/2}(M \setminus \mathcal{B}_r(x_0))} \leq C(r).$$

Now, it suffices to establish the estimate (65a) for $x \in B_r(x_0)$. For r suitably small, choose a Fermi coordinate system $\{y_1, \dots, y_{n+1}\}$ centered at x_0 . Then $u_k(y)$ satisfies

$$\begin{cases} \partial_i(\rho^{1-2\sigma} \sqrt{\det g} g^{ij} \partial_j u_k) = 0, & \text{in } \mathcal{B}_{2r}^+, \\ -\lim_{y_{n+1} \rightarrow 0} \rho^{1-2\sigma} \sqrt{\det g} \frac{\partial u_k}{\partial y_{n+1}} = f_k, & \text{on } \partial' \mathcal{B}_{2r}^+. \end{cases}$$

Let v_k be the unique weak solution of

$$\begin{cases} \partial_i(\rho^{1-2\sigma} \sqrt{\det g} g^{ij} \partial_j v_k) = 0, & \text{in } \mathcal{B}_{2r}^+, \\ -\lim_{y_{n+1} \rightarrow 0} \rho^{1-2\sigma} \sqrt{\det g} \frac{\partial v_k}{\partial y_{n+1}} = -\frac{1}{|\partial M|}, & \text{on } \partial' \mathcal{B}_{2r}^+, \\ v_k = u_k & \text{on } \partial'' \mathcal{B}_{2r}^+. \end{cases}$$

in $H^1(\rho^{1-2\sigma}, M)$. In view of (66), $\|v_k\|_{L^\infty(\mathcal{B}_{2r})} \leq C(r)$ and hence $\|v_k\|_{C^\alpha(\mathcal{B}_r^+)} \leq C(r)$. Moreover, $w_k := u_k - v_k \in H^1(\rho^{1-2\sigma}, M)$ satisfies

$$\begin{cases} \partial_i(\rho^{1-2\sigma} \sqrt{\det g} g^{ij} \partial_j w_k) = 0, & \text{in } \mathcal{B}_{2r}^+, \\ -\lim_{y_{n+1} \rightarrow 0} \rho^{1-2\sigma} \sqrt{\det g} \frac{\partial w_k}{\partial y_{n+1}} = f_k + \frac{1}{|\partial M|}, & \text{on } \partial' \mathcal{B}_{2r}^+, \\ w_k = 0 & \text{on } \partial'' \mathcal{B}_{2r}^+. \end{cases}$$

Recall that $g^{i,n+1} = 0$ for $i < n+1$ on $\partial' \mathcal{B}_{2r}^+$. Let \bar{w}_k be the even extension of w_k in \mathcal{B}_{2r} , i.e.,

$$\bar{w}_k = \begin{cases} w_k(y', y_{n+1}), & y_{n+1} \geq 0, \\ w_k(y', -y_{n+1}), & y_{n+1} \leq 0. \end{cases}$$

We also evenly extend g and ρ to be \bar{g} and $\bar{\rho}$, respectively. It is easy to verify that the weak limit w of \bar{w}_k in $L^{1+\varepsilon_0}(\rho^{1-2\sigma}, \mathcal{B}_{2r})$ is the *weak solution vanishing on $\partial \mathcal{B}_{2r}$* (see page 162 of [16]) of

$$\partial_i(\bar{\rho}^{1-2\sigma} \sqrt{\det \bar{g}} \bar{g}^{ij} \partial_j w) = -2\delta_0 \quad \text{in } \mathcal{B}_{2r}.$$

It follows from Theorem 3.3 of [16] that w satisfies the estimates (65a) in $\mathcal{B}_r(x_0)$. Thus, u satisfies (65a). Finally, (65b) and (65c) follows from (65a), Theorem A.3, Proposition A.1 and some scaling arguments. \square

References

- [1] Adimurthi; Yadava, S. L.: *Some remarks on Sobolev type inequalities*, Calc. Var. Partial Differential Equations **2** (1994), 427–442.
- [2] Aubin, T.: Problèmes isopérimétriques et espaces de Sobolev, J. Differential Geom. **11** (1976), no. 4, 573–598.
- [3] Aubin, T.: *Some nonlinear problems in Riemannian geometry*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1998. xviii+395 pp. ISBN: 3-540-60752-8.
- [4] Aubin, T.; Li, Y.Y.: *On the best Sobolev inequality*, J. Math. Pures Appl., **78** (1999), 353–387.
- [5] Beckner, W.: *Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality*, Ann. of Math. (2) **138** (1993), 213–242.
- [6] Boccardo, L.; Gallouët, T.: *Nonlinear elliptic and parabolic equations involving measure data*, J. Funct. Anal. **87** (1989), 149–169.
- [7] Brezis, H.; Strauss, W. A.: *Semi-linear second-order elliptic equations in L^1* , J. Math. Soc. Japan **25** (1973), 565–590.
- [8] Cabre, X.; Sire, Y.: *Nonlinear equations for fractional Laplacians I: Regularity, maximum principles, and Hamiltonian estimates*, arXiv:1012.0867.
- [9] Caffarelli, L.; Silvestre, L.: *An extension problem related to the fractional Laplacian*, Comm. Partial. Diff. Equ., **32** (2007), 1245–1260.
- [10] Chang, S.-Y.; González, M.: *Fractional Laplacian in conformal geometry*, Adv. Math. **226** (2011), 1410–1432.
- [11] Druet, O.: *The best constants problem in Sobolev inequalities*, Math. Ann., **314** (1999), 327–346.
- [12] Druet, O.: *Isoperimetric inequalities on compact manifolds*, Geom. Dedicata, **90** (2002), 217–236.
- [13] Druet, O.; Hebey, E.: *The AB program in geometric analysis: sharp Sobolev inequalities and related problems*, Mem. Amer. Math. Soc. **160** (2002), no. 761.
- [14] Escobar, J. F.: *Sharp constant in a Sobolev trace inequality*, Indiana Univ. Math. J. **37** (1988), 687–698.
- [15] —: *Conformal deformation of a Riemannian metric to a scalar flat metric with constant mean curvature on the boundary*, Ann. of Math. (2) **136** (1992), no. 1, 1–50.
- [16] Fabes, E.; Jerison, D.; Kenig, C.: *The Wiener test for degenerate elliptic equations*, Annales de l’institut Fourier, **32** (1982), 151–182.
- [17] Fabes, E.; Kenig, C. and Serapioni, R.: *The local regularity of solutions of degenerate elliptic equations*, Comm. Partial Differential Equations **7** (1982), 77–116.
- [18] Gallouët, T.; Sire, Y.: *Some possibly degenerate elliptic problems with measure data and non linearity on the boundary*, arXiv:1002.4982v1.
- [19] González, M.: *Gamma convergence of an energy functional related to the fractional Laplacian*, Calc. Var. Partial Differential Equations **36** (2009), 173–210.
- [20] González, M.; Mazzeo, R.; Sire, Y.: *Singular Solutions of Fractional Order Conformal Laplacians*, to appear in J. Geom. Anal..
- [21] González, M.; Qing, J.: *Fractional conformal Laplacians and fractional Yamabe problems*, preprint, arXiv:1012.0579v1.
- [22] Graham, C.R.; Zworski, M.: *Scattering matrix in conformal geometry*, Invent. Math. **152** (2003), 89–118.
- [23] Gurka, P.; Opic, B.: *Continuous and compact imbeddings of weighted Sobolev spaces II*, Czechoslovak Math. J. **39** (1989), 78–94.

- [24] Hebey, E.: “Nonlinear analysis on manifolds: Sobolev spaces and inequalities,” Courant Lecture Notes in Math., 5. New York University, Courant Institute of Mathematical Sciences, New York; Amer. Math. Soc., Providence, RI, 1999.
- [25] Hebey, E.; Vaugon, M.: *Meilleures constantes dans le théorème d’inclusion de Sobolev*, Ann. Inst. H. Poincaré, **13** (1996), 57–93.
- [26] Jin, T.; Li, Y.Y.; Xiong, J.: *On a fractional Nirenberg problem, part I: blow up analysis and compactness of solutions*, arXiv:1111.1332v1.
- [27] Jin, T.; Li, Y.Y.; Xiong, J.: *On a fractional Nirenberg problem, part II: existence of solutions*, in preparation, 2011.
- [28] Jin, T.; Xiong, J.: *A fractional Yamabe flow and some applications*, arXiv:1110.5664v1.
- [29] Kenig, C.; Pipher, J.: *The Neumann problem for elliptic equations with non-smooth coefficients*, Invent. Math. **113** (1993), 447–509.
- [30] Kufner, A.: *Weighted sobolev spaces*, John Wiley & Sons, Inc., New York, 1985.
- [31] Li, Y.Y.; Ricciardi, T.: *A sharp Sobolev inequality on Riemannian manifolds*, Comm. Pure Appl. Anal. **2** (2003), 1–31.
- [32] Li, Y.Y.; Zhu, M.: *Sharp Sobolev trace inequalities on Riemannian manifolds with boundaries*, Comm. Pure Appl. Math. **50** (1997), 449–487.
- [33] Li, Y.Y.; Zhu, M.: *Sharp Sobolev inequalities involving boundary terms*, Geom. Funct. Anal. **8** (1998), 59–87.
- [34] Lieb, E.H.: *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*, Ann. of Math. (2) **118** (1983), 349–374.
- [35] Lions, P.-L.: *The concentration-compactness principle in the calculus of variations, The limit case, II*, Rev. Mat. Iberoamericana **1** (1985), no.2, 45–121.
- [36] Maz’ya, V.: *Sobolev spaces with applications to elliptic partial differential equations*. Second, revised and augmented edition. Springer, Heidelberg, 2011.
- [37] Mazzeo, R.: *The Hodge cohomology of a conformally compact metric*, J. Differential Geom. **28** (1988), 309–339.
- [38] Mazzeo, R.; Melrose, R.: *Meromorphic extension of the resolvent on complete spaces with asymptotically constant negative curvature*, J. Funct. Anal. **75** (1987), 260–310.
- [39] Nekvinda, A.: *Characterization of traces of the weighted Sobolev space $W^{1,p}(\Omega, d_M^\varepsilon)$ on M* , Czechoslovak Math. J., **43** (1993), 695–711.
- [40] Palatucci, G.; Sire, Y.: *Γ -convergence of some super quadratic functionals with singular weights*, Math. Z. **266** (2010), 533–560.
- [41] Qing, J.; Raske, D.: *On positive solutions to semilinear conformally invariant equations on locally conformally flat manifolds*, Int. Math. Res. Not. 2006, Art. ID 94172, 20 pp.
- [42] Schoen, R.; Yau, S. T.: *Lectures on differential Geometry*. International Press, Cambridge, MA, 1994.
- [43] Tan, J.; Xiong, J.: *A Harnack inequality for fractional Laplace equations with lower order terms*, DCDS-A **31** (2011), 975–983.

Tianling Jin
 Department of Mathematics, Rutgers University
 110 Frelinghuysen Road, Piscataway, NJ 08854, USA
 Email: kingbull@math.rutgers.edu
 Jingang Xiong

School of Mathematical Sciences, Beijing Normal University
Beijing 100875, China

and

Department of Mathematics, Rutgers University
110 Frelinghuysen Road, Piscataway, NJ 08854, USA
Email: jxiong@mail.bnu.edu.cn/jxiong@math.rutgers.edu