

A note on property (gb) and perturbations ^{*}

Qingping ZENG[†] Huaijie ZHONG

(School of Mathematics and Computer Science, Fujian Normal University, Fuzhou 350007, P.R. China)

Abstract: An operator $T \in \mathcal{B}(X)$ defined on a Banach space X satisfies property (gb) if the complement in the approximate point spectrum $\sigma_a(T)$ of the upper semi-B-Weyl spectrum $\sigma_{SBF_+^-}(T)$ coincides with the set $\Pi(T)$ of all poles of the resolvent of T . In this note we continue to study property (gb) and the stability of it, for a bounded linear operator T acting on a Banach space, under perturbations by nilpotent operators, by finite rank operators, by quasi-nilpotent operators commuting with T . Two counterexamples show that property (gb) in general is not preserved under commuting quasi-nilpotent perturbations or commuting finite rank perturbations.

2010 Mathematics Subject Classification: primary 47A10, 47A11; secondary 47A53, 47A55

Key words: Generalized a-Browder's theorem; property (gb) ; eventual topological uniform descent; commuting perturbation.

1 Introduction

Throughout this note, let $\mathcal{B}(X)$ denote the Banach algebra of all bounded linear operators acting on an infinite dimensional complex Banach space X , and $\mathcal{F}(X)$ denote its ideal of finite rank operators on X . For an operator $T \in \mathcal{B}(X)$, let T^* denote its dual, $\mathcal{N}(T)$ its kernel, $\alpha(T)$ its nullity, $\mathcal{R}(T)$ its range, $\beta(T)$ its defect, $\sigma(T)$ its spectrum and $\sigma_a(T)$ its approximate point spectrum. If the range $\mathcal{R}(T)$ is closed and $\alpha(T) < \infty$ (resp. $\beta(T) < \infty$), then T is said to be *upper semi-Fredholm* (resp. *lower semi-Fredholm*). If $T \in \mathcal{B}(X)$ is both upper and lower semi-Fredholm, then T is said to be *Fredholm*. If $T \in \mathcal{B}(X)$ is either upper or lower semi-Fredholm, then T is said to be *semi-Fredholm*, and its index is defined by $\text{ind}(T) = \alpha(T) - \beta(T)$. The *upper semi-Weyl operators* are defined as the class of upper semi-Fredholm operators with index less than or equal to zero, while *Weyl operators* are defined as the class of Fredholm operators of index zero. These classes of operators generate the following spectra: the *Weyl spectrum* defined

^{*}This work has been supported by National Natural Science Foundation of China (11171066), Specialized Research Fund for the Doctoral Program of Higher Education (2010350311001, 20113503120003), Natural Science Foundation of Fujian Province (2009J01005, 2011J05002) and Foundation of the Education Department of Fujian Province (JB10042, JA08036).

[†]Email address: zqping2003@163.com.

by

$$\sigma_W(T) := \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a Weyl operator}\},$$

the *upper semi-Weyl spectrum* (in literature called also *Weyl essential approximate point spectrum*) defined by

$$\sigma_{SF_+^-}(T) := \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a upper semi-Weyl operator}\}.$$

Recall that the *descent* and the *ascent* of $T \in \mathcal{B}(X)$ are $dsc(T) = \inf\{n \in \mathbb{N} : \mathcal{R}(T^n) = \mathcal{R}(T^{n+1})\}$ and $asc(T) = \inf\{n \in \mathbb{N} : \mathcal{N}(T^n) = \mathcal{N}(T^{n+1})\}$, respectively (the infimum of an empty set is defined to be ∞). If $asc(T) < \infty$ and $\mathcal{R}(T^{asc(T)+1})$ is closed, then T is said to be *left Drazin invertible*. If $dsc(T) < \infty$ and $\mathcal{R}(T^{dsc(T)})$ is closed, then T is said to be *right Drazin invertible*. If $asc(T) = dsc(T) < \infty$, then T is said to be *Drazin invertible*. Clearly, $T \in \mathcal{B}(X)$ is both left and right Drazin invertible if and only if T is Drazin invertible. An operator $T \in \mathcal{B}(X)$ is called *upper semi-Browder* if it is a upper semi-Fredholm operator with finite ascent, while T is called *Browder* if it is a Fredholm operator of finite ascent and descent. The *Browder spectrum* of $T \in \mathcal{B}(X)$ is defined by

$$\sigma_B(T) := \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a Browder operator}\},$$

the *upper semi-Browder spectrum* (in literature called also *Browder essential approximate point spectrum*) is defined by

$$\sigma_{UB}(T) := \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a upper semi-Browder operator}\}.$$

An operator $T \in \mathcal{B}(X)$ is called *Riesz* if its essential spectrum $\sigma_e(T) := \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Fredholm}\} = \{0\}$.

Suppose that $T \in \mathcal{B}(X)$ and that $R \in \mathcal{B}(X)$ is a Riesz operator commuting with T . Then it follows from [27, Proposition 5] and [25, Theorem 1] that

$$\sigma_{SF_+^-}(T + R) = \sigma_{SF_+^-}(T); \quad (1.1)$$

$$\sigma_W(T + R) = \sigma_W(T); \quad (1.2)$$

$$\sigma_{UB}(T + R) = \sigma_{UB}(T); \quad (1.3)$$

$$\sigma_B(T + R) = \sigma_B(T). \quad (1.4)$$

For each integer n , define T_n to be the restriction of T to $\mathcal{R}(T^n)$ viewed as the map from $\mathcal{R}(T^n)$ into $\mathcal{R}(T^n)$ (in particular $T_0 = T$). If there exists $n \in \mathbb{N}$ such that $\mathcal{R}(T^n)$ is closed and T_n is upper semi-Fredholm, then T is called *upper semi-B-Fredholm*. It follows from [8, Proposition 2.1] that if there exists $n \in \mathbb{N}$ such that $\mathcal{R}(T^n)$ is closed and T_n is upper semi-Fredholm, then $\mathcal{R}(T^m)$ is closed, T_m is upper semi-Fredholm and $\text{ind}(T_m) = \text{ind}(T_n)$ for all $m \geq n$. This enables us to define the index of a upper semi-B-Fredholm operator T as the index of the upper semi-Fredholm operator T_n , where n is an integer satisfying $\mathcal{R}(T^n)$ is closed and T_n is upper semi-Fredholm. An operator $T \in \mathcal{B}(X)$ is called *upper semi-B-Weyl* if T is upper semi-B-Fredholm and $\text{ind}(T) \leq 0$.

For $T \in \mathcal{B}(X)$, let us define the *left Drazin spectrum*, the *Drazin spectrum* and the *upper semi-B-Weyl spectrum* of T as follows respectively:

$$\sigma_{LD}(T) := \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a left Drazin invertible operator}\};$$

$$\sigma_D(T) := \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a Drazin invertible operator}\};$$

$$\sigma_{SBF_+^-}(T) := \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a upper semi-B-Weyl operator}\}.$$

Let $\Pi(T)$ denote the set of all poles of T . We say that $\lambda \in \sigma_a(T)$ is a left pole of T if $T - \lambda I$ is left Drazin invertible. Let $\Pi_a(T)$ denote the set of all left poles of T . It is well known that $\Pi(T) = \sigma(T) \setminus \sigma_D(T) = \text{iso}\sigma(T) \setminus \sigma_D(T)$ and $\Pi_a(T) = \sigma_a(T) \setminus \sigma_{LD}(T) = \text{iso}\sigma_a(T) \setminus \sigma_{LD}(T)$. Here and henceforth, for $A \subseteq \mathbb{C}$, $\text{iso}A$ is the set of isolated points of A . An operator $T \in \mathcal{B}(X)$ is called *a-polaroid* if $\text{iso}\sigma_a(T) = \emptyset$ or every isolated point of $\sigma_a(T)$ is a left pole of T .

Following Harte and Lee [19] we say that $T \in \mathcal{B}(X)$ satisfies Browder's theorem if $\sigma_W(T) = \sigma_B(T)$. While, according to Djordjević and Han [12], we say that T satisfies a-Browder's theorem if $\sigma_{SF_+^-}(T) = \sigma_{UB}(T)$.

The following two variants of Browder's theorem have been introduced by Berkani and Zariouh [9] and Berkani and Koliha [7], respectively.

Definition 1.1. An operator $T \in \mathcal{B}(X)$ is said to possess property *(gb)* if

$$\sigma_a(T) \setminus \sigma_{SBF_+^-}(T) = \Pi(T).$$

While $T \in \mathcal{B}(X)$ is said to satisfy generalized a-Browder's theorem if

$$\sigma_a(T) \setminus \sigma_{SBF_+^-}(T) = \Pi_a(T).$$

From formulas (1.1)–(1.4), it follows immediately that Browder's theorem and a-Browder's theorem are preserved under commuting Riesz perturbations. It is proved in [4, Theorem 2.2] that generalized a-Browder's theorem is equivalent to a-Browder's theorem. Hence, generalized a-Browder's theorem is stable under commuting Riesz perturbations. That is, if $T \in \mathcal{B}(X)$ satisfies generalized a-Browder's theorem and R is a Riesz operator commuting with T , then $T + R$ satisfies generalized a-Browder's theorem.

The single-valued extension property was introduced by Dunford in [13, 14] and has an important role in local spectral theory and Fredholm theory, see the recent monographs [1] by Aiena and [22] by Laursen and Neumann.

Definition 1.2. An operator $T \in \mathcal{B}(X)$ is said to have the single-valued extension property at $\lambda_0 \in \mathbb{C}$ (SVEP at λ_0 for brevity), if for every open neighborhood U of λ_0 the only analytic function $f : U \rightarrow X$ which satisfies the equation $(\lambda I - T)f(\lambda) = 0$ for all $\lambda \in U$ is the function $f(\lambda) \equiv 0$.

Let $S(T) := \{\lambda \in \mathbb{C} : T \text{ does not have the SVEP at } \lambda\}$. An operator $T \in \mathcal{B}(X)$ is said to have SVEP if $S(T) = \emptyset$.

In this note we continue the study of property *(gb)* which is studied in some recent papers [9, 10, 26]. We show that property *(gb)* is satisfied by an operator T satisfying

$S(T^*) \subseteq \sigma_{SBF_+^-}(T)$. We give a revised proof of [26, Theorem 3.10] to prove that property (gb) is preserved under commuting nilpotent perturbations. We show also that if $T \in \mathcal{B}(X)$ satisfies $S(T^*) \subseteq \sigma_{SBF_+^-}(T)$ and F is a finite rank operator commuting with T , then $T + F$ satisfies property (gb) . We show that if $T \in \mathcal{B}(X)$ is a a -polaroid operator satisfying property (gb) and Q is a quasi-nilpotent operator commuting with T , then $T + Q$ satisfies property (gb) . Two counterexamples are also given to show that property (gb) in general is not preserved under commuting quasi-nilpotent perturbations or commuting finite rank perturbations. These results improve and revise some recent results of Rashid in [26].

2 Main results

We begin with the following lemmas.

Lemma 2.1. ([9, Corollary 2.9]) *An operator $T \in \mathcal{B}(X)$ possesses property (gb) if and only if T satisfies generalized a -Browder's theorem and $\Pi(T) = \Pi_a(T)$.*

Lemma 2.2. *If the equality $\sigma_{SBF_+^-}(T) = \sigma_D(T)$ holds for $T \in \mathcal{B}(X)$, then T possesses property (gb) .*

Proof. Suppose that $\sigma_{SBF_+^-}(T) = \sigma_D(T)$. If $\lambda \in \sigma_a(T) \setminus \sigma_{SBF_+^-}(T)$, then $\lambda \in \sigma_a(T) \setminus \sigma_D(T) \subseteq \Pi(T)$. This implies that $\sigma_a(T) \setminus \sigma_{SBF_+^-}(T) = \Pi(T)$. Since $\Pi(T) \subseteq \sigma_a(T) \setminus \sigma_{SBF_+^-}(T)$ is always true, $\sigma_a(T) \setminus \sigma_{SBF_+^-}(T) = \Pi(T)$, i.e. T possesses property (gb) . \square

Lemma 2.3. *If $T \in \mathcal{B}(X)$, then $\sigma_{SBF_+^-}(T) \cup S(T^*) = \sigma_D(T)$.*

Proof. Let $\lambda \notin \sigma_{SBF_+^-}(T) \cup S(T^*)$. Then $T - \lambda$ is a upper semi-Weyl operator and T^* has SVEP at λ . Thus $T - \lambda$ is a upper semi-B-Fredholm operator and $\text{ind}(T - \lambda) \leq 0$. Hence there exists $n \in \mathbb{N}$ such that $\mathcal{R}((T - \lambda)^n)$ is closed, $(T - \lambda)_n$ is a upper semi-Fredholm operator and $\text{ind}(T - \lambda)_n \leq 0$. By [2, Theorem 2.11], $dsc(T - \lambda) < \infty$. Thus $dsc(T - \lambda)_n < \infty$, by [1, Theorem 3.4(ii)], $\text{ind}(T - \lambda)_n \geq 0$. By [1, Theorem 3.4(iv)], $asc(T - \lambda)_n = dsc(T - \lambda)_n < \infty$. Consequently, $(T - \lambda)_n$ is a Browder operator. Thus by [3, Theorem 2.9] we then conclude that $T - \lambda$ is Drazin invertible, i.e. $\lambda \notin \sigma_D(T)$. Hence $\sigma_D(T) \subseteq \sigma_{SBF_+^-}(T) \cup S(T^*)$. Since the reverse inclusion obviously holds, we get $\sigma_{SBF_+^-}(T) \cup S(T^*) = \sigma_D(T)$. \square

Theorem 2.4. *If $T \in \mathcal{B}(X)$ satisfies $S(T^*) \subseteq \sigma_{SBF_+^-}(T)$, then T possesses property (gb) . In particular, if T^* has SVEP, then T possesses property (gb) .*

Proof. Suppose that $S(T^*) \subseteq \sigma_{SBF_+^-}(T)$. Then by Lemma 2.3, we get $\sigma_{SBF_+^-}(T) = \sigma_D(T)$. Consequently, by Lemma 2.2, T possesses property (gb) . If T^* has SVEP, then $S(T^*) = \emptyset$, the conclusion follows immediately. \square

The following example shows that the converse of Theorem 2.4 is not true.

Example 2.5. Let X be the Hilbert space $l_2(\mathbb{N})$ and let $T : l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be the unilateral right shift operator defined by

$$T(x_1, x_2, \dots) = (0, x_1, x_2, \dots) \quad \text{for all } (x_n) \in l_2(\mathbb{N}).$$

Then,

$$\begin{aligned} \sigma_a(T) &= \{\lambda \in \mathbb{C} : |\lambda| = 1\}, \\ \sigma_{SBF_+^-}(T) &= \{\lambda \in \mathbb{C} : |\lambda| = 1\} \end{aligned}$$

and

$$\Pi(T) = \emptyset.$$

Hence $\sigma_a(T) \setminus \sigma_{SBF_+^-}(T) = \Pi(T)$, i.e. T possesses property (gb) .

$$\text{But } S(T^*) = \{\lambda \in \mathbb{C} : 0 \leq |\lambda| < 1\} \not\subseteq \{\lambda \in \mathbb{C} : |\lambda| = 1\} = \sigma_{SBF_+^-}(T).$$

The next theorem had been established in [26, Theorem 3.10], but its proof was not so clear. Hence we give a revised proof of it.

Theorem 2.6. *If $T \in \mathcal{B}(X)$ satisfies property (gb) and N is a nilpotent operator that commutes with T , then $T + N$ satisfies property (gb) .*

Proof. Suppose that $T \in \mathcal{B}(X)$ satisfies property (gb) and N is a nilpotent operator that commutes with T . By Lemma 2.1, T satisfies generalized a-Browder's theorem and $\Pi(T) = \Pi_a(T)$. Hence $T + N$ satisfies generalized a-Browder's theorem. By [24], $\sigma(T + N) = \sigma(T)$ and $\sigma_a(T + N) = \sigma_a(T)$. Hence, by [21, Theorem 2.2] and [16, Theorem 3.2], we have that $\Pi(T + N) = \sigma(T + N) \setminus \sigma_D(T + N) = \sigma(T) \setminus \sigma_D(T) = \Pi(T) = \Pi_a(T) = \sigma_a(T) \setminus \sigma_{LD}(T) = \sigma_a(T + N) \setminus \sigma_{LD}(T + N) = \Pi_a(T + N)$. By Lemma 2.1 again, $T + N$ satisfies property (gb) . \square

The following example, which is a revised version of [26, Example 3.11], shows that the hypothesis of commutativity in Theorem 2.6 is crucial.

Example 2.7. Let $T : l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be the unilateral right shift operator defined by

$$T(x_1, x_2, \dots) = (0, x_1, x_2, \dots) \quad \text{for all } (x_n) \in l_2(\mathbb{N}).$$

Let $N : l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be a nilpotent operator with rank one defined by

$$N(x_1, x_2, \dots) = (0, -x_1, 0, \dots) \quad \text{for all } (x_n) \in l_2(\mathbb{N}).$$

Then $TN \neq NT$. Moreover,

$$\begin{aligned} \sigma(T) &= \{\lambda \in \mathbb{C} : 0 \leq |\lambda| \leq 1\}, \\ \sigma_a(T) &= \{\lambda \in \mathbb{C} : |\lambda| = 1\}, \\ \sigma(T + N) &= \{\lambda \in \mathbb{C} : 0 \leq |\lambda| \leq 1\} \end{aligned}$$

and

$$\sigma_a(T + N) = \{\lambda \in \mathbb{C} : |\lambda| = 1\} \cup \{0\}.$$

It follows that $\Pi_a(T) = \Pi(T) = \emptyset$ and $\{0\} = \Pi_a(T + N) \neq \Pi(T + N) = \emptyset$. Hence by Lemma 2.1, $T + N$ does not satisfy property (gb). But since T has SVEP, T satisfies a-Browder's theorem or equivalently, by [4, Theorem 2.2], T satisfies generalized a-Browder's theorem. Therefore by Lemma 2.1 again, T satisfies property (gb).

To continue the discussion of this note, we recall some classical definitions. Using the isomorphism $X/\mathcal{N}(T^d) \approx \mathcal{R}(T^d)$ and following [18], a topology on $\mathcal{R}(T^d)$ is defined as follows.

Definition 2.8. Let $T \in \mathcal{B}(X)$. For every $d \in \mathbb{N}$, the operator range topological on $\mathcal{R}(T^d)$ is defined by the norm $\|\cdot\|_{\mathcal{R}(T^d)}$ such that for all $y \in \mathcal{R}(T^d)$,

$$\|y\|_{\mathcal{R}(T^d)} := \inf\{\|x\| : x \in X, y = T^d x\}.$$

For a detailed discussion of operator ranges and their topologies, we refer the reader to [15] and [17].

Definition 2.9. Let $T \in \mathcal{B}(X)$ and let $d \in \mathbb{N}$. Then T has *uniform descent* for $n \geq d$ if $k_n(T) = 0$ for all $n \geq d$. If in addition $\mathcal{R}(T^n)$ is closed in the operator range topology of $\mathcal{R}(T^d)$ for all $n \geq d$, then we say that T has *eventual topological uniform descent*, and, more precisely, that T has *topological uniform descent* for $n \geq d$.

Operators with eventual topological uniform descent are introduced by Grabiner in [18]. It includes many classes of operators introduced in the Introduction of this note, such as upper semi-B-Fredholm operators, left Drazin invertible operators, Drazin invertible operators, and so on. It also includes many other classes of operators such as operators of Kato type, quasi-Fredholm operators, operators with finite descent and operators with finite essential descent, and so on. A very detailed and far-reaching account of these notations can be seen in [1, 5, 24]. Especially, operators which have topological uniform descent for $n \geq 0$ are precisely the *semi-regular* operators studied by Mbekhta in [23]. Discussions of operators with eventual topological uniform descent may be found in [6, 11, 18, 20, 28].

Lemma 2.10. If $T \in \mathcal{B}(X)$ and F is a finite rank operator commuting with T , then

- (1) $\sigma_{SBF_+^-}(T + F) = \sigma_{SBF_+^-}(T)$;
- (2) $\sigma_D(T + F) = \sigma_D(T)$.

Proof. (1) Without loss of generality, we need only to show that $0 \notin \sigma_{SBF_+^-}(T + F)$ if and only if $0 \notin \sigma_{SBF_+^-}(T)$. By symmetry, it suffices to prove that $0 \notin \sigma_{SBF_+^-}(T + F)$ if $0 \notin \sigma_{SBF_+^-}(T)$.

Suppose that $0 \notin \sigma_{SBF_+^-}(T)$. Then T is a upper semi-B-Fredholm operator and $\text{ind}(T) \leq 0$. Hence it follows from [5, Theorem 3.6] and [16, Theorem 3.2] that $T + F$ is also a upper semi-B-Fredholm operator. Thus by [18, Theorem 5.8], $\text{ind}(T + F) = \text{ind}(T) \leq 0$. Consequently, $T + F$ is a semi-B-Weyl operator, i.e. $0 \notin \sigma_{SBF_+^-}(T)$, and this completes the proof of (1).

(2) Noting that an operator is Drazin invertible if and only if it is of finite ascent and finite descent, the conclusion follows from [21, Theorem 2.2]. \square

Theorem 2.11. *If $T \in \mathcal{B}(X)$ satisfies $S(T^*) \subseteq \sigma_{SBF_+^-}(T)$ and F is a finite rank operator commuting with T , then $T + F$ satisfies property (gb).*

Proof. Since F is a finite rank operator commuting with T , by Lemma 2.10, $\sigma_{SBF_+^-}(T + F) = \sigma_{SBF_+^-}(T)$ and $\sigma_D(T + F) = \sigma_D(T)$. Since $S(T^*) \subseteq \sigma_{SBF_+^-}(T)$, by Lemma 2.3, $\sigma_{SBF_+^-}(T) = \sigma_D(T)$. Thus, $\sigma_{SBF_+^-}(T + F) = \sigma_D(T + F)$. By Lemma 2.2, $T + F$ satisfies property (gb). \square

The following example illustrates that property (gb) in general is not preserved under commuting finite rank perturbations.

Example 2.12. Let $U : l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be the unilateral right shift operator defined by

$$U(x_1, x_2, \dots) = (0, x_1, x_2, \dots) \quad \text{for all } (x_n) \in l_2(\mathbb{N}).$$

For fixed $0 < \varepsilon < 1$, let $F_\varepsilon : l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be a finite rank operator defined by

$$F_\varepsilon(x_1, x_2, \dots) = (-\varepsilon x_1, 0, 0, \dots) \quad \text{for all } (x_n) \in l_2(\mathbb{N}).$$

We consider the operators T and F defined by $T = U \oplus I$ and $F = 0 \oplus F_\varepsilon$, respectively. Then F is a finite rank operator and $TF = FT$. Moreover,

$$\sigma(T) = \sigma(U) \cup \sigma(I) = \{\lambda \in \mathbb{C} : 0 \leq |\lambda| \leq 1\},$$

$$\sigma_a(T) = \sigma_a(U) \cup \sigma_a(I) = \{\lambda \in \mathbb{C} : |\lambda| = 1\},$$

$$\sigma(T + F) = \sigma(U) \cup \sigma(I + F_\varepsilon) = \{\lambda \in \mathbb{C} : 0 \leq |\lambda| \leq 1\}$$

and

$$\sigma_a(T + F) = \sigma_a(U) \cup \sigma_a(I + F_\varepsilon) = \{\lambda \in \mathbb{C} : |\lambda| = 1\} \cup \{1 - \varepsilon\}.$$

It follows that $\Pi_a(T) = \Pi(T) = \emptyset$ and $\{1 - \varepsilon\} = \Pi_a(T + F) \neq \Pi(T + F) = \emptyset$. Hence by Lemma 2.1, $T + F$ does not satisfy property (gb). But since T has SVEP, T satisfies a-Browder's theorem or equivalently, by [4, Theorem 2.2], T satisfies generalized a-Browder's theorem. Therefore by Lemma 2.1 again, T satisfies property (gb).

Rashid gives in [26, Theorem 3.15] that if $T \in \mathcal{B}(X)$ and Q is a quasi-nilpotent operator that commute with T , then

$$\sigma_{SBF_+^-}(T + Q) = \sigma_{SBF_+^-}(T).$$

The next example show that this equality does not hold in general.

Example 2.13. Let Q denote the Volterra operator on the Banach space $C[0, 1]$ defined by

$$(Qf)(t) = \int_0^t f(s) ds \quad \text{for all } f \in C[0, 1] \text{ and } t \in [0, 1].$$

Q is injective and quasi-nilpotent. Hence it is easy to see that $\mathcal{R}(Q^n)$ is not closed for every $n \in \mathbb{N}$. Let $T = 0 \in \mathcal{B}(C[0, 1])$. It is easy to see that $TQ = 0 = QT$ and $0 \notin \sigma_{SBF_+^-}(0) = \sigma_{SBF_+^-}(T)$, but $0 \in \sigma_{SBF_+^-}(Q) = \sigma_{SBF_+^-}(0 + Q) = \sigma_{SBF_+^-}(T + Q)$. Hence $\sigma_{SBF_+^-}(T + Q) \neq \sigma_{SBF_+^-}(T)$.

Rashid claims in [26, Theorem 3.16] that property (gb) is stable under commuting quasi-nilpotent perturbations, but its proof is rely on [26, Theorem 3.15] which, by Example 2.11, is not always true. The following example shows that property (gb) in general is not preserved under commuting quasi-nilpotent perturbations.

Example 2.14. Let $U : l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be the unilateral right shift operator defined by

$$U(x_1, x_2, \dots) = (0, x_1, x_2, \dots) \quad \text{for all } (x_n) \in l_2(\mathbb{N}).$$

Let $V : l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be a quasi-nilpotent operator defined by

$$V(x_1, x_2, \dots) = (0, x_1, 0, \frac{x_3}{3}, \frac{x_4}{4} \dots) \quad \text{for all } (x_n) \in l_2(\mathbb{N}).$$

Let $N : l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be a quasi-nilpotent operator defined by

$$N(x_1, x_2, \dots) = (0, 0, 0, -\frac{x_3}{3}, -\frac{x_4}{4} \dots) \quad \text{for all } (x_n) \in l_2(\mathbb{N}).$$

It is easy to verify that $VN = NV$. We consider the operators T and Q defined by $T = U \oplus V$ and $Q = 0 \oplus N$, respectively. Then Q is quasi-nilpotent and $TQ = QT$. Moreover,

$$\sigma(T) = \sigma(U) \cup \sigma(V) = \{\lambda \in \mathbb{C} : 0 \leq |\lambda| \leq 1\},$$

$$\sigma_a(T) = \sigma_a(U) \cup \sigma_a(V) = \{\lambda \in \mathbb{C} : |\lambda| = 1\} \cup \{0\},$$

$$\sigma(T + Q) = \sigma(U) \cup \sigma(V + N) = \{\lambda \in \mathbb{C} : 0 \leq |\lambda| \leq 1\}$$

and

$$\sigma_a(T + Q) = \sigma_a(U) \cup \sigma_a(V + N) = \{\lambda \in \mathbb{C} : |\lambda| = 1\} \cup \{0\}.$$

It follows that $\Pi_a(T) = \Pi(T) = \emptyset$ and $\{0\} = \Pi_a(T + Q) \neq \Pi(T + Q) = \emptyset$. Hence by Lemma 2.1, $T + Q$ does not satisfy property (gb) . But since T has SVEP, T satisfies a-Browder's theorem or equivalently, by [4, Theorem 2.2], T satisfies generalized a-Browder's theorem. Therefore by Lemma 2.1 again, T satisfies property (gb) .

Theorem 2.15. Suppose that $T \in \mathcal{B}(X)$ obeys property (gb) and that $Q \in \mathcal{B}(X)$ is a quasi-nilpotent operator commuting with T . If T is a-polaroid, then $T + Q$ obeys (gb) .

Proof. Since T satisfies property (gb) , by Lemma 2.1, T satisfies generalized a-Browder's theorem and $\Pi(T) = \Pi_a(T)$. Hence $T + Q$ satisfies generalized a-Browder's theorem. In order to show that $T + Q$ satisfies property (gb) , by Lemma 2.1 again, it suffices to show that $\Pi(T + Q) = \Pi_a(T + Q)$. Since $\Pi(T + Q) \subseteq \Pi_a(T + Q)$ is always true, one needs only to show that $\Pi_a(T + Q) \subseteq \Pi(T + Q)$.

Let $\lambda \in \Pi_a(T + Q) = \sigma_a(T + Q) \setminus \sigma_{LD}(T + Q) = \text{iso}\sigma_a(T + Q) \setminus \sigma_{LD}(T + Q)$. Then by [24], $\lambda \in \text{iso}\sigma_a(T)$. Since T is a-polaroid, $\lambda \in \Pi_a(T) = \Pi(T)$. Thus by [28, Theorem 3.12], $\lambda \in \Pi(T + Q)$. Therefore $\Pi_a(T + Q) \subseteq \Pi(T + Q)$, and this completes the proof. \square

References

- [1] *P. Aiena*, Fredholm and local spectral Theory, with application to Multipliers, Kluwer Academic Publishers, Dordrecht 2004.
- [2] *P. Aiena*, Quasi-Fredholm operators and localized SVEP, Acta Sci. Math. (Szeged) **73** 2007, 251–263.
- [3] *P. Aiena*, *M. T. Biondi* and *C. Carpintero*, On Drazin invertibility, Proc. Amer. Math. Soc. **136** 2008, 2839–2848.
- [4] *M. Amouch* and *H. Zguitti*, On the equivalence of Browder’s and generalized Browder’s theorem, Glasgow Math. J. **48** 2006, 179–185.
- [5] *M. Berkani*, Restriction of an operator to the range of its powers, Studia Math. **140** 2000, 163–175.
- [6] *M. Berkani*, *N. Castro*, and *S. Djordjević*, Single valued extension property and generalized Weyl’s theorem, Math. Bohem. **131**(1) 2006, 29–38.
- [7] *M. Berkani* and *J.J. Koliha*, Weyl type theorems for bounded linear operators, Acta Sci. Math. (Szeged) **69** 2003, 359–376.
- [8] *M. Berkani* and *M. Sarik*, On semi B-Fredholm operators, Glasgow Math. J. **43** 2001, 457–465.
- [9] *M. Berkani* and *H. Zariouh*, Extended Weyl type theorems, Math. Bohem. **134**(4) 2009, 369–378.
- [10] *M. Berkani* and *H. Zariouh*, New extended Weyl type theorems, Mat. Vesnik. **62**(2) 2010, 145–154.
- [11] *X.H. Cao*, Topological uniform descent and Weyl type theorem, Linear Algebra Appl. **420** 2007, 175–182.
- [12] *S.V. Djordjević* and *Y.M. Han*, Browder’s theorems and spectral continuity, Glasgow Math. J. **42** 2000, 479–486.
- [13] *N. Dunford*, Spectral theory II. Resolutions of the identity, Pacific J. Math. **2** 1952, 559–614.
- [14] *N. Dunford*, Spectral operators, Pacific J. Math. **4** 1954, 321–354.
- [15] *P.A. Fillmore* and *J.P. Williams*, On operator ranges. Advances in Math. **7**(3) 1971, 254–281.
- [16] *O. Bel Hadj Fredj*, *M. Burgos*, and *M. Oudghiri*, Ascent spectrum and essential ascent spectrum, Studia Math. **187** 2008, 59–73.
- [17] *S. Grabiner*, Ranges of products of operators. Canad. J. Math. **XXVI**(6) 1974, 1430–1441.

- [18] *S. Grabiner*, Uniform ascent and descent of bounded operators, J. Math. Soc. Japan **34** 1982, 317–337.
- [19] *R.E. Harte* and *W.Y. Lee*, Another note on Weyl’s theorem, Trans. Amer. Math. Soc. **349** 1997, 2115–2124.
- [20] *Q.F. Jiang*, *H.J. Zhong* and *Q.P. Zeng*, Topological uniform descent and localized SVEP, J. Math. Anal. Appl. **390** 2012, 355–361.
- [21] *M.A. Kaashoek* and *D.C. Lay*, Ascent, descent, and commuting perturbations, Trans. Amer. Math. Soc. **169** 1972, 35–47.
- [22] *K.B. Laursen* and *M.M. Neumann*, An Introduction to Local Spectral Theory, Oxford University Press, 2000.
- [23] *M. Mbekhta*, Résolvant généralisé et théorie spectrale, J. Operator Theory. **21** 1989, 69–105.
- [24] *M. Mbekhta* and *V. Müller*, On the axiomatic theory of spectrum II, Studia Math. **119** 1996, 129–147.
- [25] *V. Rakocëvić*, Semi-Browder operators and perturbations, Studia Math. **122** 1997, 131–137.
- [26] *M.H.M. Rashid*, Property (gb) and perturbations, J. Math. Anal. Appl. **383** 2011, 82–94.
- [27] *H.-O. Tylli*, On the asymptotic behaviour of some quantities related to semi-Fredholm operators, J. London Math. Soc. **31** 1985, 340–348.
- [28] *Q.P. Zeng*, *H.J. Zhong* and *Z.Y. Wu*, Small essential spectral radius perturbations of operators with topological uniform descent, Bull. Aust. Math. Soc. **85** 2012, 26–45.