

# Two remarks on $C^\infty$ Anosov diffeomorphisms

Shigenori Matsumoto

ABSTRACT. Let  $M$  be a closed oriented  $C^\infty$  manifold and  $f$  a  $C^\infty$  Anosov diffeomorphism on  $M$ . We show that if  $M$  is the two torus  $T^2$ , then  $f$  is conjugate to a hyperbolic automorphism of  $T^2$ , either by a  $C^\infty$  diffeomorphism or by a singular homeomorphism. We also show that for general  $M$ , if  $f$  admits an absolutely continuous invariant measure  $\mu$ , then  $\mu$  is a  $C^\infty$  volume. The proofs are concatenations of well known results in the field.

## 1. Conjugacy

Let  $f$  be a  $C^\infty$  Anosov diffeomorphism on the two torus  $T^2$ . Then

$$A = f_* \in \text{Aut}(H_1(T^2, \mathbb{Z})) = \text{SL}(2, \mathbb{Z})$$

defines a hyperbolic automorphism of the abelian Lie group  $T^2$ , and  $f$  is isotopic to  $A$ . It is known [F, M] that  $f$  is conjugate to  $A$  by a homeomorphism  $h$  which is isotopic to the identity:  $h \circ A = f \circ h$ . It is well known that the conjugacy  $h$  is a bi-Hölder homeomorphism. Also it is easy to show that  $h$  is unique. Let us denote by  $m$  the normalized Haar measure of  $T^2$ . A homeomorphism  $h$  of  $T^2$  is said to be *singular* if there is an  $m$ -conull Borel set  $E$  such that  $h(E)$  is  $m$ -null. Our first result is the following.

**THEOREM 1.** *The conjugacy  $h$  is either a  $C^\infty$  diffeomorphism or a singular homeomorphism.*

**PROOF.** Let  $TT^2 = E^u \oplus E^s$  be the hyperbolic splitting associated with  $f$ . By a dimensional reason, it is a  $C^1$  splitting [MM1]. Fix a translation invariant  $C^\infty$  Riemannian metric  $g$  on  $T^2$ . The derivative of  $f$  along  $E^u$  (resp.  $E^s$ ) measured with respect to  $g$  is denoted by  $J^s f$  (resp.  $J^u f$ ). These are  $C^1$  functions. The Gibbs measure [S, B] for the potential  $-\log|J^u f|$  (resp.  $\log|J^s f|$ ) is denoted by  $\mu_+$  (resp.  $\mu_-$ ).

---

1991 *Mathematics Subject Classification.* 37D20.

*Key words and phrases.* Anosov diffeomorphism, topological conjugacy, Gibbs measure, absolutely continuous invariant measure.

The author is partially supported by Grant-in-Aid for Scientific Research (C) No. 20540096.

Let  $f$ ,  $A$  and  $h$  be as above. First consider the case where  $f$  does not admit an a. c. i. m. (absolutely continuous invariant measure). Then the  $f$ -invariant measure  $h_*m$  is singular to  $m$ .

To show this, notice that  $h_*m$  is decomposed into two parts; one absolutely continuous and the other singular. Since  $h$  is a  $C^\infty$  diffeomorphism, it leaves each part invariant. But the absolutely continuous part must be zero since by the assumption there is no a. c. i. m. for  $f$ .

Thus  $h$  maps the measure  $m$  to a singular measure  $h_*m$ . Since  $h_*m$  is singular, there is an  $m$ -null set  $E'$  such that  $h_*m(E') = m(h^{-1}E') = 1$ . The conjugacy  $h$  maps the  $m$ -conull set  $h^{-1}(E')$  to the  $m$ -null set  $E'$ , showing that  $h$  is a singular homeomorphism.

Next consider the case where  $f$  admits an a. c. i. m.  $\mu$ . Then we have  $\mu = \mu_+ = \mu_-$  (Proof of Corollary 1 of [S], Corollary 4.13 of [B]). In particular an a. c. i. m.  $\mu$  is unique and ergodic. The induced measure  $h_*m$  is also ergodic. Therefore either  $\mu$  and  $h_*m$  are mutually singular or coincide. In the former case, we argue just as before, to conclude that the conjugacy  $h$  is a singular homeomorphism.

Finally assume that  $\mu_+ = \mu_- = h_*m$ . These are the Gibbs measures of three potentials,  $-\log|J^u f|$ ,  $\log|J^s f|$  and a constant. By Section 3.4 of [S], these three functions, with the identical Gibbs measure, are mutually cohomologous modulo constant. That is, there are continuous functions  $v_1$ ,  $v_2$  and constants  $c_1$ ,  $c_2$  such that

$$\begin{aligned} -\log|J^u f| &= v_1 \circ f - v_1 + c_1, \\ \log|J^s f| &= v_2 \circ f - v_2 + c_2. \end{aligned}$$

This shows that the Lyapunov exponents of all periodic orbits are the same. By Theorem 1 of [MM2], the conjugacy  $h$  is a  $C^\infty$  diffeomorphism. The proof of Theorem 1 is complete.  $\square$

## 2. Absolutely continuous invariant measure

Let  $M$  be a closed oriented  $n$ -dimensional  $C^\infty$  manifold and  $f$  a  $C^\infty$  Anosov diffeomorphism on  $M$ . Let  $g$  be a  $C^\infty$  Riemannian metric on  $M$ , and  $m$  the normalized measure given by the volume form associated with  $g$ .

**THEOREM 2.** *Assume  $f$  admits an a. c. i. m.  $\mu$  with density  $\varphi$ :  $\mu = \varphi m$ ,  $\varphi \in L^1(m)$ . Then the density  $\varphi$  is a positive  $C^\infty$  function.*

**PROOF.** Let  $TM = E^u \oplus E^s$  be the hyperbolic splitting associated with  $f$ . Denote the Jacobian along  $E^u$  (resp.  $E^s$ ) measured with respect to  $g$  by  $J^u f$  (resp.  $J^s f$ ). The total Jacobian measured with respect to  $g$  is denoted by  $Jf$ . All these are continuous real valued functions on  $M$ .

Define another continuous Riemannian metric  $g'$  by  $g' = g|_{E^u} \oplus g|_{E^s}$ . Thus  $E^u$  and  $E^s$  are perpendicular with respect to  $g'$ . Let  $m'$  be the normalized measure given by the volume form associated with  $g'$ . We have  $m' = e^a m$  for a continuous function  $a$ .

Denote by  $J'f$  the total Jacobian with respect to  $g'$ . Then we have

$$(2.1) \quad \log|J'f| = \log|J^u f| + \log|J^s f|.$$

By [S, B], we have  $\mu = \mu_+ = \mu_-$ , where  $\mu_+$  (resp.  $\mu_-$ ) is the Gibbs measure for the potential  $-\log|J^u f|$  (resp.  $\log|J^s f|$ ). Then by [S],  $\log|J^u f| + \log|J^s f|$  is cohomologous to a constant. Thus by (2.1), we have

$$(2.2) \quad \log|J'f| = b \circ f - b + C.$$

for a continuous function  $b$  and a constant  $C$ .

On the other hand, by the invariance of the a. c. i. m.  $\mu = \varphi e^{-a} m'$ , we have  $\mu$ -almost everywhere

$$(2.3) \quad \log|J'f| = (a - \log \varphi) \circ f - (a - \log \varphi)$$

Now by (2.3), we have  $\mu(\log|J'f|) = 0$ . This implies that  $C = 0$  in (2.2). Then (2.2) implies the invariance of the measure  $e^{-b} m' = e^{-b+a} m$ . Moreover, adding an appropriate constant to  $b$ , we may assume that  $e^{-b+a} m$  is a probability measure. By the uniqueness of the a. c. i. m., we have  $\mu = e^{-b+a} m$ . That is, the density of  $\mu$  is positive and continuous. Then by Corollary 2.1 of [LMM], we obtain that  $e^{-b+a}$  is a  $C^\infty$  function. The proof is complete.  $\square$

## References

- [B] R. Bowen, *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*, Springer Lect. Notes Series in Math. Vol. 470, 2nd Edition Ed. J.-R. Chazottes, 2008, Springer-Verlag, Berlin Heidelberg.
- [F] J. Franks, *Anosov diffeomorphisms on tori*, Trans. A. M. S. **145**(1969), 117–124.
- [LMM] R. de la Llave, J. M. Marco and R. Moriyo, *Canonical perturbation theory of Anosov systems and regularity results for the Livsic cohomology equation*, Ann. Math. **123**(1986), 537–611.
- [M] A. Manning, *There are no new Anosov diffeomorphisms on tori*, Amer. J. Math. **96** No. 3(1974) 422–429.
- [MM1] J. M. Marco and R. Moriyo, *Invariants for smooth conjugacy of hyperbolic dynamical Systems, I*, Commun. Math. Phys. /bf 109(1987), 681–689.
- [MM2] J. M. Marco and R. Moriyo, *Invariants for smooth conjugacy of hyperbolic dynamical Systems, III*, Commun. Math. Phys. /bf 112(1987), 317–333.
- [S] Ya. Sinai, *Gibbs measures in ergodic theory*, Russ. Math. Surveys **27** No. 4(1972) 21–69.

DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE AND TECHNOLOGY, NIHON UNIVERSITY, 1-8-14 KANDA, SURUGADAI, CHIYODA-KU, TOKYO, 101-8308 JAPAN  
*E-mail address:* matsumo@math.cst.nihon-u.ac.jp