A simple proof of the Fundamental Theorem of Calculus for the Lebesgue integral

March, 2012

Rodrigo López Pouso Departamento de Análise Matemática Facultade de Matemáticas, Universidade de Santiago de Compostela, Campus Sur 15782 Santiago de Compostela, Spain.

Abstract

This paper contains a new elementary proof of the Fundamental Theorem of Calculus for the Lebesgue integral. The hardest part of our proof simply concerns the convergence in L^1 of a certain sequence of step functions, and we prove it using only basic elements from Lebesgue integration theory.

1 Introduction

Let $f : [a, b] \longrightarrow \mathbb{R}$ be absolutely continuous on [a, b], i.e., for every $\varepsilon > 0$ there exists $\delta > 0$ such that if $\{(a_j, b_j)\}_{j=1}^n$ is a family of pairwise disjoint subintervals of [a, b] satisfying

$$\sum_{j=1}^{n} \left(b_j - a_j \right) < \delta$$

then

$$\sum_{j=1}^{n} |f(b_j) - f(a_j)| < \varepsilon.$$

Classical results ensure that f has a finite derivative almost everywhere in I = [a, b], and that $f' \in L^1(I)$, see [3] or [8, Corollary 6.83]. These results, which we shall use in this paper, are the first steps in the proof of the main connection between absolute continuity and Lebesgue integration: the Fundamental Theorem of Calculus for the Lebesgue integral.

Theorem 1.1 If $f: I = [a, b] \longrightarrow \mathbb{R}$ is absolutely continuous on I then

$$f(b) - f(a) = \int_{a}^{b} f'(x) dx$$
 in Lebesgue's sense.

In this note we present a new elementary proof to Theorem 1.1 which seems more natural and easy than the existing ones. Indeed, our proof can be sketched simply as follows:

1. We consider a well-known sequence of step functions $\{h_n\}_{n\in\mathbb{N}}$ which tends to f' almost everywhere in I and, moreover,

$$\int_{a}^{b} h_{n}(x) \, dx = f(b) - f(a) \quad \text{for all } n \in \mathbb{N}.$$

2. We prove, by means of elementary arguments, that

$$\lim_{n \to \infty} \int_a^b h_n(x) \, dx = \int_a^b f'(x) \, dx.$$

More precise comparison with the literature on Theorem 1.1 and its several proofs will be given in Section 3.

In the sequel m stands for the Lebesgue measure in \mathbb{R} .

$\mathbf{2}$ Proof of Theorem 1.1

For each $n \in \mathbb{N}$ we consider the partition of the interval I = [a, b] which divides it into 2^n subintervals of length $(b-a)2^{-n}$, namely

$$x_{n,0} < x_{n,1} < x_{n,2} < \dots < x_{n,2^n},$$

where $x_{n,i} = a + i(b-a)2^{-n}$ for $i = 0, 1, 2, ..., 2^n$. Now we construct a step function $h_n : [a, b) \longrightarrow \mathbb{R}$ as follows: for each $x \in [a, b)$ there is a unique $i \in \{0, 1, 2, \dots, 2^n - 1\}$ such that

$$x \in [x_{n,i}, x_{n,i+1})$$

and we define

$$h_n(x) = \frac{f(x_{n,i+1}) - f(x_{n,i})}{x_{n,i+1} - x_{n,i}} = \frac{2^n}{b-a} [f(x_{n,i+1}) - f(x_{n,i})].$$

On the one hand, the construction of $\{h_n\}_{n\in\mathbb{N}}$ implies that

$$\lim_{n \to \infty} h_n(x) = f'(x) \quad \text{for all } x \in [a, b) \setminus N,$$
(2.1)

where $N \subset I$ is a null-measure set such that f'(x) exists for all $x \in I \setminus N$. On the other hand, for each $n \in \mathbb{N}$ we compute

$$\int_{a}^{b} h_{n}(x)dx = \sum_{i=0}^{2^{n}-1} \int_{x_{n,i}}^{x_{n,i+1}} h_{n}(x)dx = \sum_{i=0}^{2^{n}-1} [f(x_{n,i+1}) - f(x_{n,i})] = f(b) - f(a)$$

and therefore it only remains to prove that

$$\lim_{n \to \infty} \int_a^b h_n(x) \, dx = \int_a^b f'(x) \, dx.$$

Let us prove that, in fact, we have convergence in $L^{1}(I)$, i.e.,

$$\lim_{n \to \infty} \int_{a}^{b} |h_{n}(x) - f'(x)| \, dx = 0.$$
(2.2)

Let $\varepsilon > 0$ be fixed and let $\delta > 0$ be one of the values corresponding to $\varepsilon/4$ in the definition of absolute continuity of f.

Since $f' \in L^1(I)$ we can find $\rho > 0$ such that for any measurable set $E \subset I$ we have

$$\int_{E} |f'(x)| \, dx < \frac{\varepsilon}{4} \quad \text{whenever } m(E) < \rho. \tag{2.3}$$

The following lemma will give us fine estimates for the integrals when $|h_n|$ is "small". We postpone its proof for better readability.

Lemma 2.1 For each $\varepsilon > 0$ there exist $k, n_k \in \mathbb{N}$ such that

$$k \cdot m\left(\left\{x \in I : \sup_{n \ge n_k} |h_n(x)| > k\right\}\right) < \varepsilon.$$

Lemma 2.1 guarantees that there exist $k, n_k \in \mathbb{N}$ such that

$$k \cdot m\left(\left\{x \in I : \sup_{n \ge n_k} |h_n(x)| > k\right\}\right) < \min\left\{\delta, \frac{\varepsilon}{4}, \rho\right\}.$$
 (2.4)

Let us denote

$$A = \left\{ x \in I : \sup_{n \ge n_k} |h_n(x)| > k \right\},$$

which, by virtue of (2.4) and (2.3), satisfies the following properties:

$$m(A) < \delta, \tag{2.5}$$

$$k \cdot m(A) < \frac{\varepsilon}{4},\tag{2.6}$$

$$\int_{A} |f'(x)| \, dx < \frac{\varepsilon}{4}. \tag{2.7}$$

We are now in a position to prove that the integrals in (2.2) are smaller than ε for all sufficiently large values of $n \in \mathbb{N}$. We start by noticing that (2.7) guarantees that for all $n \in \mathbb{N}$ we have

$$\int_{I} |h_{n}(x) - f'(x)| \, dx = \int_{I \setminus A} |h_{n}(x) - f'(x)| \, dx + \int_{A} |h_{n}(x) - f'(x)| \, dx$$
$$< \int_{I \setminus A} |h_{n}(x) - f'(x)| \, dx + \int_{A} |h_{n}(x)| \, dx + \frac{\varepsilon}{4}. \quad (2.8)$$

The definition of the set A implies that for all $n \in \mathbb{N}$, $n \ge n_k$, we have

$$|h_n(x) - f'(x)| \le k + |f'(x)|$$
 for almost all $x \in I \setminus A$,

so the Dominated Convergence Theorem yields

$$\lim_{n \to \infty} \int_{I \setminus A} |h_n(x) - f'(x)| \, dx = 0.$$
(2.9)

From (2.8) and (2.9) we deduce that there exists $n_{\varepsilon} \in \mathbb{N}$, $n_{\varepsilon} \ge n_k$, such that for all $n \in \mathbb{N}$, $n \ge n_{\varepsilon}$, we have

$$\int_{I} |h_{n}(x) - f'(x)| \, dx < \frac{\varepsilon}{2} + \int_{A} |h_{n}(x)| \, dx.$$
(2.10)

Finally, we estimate $\int_A |h_n|$ for each fixed $n \in \mathbb{N}$, $n \ge n_{\varepsilon}$. First, we decompose $A = B \cup C$, where

$$B = \{x \in A : |h_n(x)| \le k\}$$
 and $C = A \setminus B$.

We immediately have

$$\int_{B} |h_n(x)| \, dx \le k \cdot m(B) \le k \cdot m(A) < \frac{\varepsilon}{4} \quad \text{by (2.6)}.$$

Obviously, $\int_C |h_n| < \varepsilon/4$ when $C = \emptyset$. Let us see that this inequality holds true when $C \neq \emptyset$. For every $x \in C = \{x \in A : |h_n(x)| > k\}$ there is a unique index $i \in \{0, 1, 2, \ldots, 2^n - 1\}$ such that $x \in [x_{n,i}, x_{n,i+1})$. Since $|h_n|$ is constant on $[x_{n,i}, x_{n,i+1})$ we deduce that $[x_{n,i}, x_{n,i+1}) \subset C$. Thus there exist indexes $i_l \in \{0, 1, 2, \ldots, 2^n - 1\}$, with $l = 1, 2, \ldots, p$ and $i_l \neq i_{\tilde{l}}$ if $l \neq \tilde{l}$, such that

$$C = \bigcup_{l=1}^{p} [x_{n,i_l}, x_{n,i_l+1}).$$

Therefore

$$\sum_{l=1}^{p} (x_{n,i_l+1} - x_{n,i_l}) = m(C) \le m(A) < \delta \quad \text{by (2.5)},$$

and then the absolute continuity of f finally comes into action:

$$\int_{C} |h_{n}(x)| \, dx = \sum_{l=1}^{p} \int_{x_{n,i_{l}}}^{x_{n,i_{l}+1}} |h_{n}(x)| \, dx$$
$$= \sum_{l=1}^{p} |f(x_{n,i_{l}+1}) - f(x_{n,i_{l}})| < \frac{\varepsilon}{4}.$$

This inequality, along with (2.10) and (2.11), guarantee that for all $n \in \mathbb{N}$, $n \geq n_{\varepsilon}$, we have

$$\int_{I} |h_n(x) - f'(x)| \, dx < \varepsilon,$$

thus proving (2.2) because ε was arbitrary.

Proof of Lemma 2.1. Let $\varepsilon > 0$ be fixed and let $\rho > 0$ be such that for every measurable set $E \subset I$ with $m(E) < \rho$ we have

$$\int_E |f'(x)| \, dx < \frac{\varepsilon}{2}.$$

Let $N \subset I$ be as in (2.1) and let $k \in \mathbb{N}$ be sufficiently large so that

$$m\left(\left\{x \in I \setminus N : |f'(x)| \ge k\right\}\right) < \rho,$$

which implies that

$$k \cdot m\left(\left\{x \in I \setminus N : |f'(x)| \ge k\right\}\right) \le \int_{\{x \in I \setminus N : |f'(x)| \ge k\}} |f'(x)| \, dx < \frac{\varepsilon}{2}.$$
(2.12)

Let us define

$$E_j = \left\{ x \in I \setminus N : \sup_{n \ge j} |h_n(x)| > k \right\} \quad (j \in \mathbb{N}).$$

Notice that $E_{j+1} \subset E_j$ for every $j \in \mathbb{N}$, and $m(E_1) < \infty$, hence

$$\lim_{j \to \infty} m(E_j) = m\left(\bigcap_{j=1}^{\infty} E_j\right).$$
(2.13)

Clearly, $\bigcap_{j=1}^{\infty} E_j \subset \{x \in I \setminus N : |f'(x)| \ge k\}$, so we deduce from (2.13) that we can find some $n_k \in \mathbb{N}$ such that

$$m(E_{n_k}) \le m\left(\left\{x \in I \setminus N : |f'(x)| \ge k\right\}\right) + \frac{\varepsilon}{2k},$$

and then (2.12) yields $k \cdot m(E_{n_k}) < \varepsilon$.

3 Final remarks

The sequence $\{h_n\}_{n\in\mathbb{N}}$ is used in other proofs of Theorem 1.1, see [1] or [10]. The novelty in this paper is our elementary and self-contained proof of (2.2). Incidentally, a revision of the proof of our Lemma 2.1 shows that it holds true for any sequence of measurable functions $\tilde{h}_n : E \subset \mathbb{R} \longrightarrow \mathbb{R}$ which converges pointwise almost everywhere to some $h \in L^1(E)$ and $m(E) < \infty$.

Our proof avoids somewhat technical results often invoked to prove Theorem 1.1. For instance, we do not use any sophisticated estimate for the measure of image sets such as [4, Theorem 7.20], [8, Lemma 6.88] or [10, Proposition 1.2], see also [6]. We do not use the following standard lemma either: an absolutely continuous function having zero derivative almost everywhere is constant, see [4, Theorem 7.16] or [8, Lemma 6.89]. It is worth having a look at [5] for a proof of that lemma using tagged partitions; see also [2] for a proof based on full covers [9]. Concise proofs of Theorem 1.1 follow from the Radon–Nikodym Theorem, see [1], [4] or [7], but this is far from being elementary. Finally, it is interesting to note that (2.2) easily follows from the Dominated Convergence Theorem when f is Lipschitz continuous on I. This fact made the author think about the following project for students in an introductory course to Lebesgue integration.

PROJECT: TWO IMPORTANT RESULTS FOR THE PRICE OF ONE.

1. Let $f: I = [a, b] \longrightarrow \mathbb{R}$ be Lispchitz continuous on I. A deep result (worth to know without proof) guarantees that f'(x) exists for almost all $x \in I$, see [4, Theorem 7.8].

Consider the sequence $\{h_n\}_{n \in \mathbb{N}}$ as defined in Section 2 and prove

- (a) $\{h_n(x)\}_{n \in \mathbb{N}}$ tends to f'(x) for almost all $x \in I$;
- (b) $\int_{I} h_n(x) dx = f(b) f(a)$ for all $n \in \mathbb{N}$;
- (c) (Use the Dominated Convergence Theorem) $f' \in L^1(I)$ and

$$f(b) - f(a) = \int_{a}^{b} f'(x) dx$$
 in Lebesgue's sense.

2. Let $g: I = [a, b] \longrightarrow \mathbb{R}$ be Riemann-integrable on I and define

$$f(x) = (R) \int_a^x g(s) \, ds \quad (x \in I),$$

where $(R) \int$ stands for the Riemann integral.

Use the information in Exercise 1 to deduce that $g \in L^1(I)$ and

(R)
$$\int_{a}^{b} g(x) dx = \int_{a}^{b} g(x) dx$$
 in Lebesgue's sense.

References

- D. Bárcenas, The Fundamental Theorem of Calculus for Lebesgue Integral, Divulg. Mat. 8 (2000), 75–85.
- [2] M. W. Botsko, The use of full covers in real analysis, Amer. Math. Monthly 96 (1989), 328-333.
- [3] M. W. Botsko, An elementary proof of Lebesgue's Differentiation Theorem, Amer. Math. Monthly 110 (2003), 834–838.

- [4] A. M. Bruckner, J. B. Bruckner and B. S. Thomson, *Real Analysis*, 2nd Edition, 2008, http://classicalrealanalysis.info.
- [5] R. A. Gordon, The use of tagged partitions in elementary real analysis, Amer. Math. Monthly 105 (1998), 107–117.
- [6] J. J. Koliha, Mean, Meaner and the Meanest Mean Value Theorem, Amer. Math. Monthly 116 (2009), 356–361.
- [7] W. Rudin, Real and Complex Analysis, 3rd Edition, McGraw-Hill, 1987.
- [8] K. R. Stromberg, An Introduction to Classical Real Analysis, Wadsworth Inc., Belmont, California, 1981.
- [9] B. S. Thomson, On full covering properties, Real Anal. Exchange 6 (1980/1981), 77–93.
- [10] C. Volintiru, A proof of the fundamental theorem of calculus using Hausdorff measures, *Real Anal. Exchange* **26** (2000/01), 381–389.