

Domain dependence of eigenvalues of elliptic type operators

Vladimir Kozlov (Linköping)*

February 19, 2022

*Department of Mathematics, Linköping University, S-581 83 Linköping,
Sweden*

E-mail address: vlkoz@mai.liu.se

Abstract. The dependence on the domain is studied for the Dirichlet eigenvalues of an elliptic operator considered in bounded domains. Their proximity is measured by a norm of the difference of two orthogonal projectors corresponding to the reference domain and the perturbed one; this allows to compare domains that have non-smooth boundaries and different topology. The main result is an asymptotic formula in which the remainder is evaluated in terms of this quantity. As an application, the stability of eigenvalues is estimated by virtue of integrals of squares of the gradients of eigenfunctions for elliptic problems in different domains. It occurs that these stability estimates imply well-known inequalities for perturbed eigenvalues.

1 Introduction

We consider eigenvalues of the Dirichlet problem for an elliptic operator in a domain in \mathbb{R}^n , $n \geq 2$. Our main aim is to study how these eigenvalues depend on the domain. The first results concerning this classical problem can be found presumably in the book [16] by Rayleigh. A general technique was proposed by Hadamard [8], [9], who studied perturbations of a domain with a smooth boundary. In his works, the boundary of the perturbed domain

*The author was supported by the Swedish Research Council (VR)

Ω_ε is described by the function $\tau = \varepsilon h(x')$, where τ is the variable along the normal to the boundary, h is a smooth function on the boundary and ε is a small parameter. Hadamard's formula for the perturbed first eigenvalue $\lambda(\Omega_\varepsilon)$ of the Dirichlet Laplacian is as follows:

$$\lambda(\Omega_\varepsilon) = \lambda(\Omega_0) + \varepsilon \int_{\partial\Omega_0} |\partial_\tau \varphi|^2 h \, dS + o(\varepsilon), \quad (1)$$

where φ is the first eigenfunction such that $\|\varphi\|_{L^2} = 1$ and dS is the surface measure on $\partial\Omega_0$. In various generalizations of this formula, perturbations are described by a family of smooth mappings, other boundary conditions are considered as well as more general elliptic operators; see [4], [1], [3], [15], [10] and references cited therein. Some non-smooth perturbations of smooth boundaries, that are described by normal shift functions, were treated in [5]–[7], [17]. On the other hand, there are many problems involving more general classes of perturbations, namely, non-smooth perturbations of non-smooth boundaries and perturbations that cannot be described by a family of isomorphisms.

In [12]–[14], another approach was proposed to studying the dependence of λ on the domain and the operator's coefficients. It is based on an abstract theorem concerning perturbation of eigenvalues for operators acting in different spaces. Here we further develop the approach used in [12]. In this paper, the main novelty is the application of a new small parameter to measure the proximity of two spectral problems. This parameter is a norm of the difference of orthogonal projectors on Sobolev spaces consisting of functions given on these domains. In Sect. 2.1, we show that the convergence with respect to this parameter (it is actually a distance) is equivalent to the convergence in the sense of Mosco or to γ -convergence, see [10] and [1]. The latter type of convergence plays an important role in proving the existence of solutions to various shape optimization problems dealing with eigenvalues. Another new point is a refined estimate of the remainder term. It allows us to extend essentially the class of domains and their perturbations for which a certain "generalized" asymptotic formula is still valid. Let us turn to a detailed description of the results. For this purpose we shall use an example of second order elliptic operator.

Let Ω_1 and Ω_2 be two bounded domains with nonempty intersection. We

consider a bilinear form

$$(u, v) = \sum_{i,j=1}^n \int_{\mathbb{R}^n} A_{ij}(x) \partial_{x_j} u \partial_{x_i} \bar{v} dx, \quad (2)$$

where A_{ij} are bounded measurable real-valued functions such that $A_{ij} = A_{ji}$ and

$$\nu |\xi|^2 \leq \sum_{i,j=1}^n A_{ij}(x) \xi_i \xi_j \leq \nu^{-1} |\xi|^2 \quad (3)$$

for all $\xi \in \mathbb{R}^n \setminus \mathcal{O}$ and $x \in \mathbb{R}^n$. The form (\cdot, \cdot) defines a new inner product in the space $H_k = \dot{W}^{1,2}(\Omega_k)$, $k = 1, 2$ (we suppose that functions belonging to both of these spaces are extended by zero to the whole \mathbb{R}^n); the corresponding norm will be denoted by $\|\cdot\|$. Let us consider the following spectral problems

$$(\varphi, v) = \lambda \langle \varphi, v \rangle \quad \text{for all } v \in H_1 \quad (4)$$

and

$$(U, V) = \mu \langle U, V \rangle \quad \text{for all } V \in H_2, \quad (5)$$

where $\langle \cdot, \cdot \rangle$ is the inner product in L^2 , $\varphi \in H_1$ and $U \in H_2$. By λ_m we denote the m th eigenvalue of problem (4); let X_m be the corresponding eigenspace, $J_m = \dim X_m$. Our aim is to describe eigenvalues of problem (5) located near the eigenvalue λ_m of (4).

If Ω_1 and Ω_2 are sub-domains of a bounded domain D , then by S_j , $j = 1, 2$, we denote the orthogonal projector defined on $\dot{W}^{1,2}(D)$ whose image belongs to H_j . For characterizing the proximity of Ω_1 and Ω_2 we use the best constant $\sigma = \sigma(H_1, H_2)$ in the inequality

$$\| (S_1 - S_2)u \|^2 \leq \sigma \|u\|^2, \quad u \in \dot{W}^{1,2}(D), \quad (6)$$

where $\|\cdot\|$ is the L^2 -norm. In Sect. 2.1, we show that the γ -convergence of domains is equivalent to their convergence in terms of the distance (6).

If σ is sufficiently small, then problem (5) has exactly J_m eigenvalues, say μ_1, \dots, μ_{J_m} , in a neighborhood of λ_m ; see Proposition 3. In order to formulate one of our main results we introduce the following notation: $T_2 u = u - S_2 u$, $\Psi = \Psi_\varphi \in H_2$ is the solution of the equation

$$(\Psi, w) = (\varphi, w) - \lambda_m \langle \varphi, w \rangle \quad \text{for all } w \in H_2 \quad (7)$$

and

$$\rho = \max_{\varphi \in X_m, \|\varphi\|=1} (\|T\varphi\|^2 + \|\Psi_\varphi\|^2 + \sigma \|\Psi_\varphi\|^2). \quad (8)$$

Now we are in a position to formulate the following.

Theorem 1. *The asymptotic formula holds:*

$$\mu_k^{-1} = \lambda_m^{-1} + \tau_k + O(\rho + |\tau_k|\sigma), \quad k = 1, \dots, J_m. \quad (9)$$

Here $\tau = \tau_k$ is an eigenvalue of the problem

$$\frac{1}{\lambda_m} \left((\Psi_\varphi, \Psi_\psi) - (T\varphi, T\psi) - (\Psi_\varphi, \psi) - (\varphi, \Psi_\psi) \right) = \tau(S\varphi, S\psi) \quad \text{for all } \psi \in X_m, \quad (10)$$

where $\varphi \in X_m$; moreover, τ_1, \dots, τ_m in (9) run through all eigenvalues of (10) counted according to their multiplicity.

In the case when $\Omega_2 \subset \Omega_1$, the function Ψ_φ vanishes and the eigenvalue problem (10) takes the form:

$$-\frac{1}{\lambda_m} (T\varphi, T\psi) = \tau(S\varphi, S\psi) \quad \text{for all } \psi \in X_m. \quad (11)$$

If $\Omega_1 \subset \Omega_2$, then $T\varphi = 0$ and (10) can be written as follows:

$$\frac{1}{\lambda_m} (\Psi_\varphi, \Psi_\psi) = \tau(S\varphi, S\psi) \quad \text{for all } \psi \in X_m. \quad (12)$$

Similar theorems were proved in [12]¹ and [14], but here, the main novelty is the use of the small parameter σ which makes the present theorem applicable to a substantially larger class of perturbations. The importance of this parameter, that serves as a distance in the set of all closed subspaces of $\mathring{W}^{1,2}(D)$, lies in the fact that this set is compact with respect to this distance (see Proposition 2). As a result one obtains solutions of various optimization problems for functionals that are continuous with respect to this distance. Another new point is that the remainder in (10) has the form, in which the small parameter σ appears explicitly together with the quantities involved in the finite dimensional spectral problem (10).

¹Two terms are lost in formula (5), [12], compare with (9) and (10). However, if $\Omega_2 \subset \Omega_1$ or $\Omega_1 \subset \Omega_2$, then formula (5), [12], is true.

Let us describe some applications of the asymptotic formula (10). We begin with the case when both Ω_1 and Ω_2 are Lipschitz domains with the Lipschitz constant less than or equal to C_* or, what is the same, uniformly Lipschitz (see [11] and [10], where the definitions of these notions are given). We assume that

$$\Omega_\varepsilon = \{x \in \Omega_1, \text{dist}(x, \partial\Omega_1) > \varepsilon\} \subset \Omega_2 \subset \Omega_\varepsilon^+ = \{x \in \mathbb{R}^n, \text{dist}(x, \Omega_1) < \varepsilon\}, \quad (13)$$

where ε is a small positive parameter. It is supposed that the constant C_* is independent of ε . Then there exists a set S , subject to the conditions

$$\Omega_1 \setminus \Omega_{2\varepsilon} \subset S \subset \Omega_1 \quad \text{and} \quad |S| \leq C_1 |\Omega_2 \setminus \Omega_1|,$$

and such that the following inequality holds:

$$|\mu_k^{-1} - \lambda_m^{-1}| \leq C_1 \max_{\varphi \in X_m, \|\varphi\|=1} \int_{(\Omega_1 \setminus \Omega_2) \cup S} |\nabla \varphi|^2 dx, \quad k = 1, \dots, J_m. \quad (14)$$

Here $|S|$ is the surface measure of S and the constants C_1 and C_2 do not depend on ε . If $\Omega_2 \subset \Omega_1$, then one has that

$$|\mu_k^{-1} - \lambda_m^{-1}| \geq c \min_{\varphi \in X_m, \|\varphi\|=1} \int_{(\Omega_1 \setminus \Omega_2)} |\nabla \varphi|^2 dx, \quad k = 1, \dots, J_m,$$

where the integral is the same as in the right-hand side of (14). If we assume that the eigenfunctions φ_k , $k = 1, \dots, J_m$, belong to $W^{1,p}(\Omega_1)$ for some $p \in (2, \infty]$ then formula (14) implies that

$$|\mu_k^{-1} - \lambda_m^{-1}| \leq C |\Omega_1 \triangle \Omega_2|^{1-\frac{2}{p}}, \quad (15)$$

where $\Omega_1 \triangle \Omega_2$ denotes the symmetric difference of Ω_1 and Ω_2 . Estimate (15) is equivalent to (1.8), [2], but the assumptions imposed on Ω_1 and Ω_2 are weaker here; in particular, all egenfunctions are required to belong to $W^{1,p}(\Omega_1)$ in [2].

Let Ω_1 be a bounded domain, $x_0 \in \partial\Omega_1$ and $\varepsilon > 0$. We assume that

$$\Omega_1 \setminus B_\varepsilon(x_0) \subset \Omega_2 \subset \Omega_1 \cup B_\varepsilon(x_0) = \Omega_\varepsilon^+(x_0), \quad (16)$$

where $B_\varepsilon(x_0)$ is the ball of the radius ε centered at x_0 . Let also for all $u \in W^{1,2}(B_{q\varepsilon}(x_0))$ such that $u = 0$ on $B_{q\varepsilon}(x_0) \setminus \Omega_\varepsilon^+(x_0)$ the following inequality

$$\int_{\Omega_\varepsilon^+(x_0) \cap B_{q\varepsilon}(x_0)} |u|^2 dx \leq C\varepsilon^2 \int_{\Omega_\varepsilon^+(x_0) \cap B_{q\varepsilon}(x_0)} |\nabla u|^2 dx \quad (17)$$

hold with some $q > 1$ and a constant C independent of ε . Then it is proved in Sect.5.1 that

$$|\mu_k^{-1} - \lambda_m^{-1}| \leq C \max_{\varphi \in X_m, \|\varphi\|=1} \int_{\Omega_1 \setminus B_{q\varepsilon}(x_0)} |\nabla \varphi|^2 dx, \quad k = 1, \dots, J_m.$$

In Sect.5.2, we consider perturbations satisfying (13). Assuming that for some $q > 1$ inequality (17) holds for all $x_0 \in \partial\Omega_1$, we prove the following estimate for the perturbed eigenvalues:

$$|\mu_k^{-1} - \lambda_m^{-1}| \leq C \max_{\varphi \in X_m, \|\varphi\|=1} \int_{\Omega_1 \setminus \Omega_{q\varepsilon}} |\nabla \varphi|^2 dx, \quad k = 1, \dots, J_m.$$

Notice that the Sobolev space $\mathring{W}^{1,2}(\Omega_k)$ can be considered as a subspace of a similar Sobolev space in a larger domain. Thus, we present an abstract approach for comparison of eigenvalues and eigenfunctions of operators acting in different subspaces of a certain Hilbert space in Sections 2–4. There, in order to measure the proximity of two subspaces, we introduce a distance σ as a norm of two projectors onto these subspaces and show that the convergence with respect to this distance is equivalent to the Mosco convergence. In Sect. 2, we formulate a proposition about the closeness of eigenvalues and eigenfunctions of two eigenvalue problems, see Proposition 3. In the same section, we present the main asymptotic theorem, see Theorem 2. Proposition 3 and Theorem 2 are proved in Sections 3 and 4, respectively. In Sect. 5, we apply our asymptotic formula to eigenvalues of the Dirichlet problem for a second-order differential operator. In particular, we consider local and global perturbations of the boundary; the case of uniformly Lipschitz boundaries is also treated. A class of domain perturbations, for which the quantity σ is small, is described for both cases; a stability estimate, which evaluates eigenvalues by integrals of squares of the gradient of eigenfunctions, is also presented.

2 Perturbation of eigenvalues. Abstract version.

2.1 Statement of the perturbation problem

Here we present an abstract approach for study of perturbation of eigenvalues to the spectral problems from Introduction keeping the same notations.

Let H and \mathcal{H} be Hilbert spaces with inner products (\cdot, \cdot) and $\langle \cdot, \cdot \rangle$ and with corresponding norms $\|\cdot\|$ and $|\cdot|$ respectively. We suppose that H is compactly imbedded in \mathcal{H} . This implies existence of $c_0 > 0$ such that

$$|u| \leq c_0 \|u\|, \quad \text{for } u \in H. \quad (18)$$

Let H_1 and H_2 be two subspaces of H of infinite dimension. We introduce the operators $K_j : H_j \rightarrow H_j$ by $(K_j u, v) = \langle u, v \rangle$, where $u, v \in H_j$, $j = 1, 2$. One can check that the operators K_1 and K_2 are self-adjoint, positive definite and compact. Let \mathcal{H}_j , $j = 1, 2$, be the closure of H_j in the space \mathcal{H} . From the definition of K_j , $j = 1, 2$, it follows that the operator can be extended to \mathcal{H}_j and

$$\|K_j\|_{\mathcal{H}_j \rightarrow H_j} \leq c_0,$$

where c_0 is the constant in (18).

We consider two spectral problems

$$K_1 \varphi = \lambda^{-1} \varphi, \quad \varphi \in H_1, \quad (19)$$

and

$$K_2 U = \mu^{-1} U, \quad U \in H_2, \quad (20)$$

We denote by λ_j^{-1} , $j = 1, \dots$, eigenvalues of K_1 , numerated according to $0 < \lambda_1 < \lambda_2 < \dots$, and by X_j corresponding eigenspaces. We put $J_j = \dim X_j$. Our goal is to study eigenvalues of (20) located in a neighborhood of λ_m for certain fixed m .²

We denote by S_j , $j = 1, 2$, the orthogonal projector in H with the image H_j . We will measure the proximity between H_1 and H_2 by the constant $\sigma = \sigma(H_1, H_2)$ in the inequality

$$|(S_1 - S_2)u|^2 \leq \sigma \|u\|^2, \quad u \in H. \quad (21)$$

The quantity $\sigma(H_1, H_2)$ is a distance in $\mathcal{S}(H)$ -the space of all closed subspaces of H . In the next proposition we prove that the Mosco convergence of subspaces (see Sect. 2.3.3 in [10] or Sect.4.5 in [1]) is equivalent to convergence of subspaces with respect to the distance σ .

²We note that the spectral problem (19) and (4), and also (20) and (5) have the same eigenvalues and eigenvectors.

Proposition 1. *Let H_j , $j = 1, \dots$, and H_* be subspaces in H and let S_j and S_* be corresponding orthogonal projectors. the following assertions are equivalent:*

- (i) *for all $u \in H$, $S_j u \rightarrow S_* u$ as $j \rightarrow \infty$;*
- (ii) *$\sigma(H_j, H_*) \rightarrow 0$ as $j \rightarrow \infty$.*

Proof. We denote by $0 < s_1 \leq s_2 \leq \dots$, eigenvalues of problem (19), when $H_1 = H$. Here, we numerate them accounting their multiplicities. Due to the compactness of the imbedding $H \subset \mathcal{H}$, $s_k \rightarrow \infty$ as $k \rightarrow \infty$. We denote by v_1, v_2, \dots , corresponding eigenvectors which form an orthogonal basis in H . We normalize them according to $\|v_k\| = 1$. It is clear that

$$(v_k, w) = s_k \langle v_k, w \rangle \quad \text{for all } w \in H.$$

Therefore, the vectors $w_k = \sqrt{s_k} v_k$, $k = 1, 2, \dots$, form the orthogonal basis in \mathcal{H} subject to $\|w_k\| = 1$.

Let us show that (i) implies (ii). We have

$$(S_j - S_*)u = \sum_{k=1}^N s_k^{-1/2} (u, (S_j - S_*)v_k) w_k + \sum_{k=N+1}^{\infty} s_k^{-1/2} ((S_j - S_*)u, v_k) w_k.$$

Therefore,

$$\|(S_j - S_*)u\|^2 \leq \left(\sum_{k=1}^N s_k^{-1} \|(S_j - S_*)v_k\|^2 + s_{N+1}^{-1} \right) \|u\|^2.$$

this implies

$$\sigma(H_j, H_*) \leq \inf_N \left(\sum_{k=1}^N s_k^{-1} \|(S_j - S_*)v_k\|^2 + s_{N+1}^{-1} \right).$$

The right hand side in the last inequality tends to zero when $j \rightarrow \infty$ because of strong convergence of S_j to S_* .

Let us prove the implication (ii) \Rightarrow (i). For $u \in H$, we have

$$((S_j - S_*)u, v_k) = s_k \langle (S_j - S_*)u, v_k \rangle \rightarrow 0 \quad \text{as } j \rightarrow \infty$$

for $k = 1, 2, \dots$. This implies that $((S_j - S_*)u, v) \rightarrow 0$ as $j \rightarrow \infty$ for all $v \in H$.

Therefore,

$$\begin{aligned} ((S_j - S_*)u, (S_j - S_*)u) &= (S_j u, u) - (S_* u, S_j u) - (S_j u, S_* u) + (S_* u, S_* u) \\ &\rightarrow (S_* u, u) - (S_* u, S_* u) - (S_* u, S_* u) + (S_* u, S_* u) = 0 \quad \text{as } j \rightarrow \infty. \end{aligned}$$

The proof is complete. \square

Using equivalence of γ convergence of domains and strong convergence of corresponding operators (see Theorem 2.3.10 in [9]), we derive from the previous assertion the following

Corollary 1. *Let Ω_j , $j = 1, \dots$, and Ω_* be domains belonging to a bounded domain D . If Ω_j γ -converges to Ω_* then $\sigma(H_j, H_*) \rightarrow 0$ as $j \rightarrow \infty$, where $H_j = \dot{W}^{1,2}(\Omega_j)$ and $H_* = \dot{W}^{1,2}(\Omega_*)$.*

Proposition 2. *The metric space $\mathcal{S}(H)$ with the distance σ is compact.*

Proof. Let S_k , $k = 1, 2, \dots$, be a family of orthogonal projectors. Let us show that one can choose a convergent subsequence. Let v_j be the same vectors as in the proof of Proposition 1. We can choose a subsequence such that

$$(S_k v_i, v_j) \rightarrow \alpha_{ij} \quad \text{as } k \rightarrow \infty \quad (22)$$

for all $i, j \geq 1$. We used the same index for the subsequence in (22). Since $\|S_k\|_{H \rightarrow H} = 1$, we derive from (22) that

$$(S_k u, v) \rightarrow \alpha(u, v) \quad \text{as } k \rightarrow \infty$$

for all $u, v \in H$, where $\alpha(u, v)$ is a certain number. One can check that the form α is linear with respect to the first argument and anti-linear with respect to the second one. Moreover, the form α is bounded. Therefore, $\alpha(u, v) = (S_* u, v)$, where S_* is an orthogonal projector. Reasoning as in the proof of Proposition 1(ii), we conclude that $S_k u \rightarrow S_* u$ as $k \rightarrow \infty$. \square

In what follows we shall use that the orthogonal projectors S_1 and S_2 possess the following symmetry property:

$$(S_2 v, w) = (v, S_1 w) \quad \text{for } v \in H_1 \text{ and } w \in H_2.$$

2.2 Formulation of results

In what follows we denote by P_m the orthogonal projector in H with the image SX_m .

Proposition 3. *There exists positive σ_0 , c and C depending on $\lambda_1, \dots, \lambda_{m+1}$ and c_0 such that for $\sigma \leq \sigma_0$ the following assertions are valid:*

(i) *The operator K_2 has exactly J_m eigenvalue in $(1/\lambda_{m+1} + c\sigma^{1/2}, 1/\lambda_{m-1} - c\sigma^{1/2})$ and all of them are located in $(1/\lambda_m - c\sigma^{1/2}, 1/\lambda_m + c\sigma^{1/2})$.*

(ii) If μ^{-1} is an eigenvalue of (20) from the interval $(1/\lambda_m - c\sigma^{1/2}, 1/\lambda_m + c\sigma^{1/2})$ and U is a corresponding eigenvector then

$$\|U - P_m U\| \leq C\sigma^{1/2}\|U\|. \quad (23)$$

We denote by μ_k^{-1} , $k = 1, \dots, J_m$, the eigenvalues of the spectral problem (20) located in the interval $(\lambda_m^{-1} - c\sigma, \lambda_m^{-1} + c\sigma)$, where c is the same positive constant as in Proposition 3. In order to formulate the main result of this paper let us introduce some more objects. We put $T_2 u = u - S_2 u$ and define the vector $\Psi = \Psi_\varphi \in H_2$ as the solution of equation (7). Let also

$$\rho = \max_{\varphi \in X_m, \|\varphi\|=1} (\sigma \|\Psi_\varphi\|^2 + \|T\varphi\|^2 + \|\Psi_\varphi\|^2). \quad (24)$$

Theorem 2. *The following asymptotic formula holds:*

$$\mu_k^{-1} = \lambda_m^{-1} + \tau_k + O(\rho + |\tau_k|\sigma), \quad k = 1, \dots, J_m, \quad (25)$$

where $\tau = \tau_k$ is an eigenvalue of the problem

$$\frac{1}{\lambda_m} \left((\Psi_\varphi, \Psi_\psi) - (T\varphi, T\psi) - (\Psi_\varphi, \psi) - (\varphi, \Psi_\psi) \right) = \tau (S\varphi, S\psi) \quad \text{for all } \psi \in X_m, \quad (26)$$

where $\varphi \in X_m$. Moreover, τ_1, \dots, τ_m in (25) run through all eigenvalues of (26) counting their multiplicities.

In Sect. 3.1 we prove that $\|\varphi\|^2 - \|S\varphi\|^2 \leq 1 - C\sigma^{1/2}$ for $\varphi \in X_m$. This fact and Theorem 2 lead to the following corollaries.

Corollary 2. *If $H_2 \subset H_1$ then $\Psi_\varphi = 0$ for all $\varphi \in X_m$. Therefore, from (26) it follows that $|\tau_k| \leq C\rho$ and hence (25) implies*

$$c \min_{\varphi \in X_m, \|\varphi\|=1} \|T\varphi\|^2 \leq |\mu_k^{-1} - \lambda_m^{-1}| \leq C \max_{\varphi \in X_m, \|\varphi\|=1} \|T\varphi\|^2,$$

where c and C are positive constants.

Corollary 3. *Let S_0 be the orthogonal projector onto $H_1 \cap H_2$ and let $T_0 u = u - S_0 u$. Then $(\Psi_\varphi, \psi) = (\Psi_\varphi, T_0 \psi)$ and therefore from (26) it follows $|\tau_k| \leq C\rho_0$, where*

$$\rho_0 = \max_{\varphi \in X_m, \|\varphi\|=1} (\|T_0 \varphi\|^2 + \|\Psi_\varphi\|^2).$$

Consequently,

$$|\mu_k^{-1} - \lambda_m^{-1}| \leq C\rho_0$$

for arbitrary H_1 and H_2 subject to (21) with small constant σ .

Remark 1. In [12] and [14] another quantity is used to measure proximity of two subspaces. It is defined as a constant $\sigma_* = \sigma_*(H_1, H_2)$ in the inequality

$$\|u\|^2 \leq \sigma_* \|u\|^2 \quad (27)$$

for all elements u from $u \in H_1 + H_2$ orthogonal to $H_1 \cap H_2$. Since

$$\|(S_1 - S_2)u\| \leq \sqrt{\sigma_*} \|(S_1 - S_2)u\| \leq 2\sqrt{\sigma_*} \|u\|,$$

the constant σ in (21) is subject to

$$\sigma \leq 4\sigma_*. \quad (28)$$

One can check that $T\varphi$ and Ψ_φ belong to $H_1 + H_2$ and both of them are orthogonal to $H_1 \cap H_2$ for $\varphi \in X_m$. Therefore, $\|T\varphi\| \leq \sigma_* \|T\varphi\|^2$ and $\|\Psi_\varphi\| \leq \sigma_* \|\Psi_\varphi\|^2$ due to (27). So, under assumption (27), we have

$$\rho \leq c\sigma_*\rho_*, \quad \rho_* = \max_{\varphi \in X_m, \|\varphi\|=1} (\|T\varphi\|^2 + \|\Psi_\varphi\|^2). \quad (29)$$

Using (29), we see that formula (25) in Theorem 2 implies the following asymptotic formula

$$\mu_k^{-1} = \lambda_m^{-1} + \tau_k + O(\sigma_*\rho_* + |\tau_k|\sigma_*), \quad k = 1, \dots, J_m, \quad (30)$$

where $\tau = \tau_k$ is an eigenvalue of the problem (10). Moreover, τ_1, \dots, τ_m in (30) run through all eigenvalues of (10) counting their multiplicities.

3 Proof of Proposition 3

In what follows, we put $\mathcal{X}_m = X_1 + \dots + X_m$ and by \mathcal{Y}_m we denote the orthogonal complement of $S\mathcal{X}_m$ in H_2 .

An important role will be played by the operator

$$B = K_2 S_2 - S_2 K_1. \quad (31)$$

In Sect 3.1, we will show in particular, that the norm of $B : H_1 \rightarrow H_2$ is estimated by a constant times $\sigma^{1/2}$.

By c, C, \dots we denote various constants depending on $\lambda_1, \dots, \lambda_{m+1}$ and c_0 .

3.1 Some inequalities

In this section we prove some important estimates, which follow from (21) and which will be used in the proofs of Proposition 3 and Theorem 2.

Estimates for S_2 . Let us start with the following inequality

$$\left(1 - \Lambda_m \sigma^{1/2}\right) \|\varphi\|^2 \leq \|S\varphi\|^2 \leq \|\varphi\|^2 \quad \text{for } \varphi \in \mathcal{X}_m, \quad (32)$$

where $\Lambda_m = \left(\sum_{k=1}^m \lambda_k\right)^{1/2}$. Since S_2 is an orthogonal projector, the right inequality is obvious. Let us prove the left one. We represent φ as $\varphi_0 + \varphi_1$, where $\varphi_0 = S_2\varphi$ and $\varphi_1 = (I - S_2)\varphi$. Since $S_2\varphi_0 = \varphi_0$ and $S_2\varphi_1 = 0$, we have $\|S\varphi\|^2 = \|\varphi_0\|^2$. Using that $S_1\varphi = \varphi$, we conclude that $\varphi_1 = (S_1 - S_2)\varphi$ and from (21) it follows that

$$\|\varphi_1\|^2 = \|(S_1 - S_2)\varphi\|^2 \leq \sigma \|\varphi\|^2. \quad (33)$$

Representing φ as $\varphi = \zeta_1 + \dots + \zeta_m$, where $\zeta_k \in X_k$, and noting that $(\varphi_1, \varphi_1) = (\varphi, \varphi_1)$, we get

$$\begin{aligned} (\varphi_1, \varphi_1) &= \sum_{k=1}^m (\zeta_k, \varphi_1) = \sum_{k=1}^m \lambda_k \langle \zeta_k, \varphi_1 \rangle \leq \sum_{k=1}^m \lambda_k \|\zeta_k\| \|\varphi_1\| \\ &\leq \sum_{k=1}^m \lambda_k^{1/2} \|\zeta_k\| \|\varphi_1\| \leq \Lambda_m \|\varphi\| \|\varphi_1\|, \end{aligned}$$

where we used that ζ_k is an eigenvector of K_1 corresponding to the eigenvalue λ_k^{-1} . Using the last inequality together with (33), we get

$$(\varphi_1, \varphi_1) \leq \Lambda_m \sigma^{1/2} (\varphi, \varphi).$$

Therefore,

$$\|S\varphi\|^2 = \|\varphi_0\|^2 = \|\varphi\|^2 - \|\varphi_1\|^2 \geq (1 - \Lambda_m \sigma^{1/2}) \|\varphi\|^2,$$

which implies the left inequality in (32).

From (32) one can derive the estimate

$$|(S\varphi, S\psi) - (\varphi, \psi)| \leq 3\Lambda_m \sigma^{1/2} (\|\varphi\|^2 + \|\psi\|^2) \quad \text{for } \varphi, \psi \in \mathcal{X}_m. \quad (34)$$

Indeed, introduce the form $b(u, v) = (S\varphi, S\psi) - (\varphi, \psi)$. By (33),

$$|b(u, u)| \leq \varepsilon \|u\|^2, \quad (35)$$

where $\varepsilon = \Lambda_m \sigma^{1/2}$. Using that

$$b(\varphi + \psi, \varphi + \psi) = b(\varphi, \varphi) + b(\psi, \psi) + b(\varphi, \psi) + b(\psi, \varphi)$$

and applying (35) for estimating quadratic terms here we obtain

$$|b(\varphi, \psi) + b(\psi, \varphi)| \leq 3\varepsilon(\|\varphi\|^2 + \|\psi\|^2).$$

Similar arguments applied to $b(\varphi + i\psi, \varphi + i\psi)$ give the estimate

$$|b(\varphi, \psi) - b(\psi, \varphi)| \leq 3\varepsilon(\|\varphi\|^2 + \|\psi\|^2),$$

which together with the previous one leads to (34).

An estimate for the operator B. Let us prove that

$$\|Bv\| \leq 2c_0\sigma^{1/2}\|v\| \quad \text{for } v \in H_1. \quad (36)$$

For $w \in H_2$ we have

$$(Bv, w) = (K_2 S_2 v, w) - (S_2 K_1 v, w) = \langle S_2 v, w \rangle - \langle v, S_1 w \rangle. \quad (37)$$

We write v and w as $v = S_2 v + (I - S_2)v = v_0 + v_1$ and $w = S_1 w + (I - S_1)w = w_0 + w_1$. Then (37) implies

$$(Bv, w) = \langle v_0, w_1 \rangle - \langle v_1, w_0 \rangle, \quad (38)$$

where we have used the equalities $S_2 v_1 = S_1 w_1 = 0$. Since $v_1 = (S_1 - S_2)v$ and $w_1 = (S_2 - S_1)w$, then by using estimate (21) for function containing index 1, we get

$$|(Bv, w)| \leq \sigma^{1/2}(\|v_0\| \|w_1\| + \|v_1\| \|w_0\|).$$

Applying here estimate (18), we get

$$|(Bv, w)| \leq 2c_0\sigma^{1/2}\|v\|\|w\|,$$

which implies (36).

An inequality for K_1 and K_2 . Finally, let us show that

$$(K_2 w, w) \leq (K_1 S_1 w, S_1 w) + (2c_0\sqrt{\sigma} + \sigma)\|w\|^2 \quad \text{for } w \in H_2, \quad (39)$$

or what is equivalent, due to the definition of K_1 and K_2 ,

$$\|w\|^2 \leq \|S_1 w\|^2 + (2c_0\sqrt{\sigma} + \sigma)\|w\|^2. \quad (40)$$

We write $w = S_1 w + (I - S_1)w = w_0 + w_1$. Since $S_1 w_1 = 0$ and $S_1 w_0 = w_0$, relation (40) takes the form

$$\|w_0 + w_1\|^2 \leq \|w_0\|^2 + (2c_0\sqrt{\sigma} + \sigma)\|w\|^2.$$

or

$$\|w_1\|^2 + 2\langle w_0, w_1 \rangle \leq (2c_0\sqrt{\sigma} + \sigma)\|w\|^2. \quad (41)$$

Using that $w_1 = (S_2 - S_1)w$ and applying (21), we estimate the left-hand side of (41) by

$$\sigma\|w\|^2 + 2\sigma^{1/2}\|w_0\|\|w\|.$$

According to (18), $\|w_0\| \leq c_0\|w_0\| \leq c_0\|w\|$, which implies (41) and hence (39).

3.2 Proof of Proposition 3(i)

1) Let $u \in \mathcal{Y}_m$. Then $S_1 u$ is orthogonal to $X_1 + \dots + X_m$ and by (39)

$$\begin{aligned} (K_2 u, u) &\leq (K_1 S_1 u, S_1 u) + c\sigma^{1/2}\|u\|^2 \leq \frac{1}{\lambda_{m+1}}(S_1 u, S_1 u) + c\sigma^{1/2}\|u\|^2 \\ &\leq \left(\frac{1}{\lambda_{m+1}} + c\sigma^{1/2} \right) \|u\|^2. \end{aligned} \quad (42)$$

From this inequality it follows that there are $\leq J_1 + \dots + J_m$ eigenvalues of K_2 counted together with their multiplicity in the interval $(1/\lambda_{m+1} + c\sigma^{1/2}, \infty)$.

2) Let $u = S_2 \varphi$, $\varphi \in \mathcal{X}_m$. Then

$$(K_2 u, u) = (S_2 K_1 \varphi, S_2 \varphi) + (B \varphi, u)$$

and using (36), we get

$$(K_2 u, u) \geq (S_2 K_1 \varphi, S_2 \varphi) - C\sigma^{1/2}\|\varphi\|\|u\|.$$

Applying (34) to the first term in the left-hand side, we obtain

$$(K_2 u, u) \geq (K_1 \varphi, \varphi) - C_1 \sigma^{1/2}(\|K_1 \varphi\|^2 + \|\varphi\|^2) - C\sigma^{1/2}\|\varphi\|\|u\|,$$

which leads to

$$(K_2 u, u) \geq (K_1 \varphi, \varphi) - C_2 \sigma^{1/2} (\|\varphi\|^2 \geq \left(\frac{1}{\lambda_{m-1}} - C_2 \sigma^{1/2} \right) \|\varphi\|^2.$$

Applying (32) for estimating the second term in the right-hand side in the last inequality, we arrive at

$$(K_2 u, u) \geq \left(\frac{1}{\lambda_m} - C_2 \sigma^{1/2} \right) \|u\|^2.$$

This implies that there are $\geq J_1 + \dots + J_m$ eigenvalues of K_2 in the interval $(1/\lambda_m - C_2 \sigma^{1/2}, \infty)$.

Combining 1) and 2) we conclude that there are exactly $J_1 + \dots + J_m$ eigenvalues of K_2 in the interval (γ_1, ∞) for $\gamma_1 \in (1/\lambda_{m+1} + C_1 \sigma^{1/2}, 1/\lambda_m - C_2 \sigma^{1/2})$.

Applying 1) and 2) with m replaced by $m - 1$ we obtain that there are exactly $J_1 + \dots + J_{m-1}$ eigenvalues of K_2 in the interval (γ_2, ∞) for $\gamma_2 \in (1/\lambda_m + C_1 \sigma^{1/2}, 1/\lambda_{m-1} - C_2 \sigma^{1/2})$. Therefore there exists positive constants c and σ_0 depending on $\lambda_1, \dots, \lambda_{m+1}$ and c_0 such that the intervals $(1/\lambda_m - c\sigma^{1/2}, 1/\lambda_m + c\sigma^{1/2})$ and $(1/\lambda_{m+1} + c\sigma^{1/2}, 1/\lambda_{m-1} - c\sigma^{1/2})$ contains exactly J_m eigenvalues of K_2 for $\sigma \leq \sigma_0$. The proof of Proposition 3(i) is complete.

3.3 Proof of Proposition 3(ii)

First, let us consider the equation

$$Q_m K_2 w - \mu w = f, \tag{43}$$

where $f, w \in Q_m H_2$, $Q_m = I - P_m$ and

$$|\mu^{-1} - \lambda_m^{-1}| \leq c\sigma^{1/2}, \tag{44}$$

where c is the same constant as in (i). Our first goal is to prove the estimate

$$\|w\| \leq c_1 \|f\| \tag{45}$$

for solutions of equation (43). Here the constant c_1 depends on $\lambda_1, \dots, \lambda_{m+1}$ and c_0 .

We represent $S_2\mathcal{X}_m$ as $S_2X_m + Y_m$ where Y_m is the orthogonal complement to S_2X_m in $S_2\mathcal{X}_m$. We introduce the orthogonal projectors R_m onto Y_m and \mathcal{T}_m onto \mathcal{Y}_m . Then $Q_m = R_m + \mathcal{T}_m$. We write $w = w_0 + w_1$ and $f = f_0 + f_1$, where $w_0, f_0 \in Y_m$ and $w_1, f_1 \in \mathcal{Y}_m$. Equation (43) can be written as the following system of equations

$$R_m K_2(w_0 + w_1) - \mu w_0 = f_0 \quad (46)$$

and

$$\mathcal{T}_m K_2 w_1 - \mu w_1 = f_1 - \mathcal{T}_m K_2 w_0. \quad (47)$$

From the second equation we get

$$(K_2 w_1, w_1) - \mu(w_1, w_1) = (f_1 - \mathcal{T}_m K_2 w_0, w_1).$$

Using estimate (42), we obtain

$$\mu(w_1, w_1) - \left(\frac{1}{\lambda_{m+1}} + c\sigma^{1/2} \right) \|w_1\|^2 \leq (\mathcal{T}_m K_2 w_0, w_1) - (f_1, w_1),$$

which implies that the operator $\mu^{-1} - Q_m K_2$ is positive definite on \mathcal{Y}_m , equation (47) is uniquely solvable and its solution satisfies

$$\left(\mu - \frac{1}{\lambda_{m+1}} - c\sigma^{1/2} \right) \|w_1\|_2 \leq \|\mathcal{T}_m K_2 w_0\| + \|f_1\|. \quad (48)$$

Furthermore, representing w_0 as $S\varphi_0$, $\varphi_0 \in \mathcal{X}_m$ we have

$$\mathcal{T}_m K_2 w_0 = \mathcal{T}_m K_2 S_2 \varphi_0 = \mathcal{T}_m (S_2 K_1 \varphi_0 + B\varphi_0) = \mathcal{T}_m B\varphi_0.$$

Therefore, by (36) and (32)

$$\|\mathcal{T}_m K_2 w_0\| \leq c\sigma^{1/2} \|\varphi_0\| \leq c_1 \sigma^{1/2} \|w_0\|. \quad (49)$$

Combining this estimate with (48), we get

$$\|w_1\| \leq C(\|f_1\| + \sigma^{1/2} \|w_0\|). \quad (50)$$

We represent φ_0 as $\varphi' + \varphi_m$, where $\varphi_m \in X_m$ and $\varphi' \in X_1 + \dots + X_{m-1}$. Since $(S_2 \varphi_0, S_2 \varphi_m) = 0$, using (34) we get

$$(\varphi_0, \varphi_m) = \|\varphi_m\|^2 \leq 3\lambda_m^{1/2} \sigma^{1/2} \|\varphi_0\|^2. \quad (51)$$

Therefore,

$$(K_2 w_0, w_0) = (K_2 S_2 \varphi_0, S_2 \varphi_0) = (S_2 K_1 \varphi_0, S_2 \varphi_0) + (B \varphi_0, S_2 \varphi_0).$$

Using (34) and (36) for estimating the first and second terms in the right-hand side of the last relation respectively we get

$$(K_2 w_0, w_0) \geq (K_1 \varphi_0, \varphi_0) - c \sigma^{1/2} \|\varphi_0\|^2, \quad (52)$$

where c depends on the eigenvalues λ_1, λ_m and the constant c_0 in (18). Since

$$(K_1 \varphi_0, \varphi_0) = (K_1 \varphi_m, \varphi_m) + (K_1 \varphi', \varphi') = \frac{1}{\lambda_m} \|\varphi_m\|^2 + (K_1 \varphi', \varphi'),$$

by using (51), we get

$$(K_1 \varphi_0, \varphi_0) \geq \left(\frac{1}{\lambda_{m-1}} - c \sigma^{1/2} \right) \|\varphi_0\|^2 \geq \left(\frac{1}{\lambda_{m-1}} - c_1 \sigma^{1/2} \right) \|w_0\|^2.$$

Here we used also (32) in order to obtain the last inequality. Applying the last estimate to the first term in the right-hand side in (52), we obtain

$$(K_2 w_0, w_0) \geq \left(\frac{1}{\lambda_{m-1}} - c_2 \sigma^{1/2} \right) \|w_0\|^2. \quad (53)$$

Using (53) and (49), we conclude that equation (46) is solvable with respect to w_0 and $\|w_0\| \leq c \|f\|$. Similar estimate for w_1 follows from (50). Conclusively, equation (43) is uniquely solvable and for its solution $w \in Q_m H_2$ estimate (45) holds.

Let μ^{-1} be an eigenvalue of problem (20) satisfying (44) and let U be a corresponding eigenvector. We represent it as $U = S_2 \varphi_m + w$ where $\varphi_m \in X_m$ and $w \in w \in Q_m H_2$. Then

$$K_2(S_2 \varphi_m + w) = \frac{1}{\mu}(S_2 \varphi_m + w).$$

We write this relation as

$$K_2 w - \frac{1}{\mu} w + S_2 K_1 \varphi_m + B \varphi_m - \frac{1}{\mu} S_2 \varphi_m = 0,$$

or, equivalently

$$K_2 w - \frac{1}{\mu} w = -B\varphi_m + \left(\frac{1}{\mu} - \frac{1}{\lambda_m}\right) S_2 \varphi_m. \quad (54)$$

We denote the left-hand side by f . Then by (36) and assumption on μ , we have

$$\|f\| \leq C\sigma^{1/2}\|\varphi_m\|.$$

Applying operator \mathcal{T}_m to both sides in (54) and using (45) together with the last estimate of f we obtain

$$\|w\| \leq C\sigma^{1/2}\|\varphi_m\| \leq C_1\sigma^{1/2}\|w\|. \quad (55)$$

3.4 Corollary of Proposition 3

Let Z_m be the space of eigenvectors of the problem (20) corresponding to eigenvalues located in the interval $(1/\lambda_m - c\sigma^{1/2}, 1/\lambda_m + c\sigma^{1/2})$, see Proposition 3(i). According to the same proposition $\dim Z_m = J_m$. Let us show that

$$|(U, V) - (P_m U, P_m V)| \leq C\sigma(\|U\|^2 + \|V\|^2) \quad \text{for all } U, V \in Z_m. \quad (56)$$

First, let us check that

$$(U, U) - (P_m U, P_m U) \leq C\sigma\|U\|^2 \quad \text{for all } U \in Z_m. \quad (57)$$

Indeed, introduce an orthonormal basis U_1, \dots, U_{J_m} in Z_m consisting of eigenvectors of problem (20). If $U \in Z_m$ we represent it as $U = a_1 U_1 + \dots + a_{J_m} U_{J_m}$. Using (23), we get

$$\|U - P_m U\| \leq \sum_{j=1}^{J_m} |a_j| \|U_j - P_m U_j\| \leq c\sigma^{1/2} \sum_{j=1}^{J_m} |a_j| \leq c_1\sigma^{1/2}\|U\|,$$

which implies (57) since $\|U - P_m U\|^2 = (U, U) - (P_m U, P_m U)$.

In order to prove (56) we introduce the quasi-linear form $b(U, V) = (U, V) - (P_m U, P_m V)$. Since $b(U, U) \geq 0$, we have

$$|b(U, V)| \leq b(U, U)^{1/2} b(V, V)^{1/2} \leq \frac{1}{2}(b(U, U) + b(V, V)),$$

which together with (57) implies (56).

4 Proof of Theorem 2

4.1 A finite dimensional reduction

We represent the function $U \in H_2$ in (5) as $U = S_2\varphi + w$, where $\varphi \in X_m$ and $w \in Q_m H_2$. Then (5) takes the form

$$(\mu^{-1} - K_2)(S_2\varphi + w) = 0.$$

By using the operator B , we can write the last relation as

$$\left(\frac{1}{\mu} - \frac{1}{\lambda_m}\right)S_2\varphi - B\varphi + (\mu^{-1} - K_2)w = 0$$

Applying operators P_m and Q_m we get

$$\left(\frac{1}{\mu} - \frac{1}{\lambda_m}\right)S_2\varphi - P_mB\varphi - P_mK_2w = 0 \quad (58)$$

and

$$Q_m(\mu^{-1} - K_2)w = Q_mB\varphi. \quad (59)$$

We assume that μ satisfies (44). Then the last equation coincides with (43) if we take there $f = -Q_mB\varphi$. Therefore equation (59) is uniquely solvable and

$$\|(\mu^{-1} - Q_mK_2Q_m)^{-1}\|_{Q_mH_2 \rightarrow Q_mH_2} \leq c_1, \quad (60)$$

where c_1 is the constant from (45). Inserting $w = (\mu^{-1} - Q_mK_2Q_m)^{-1}Q_mB\varphi$ in (58), we obtain

$$\begin{aligned} \left(\frac{1}{\mu} - \frac{1}{\lambda_m}\right)S_2\varphi - P_mB\varphi - R(\lambda, \varphi) &= 0, \\ R(\lambda, \varphi) &= P_mK_2Q_m(\mu^{-1} - Q_mK_2Q_m)^{-1}Q_mB\varphi \end{aligned} \quad (61)$$

We represent R as $R(\lambda, \varphi) = \mu P_mK_2Q_mB\varphi + R_1(\lambda, \varphi)$, where

$$R_1(\lambda, \varphi) = P_mK_2Q_mK_2Q_m(\mu^{-1} - Q_mK_2Q_m)^{-1}Q_mB\varphi.$$

Equation (61) becomes

$$\left(\frac{1}{\mu} - \frac{1}{\lambda_m}\right)S_2\varphi - P_mB\varphi - \mu P_mK_2Q_mB\varphi - R_1(\lambda, \varphi) = 0. \quad (62)$$

Taking the inner product of the left-hand side with $S_2\psi$ in H_2 , where $\psi \in X_m$, we get

$$\left(\frac{1}{\mu} - \frac{1}{\lambda_m}\right)(S_2\varphi, S_2\psi)_2 - (B\varphi, S_2\psi) - \mu(Q_m B\varphi, K_2 S_2\psi) - (R_1(\lambda, \varphi), S_2\psi) = 0$$

Using that $Q_m K_2 S_2\psi = Q_m B\psi$, we arrive at

$$\left(\frac{1}{\mu} - \frac{1}{\lambda_m}\right)(S_2\varphi, S_2\psi)_2 - (B\varphi, S_2\psi)_2 - \mu(Q_m B\varphi, B\psi)_2 - (L(\mu)B\varphi, B\psi)_2 = 0, \quad (63)$$

where $L(\mu) = Q_m K_2 Q_m (\mu^{-1} - Q_m K_2 Q_m)^{-1} Q_m$. By (60), one can check the following estimate for the last operator

$$|(L(\mu)B\psi, B\psi)| \leq C(K_2 Q_m B\psi, Q_m B\psi)_2 = C\langle Q_m B\psi, Q_m B\psi \rangle. \quad (64)$$

4.2 Representation of $(B\varphi, S\psi)$

For $\varphi, \psi \in X_m$, we use the representations

$$\varphi = S_2\varphi + T_2\varphi, \quad \psi = S_2\psi + T_2\psi. \quad (65)$$

Then

$$\begin{aligned} (B\varphi, S_2\psi) &= \langle S_2\varphi, S_2\psi \rangle - \lambda_m^{-1}(S_2\varphi, S_2\psi) \\ &= \langle \varphi - T_2\varphi, \psi - T_2\psi \rangle - \lambda_m^{-1}(\varphi - T_2\varphi, \psi - T_2\psi). \end{aligned} \quad (66)$$

Using the relation

$$(\Psi_\varphi, S_2\psi) = (\Psi_\varphi, \psi) = (\varphi, S_2\psi) - \lambda_m \langle \varphi, S_2\psi \rangle$$

and similar equality with exchanged φ and ψ , we obtain

$$\begin{aligned} (\varphi - T_2\varphi, \psi - T_2\psi) &= (T_2\varphi, T_2\psi) - (\varphi, \psi) + (\varphi, S_2\psi) + (S_2\varphi, \psi) \\ &= (T_2\varphi, T_2\psi) - \lambda_m \langle \varphi, \psi \rangle + (\Psi_\varphi, \psi) + \lambda_m \langle \varphi, S_2\psi \rangle \\ &\quad + (\varphi, \Psi_\psi) + \lambda_m \langle S_2\varphi, \psi \rangle. \end{aligned}$$

Replacing the last term in (66) according to this formula, we derive from (66)

$$(B\varphi, S_2\psi) = \langle T_2\varphi, T_2\psi \rangle - \lambda_m^{-1} \left((T_2\varphi, T_2\psi) + (\Psi_\varphi, \psi) + (\varphi, \Psi_\psi) \right). \quad (67)$$

Corollary 4. *Let $H_2 \subset H_1$ and let λ_1 be the first eigenvalue of (19) and φ be a corresponding eigenfunction. Then $\Psi_\varphi = \Psi_\psi = 0$ and from (63) and (67) it follows that*

$$1 - \frac{\lambda_1}{\mu_1} \leq \frac{\|T\varphi\|^2}{\|S\varphi\|^2}, \quad (68)$$

where μ_1 is the first eigenvalue of (20).

4.3 Representation of $(Q_m B\varphi, B\psi)$

For $\varphi \in X_M$ and $w \in H_2$, we have the following representation

$$\begin{aligned} (B\varphi, w) &= (K_2 S_2 \varphi, w) - \frac{1}{\lambda_m} (S_2 \varphi, w) = \langle \varphi - T_2 \varphi, w \rangle - \frac{1}{\lambda_m} (\varphi, w) \\ &= -\frac{1}{\lambda_m} (\Psi_\varphi, w) - \langle T_2 \varphi, w \rangle. \end{aligned}$$

This implies

$$B\varphi = K_2 \Phi_\varphi - \lambda_m^{-1} \Psi_\varphi, \quad (69)$$

where Φ_φ is the orthogonal with respect to the inner product in \mathcal{H} projection of $T_2 \varphi$ onto \mathcal{H}_2 . Using (69), we get the desired representation

$$(Q_m B\varphi, B\psi) = \frac{1}{\lambda_m^2} (\Psi_\varphi, \Psi_\psi) + b_0(\varphi, \psi), \quad (70)$$

where

$$\begin{aligned} b_0(\varphi, \psi) &= -\frac{1}{\lambda_m^2} (P_m \Psi_\varphi, \Psi_\psi) + (Q_m K_2 \Phi_\varphi, K_2 \Phi_\psi) \\ &\quad - \frac{1}{\lambda_m} (\langle \Phi_\varphi, Q_m \Psi_\psi \rangle + \langle Q_m \Psi_\varphi, \Phi_\psi \rangle). \end{aligned} \quad (71)$$

Let us estimate the form b_0 . Since

$$\|K_2 \Phi_\varphi\| \leq C \|\Phi_\varphi\| \leq C \|T_2 \varphi\|,$$

it follows from (69) and (36) that

$$\|\Psi_\varphi\| \leq C(\sigma^{1/2} \|\varphi\| + \|T_2 \varphi\|) \quad \text{for } \varphi \in X_m. \quad (72)$$

Using that $T_2 \varphi = (S_1 - S_2)\varphi$, we have that $\|T_2 \varphi\|^2 \leq \sigma \|\varphi\|^2$. Therefore, (72) implies

$$\|\Psi_\varphi\| \leq C\sigma^{1/2} \|\varphi\| \quad \text{for } \varphi \in X_m. \quad (73)$$

Let $\Upsilon_k, k = 1, \dots, J_m$, be an orthogonal basis in SX_m and let $\Upsilon_k = S\varphi_k$, $\varphi_k \in X_m$. Then

$$(P_m \Psi_\varphi, \Psi_\psi) = \sum_{k=1}^{J_m} (\Psi_\varphi, S\varphi_k) (S\varphi_k, \Psi_\psi). \quad (74)$$

Using definitions of S , K_2 and Ψ_φ , one can verify that

$$S\varphi = \lambda_m K_2 \varphi^* + \Psi_\varphi \quad \text{for } \varphi \in X_m, \quad (75)$$

where φ^* is orthogonal in \mathcal{H} projection of φ on \mathcal{H}_2 . Therefore,

$$(\Psi_\varphi, S\varphi_k) = \lambda_m \langle \Psi_\varphi, \varphi_k^* \rangle + (\Psi_\varphi, \Psi_{\varphi_k}).$$

Applying inequalities (73) for estimation of $\|\Psi_{\varphi_k}\|$, we get

$$|(\Psi_\varphi, S\varphi_k)| \leq C (\|\Psi_\varphi\| + \sigma^{1/2} \|\Psi_\varphi\|).$$

Therefore, it follows from (74) that

$$|(P_m \Psi_\varphi, \Psi_\psi)| \leq C (\sigma (\|\Psi_\varphi\|^2 + \|\Psi_\psi\|^2) + \|\Psi_\varphi\|^2 + \|\Psi_\psi\|^2). \quad (76)$$

Since K_2 is a bounded operator from \mathcal{H}_2 to H_2 , we have

$$(Q_m K_2 \Phi_\varphi, K_2 \Phi_\varphi) \leq (K_2 \Phi_\varphi, K_2 \Phi_\varphi) \leq C \langle \Phi_\varphi, \Phi_\varphi \rangle \leq C \|T\varphi\|^2. \quad (77)$$

Similarly,

$$\left| \langle \Phi_\varphi, \Psi_\psi \rangle + \langle \Psi_\varphi, \Phi_\psi \rangle \right| \leq \|T\varphi\|^2 + \|T\psi\|^2 + \|\Psi_\varphi\|^2 + \|\Psi_\psi\|^2. \quad (78)$$

Combining (76)–(78), we obtain

$$b_0(\varphi, \psi) \leq C(\wp(\varphi) + \wp(\psi)), \quad (79)$$

where

$$\wp(\varphi) = \|T_2\varphi\|^2 + \|\Psi_\varphi\|^2 + \sigma \|\Psi_\varphi\|^2.$$

We note that from the definition of ρ , see (24), it follows that

$$\rho = \max_{\varphi \in X_m, \|\varphi\|=1} \wp(\varphi). \quad (80)$$

4.4 Estimate of $\langle Q_m B\varphi, Q_m B\psi \rangle$

Let us consider first the term $\langle P_m B\varphi, B\psi \rangle$. Using the basis $\Upsilon_k = S\varphi_k$, $k = 1, \dots, J_m$, introduced in Sect. 4.3, we have

$$\langle P_m B\varphi, B\psi \rangle = \sum_{k=1}^{J_m} (B\varphi, S\varphi_k) \langle S\varphi_k, B\psi \rangle. \quad (81)$$

Applying representation (75), we get

$$(B\varphi, S\varphi_k) = \lambda_m \langle B\varphi, \varphi_k^* \rangle + (B\varphi, \Psi_{\varphi_k}),$$

which together with (73) and (69) leads

$$(B\varphi, S\varphi_k) \leq C(|T_2\varphi| + |\Psi_\varphi| + \sigma^{1/2}||\Psi_\varphi||).$$

Similarly,

$$|\langle S\varphi_k, B\psi \rangle| \leq C(|T_2\psi| + |\Psi_\psi| + \sigma^{1/2}||\Psi_\psi||).$$

Applying these for estimating the right-hand side of (81), we get

$$|\langle P_m B\varphi, B\psi \rangle| \leq C(\wp(\varphi) + \wp(\psi)). \quad (82)$$

Similar considerations give the estimate

$$|\langle P_m B\varphi, P_m B\psi \rangle| \leq C(\wp(\varphi) + \wp(\psi)). \quad (83)$$

Furthermore, using (69), we get

$$|\langle B\varphi, B\psi \rangle| \leq C(|T_2\varphi|^2 + |T_2\psi|^2 + |\Psi_\varphi|^2 + |\Psi_\psi|^2). \quad (84)$$

Applying (82)–(84) for estimating the right-hand side in (64), we get

$$|(L(\mu)B\varphi, B\psi)| \leq C(\wp(\varphi) + \wp(\psi)). \quad (85)$$

4.5 Proof of Theorem 2

Let μ satisfies (44). We put $\hat{\tau} = \mu^{-1} - \lambda_m^{-1}$. Then $\mu = \lambda_m - \lambda_m^2 \hat{\tau} + O(\sigma)$. Therefore, from (70) and (79) it follows that

$$\mu(Q_m B\varphi, B\psi) = \frac{1 - \lambda_m \hat{\tau}}{\lambda_m} (\Psi_\varphi, \Psi_\psi) + b_2(\varphi, \psi), \quad (86)$$

where

$$b_2(\varphi, \psi) = \mu b_0(\varphi, \psi) + \hat{\tau} \frac{\lambda_m - \mu}{\lambda_m} (\Psi_\varphi, \Psi_\psi)$$

and b_0 is given by (71). Using (79) and (44), we get

$$b_2(\varphi, \psi) \leq C(\wp(\varphi) + \wp(\psi)). \quad (87)$$

Using (67), (86) and (87) together with (85), we derive from (63)

$$\begin{aligned} \hat{\tau}((S\varphi, S\psi) + (\Psi_\varphi, \Psi_\psi)) &= \frac{1}{\lambda_m} \left((\Psi_\varphi, \Psi_\psi) - (T\varphi, T\psi) \right. \\ &\quad \left. - (\Psi_\varphi, \psi) - (\varphi, \Psi_\psi) \right) + b(\varphi, \psi) \quad \text{for all } \psi \in X_m, \end{aligned} \quad (88)$$

where

$$b(\varphi, \psi) = b_2(\varphi, \psi) + (L(\mu)B\varphi, B\psi) + \langle T_2\varphi, T_2\psi \rangle.$$

Due to (87) and (85), the form b is subject to

$$|b(\varphi, \psi)| \leq C(\wp(\varphi) + \wp(\psi)). \quad (89)$$

Moreover, by (73)

$$|(\Psi_\varphi, \Psi_\psi)| \leq c\sigma(\|\varphi\|^2 + \|\psi\|^2).$$

From (32) it follows that $\|\varphi\|^2 \leq (1 + c\sigma^{1/2})\|S\varphi\|^2$. Therefore

$$|(\Psi_\varphi, \Psi_\psi)| \leq c\sigma(\|S\varphi\|^2 + \|S\psi\|^2). \quad (90)$$

Let $\mu_1^{-1}, \dots, \mu_{J_m}^{-1}$ be the eigenvalues of the problem (20) and let U_1, \dots, U_{J_m} be corresponding eigenvectors. We assume that the eigenvectors are chosen to satisfy $(U_j, U_k) = \delta_{j,k}$, where $\delta_{j,k}$ is the Kronecker delta. Since P_m is the orthogonal projector with the image SX_m , we can represent $P_m U_j$ as $P_m U_j = SV_j$, where $V_j \in X_m$. According to (56),

$$(SV_j, SV_k) = \delta_{jk} + \mathcal{O}(\sigma) \quad \text{for } j, k = 1, \dots, J_m. \quad (91)$$

Moreover according to the reduction from Sect. 4.1, $\hat{\tau}_j = \mu_j^{-1} - \lambda_m^{-1}$, $j = 1, \dots, J_m$, is an eigenvalue to problem (88) and $\varphi = V_j$ is the corresponding eigenvector.

Denote by τ_k , $k = 1, \dots, J_m$, the eigenvalue of the finite-dimensional problem (26) and by $\Phi_1, \dots, \Phi_{J_m}$ corresponding eigenvectors from X_m , which satisfies the bi-orthogonality condition

$$(S\Phi_j, S\Phi_k) = \delta_{jk} \quad \text{for } j, k = 1, \dots, J_m. \quad (92)$$

Let us show that for each $j = 1, \dots, J_m$ there exists $k = k(j)$, $1 \leq k \leq \dots, J_m$ such that

$$(SV_j, S\Phi_{k(j)}) \geq c_*, \quad (93)$$

where c_* is a positive constant depending on J_m . Moreover the function $k(j)$ is isomorphism. In order to prove these facts we consider the matrix $A = \{A_{jk}\}$, where $A_{jk} = (SV_j, S\Phi_k)$, $j, k = 1, \dots, J_m$. Let us show that

$$\det A = 1 + \mathcal{O}(\sigma^{1/2}) \quad (94)$$

Indeed, let ν be an eigenvalue of the matrix A and $\mathbf{a} = (a_1, \dots, a_{J_m})$ be corresponding eigenvector with the norm 1. Then

$$\sum_k (SV_j, S\Phi_k) a_k = \nu a_j \quad \text{for } j = 1, \dots, J_m,$$

or equivalently $(SV_j, S\Phi) = \nu a_j$, where $\Phi = \bar{a}_1 \Phi_1 + \dots + \bar{a}_{J_m} \Phi_{J_m}$. Clearly, $\|S\Phi\| = 1$. We chose the constants b_j , $j = 1, \dots, J_m$, such that $\sum b_j V_j = \Phi$. Clearly, the norm of the vector $\mathbf{b} = (b_1, \dots, b_{J_m})$ is equal to $1 + \mathcal{O}(\sigma)$. Then

$$\sum_{j=1}^{J_m} b_j (SV_j, S\Phi) = (S\Phi, S\Phi) = \nu \sum_{j=1}^{J_m} a_j b_j.$$

This implies $1 \leq |\nu|(1 + \mathcal{O}(\sigma))$. Since the last relation is valid for all eigenvalues of A we obtain (94). Therefore, there exists an isomorphism $k(j)$ such that the equality (93) is valid for $j = 1, \dots, J_m$. After the re-numeration of eigenvalues τ_j and corresponding eigenvectors we can assume that the relations

$$(SV_j, S\Phi_j) \geq c_*, \quad j = 1, \dots, J_m, \quad (95)$$

hold.

Choosing $\varphi = V_j$ and $\psi = \Phi_j$ in (88) we obtain

$$(\widehat{\tau}_k - \tau_k)(SV_j, S\Phi_j) = b(V_j, \Phi_j) - \widehat{\tau}_j(\Psi_{V_j}, \Psi_{\Phi_j}). \quad (96)$$

Using relations (95), (89) and (73) together with definition (24) of ρ , we derive from (96)

$$|\widehat{\tau}_k - \tau_k| \leq C \left(\max_{\psi \in X_m, \|\psi\|=1} \wp(\psi) + |\widehat{\tau}_k| \sigma \right). \quad (97)$$

By (80), we get

$$|\widehat{\tau}_k - \tau_k| \leq C(\rho + |\widehat{\tau}_k|\sigma). \quad (98)$$

This implies

$$|\widehat{\tau}_k - \tau_k| \leq C(\rho + |\tau_k|\sigma) \quad (99)$$

and hence (25).

5 Application to a second order elliptic equation

Here we consider the spectral problems (4) and (5) generated by the bi-linear form (2). Instead of (21) it is sufficient to check inequality (27).

5.1 Local perturbation of the boundary

Constant σ_ .* Let ε be a small positive number. We assume that there exists a point x_0 in $\partial\Omega_1$ such that

$$\Omega_1 \setminus B_\varepsilon(x_0) \subset \Omega_2 \subset \Omega_1 \cup B_\varepsilon(x_0). \quad (100)$$

We denote the domain $\Omega_1 \cup B_\varepsilon(x_0)$ by $\Omega_\varepsilon^+(x_0)$. We also assume that for $u \in \mathring{W}^{1,2}(\Omega_\varepsilon^+(x_0))$

$$\int_{B_{q\varepsilon}(x_0) \cap \Omega_\varepsilon^+(x_0)} |u|^2 dx \leq C\varepsilon^2 \int_{B_{q\varepsilon}(x_0) \cap \Omega_\varepsilon^+(x_0)} |\nabla u|^2 dx \quad (101)$$

for a certain $q > 1$ independent of ε . Here the constant C may depend on Ω_1 , n , q and the ellipticity constant ν .

Let $\mathcal{Z}_\varepsilon(x_0)$ be subspace of function in $\mathring{W}^{1,2}(\Omega_\varepsilon^+)$ subject to

$$\sum_{i,j=1}^n \int A_{ij}(x) \partial_{x_j} u \partial_{x_i} w dx = 0 \quad \text{for all } w \in \mathring{W}^{1,2}(\Omega_1 \setminus B_\varepsilon(x_0)). \quad (102)$$

In what follows we will omit the summation sign in formulas like (102).

Lemma 1. (i) *Let $u \in \mathcal{Z}_{q^{1/2}\varepsilon}(x_0)$. Then*

$$\int_{\Omega_\varepsilon^+(x_0)} |\nabla u|^2 dx \leq C_1 \int_{B_{q\varepsilon}(x_0) \cap (\Omega_\varepsilon^+(x_0))} |\nabla u|^2 dx. \quad (103)$$

(ii) Let $q_* \in [1, q^{1/2}]$ and let $T_{q_*\varepsilon}(x_0)$ be orthogonal projector from $\mathring{W}^{1,2}(\Omega_\varepsilon^+(x_0))$ onto $\mathcal{Z}_{q_*\varepsilon}(x_0)$. Then

$$\int_{\Omega_\varepsilon^+(x_0)} |\nabla(T_{q_*\varepsilon}(x_0)u)|^2 dx \leq C_1 \int_{B_{q\varepsilon}(x_0) \cap (\Omega_\varepsilon^+(x_0))} |\nabla u|^2 dx. \quad (104)$$

The constant C_1 in (i) and (ii) may depend on n, ν, q and Ω_1 .

Proof. We introduce a smooth function $\eta = \eta(t)$ which is equal to 1 for $t < q^{1/2}$ and to 0 for $t > q$ and let $\eta_\varepsilon(x) = \eta(|x - x_0|/\varepsilon)$.

(i) Put $u_\varepsilon = u - \eta_\varepsilon u$. One can check that $u_\varepsilon \in \mathring{W}^{1,2}(\Omega_1 \setminus B_{q^{1/2}\varepsilon}(x_0))$. From (102) it follows that

$$\int A_{ij} \partial_{x_j} u_\varepsilon \partial_{x_j} w dx = - \int A_{ij} \partial_{x_j} (\eta_\varepsilon u) \partial_{x_i} w dx \quad \text{for all } w \in \mathring{W}^{1,2}(\Omega_1 \setminus B_{q^{1/2}\varepsilon}(x_0)). \quad (105)$$

Taking $w = u_\varepsilon$ in (105) we obtain

$$\int A_{ij} \partial_{x_j} u_\varepsilon \partial_{x_i} u_\varepsilon dx = - \int A_{ij} \partial_{x_j} (\eta_\varepsilon u) \partial_{x_i} u_\varepsilon dx, \quad (106)$$

which together with (3) implies

$$\int |\nabla u_\varepsilon|^2 dx \leq \nu^{-4} \int |\nabla(\eta_\varepsilon u)|^2 dx.$$

Using (101), we derive the following estimate from the last inequality:

$$\int_{\Omega_1} |\nabla u_\varepsilon|^2 dx \leq C_1 \int_{B_{q\varepsilon}(x_0) \cap (\Omega_\varepsilon^+(x_0))} |\nabla u|^2 dx.$$

Similar estimate for $\eta_\varepsilon u$ follows from (101), which together with the estimate for u_ε leads to (103).

(ii) Let $U_\varepsilon = T_{q_*\varepsilon}(x_0)u - \eta_\varepsilon u$. Then $U_\varepsilon(x) \in \mathring{W}^{1,2}(\Omega_1 \setminus B_{q_*\varepsilon}(x_0))$ and $U_\varepsilon + \eta_\varepsilon u \in \mathcal{Z}_{q_*\varepsilon}(x_0)$. Therefore,

$$\int A_{ij} \partial_{x_j} U_\varepsilon \partial_{x_j} w dx = - \int A_{ij} \partial_{x_j} (\eta_\varepsilon u) \partial_{x_j} w dx \quad \text{for all } w \in \mathring{W}^{1,2}(\Omega_1 \setminus B_{q_*\varepsilon}(x_0)).$$

Taking here $w = U_\varepsilon$ and using Hölder inequality along with inequalities (3), we obtain

$$\int |\nabla U_\varepsilon|^2 dx \leq \nu^{-4} \int |\nabla(\eta_\varepsilon u)|^2 dx \leq C \int_{B_{q\varepsilon}(x_0) \cap (\Omega_\varepsilon^+(x_0))} |\nabla u|^2 dx.$$

To get the last inequality we applied (101). The estimate of $\eta_\varepsilon u$ by the right-hand side of the last inequality follows from (101). Since $T_{q^*\varepsilon}(x_0)u = \eta_\varepsilon u + U_\varepsilon$, the above two estimates give (104). The proof is complete. \square

Now we are in position to prove the following

Proposition 4. *There exists a function $\sigma_* = \sigma_*(\varepsilon)$ such that $\sigma_*(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$ and for all $u \in \mathcal{Z}_\varepsilon(x_0)$*

$$\int_{\Omega_\varepsilon^+(x_0)} |u|^2 dx \leq \sigma \int_{\Omega_\varepsilon^+(x_0)} |\nabla u|^2 dx. \quad (107)$$

Proof. Let λ_k , $k = 1, \dots$, be eigenvalues of the problem (4) and let φ_k be corresponding eigenfunctions normalized by $\|\varphi\|_{W^{1,2}(\Omega_1)} = 1$. We represent u as $u = \eta_{q^{-1/2}\varepsilon} u + u_\varepsilon$. Then $u_\varepsilon \in \dot{W}^{1,2}(\Omega_1)$ and $T_{q^{1/2}\varepsilon} u_\varepsilon = u_\varepsilon$. Therefore we may represent u as

$$u_\varepsilon = \sum_{k=N+1}^{\infty} (u_\varepsilon, \varphi_k) \varphi_k + \sum_{k=1}^N (u_\varepsilon, T_{q^{1/2}\varepsilon} \varphi_k) \varphi_k.$$

Since $\{\varphi_k\}_{k \geq 1}$ is an orthogonal basis in $L^2(\Omega_1)$ also and $\|\varphi_k\|_{L^2(\Omega_1)}^2 = \lambda_k^{-1}$, we have

$$\begin{aligned} \|u_\varepsilon\|_{L^2(\Omega_1)}^2 &\leq \frac{1}{\lambda_{N+1}} \|u_\varepsilon\|_{W^{1,2}(\Omega_1)}^2 + \sum_{k=1}^N \frac{1}{\lambda_k} |(u_\varepsilon, T_{q^{1/2}\varepsilon} \varphi_k)|^2 \\ &\leq \left(\frac{1}{\lambda_{N+1}} + \sum_{k=1}^N \frac{1}{\lambda_k} \|T_{q^{1/2}\varepsilon} \varphi_k\|_{W^{1,2}(\Omega_1)}^2 \right) \|u_\varepsilon\|_{W^{1,2}(\Omega_1)}^2. \end{aligned}$$

Using (103) we get

$$\|u_\varepsilon\|_{L^2(\Omega_1)}^2 \leq \left(\frac{1}{\lambda_{N+1}} + C \sum_{k=1}^N \frac{1}{\lambda_k} \int_{\Omega_1 \setminus \Omega_{q\varepsilon}} |\nabla \varphi_k|^2 dx \right) \|u_\varepsilon\|_{W^{1,2}(\Omega_1)}^2$$

Let

$$\sigma_1 = \inf \left(\frac{1}{\lambda_{N+1}} + C \sum_{k=1}^N \frac{1}{\lambda_k} \int_{\Omega_1 \setminus \Omega_{q\varepsilon}} |\nabla \varphi_k|^2 dx \right).$$

then $\sigma_1(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$ and by the last inequality for u_ε combined with (101), we obtain

$$\|u_\varepsilon\|_{L^2(\Omega_1)}^2 \leq \sigma_1 \|\nabla(u - \eta_{q^{-1/2}\varepsilon}u)\|_{W^{1,2}(\Omega_\varepsilon^+)} \leq C_1(\sigma_1 + \varepsilon^2) \|\nabla u\|_{W^{1,2}(\Omega_\varepsilon^+)}^2.$$

Using (101) again, we get

$$\|\eta_{q^{-1/2}\varepsilon}u\|_{L^2(\Omega_\varepsilon^+(x_0))}^2 \leq C_2\varepsilon^2 \int_{\Omega_\varepsilon^+(x_0) \setminus B_{q\varepsilon}(x_0)} |\nabla u|^2 dx.$$

Now setting $\sigma = 2C_1(\sigma_1 + \varepsilon) + 2C_2\varepsilon^2$ and using the representation $u = \eta_{q^{-1/2}\varepsilon}u + u_\varepsilon$ we arrive at (107). \square

Remark 2. In the case $\Omega_2 \subset \Omega_1$ Lemma 1 and Proposition 4 are valid if $\Omega_\varepsilon(x_0)$ is replaced by Ω_1 in their formulations and in (101).

Remark 3. Let several points x_1, \dots, x_m be given on the boundary $\partial\Omega_1$. Let also ε be a small positive number, $q > 1$ and inequality (101) be valid for all points x_1, \dots, x_m . Then Lemma 1 and Proposition 4 remain true for perturbation Ω_2 subject to

$$\Omega_1 \setminus \bigcup_{j=1}^m B_\varepsilon(x_j) \subset \Omega_2 \subset \Omega_1 \cup \bigcup_{j=1}^m B_\varepsilon(x_j)$$

if we replace $\Omega_\varepsilon^+(x_0)$ by $\Omega_1 \cup \bigcup_{j=1}^m B_\varepsilon(x_j)$ and $B_{q\varepsilon}(x_0)$ by $\bigcup_{j=1}^m B_{q\varepsilon}(x_j)$ in their formulations.

Estimates of the function $u = T\varphi$, $\varphi \in X_m$. Since $TT_\varepsilon = T_\varepsilon T = T$, we have that

$$\int A_{ij} \partial_{x_j}(T\varphi) \partial_{x_i}(T\varphi) dx \leq \int A_{ij} \partial_{x_j}(T_\varepsilon\varphi) \partial_{x_i}(T_\varepsilon\varphi) dx. \quad (108)$$

This together with Lemma 1(ii) gives

$$\|\nabla(T\varphi)\|_{L^2(\Omega_2)} \leq c \|\nabla(T\varphi)\|_{L^2(B_{q\varepsilon}(x_0) \cap \Omega_\varepsilon^+)}. \quad (109)$$

Estimate of the function $\Psi = \Psi_\varphi$. We seek Ψ in the form $\Psi = \eta_\varepsilon(x) + v$, where $\eta = \eta(t)$ is a smooth function equals 1 for $t < q_1 = (1+q)/2$ and 0 for $t > q_2 = (1+q_1)/2$. Then the function v belongs to $W^{1,2}(\Omega_2)$ and satisfies

$$(v, w) = -\lambda_m \langle \varphi, \eta_\varepsilon w \rangle + \int_{\Omega_2} A_{ij} \partial_{x_j} v w \partial_{x_i} \eta_\varepsilon dx - \int_{\Omega_2} v A_{ij} \partial_{x_j} \eta_\varepsilon \partial_{x_i} w dx. \quad (110)$$

Applying Hölder's inequality to the left-hand side of (110) and using then (101) we arrive at

$$\|\nabla v\|_{L^2(\Omega_2)} \leq c \|\nabla \varphi\|_{L^2(\Omega_1 \cap B_{q\delta}(x_2))}. \quad (111)$$

Applying (101), we get an estimate of η_ε by the right-hand side of (111). Combining these two estimates, we obtain

$$\|\nabla \Psi\|_{L^2(\Omega_2)} \leq c \|\nabla \varphi\|_{L^2(\Omega_1 \cap B_{q\varepsilon}(x_0))}. \quad (112)$$

Using estimates (109), (112) and Corollary 3, we arrive at

Corollary 5. *Under the assumptions on Ω_1 and Ω_2 of this section, the following estimate for the eigenvalues of the problems (4) and (5) holds*

$$|\lambda_m^{-1} - \mu_k^{-1}| \leq C \max_{\varphi \in X_m, \|\varphi\|=1} \|\nabla \varphi\|_{L^2(\Omega_1 \cap B_{q\varepsilon}(x_0))}^2,$$

where μ_1, \dots, μ_{J_m} are eigenvalues of (5) located near λ_m^{-1} , see Proposition 3.

5.2 Global perturbation of the boundary

Here we consider perturbations of Ω_1 located near the boundary. Let ε be a small positive number. We introduce the sets

$$\Omega_\varepsilon = \{x \in \Omega_1 : \text{dist}(x, \partial\Omega_1) > \varepsilon\} \text{ and } \Omega_\varepsilon^+ = \{y \in \mathbb{R}^n : \text{dist}(y, \Omega_1) < \varepsilon\}.$$

We assume that

$$\Omega_\varepsilon \subset \Omega_2 \subset \Omega_\varepsilon^+. \quad (113)$$

and that for all $x_0 \in \partial\Omega_1$ and $u \in \mathring{W}^{1,2}(\Omega_\varepsilon^+)$ the inequality

$$\int_{B_{q\varepsilon}(x_0) \cap \Omega_\varepsilon^+} |u|^2 dx \leq C\varepsilon^2 \int_{B_{q\varepsilon}(x_0) \cap \Omega_\varepsilon^+} |\nabla u|^2 dx \quad (114)$$

holds with a certain $q > 1$ independent of ε .

Let \mathcal{Z}_δ be subspace of function in $\mathring{W}^{1,2}(\Omega_\varepsilon^+)$ subject to

$$\sum_{i,j=1}^n \int A_{ij}(x) \partial_{x_j} u \partial_{x_i} w dx = 0 \text{ for all } w \in \mathring{W}^{1,2}(\Omega_\varepsilon^+ \setminus \Omega_\delta). \quad (115)$$

Lemma 2. (i) Let $q_1 \in (1, q)$ and $u \in \mathring{W}^{1,2}(\Omega_\varepsilon^+)$. Then

$$\int_{\Omega_\varepsilon^+ \setminus \Omega_{q_1\varepsilon}} |\nabla u|^2 dx \leq C_1 \varepsilon^2 \int_{\Omega_\varepsilon^+ \setminus \Omega_{q\varepsilon}} |\nabla u|^2 dx. \quad (116)$$

(ii) Let $u \in \mathcal{Z}_{q_0\varepsilon}$ with $q_0 \in (1, q)$. Then

$$\int_{\Omega_\varepsilon^+} |\nabla u|^2 dx \leq C_1 \int_{\Omega_\varepsilon^+ \setminus \Omega_{q\varepsilon}} |\nabla u|^2 dx. \quad (117)$$

(iii) Let $q_* \in [1, q)$ and let $T_{q_*\varepsilon}$ be orthogonal projector from $\mathring{W}^{1,2}(\Omega_\varepsilon^+)$ onto $\mathcal{Z}_{q_*\varepsilon}$. Then

$$\int_{\Omega_\varepsilon^+} |\nabla(T_{q_*\varepsilon} u)|^2 dx \leq C_1 \int_{\Omega_\varepsilon^+ \setminus \Omega_{q\varepsilon}} |\nabla u|^2 dx, \quad (118)$$

The constant C_1 in (i) –(ii) may depend on n, ν, q, q_0, q_1, q_* and Ω_1 .

Proof. (i) Let us construct a set of points on $\partial\Omega_1$ satisfying the properties a) and b) below. We put $\alpha = \sqrt{q^2 - q_1^2}$ and choose points x_1, \dots, x_N in the following way. We take an arbitrary point $x_1 \in \partial\Omega_1$. Let the points x_1, \dots, x_m have been chosen. If there is a point on the boundary, say x_* , such that $|x - x_*| > \alpha\varepsilon$ then we put $x_{m+1} = x_*$. If there are no such point the the required set is constructed and we take $N = m$. The above procedure leads to a finite set of points with the following properties:

- a). $\Omega_\varepsilon^+ \setminus \Omega_{q_1\varepsilon} \subset \bigcup_{k=1}^N B_{q\varepsilon}(x_k)$,
- b). There is a integer M , depending only on n and q_1, q , such that every $x \in \mathbb{R}^n$ may belong at most to M balls $B_{q\varepsilon}(x_k)$, $k = 1, \dots, N$.

Using a), (101) and then b), we get

$$\begin{aligned} \int_{\Omega_\varepsilon^+ \setminus \Omega_{q_1\varepsilon}} |u|^2 dx &\leq \int_{\bigcup_{k=1}^N B_{q\varepsilon}(x_k)} |u|^2 dx \\ &\leq C\varepsilon^2 \int_{\bigcup_{k=1}^N B_{q\varepsilon}(x_k)} |\nabla u|^2 dx \leq CM \int_{\Omega_\varepsilon^+ \setminus \Omega_{q\varepsilon}} |\nabla u|^2 dx, \end{aligned}$$

which leads to (116).

(ii). Let $\eta = \eta(t)$ be a smooth function which is equal to 1 for $t < q_0$ and 0 for $t > q_1 = (q_0 + q)/2$. It is clear that $1 < q_0 < q_1 < q$. Let

also $\zeta_\varepsilon(x) = \eta(d(x)/\varepsilon)$, where $d(x) = \max_k |x - x_k|$. We represent u as $u = \zeta_\varepsilon u + u_\varepsilon$. Then u_ε belongs to $\dot{W}^{1,2}(\Omega_\varepsilon^+ \setminus \Omega_{q_1\varepsilon})$ and satisfies

$$\sum_{i,j=1}^n \int A_{ij}(x) \partial_{x_j}(u_\varepsilon + \zeta_\varepsilon u) \partial_{x_i} w dx = 0 \quad \text{for all } w \in \dot{W}^{1,2}(\Omega_\varepsilon^+ \setminus \Omega_{q_0\varepsilon}).$$

Taking here $w = u_\varepsilon$, moving term with u to the right-hand side, using then Hölder inequality and (3), we obtain

$$\int |u_\varepsilon|^2 dx \leq \nu^{-4} \int |\nabla(\zeta_\varepsilon u)|^2 dx$$

which implies, due to (116),

$$\int_{\Omega_\varepsilon^+} |\nabla u_\varepsilon|^2 dx \leq C_1 \int_{\Omega_\varepsilon^+ \setminus \Omega_{q\varepsilon}} |\nabla u|^2 dx. \quad (119)$$

The estimate of $\zeta_\varepsilon u$ by the left-hand side of (119) follows from (116), which together with (119) gives (117).

(iii) Let $\eta = \eta(t)$ be a smooth function which is equal to 1 for $t < q_0 = (q_* + q)/2$ and 0 for $t > q_1 = (q_0 + q)/2$. One can check that $1 < q_0 < q_1 < q$. Let also $\zeta_\varepsilon(x) = \eta(d(x)/\varepsilon)$. We represent $T_{q_*\varepsilon}u$ as $T_{q_*\varepsilon}u = \eta_\varepsilon u + u_\varepsilon$. Since $T_{q_*\varepsilon}u(x) = u(x)$ for $x \in \Omega_\varepsilon^+ \setminus \Omega_{q_*\varepsilon}$ and

$$(T_{q_*\varepsilon}u, w) = 0 \quad \text{for all } w \in \dot{W}^{1,2}(\Omega_{q_*\varepsilon}),$$

we have that $u_\varepsilon \in \dot{W}^{1,2}(\Omega_{q_*\varepsilon})$ and

$$(u_\varepsilon, w) = -(\zeta_\varepsilon u, w) \quad \text{for all } w \in \dot{W}^{1,2}(\Omega_\varepsilon).$$

We choose here $w = u_\varepsilon$ and obtain

$$||u_\varepsilon||^2 = -(\zeta_\varepsilon u, u_\varepsilon).$$

This implies

$$||u_\varepsilon||_1^2 \leq ||\zeta_\varepsilon u||_1^2,$$

which leads to

$$\int_{\Omega_\varepsilon^+} |\nabla u_\varepsilon|^2 dx \leq C_1 \int_{\Omega_\varepsilon^+ \setminus \Omega_{q\varepsilon}} |\nabla u|^2 dx,$$

where we used (114). Similar estimate of $\eta_\varepsilon u$ by the right-hand side of the last inequality follows from (114). These two estimates give (118). \square

Proposition 5. *There exists a function $\sigma = \sigma(\varepsilon)$ such that $\sigma(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$ and for all $u \in \mathring{W}^{1,2}(\Omega_\varepsilon^+)$ satisfying (115)*

$$\int_{\Omega_\varepsilon^+} |u|^2 dx \leq \sigma \int_{\Omega_\varepsilon^+} |\nabla u|^2. \quad (120)$$

Proof. Let $\lambda_k, k = 1, \dots$, be eigenvalues of the problem (4) and let φ_k be corresponding eigenfunctions normalized by $\|\varphi\|_1 = 1$. We introduce a smooth $\eta = \eta(t)$ which is equal to 1 for $t < q_0 = (1+q)/2$ and 0 for $t > q_1 = (q_0+q)/2$. One can check that $1 < q_0 < q_1 < q$. Let also $\zeta_\varepsilon(x) = \eta(d(x)/\varepsilon)$. We represent u as $u = \eta_\varepsilon u + u_\varepsilon$. Then $u_\varepsilon(x) \in \mathring{W}^{1,2}(\Omega_{q_0\varepsilon})$ and $T_{q_1\varepsilon} u_\varepsilon = u_\varepsilon$. Therefore, we may represent u_ε as

$$u_\varepsilon = \sum_{k=N+1}^{\infty} (u_\varepsilon, \varphi_k) \varphi_k + \sum_{k=1}^N (u_\varepsilon, T_{q_1\varepsilon} \varphi_k) \varphi_k.$$

Since $\{\varphi_k\}_{k \geq 1}$ is an orthogonal basis in $L^2(\Omega_1)$ also and $\|\varphi_k\|_{L^2(\Omega_1)}^2 = \lambda_k^{-1}$, we have

$$\begin{aligned} \|u_\varepsilon\|_{L^2(\Omega_1)}^2 &\leq \frac{1}{\lambda_{N+1}} \|u_\varepsilon\|_{W^{1,2}(\Omega_1)}^2 + \sum_{k=1}^N \frac{1}{\lambda_k} |(u_\varepsilon, T_{q_1\varepsilon} \varphi_k)|^2 \\ &\leq \left(\frac{1}{\lambda_{N+1}} + \sum_{k=1}^N \frac{1}{\lambda_k} \|T_{q_1\varepsilon} \varphi_k\|_{W^{1,2}(\Omega_1)}^2 \right) \|u_\varepsilon\|_{W^{1,2}(\Omega_1)}^2. \end{aligned}$$

Using Lemma 2(ii) we get

$$\|u_\varepsilon\|_{L^2(\Omega_1)}^2 \leq \left(\frac{1}{\lambda_{N+1}} + C \sum_{k=1}^N \frac{1}{\lambda_k} \int_{\Omega_1 \setminus \Omega_{q\varepsilon}} |\nabla \varphi_k|^2 dx \right) \|u_\varepsilon\|_{W^{1,2}(\Omega_1)}^2$$

Let

$$\sigma_1 = 2 \inf \left(\frac{1}{\lambda_{N+1}} + C \sum_{k=1}^N \frac{1}{\lambda_k} \int_{\Omega_1 \setminus \Omega_{q\varepsilon}} |\nabla \varphi_k|^2 dx \right).$$

then $\sigma_1(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$ and by the last inequality for u_ε , we obtain

$$\|u_\varepsilon\|_{L^2(\Omega_1)}^2 \leq \sigma_1 \|\nabla u_\varepsilon\|_{W^{1,2}(\Omega_\varepsilon^+)}.$$

Since $u_\varepsilon = (1 - \eta_\varepsilon)u$, we get

$$\|u_\varepsilon\|_{L^2(\Omega_\varepsilon^+)}^2 \leq C \|\nabla u\|_{W^{1,2}(\Omega_\varepsilon^+)}.$$

Therefore

$$\|u_\varepsilon\|_{L^2(\Omega_1)}^2 \leq C\sigma_1 \|\nabla u_\varepsilon\|_{W^{1,2}(\Omega_\varepsilon^+)}. \quad (119)$$

From Lemma 2(i) it follows that

$$\|\eta_\varepsilon u\|_{L^2(\Omega_\varepsilon^+)}^2 \leq C_1 \varepsilon^2 \int_{\Omega_\varepsilon \setminus \Omega_{q\varepsilon}^+} |\nabla u|^2 dx.$$

Now setting $\sigma = C\sigma_1 + C_1\varepsilon^2$ and using the representation $u = \eta_\varepsilon u + u_\varepsilon$ we arrive at (120). \square

Estimates for $T\varphi$. Since $TT_\varepsilon = T_\varepsilon T = T$, we conclude that T and T_ε satisfy (108). Now, Lemma 2 and (108) lead to the estimate

$$\|T\varphi\|_{H^1(\Omega_2)} \leq C \|\nabla \varphi\|_{L^2(\Omega_1 \setminus \Omega_{q\varepsilon})}. \quad (121)$$

Estimate for Ψ_φ , $\varphi \in X_m$. Let $d = d(x)$ and $\eta_\varepsilon(x) = \eta(d(x)/\varepsilon)$ be the same as before. We are looking for the solution $\Psi = \Psi_\varphi$ to equation (7) in the form

$$\Psi = \eta_\varepsilon(x)\varphi(x) + v(x). \quad (122)$$

Then $v \in \dot{W}^{1,2}(\Omega_2)$ and satisfies

$$(v, w) = -\lambda \langle \eta_\varepsilon \varphi, w \rangle + \int_{\Omega_2} (A_{ij} \partial_{x_j} \varphi w \partial_{x_i} \eta_\varepsilon) dx - \int_{\Omega_2} \varphi A_{ij} \partial_{x_j} \eta_\varepsilon \partial_{x_i} w dx.$$

Choosing $w = v$ and using Hölder's inequality together with Lemma 2(i), we obtain

$$\|\nabla v\|_{L^2(\Omega_2)} \leq C \|\nabla \varphi\|_{L^2(\Omega_1 \setminus \Omega_{q\varepsilon})}$$

Taking into account (122), we get

$$\|\Psi\|_{L^2(\Omega_2)} \leq C \|\nabla \varphi\|_{L^2(\Omega_1 \setminus \Omega_{q\varepsilon})}. \quad (123)$$

Using (121), (123) and Corollary 3, we arrive at

Corollary 6. *Under the assumptions on Ω_1 and Ω_2 of this section, the following estimate for the eigenvalues of the problems (4) and (5) holds*

$$|\lambda_m^{-1} - \mu_k^{-1}| \leq C \max_{\varphi \in X_m, \|\varphi\|=1} \|\nabla \varphi\|_{L^2(\Omega_1 \setminus \Omega_{q\varepsilon})}^2,$$

where μ_1, \dots, μ_{J_m} are eigenvalues of (5) located near λ_m^{-1} , see Proposition 3.

5.3 The case of Lipschitz domains

We recall the domain Ω is called Lipschitz with Lipschitz constant less than or equal to C_* if the boundary $\partial\Omega$ can be covered by a finite number of balls B such that in an appropriate orthogonal system of coordinate $B \cap \Omega = B \cap \{y = (y', y_n) : y_n > h(y')\}$, where h satisfies $|h(y') - h(z')| \leq C_*|y' - z'|$ and $h(0) = 0$ and B has the center at the origin.

We assume in this section that Ω_1 is a bounded Lipschitz domain and we denote by B_k , $k = 1, \dots, M$, the balls from the covering of the boundary and by h_k corresponding Lipschitz functions. Then there exists a positive δ such that the set $\mathcal{V}_\delta = \{x \in \mathbb{R}^n : \text{dist}(x, \partial\Omega) \leq \delta\}$ contains in $\bigcup_{k=1}^M B_k$.

Concerning the domain Ω_2 we assume that

$$\Omega_\varepsilon \subset \Omega_2 \subset \Omega_\varepsilon^+,$$

where ε is a sufficiently small positive number. Moreover, we suppose that the domain Ω_2 is Lipschitz and

$$B_k \cap \Omega_2 = B_k \cap \{y = (y', y_n) : y_n > g_k(y')\} \text{ for } k = 1, \dots, M,$$

where g_k are Lipschitz functions with the Lipschitz constant $\leq C_*$.

Proposition 6. *The following inequality*

$$\|T_0\varphi\|^2 \leq C \int_{\Omega_1 \setminus \Omega_2} |\nabla\varphi|^2 dx \quad (124)$$

holds for all $\varphi \in X_m$. Here $T_0 = I - S_0$ and S_0 is the orthogonal projector onto $\mathring{W}^{1,2}(\Omega_1 \cap \Omega_2)$, compare with Corollary 3.

Proof. Since $\varphi \in \mathring{W}^{1,2}(\mathbb{R}^n \setminus (\Omega_1 \cap \Omega_2))$ the trace of this function on $\partial(\Omega_1 \cap \Omega_2)$ belongs to $W^{1/2,2}(\partial(\Omega_1 \cap \Omega_2))$ and

$$\|\varphi\|_{W^{1/2,2}(\partial(\Omega_1 \cap \Omega_2))} \leq C \|\varphi\|_{W^{1,2}(\Omega_1 \setminus \Omega_2)}.$$

Since $T_0\varphi$ is harmonic in $\Omega_1 \cap \Omega_2$ with the Dirichlet boundary condition $u = \varphi$ on $\partial\Omega_2$ we have that

$$\|\nabla T\varphi\|_{L^2(\Omega_2)} \leq c \|\varphi\|_{W^{1/2,2}(\partial\Omega_2)} \leq C_1 \|\nabla\varphi\|_{L^2(\Omega_1 \setminus \Omega_2)}.$$

This estimate together with $T\varphi = \varphi$ outside Ω_2 implies (124). \square

The estimate obtained in the last proposition together with Corollary 2 and $T_0\varphi = \varphi$ outside $\Omega_1 \cap \Omega_2$, implies the following

Corollary 7. *Let $\Omega_2 \subset \Omega_1$. Then for $j = 1, \dots, J_m$,*

$$c \min \|\nabla \varphi\|_{L^2(\Omega_1 \setminus \Omega_2)}^2 \leq \left| \mu_{m_j}^{-1} - \lambda_m^{-1} \right| \leq C \max \|\nabla \varphi\|_{L^2(\Omega_1 \setminus \Omega_2)}^2,$$

where minimum and maximum are taken over all $\varphi \in X_m$ with $\|\varphi\| = 1$.

Let us turn to estimating of the function $\Psi = \Psi_\varphi$. We have the following

Proposition 7. *There exists a domain S such that $\Omega_2 \setminus \Omega_1 \subset S \subset \Omega_2$, $|S| \leq C|(\Omega_2 \setminus \Omega_1)|$ and the following inequality*

$$\|\nabla \Psi_\varphi\|_{L^2(\Omega_2)}^2 \leq C_1 \max \int_S |\nabla \varphi|^2 dx \quad (125)$$

holds, where max is taken over all $\varphi \in X_m$ satisfying $\|\varphi\| = 1$. Here by $|S|$ is denote the area of S and the constant C is independent of ε .

Proof. There exists a set of smooth functions $\psi_k \in C^1(B_k)$ with compact support such that $\psi_1(x) + \dots + \psi_N(x) = 1$ on \mathcal{V}_δ .

We take a smooth function $\eta = \eta(t)$ which is equal to 1 for $t < 3/2$ and to 0 for $t > 2$ and introduce the function $\zeta_k(x)$ in B_k by

$$\zeta_k(x) = \eta\left(\frac{y_n - g_k(y')}{h_k(y') - g_k(y')}\right)$$

if $h_k(y') > g_k(y')$ and $\zeta_k(x) = 0$ otherwise. Here the local variable y is considered as a function of x . We define also

$$\zeta(x) = \sum_{k=1}^N \zeta_k(x) \psi_k(x).$$

One can verify that $\zeta(x) = 1$ if $x \in \Omega_2$ and $y_n - g_k(y') < 3(h_k(y') - g_k(y'))/2$. Moreover, the following inequality holds:

$$\int |(\nabla \zeta)u(x)|^2 dx \leq C \int_{\text{supp} \zeta} |\nabla u|^2 dx \quad (126)$$

for $u \in \mathring{W}^{1,2}(\Omega_2)$. It is sufficient to prove a local version of this inequality, i.e.

$$\int_{B_k} |\nabla_y \zeta_k u|^2 dy \leq c \int_{B_k, 0 < y_n - g_k(y') < 2|h_k(y') - g_k(y')|} |\nabla u|^2 dy \quad (127)$$

for smooth functions u equals zero for $y_n < g_k(y')$. Estimate (127) follows from one dimensional Hardy inequality.

Now, we are looking for the function Ψ in the form $\Psi = \zeta\varphi + v$. Then v satisfies the equation

$$(v, w) = -\langle \varphi A_{ij} \partial_{x_j} \zeta, \partial_{x_i} w \rangle + \langle A_{ij} \partial_{x_j} \zeta \partial_{x_i} \varphi, w \rangle - \lambda_m \langle \zeta \varphi, w \rangle.$$

Putting here $w = v$ and using (126), we get

$$(v, v) \leq C \int_{\text{supp} \zeta} |\nabla \varphi|^2 dx,$$

which implies (125). □

From Propositions 6, 7 and Corollary 3 it follows

Corollary 8. *There exists a domain S such that $\Omega_2 \setminus \Omega_1 \subset S \subset \Omega_2$, $|S| \leq C|\Omega_2 \setminus \Omega_1|$ and the following inequality*

$$|\mu_k^{-1} - \lambda_m^{-1}| \leq C_1 \max \int_{(\Omega_1 \setminus \Omega_2) \cup S} |\nabla \varphi|^2 dx$$

holds, where max is taken over all $\varphi \in X_m$ satisfying $\|\varphi\| = 1$.

References

- [1] Bucur, Dorin; Buttazzo, Giuseppe, Variational methods in shape optimization problems. Progress in Nonlinear Differential Equations and their Applications, 65. Birkhäuser Boston, Inc., Boston, MA, 2005.
- [2] Burenkov, Victor I.; Lamberti, Pier Domenico Spectral stability of Dirichlet second order uniformly elliptic operators. J. Differential Equations **244** (2008), no. 7, 1712–1740.
- [3] Delfour, M. C.; Zolésio, J.-P. Shapes and geometries. Analysis, differential calculus, and optimization. Advances in Design and Control, 4. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2001.

- [4] P.R. Garabedian, M. Schiffer, Convexity of domain functionals, *J. Anal. Math.* **2** (1953), 281-368.
- [5] P. Grinfeld, G. Strang, Laplace eigenvalues on polygons, *Comput. Math. Appl.* **48** (2004), 1121-1133.
- [6] Grinfeld, P. Hadamard's formula inside and out. *J. Optim. Theory Appl.* **146** (2010), no. 3, 654-690.
- [7] Grinfeld, Pavel; Strang, Gilbert Laplace eigenvalues on regular polygons: a series in $1/N$. *J. Math. Anal. Appl.* **385** (2012), no. 1, 135-149,
- [8] Hadamard, J., Mémoire sur le problème d'analyse relatif à l'équilibre des plaques élastiques encastrées, *Mémoires des Savants Étrangers* **33** (1908).
- [9] Hadamard, J., Mémoire sur le problème d'analyse relatif à l'équilibre des plaques élastiques encastrées, *Œuvres*, **2** (1968), 515-631.
- [10] Henrot, Antoine, Extremum problems for eigenvalues of elliptic operators. *Frontiers in Mathematics*. Birkhäuser Verlag, Basel, 2006.
- [11] Kenig, Carlos E., Harmonic analysis techniques for second order elliptic boundary value problems. *CBMS-AMS Regional Conference Series in Mathematics*, No. 83, 1994.
- [12] Kozlov, Vladimir, On the Hadamard formula for nonsmooth domains. *J. Differential Equations* **230** (2006), no. 2, 532–555.
- [13] Kozlov, Vladimir, L^q -perturbations of leading coefficients of elliptic operators: asymptotics of eigenvalues. *Abstr. Appl. Anal.* 2006, Art. ID 26845, 15 pp.
- [14] Kozlov, V. and Nazarov, S., On the Hadamard formula for second order systems in non-smooth domains, to appear in *CPDE*.
- [15] Kawohl, B., Some nonconvex shape optimization problems. *Optimal shape design (Tria, 1998)*, 7–46, *Lecture Notes in Math.*, 1740, Springer, Berlin, 2000.
- [16] J.W.S. Rayleigh, *The Theory of Sound* vol. I, Macmillan and Co, London, 1894.

- [17] M. van den Berg, On Rayleigh's formula for the first Dirichlet eigenvalue of a radial perturbation of a ball. To appear in Journal of Geometric Analysis.