

ON CONSTRAINED MARKOV-NIKOLSKII TYPE INEQUALITY FOR k -ABSOLUTELY MONOTONE POLYNOMIALS

OLEKSIY KLURMAN

ABSTRACT. We consider a classical problem of estimating norms of higher order derivatives of an algebraic polynomial via the norms of the polynomial itself. The corresponding extremal problem for general polynomials in the uniform norm was solved by V. A. Markov. In 1926, Bernstein found the exact constant in the Markov inequality for monotone polynomials. It was shown in [3] that the order of the constants in constrained Markov-Nikolskii inequality for k -absolutely monotone polynomials is the same as in the classical one in case $0 < p \leq q \leq \infty$. In this paper, we find the exact order for all values of $0 < p, q \leq \infty$. It turns out that for the case $q < p$ constrained Markov-Nikolskii inequality is significantly better than the unconstrained one.

1. INTRODUCTION

For $n \geq m \geq 0$, we denote

$$M_{q,p}(n, m) := \sup_{P_n \in \mathbb{P}_n} \frac{\|P_n^{(m)}\|_{L_q[-1,1]}}{\|P_n\|_{L_p[-1,1]}}.$$

In [4], complete information about the orders of $M_{q,p}(n, m)$ for all values $p, q > 0$ is given.

Theorem 1.1. *For $0 < p, q \leq \infty$ we have:*

$$(1) \quad M_{q,p}(n, m) \asymp \begin{cases} n^{2m+2/p-2/q}, & \text{if } m > 2/q - 2/p, \\ n^m (\log n)^{1/q-1/p}, & \text{if } m = 2/q - 2/p, \\ n^m, & \text{if } m < 2/q - 2/p. \end{cases}$$

The asymptotic is taken when m is fixed, so the constants may depend on (m, p, q) .

For each $f \in C[-1, 1]$ we denote

$$\|f\|_{C[-1,1]} := \|f\|.$$

By Δ_n we denote the set of all monotone polynomials of degree n on $[-1, 1]$. In 1926, S. Bernstein [1] pointed out that Markov's inequality for monotone polynomials is not essentially better than for all polynomials, in the sense, that the order of $\sup_{P_n \in \Delta_n} \|P'_n\|/\|P_n\|$ is n^2 . He proved his result only for odd n . In 2001, Qazi [6]

2000 *Mathematics Subject Classification.* 41A17.

Key words and phrases. Markov inequality, Nikolskii inequality, k -monotone polynomial.

extended Bernstein's idea to include polynomials of even degree. Next theorem contains their results:

Theorem 1.2 (Bernstein [1], Qazi [6]).

$$\sup_{P_n \in \Delta_n} \frac{\|P'_n\|}{\|P_n\|} = \begin{cases} \frac{(n+1)^2}{4}, & \text{if } n = 2k + 1, \\ \frac{n(n+2)}{4}, & \text{if } n = 2k. \end{cases}$$

A natural generalization of the concept of monotonicity is k -absolute monotonicity.

Definition 1.3. The function $f : [a, b] \rightarrow \mathbb{R}$ is *absolutely monotone of order k* if, for all $x \in [a, b]$,

$$f^{(m)}(x) \geq 0,$$

for all $0 \leq m \leq k$, and denote by $\Delta_n^{(k)}$ the set of all absolutely monotone polynomials of order k on $[-1, 1]$.

For example, absolutely monotone functions of order zero are just nonnegative functions on $[a, b]$, and $\Delta_n^{(1)} = \Delta_n \cap \Delta_n^{(0)}$ is the set of all nonnegative monotone polynomials on $[-1, 1]$.

A natural modification of $M_{q,p}(n, k)$ for $\Delta_n^{(k)}$ is

$$M_{q,p}^{(k)}(n, m) = \sup_{P_n \in \Delta_n^{(k)}} \frac{\|P_n^{(m)}\|_{L_q[-1,1]}}{\|P_n\|_{L_p[-1,1]}},$$

for $0 \leq m \leq n$, $0 \leq k \leq n$.

In 2009, A. Kroó and J. Szabados [5] found the exact constants for Markov-Nikolskii inequalities in L_1 and L_∞ . Note, that J. Szabados and A. Kroó referred to absolutely monotone polynomials of order k as “ k -monotone polynomials.”

The next theorem contains their results:

Theorem 1.4 (Kroó and Szabados [5], 2009). *For $2 \leq k \leq n$, $m = \lfloor \frac{n-k}{2} \rfloor + 1$, $\beta = \frac{1-(-1)^{n-k}}{2}$:*

$$M_{\infty,\infty}^{(k)}(n, 1) = \frac{k-1}{1-x_{1,m}^{(k-2,\beta)}},$$

$$M_{1,1}^{(k)}(n, 1) = M_{\infty,\infty}^{(k+1)}(n+1, 1),$$

where $x_{1,m}^{(k-2,\beta)}$ is the largest zero of the Jacobi polynomial $J_m^{(k-2,\beta)}$, associated with the weight $(1-x)^{k-2}(1+x)^\beta$.

T. Erdélyi [3] found the order of $M_{q,p}^{(k)}(n, m)$ in the case $q \geq p$. He was interested in how this order depends on k .

Theorem 1.5 (Erdélyi [3], 2009). *For $0 \leq m \leq k/2$, $1 \leq k \leq n$, $0 < p \leq q \leq \infty$, we have*

$$M_{q,p}^{(k)}(n, m) \asymp (n^2/k)^{m+1/p-1/q} \asymp M_{q,p}(n, m).$$

First asymptotic is taken when both $n, k \rightarrow \infty$, so the constants depend on (p, q) only. Second asymptotic is taken when k is fixed.

It follows from Theorem 1.5 that whenever $q \geq p$ the order of constants in constrained Markov-Nikolskii inequality remains the same as in the classical case. In this paper, we find exact order for all values of $0 < p, q \leq \infty$. In particular, the results imply that the order can be significantly improved when $q < p$. Our main result is:

Theorem 1.6. *For $0 < p, q \leq \infty$ and $p \neq \infty$, $0 < m \leq k \leq n$,*

$$M_{q,p}^{(k)}(n, m) \asymp \begin{cases} n^{2m+2/p-2/q}, & \text{if } m > 1/q - 1/p, \\ \log^m n, & \text{if } m = 1/q - 1/p, \\ 1, & \text{if } m < 1/q - 1/p. \end{cases}$$

If $p = \infty$, $0 < q \leq \infty$ and $0 < m \leq k \leq n$, then

$$M_{q,\infty}^{(k)}(n, m) \asymp \begin{cases} n^{2m-2/q}, & \text{if } m > \frac{1}{q}, \\ \log^{m-1} n, & \text{if } m = \frac{1}{q}, \\ 1, & \text{if } m < \frac{1}{q}. \end{cases}$$

The asymptotic is taken when k is fixed and so the constants may depend on (p, q, k) .

2. PROOF OF THE MAIN RESULT

Proof of the upper bound in Theorem 1.6. We are going to show that for $0 < p, q \leq \infty$, $0 < m \leq k \leq n$ and $p \neq \infty$,

$$(2) \quad M_{q,p}^{(k)}(n, m) \leq C(k, p, q) \begin{cases} n^{2m+2/p-2/q}, & \text{if } m > 1/q - 1/p, \\ \log^m n, & \text{if } m = 1/q - 1/p, \\ 1, & \text{if } m < 1/q - 1/p. \end{cases}$$

Consider the case $k = 1$. We distinguish between two cases.

Case 1. $q \geq 1$. Clearly, $\frac{1}{q} - \frac{1}{p} < 1$. Without loss of generality, we can assume that $P_n(-1) = 0$. Note, that for each $P_n \in \Delta_n^{(1)}$ we have $\|P'_n\|_{L_1[-1,1]} = \|P_n\|$. By Nikolskii inequality

$$\|P'_n\|_{L_q[-1,1]} \leq C_1(q) n^{2-\frac{2}{q}} \|P'_n\|_{L_1[-1,1]},$$

and

$$\|P'_n\|_{L_1[-1,1]} \leq \|P_n\| \leq C_1(p) n^{\frac{2}{p}} \|P_n\|_{L_p[-1,1]},$$

so

$$\|P'_n\|_{L_q[-1,1]} \leq C_2(q, p) n^{2-\frac{2}{q}+\frac{2}{p}} \|P_n\|_{L_p[-1,1]}.$$

Case 2. Let $q < 1$. We first prove, that for all $P_n \in \Delta_n^1$, $P_n(-1) = 0$ the following inequality holds:

$$\int_{-1}^1 P_n'^q(x) dx \leq \frac{1}{q} \int_{-1}^1 \frac{P_n^q(x)}{(1-x)^q} dx.$$

Indeed, integration by parts yields

$$S = \int_{-1}^1 \frac{P_n^q(x)}{(1-x)^q} dx = \frac{1}{q-1} P_n^q(x)(1-x)^{1-q} \Big|_{-1}^1 + \frac{q}{1-q} \int_{-1}^1 P_n'(x) P_n^{q-1}(x) (1-x)^{1-q} dx.$$

Since $P_n(-1) = 0$, we have

$$S_1 = \frac{1-q}{q} S = \int_{-1}^1 P_n'(x) P_n^{q-1}(x) (1-x)^{1-q} dx.$$

We now estimate $S_1 + S$ to get the result:

$$S_1 + S = \frac{1}{q} S = \int_{-1}^1 \left[\frac{P_n^q}{(1-x)^q} + P_n'(x) P_n^{q-1}(x) (1-x)^{1-q} \right] dx \geq \int_{-1}^1 [P_n'(x)]^q dx$$

since

$$\frac{P_n^q(x)}{(1-x)^q} + P_n'(x) P_n^{q-1}(x) (1-x)^{1-q} \geq [P_n'(x)]^q$$

pointwise. Indeed, if

$$\frac{P_n^q(x)}{(1-x)^q} \geq [P_n'(x)]^q$$

the inequality clearly holds. In the other case, if

$$\frac{P_n^q(x)}{(1-x)^q} < [P_n'(x)]^q,$$

then

$$[P_n'(x)]^{q-1} < (1-x)^{1-q} P_n^{q-1}(x)$$

and the second term dominates the right-hand side.

Next we show that it is possible to stay bounded away from the endpoints of the interval in the sense, that

$$\int_{-1}^1 [P_n'(x)]^q dx \leq C_3(q) \int_{-1}^{1-c/n^2} \frac{P_n^q(x)}{(1-x)^q} dx.$$

To prove the last inequality, we estimate

$$\begin{aligned} \int_{1-c/n^2}^1 \frac{P_n^q(x)}{(1-x)^q} dx &\leq P_n^q(1) \int_{1-c/n^2}^1 (1-x)^{-q} dx = \frac{1}{1-q} c^{1-q} n^{2q-2} \|P_n'\|_{L_1[-1,1]}^q \\ &\leq c^{1-q} C(q) \|P_n'\|_{L_q[-1,1]}^q, \end{aligned}$$

where the constant $C_1(q)$ comes from the classical Nikoskii inequality for polynomial P_n' and spaces $L_1[-1,1]$ and $L_q[-1,1]$ respectively. Taking c to be sufficiently small, we can make $c^{1-q} C(q) \leq \frac{q}{2}$. For such defined $c = c(p, q)$ we have

$$(3) \quad \int_{-1}^1 [P_n'(x)]^q dx \leq \frac{2}{q} \int_{-1}^{1-c/n^2} \frac{P_n^q(x)}{(1-x)^q} dx.$$

We are ready to prove bounds from above for $k = 1$. For $q \geq p$ the result follows from the classical Markov-Nikolskii inequality. Let $\frac{1}{q} = \frac{1}{p} + \frac{1}{r}$ and $r > 0$. Combining (3) with Young's inequality we get

$$\begin{aligned} \frac{\|P'_n\|_{L_q[-1,1]}}{\|P_n\|_{L_p[-1,1]}} &\leq \frac{2}{q} \frac{\|P_n(x)(1-x)^{-1}\|_{L_q[-1,1-c/n^2]}}{\|P_n\|_{L_p[-1,1-c/n^2]}} \\ &\leq \frac{2}{q} \|(1-x)^{-1}\|_{L_r[-1,1-c/n^2]}. \end{aligned}$$

The only thing left is to observe that

$$\|(1-x)^{-1}\|_{L_r[-1,1-c/n^2]} \asymp \begin{cases} n^{2+2/p-2/q}, & \text{if } 1 > 1/q - 1/p, \\ \log n, & \text{if } 1 = 1/q - 1/p, \\ 1, & \text{if } 1 < 1/q - 1/p. \end{cases}$$

We prove the upper bound of the theorem for all k by induction. The base case has been proved above. Let us assume that for each $P_n \in \Delta_n^{k-1}$, $k \geq 2$, $1 \leq m \leq k-1$ we have

$$\frac{\|P_n^{(m)}\|_{L_q[-1,1]}}{\|P_n\|_{L_p[-1,1]}} \leq C(k-1, q, p) \begin{cases} n^{2(m-1)+2/p-2/q}, & \text{if } m-1 > 1/q - 1/p, \\ \log^{m-1} n, & \text{if } m-1 = 1/q - 1/p, \\ 1, & \text{if } m-1 < 1/q - 1/p. \end{cases}$$

Take $P_n \in \Delta_n^k$. If $\frac{1}{q} - \frac{1}{p} = m$, then $\frac{1}{q} - \frac{1}{p/p+1} = m-1$. Using induction hypothesis for $Q_n = P'_n \in \Delta_n^{k-1}$, we get

$$\begin{aligned} \frac{\|P_n^{(m)}\|_{L_q[-1,1]}}{\|P_n\|_{L_p[-1,1]}} &\leq C(k-1, q, p/p+1) \log^{m-1} n \frac{\|P'_n\|_{L_{p/p+1}[-1,1]}}{\|P_n\|_{L_p[-1,1]}} \\ &\leq C(k, q, p) \log^m n. \end{aligned}$$

Following the same lines, if $\frac{1}{q} - \frac{1}{p} > m$, take $r < \frac{p}{p+1}$ such that $\frac{1}{q} - \frac{1}{r} > m-1$ and use induction hypothesis to arrive at

$$\begin{aligned} \frac{\|P_n^{(m)}\|_{L_q[-1,1]}}{\|P_n\|_{L_p[-1,1]}} &\leq C(k-1, q, r) \frac{\|P'_n\|_{L_r[-1,1]}}{\|P_n\|_{L_p[-1,1]}} \\ &\leq C(k, q, p). \end{aligned}$$

If $\frac{1}{q} - \frac{1}{p} < m$, take $r > \frac{p}{p+1}$ such that $\frac{1}{q} - \frac{1}{r} < m-1$ and use induction hypothesis to get

$$\begin{aligned} \frac{\|P_n^{(m)}\|_{L_q[-1,1]}}{\|P_n\|_{L_p[-1,1]}} &\leq C(k-1, q, r) n^{2(m-1)+2/p-2/q} \frac{\|P'_n\|_{L_r[-1,1]}}{\|P_n\|_{L_p[-1,1]}} \\ &\leq C(k, q, p) n^{2m+2/p-2/q}. \end{aligned}$$

The proof of an upper bound is now complete.

We treat the case $p = \infty$ separately.

Lemma 2.1.

$$M_{q,\infty}^{(k)}(n, m) \leq C(k, q) \begin{cases} n^{2m-2/q}, & \text{if } m > \frac{1}{q}, \\ \log^{m-1} n, & \text{if } m = \frac{1}{q}, \\ 1, & \text{if } m < \frac{1}{q}. \end{cases}$$

Proof. Since $\|P_n\| \geq \|P'_n\|_{L_1[-1,1]}$ the result immediately follows from (2). \square

To prove the lower bounds we begin with the following two lemmas:

Lemma 2.2. *Consider*

$$Q_n(x) = \sum_{k=1}^n \frac{\alpha(\alpha+1)\dots(\alpha+k-1)}{k!} x^k$$

for $\alpha = \frac{1}{m}$ and integer $m \geq 1$. Then

$$\int_0^1 Q_n^{\frac{1}{\alpha}}(x) dx \geq C(\alpha) \log n.$$

Proof. For $\alpha = 1$ the result immediately follows from direct integration. For $\alpha \neq 1$, we first note that

$$\frac{\alpha(\alpha+1)\dots(\alpha+k-1)}{k!} \sim k^{\alpha-1}.$$

Introducing

$$B_n(x) = 1 + \sum_{k=1}^n x^k k^{\alpha-1},$$

we are left to show that

$$\int_0^1 B_n^m(x) dx \geq C(\alpha) \log n.$$

Using generalized binomial theorem the coefficient of x^l , $1 \leq l \leq n$, is equal to

$$\sum_{l_1+l_2+\dots+l_m=l} (l_1 l_2 \dots l_m)^{\alpha-1}.$$

Therefore, since the number of ways to represent l as a sum of m positive integers is equal to $\binom{m+l-1}{m-1} \sim l^{m-1}$ and each term

$$(l_1 l_2 \dots l_m)^{\alpha-1} \geq \left(\frac{l}{m}\right)^{(\alpha-1)m} \sim l^{1-m},$$

we get

$$\sum_{l_1+l_2+\dots+l_m=l} (l_1 l_2 \dots l_m)^{\alpha-1} \geq C(\alpha).$$

Therefore

$$\int_0^1 B_n^m(x) dx \geq C(\alpha) \int_0^1 \sum_{k=0}^n x^k dx \geq C(\alpha) \log n.$$

\square

Lemma 2.3. *Let $n \in \mathbb{N}$ and*

$$Q_{n,\alpha}(x) = \sum_{k=1}^n \frac{\alpha(\alpha+1)\dots(\alpha+k-1)}{k!} x^k,$$

and $\alpha < 1$. Then

$$|Q_{n,\alpha}(x)| \leq C_1(\alpha),$$

for all $x \in [-1, 0]$.

Proof. Using Abel's summation formula, we arrive at

$$\begin{aligned} Q_{n,\alpha}(x) &= \sum_{k=1}^{n-1} \left(\frac{\alpha(\alpha+1)\dots(\alpha+k-1)}{k!} - \frac{\alpha(\alpha+1)\dots(\alpha+k)}{(k+1)!} \right) \left(\sum_{i=1}^k x^i \right) \\ &\quad + \frac{\alpha(\alpha+1)\dots(\alpha+n-1)}{n!} \sum_{i=1}^n x^i \end{aligned}$$

Observe, that each sum of the form $\sum_{i=1}^k x^i = \frac{1-x^{k+1}}{1-x} \leq 2$ for $x \in [-1, 0]$ and

$$\frac{\alpha(\alpha+1)\dots(\alpha+k-1)}{k!} - \frac{\alpha(\alpha+1)\dots(\alpha+k)}{(k+1)!} \sim k^{\alpha-2}.$$

The only thin left is to observe that the sum

$$\sum_{k=1}^{\infty} k^{\alpha-2},$$

converges for $\alpha < 1$.

□

Proof of the lower bound in Theorem 1.6. We show that for $0 < p, q \leq \infty$, $p \neq \infty$ and $0 < m \leq k \leq n$

$$(4) \quad M_{q,p}^{(k)}(n, m) \geq C(k, p, q) \begin{cases} n^{2m+2/p-2/q}, & \text{if } m > 1/q - 1/p, \\ \log^m n, & \text{if } m = 1/q - 1/p, \\ 1, & \text{if } m < 1/q - 1/p. \end{cases}$$

Note, that in the case when our order is $n^{2m+2/p-2/q}$ the polynomial was constructed by Erdélyi, more precisely, the construction in Theorem 2.1 in [3] is valid for all $0 < p, q \leq \infty$. So of interest is to construct a polynomial $P_n \in \Delta_n^k$ such that for all $0 < m \leq k$

$$M_{q,p}^{(k)}(n, m) \asymp \log^m n.$$

By continuity, we can assume that $p, q \in \mathbb{Q}$. Take

$$Q_n(x) = \sum_{k=1}^n \frac{\frac{1}{2mq}(\frac{1}{2mq} + 1)\dots(\frac{1}{2mq} + k - 1)}{k!} x^k$$

and consider

$$P_\nu(y) = P_{2mn+k}(y) = \int_{-1}^y Q_n^{2m}(x)(y-x)^{k-1} dx.$$

Clearly, $\nu = 2mn + k$ and $\deg P_\nu = 2mn + k$. It is easy to see, that $P_\nu \in \Delta_\nu^k$ and $P_\nu^{(k)}(x) = Q_n^{2m}(x)$. Lemma 2.2 implies $\|P_\nu^{(k)}\|_{L_q[-1,1]} \geq C(k, q) \log^{1/q} n$. Thus, we are left to prove that $\|P_\nu\|_{L_p[-1,1]} \leq C(k, p) \log^{1/p} n$. Since $P_\nu \in \Delta_n^k$ Remez inequality (see [2]) implies that for sufficiently small $c = c(p)$

$$\|P_\nu\|_{L_p[-1,1]}^p \leq 2 \int_0^1 P_\nu^p(x) dx \leq C(p) \int_0^{1-c/n^2} P_\nu^p(x) dx.$$

Now, for $y > 0$ using that $|a+b|^p \leq C(p)(|a|^p + |b|^p)$, $y-x \leq 1-x$ and $0 < Q_n(x) \leq (1-x)^{-1/2mq}$ for $x \geq 0$ we can estimate

$$\begin{aligned} \int_0^{1-c/n^2} P_\nu^p(x) dx &= \int_0^{1-c/n^2} \left(\int_{-1}^y Q_n^{2m}(x)(y-x)^{k-1} dx \right)^p dy \\ &\leq C(p) \int_0^{1-c/n^2} \left(\int_0^y Q_n^{2m}(x)(y-x)^{k-1} dx \right)^p dy \\ &\quad + C(p) \int_0^{1-c/n^2} \left(\int_{-1}^0 Q_n^{2m}(x)(y-x)^{k-1} dx \right)^p dy \\ &\leq C(p) \int_0^{1-c/n^2} \left(\int_0^y Q_n^{2m}(x)(y-x)^{k-1} dx \right)^p dy \\ &\quad + 2^k C(p) \left(\int_{-1}^0 Q_n^{2m}(x)(y-x)^{k-1} dx \right)^p \\ &\leq \int_0^{1-c/n^2} \left(\int_{-1}^y (1-x)^{-1/q} (1-x)^{k-1} dx \right)^p dy + C_1(p, k, m) \\ &\leq C_2(p) \log n. \end{aligned}$$

It is now straightforward to get a sharp result for all intermediate derivatives of k -absolutely monotone polynomials by using (2) and (4).

The result for the case $p = \infty$ immediately follows from the construction described above and the fact $\|P_n\| = \|P'_n\|_{L_1[-1,1]}$.

Acknowledgment

The author would like to thank Professor K. Kopotun for all his support and encouragement and to anonymous referee for useful suggestions.

REFERENCES

- [1] Serge Bernstein. Sur l'extension du théorème limite du calcul des probabilités aux sommes de quantités dépendantes. *Math. Ann.*, 97(1):1–59, 1927.
- [2] Peter Borwein and Tamás Erdélyi. *Polynomials and polynomial inequalities*, volume 161 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [3] Tamás Erdélyi. A Markov Nikolskii type inequality for absolutely monotone polynomials of order K . *J. Anal. Math.*, 112:369–381, 2010.

- [4] P. Yu. Glazyrina. The sharp Markov-Nikolskii inequality for algebraic polynomials in the spaces L_q and L_0 on an interval. *Mat. Zametki*, 84(1):3–22, 2008.
- [5] A. Kroó and J. Szabados. On the exact Markov inequality for k -monotone polynomials in uniform and L_1 -norms. *Acta Math. Hungar.*, 125(1-2):99–112, 2009.
- [6] Mohammed A. Qazi. On polynomials monotonic on the unit interval. *Analysis (Munich)*, 21(2):129–134, 2001.

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, CP
6128 SUCC. CENTRE-VILLE, MONTRÉAL QC H3C 3J7, CANADA CANADA

E-mail address: lklurman@gmail.com