

A symmetric Nörlund sum with application to inequalities

Artur M. C. Brito da Cruz, Natália Martins and Delfim F. M. Torres

[Submitted 18-Oct-2011; accepted 09-March-2012; to Proceedings of International Conference on Differential & Difference Equations and Applications, in honour of Professor Ravi P. Agarwal, to be published by Springer in the series *Proceedings in Mathematics* (PROM)]

Abstract Properties of an α, β -symmetric Nörlund sum are studied. Inspired in the work by Agarwal et al., α, β -symmetric quantum versions of Hölder, Cauchy–Schwarz and Minkowski inequalities are obtained.

1 Introduction

The symmetric derivative of function f at point x is defined as $\lim_{h \rightarrow 0} (f(x+h) - f(x-h))/(2h)$. The notion of symmetrically differentiable is interesting because if a function is differentiable at a point then it is also symmetrically differentiable, but the converse is not true. The best known example of this fact is the absolute value function: $f(x) = |x|$ is not differentiable at $x = 0$ but is symmetrically differentiable at $x = 0$ with symmetric derivative zero [6].

Quantum calculus is, roughly speaking, the equivalent to traditional infinitesimal calculus but without limits [4]. Therefore, one can introduce the symmetric quantum derivative of f at x by $(f(x+h) - f(x-h))/(2h)$. As in any calculus, it is then natural to develop a corresponding integration theory, looking to such integral as the inverse operator of the derivative.

Artur M. C. Brito da Cruz^{1,2}
e-mail: artur.cruz@estsetubal.ips.pt

Natália Martins²
e-mail: natalia@ua.pt

Delfim F. M. Torres²
e-mail: delfim@ua.pt

¹Escola Superior de Tecnologia de Setúbal, Estefanilha, 2910-761 Setúbal, Portugal

²Center for Research and Development in Mathematics and Applications
Department of Mathematics, University of Aveiro, 3810-193 Aveiro, Portugal

The main goal of this paper is to study the properties of a general symmetric quantum integral that we call, due to the so-called Nörlund sum [4], the α, β -symmetric Nörlund sum.

The paper is organized as follows. In Section 2 we define the forward and backward Nörlund sums. Then, in Section 3, we introduce the α, β -symmetric Nörlund sum and give some of its properties. We end with Section 4, proving α, β -symmetric versions of Hölder's, Cauchy–Schwarz's and Minkowski's inequalities.

2 Forward and backward Nörlund sums

This section is dedicated to the inverse operators of the α -forward and β -backward differences, $\alpha > 0, \beta > 0$, defined respectively by

$$\Delta_\alpha[f](t) := \frac{f(t+\alpha) - f(t)}{\alpha}, \quad \nabla_\beta[f](t) := \frac{f(t) - f(t-\beta)}{\beta}.$$

Definition 1. Let $I \subseteq \mathbb{R}$ be such that $a, b \in I$ with $a < b$ and $\sup I = +\infty$. For $f : I \rightarrow \mathbb{R}$ and $\alpha > 0$ we define the Nörlund sum (the α -forward integral) of f from a to b by

$$\int_a^b f(t) \Delta_\alpha t = \int_a^{+\infty} f(t) \Delta_\alpha t - \int_b^{+\infty} f(t) \Delta_\alpha t,$$

where $\int_x^{+\infty} f(t) \Delta_\alpha t = \alpha \sum_{k=0}^{+\infty} f(x+k\alpha)$, provided the series converges at $x = a$ and $x = b$. In that case, f is said to be α -forward integrable on $[a, b]$. We say that f is α -forward integrable over I if it is α -forward integrable for all $a, b \in I$.

Until Definition 2 (the backward/nabla case), we assume that I is an interval of \mathbb{R} such that $\sup I = +\infty$. Note that if $f : I \rightarrow \mathbb{R}$ is a function such that $\sup I < +\infty$, then we can extend function f to $\tilde{f} : \tilde{I} \rightarrow \mathbb{R}$, where \tilde{I} is an interval with $\sup \tilde{I} = +\infty$, in the following way: $\tilde{f}|_I = f$ and $\tilde{f}|_{\tilde{I} \setminus I} = 0$.

Using the techniques of Aldwoah in his Ph.D. thesis [2], it can be proved that the α -forward integral has the following properties:

Theorem 1. If $f, g : I \rightarrow \mathbb{R}$ are α -forward integrable on $[a, b]$, $c \in [a, b]$, $k \in \mathbb{R}$, then

1. $\int_a^a f(t) \Delta_\alpha t = 0$;
2. $\int_a^b f(t) \Delta_\alpha t = \int_a^c f(t) \Delta_\alpha t + \int_c^b f(t) \Delta_\alpha t$, when the integrals exist;
3. $\int_a^b f(t) \Delta_\alpha t = - \int_b^a f(t) \Delta_\alpha t$;
4. kf is α -forward integrable on $[a, b]$ and $\int_a^b kf(t) \Delta_\alpha t = k \int_a^b f(t) \Delta_\alpha t$;
5. $f + g$ is α -forward integrable on $[a, b]$ and

$$\int_a^b (f+g)(t) \Delta_\alpha t = \int_a^b f(t) \Delta_\alpha t + \int_a^b g(t) \Delta_\alpha t;$$

6. if $f \equiv 0$, then $\int_a^b f(t) \Delta_\alpha t = 0$.

Theorem 2. Let $f : I \rightarrow \mathbb{R}$ be α -forward integrable on $[a, b]$. If $g : I \rightarrow \mathbb{R}$ is a non-negative α -forward integrable function on $[a, b]$, then fg is α -forward integrable on $[a, b]$.

Proof. Since g is α -forward integrable, then both series $\alpha \sum_{k=0}^{+\infty} g(a+k\alpha)$ and $\alpha \sum_{k=0}^{+\infty} g(b+k\alpha)$ converge. We want to study the nature of series $\alpha \sum_{k=0}^{+\infty} fg(a+k\alpha)$ and $\alpha \sum_{k=0}^{+\infty} fg(b+k\alpha)$. Since there exists an order $N \in \mathbb{N}$ such that $|fg(b+k\alpha)| \leq g(b+k\alpha)$ and $|fg(a+k\alpha)| \leq g(a+k\alpha)$ for all $k > N$, then both $\alpha \sum_{k=0}^{+\infty} fg(a+k\alpha)$ and $\alpha \sum_{k=0}^{+\infty} fg(b+k\alpha)$ converge absolutely. The intended conclusion follows.

Theorem 3. Let $f : I \rightarrow \mathbb{R}$ and $p > 1$. If $|f|$ is α -forward integrable on $[a, b]$, then $|f|^p$ is also α -forward integrable on $[a, b]$.

Proof. There exists $N \in \mathbb{N}$ such that $|f(b+k\alpha)|^p \leq |f(b+k\alpha)|$ and $|f(a+k\alpha)|^p \leq |f(a+k\alpha)|$ for all $k > N$. Therefore, $|f|^p$ is α -forward integrable on $[a, b]$.

Theorem 4. Let $f, g : I \rightarrow \mathbb{R}$ be α -forward integrable on $[a, b]$. If $|f(t)| \leq g(t)$ for all $t \in \{a+k\alpha : k \in \mathbb{N}_0\}$, then for $b \in \{a+k\alpha : k \in \mathbb{N}_0\}$ one has

$$\left| \int_a^b f(t) \Delta_\alpha t \right| \leq \int_a^b g(t) \Delta_\alpha t.$$

Proof. Since $b \in \{a+k\alpha : k \in \mathbb{N}_0\}$, there exists k_1 such that $b = a+k_1\alpha$. Thus,

$$\begin{aligned} \left| \int_a^b f(t) \Delta_\alpha t \right| &= \left| \alpha \sum_{k=0}^{+\infty} f(a+k\alpha) - \alpha \sum_{k=0}^{+\infty} f(a+(k_1+k)\alpha) \right| \\ &= \left| \alpha \sum_{k=0}^{+\infty} f(a+k\alpha) - \alpha \sum_{k=k_1}^{+\infty} f(a+k\alpha) \right| = \left| \alpha \sum_{k=0}^{k_1-1} f(a+k\alpha) \right| \\ &\leq \alpha \sum_{k=0}^{k_1-1} |f(a+k\alpha)| \leq \alpha \sum_{k=0}^{k_1-1} g(a+k\alpha) \\ &= \alpha \sum_{k=0}^{+\infty} g(a+k\alpha) - \alpha \sum_{k=k_1}^{+\infty} g(a+k\alpha) = \int_a^b g(t) \Delta_\alpha t. \end{aligned}$$

Corollary 1. Let $f, g : I \rightarrow \mathbb{R}$ be α -forward integrable on $[a, b]$ with $b = a+k\alpha$ for some $k \in \mathbb{N}_0$.

1. If $f(t) \geq 0$ for all $t \in \{a+k\alpha : k \in \mathbb{N}_0\}$, then $\int_a^b f(t) \Delta_\alpha t \geq 0$.
2. If $g(t) \geq f(t)$ for all $t \in \{a+k\alpha : k \in \mathbb{N}_0\}$, then $\int_a^b g(t) \Delta_\alpha t \geq \int_a^b f(t) \Delta_\alpha t$.

We can now prove the following fundamental theorem of the α -forward calculus.

Theorem 5 (Fundamental theorem of Nörlund calculus). *Let $f : I \rightarrow \mathbb{R}$ be α -forward integrable over I . Let $x \in I$ and define $F(x) := \int_a^x f(t) \Delta_\alpha t$. Then, $\Delta_\alpha [F](x) = f(x)$. Conversely, $\int_a^b \Delta_\alpha [f](t) \Delta_\alpha t = f(b) - f(a)$.*

Proof. If $G(x) = -\int_x^{+\infty} f(t) \Delta_\alpha t$, then

$$\begin{aligned} \Delta_\alpha [G](x) &= \frac{G(x+\alpha) - G(x)}{\alpha} = \frac{-\alpha \sum_{k=0}^{+\infty} f(x+\alpha+k\alpha) + \alpha \sum_{k=0}^{+\infty} f(x+k\alpha)}{\alpha} \\ &= \sum_{k=0}^{+\infty} f(x+k\alpha) - \sum_{k=0}^{+\infty} f(x+(k+1)\alpha) = f(x). \end{aligned}$$

Therefore, $\Delta_\alpha [F](x) = \Delta_\alpha \left(\int_a^{+\infty} f(t) \Delta_\alpha t - \int_x^{+\infty} f(t) \Delta_\alpha t \right) = f(x)$. Using the definition of α -forward difference operator, the second part of the theorem is also a consequence of the properties of Mengoli's series. Since

$$\begin{aligned} \int_a^{+\infty} \Delta_\alpha [f](t) \Delta_\alpha t &= \alpha \sum_{k=0}^{+\infty} \Delta_\alpha [f](a+k\alpha) = \alpha \sum_{k=0}^{+\infty} \frac{f(a+k\alpha+\alpha) - f(a+k\alpha)}{\alpha} \\ &= \sum_{k=0}^{+\infty} \left(f(a+(k+1)\alpha) - f(a+k\alpha) \right) = -f(a) \end{aligned}$$

and $\int_b^{+\infty} \Delta_\alpha [f](t) \Delta_\alpha t = -f(b)$, it follows that

$$\int_a^b \Delta_\alpha [f](t) \Delta_\alpha t = \int_a^{+\infty} f(t) \Delta_\alpha t - \int_b^{+\infty} f(t) \Delta_\alpha t = f(b) - f(a).$$

Corollary 2 (α -forward integration by parts). *Let $f, g : I \rightarrow \mathbb{R}$. If fg and $f \Delta_\alpha [g]$ are α -forward integrable on $[a, b]$, then*

$$\int_a^b f(t) \Delta_\alpha [g](t) \Delta_\alpha t = f(t)g(t) \Big|_a^b - \int_a^b \Delta_\alpha [f](t) g(t+\alpha) \Delta_\alpha t$$

Proof. Since $\Delta_\alpha [fg](t) = \Delta_\alpha [f](t)g(t+\alpha) + f(t)\Delta_\alpha [g](t)$, then

$$\begin{aligned} \int_a^b f(t) \Delta_\alpha [g](t) \Delta_\alpha t &= \int_a^b \left(\Delta_\alpha [fg](t) - \Delta_\alpha [f](t)g(t+\alpha) \right) \Delta_\alpha t \\ &= \int_a^b \Delta_\alpha [fg](t) \Delta_\alpha t - \int_a^b \Delta_\alpha [f](t)g(t+\alpha) \Delta_\alpha t \\ &= f(t)g(t) \Big|_a^b - \int_a^b \Delta_\alpha [f](t)g(t+\alpha) \Delta_\alpha t. \end{aligned}$$

Remark 1. Our study of the Nörlund sum is in agreement with the Hahn quantum calculus [2, 3, 5]. In [4] $\int_a^b f(t) \Delta_\alpha t = \alpha [f(a) + f(a+\alpha) + \dots + f(b-\alpha)]$ for $a < b$ such that $b-a \in \alpha\mathbb{Z}$, $\alpha \in \mathbb{R}^+$. In contrast with [4], our definition is valid for any two real points a, b and not only for those points belonging to the time scale $\alpha\mathbb{Z}$. The definitions (only) coincide if function f is α -forward integrable on $[a, b]$.

Similarly, we introduce the β -backward integral.

Definition 2. Let I be an interval of \mathbb{R} such that $a, b \in I$ with $a < b$ and $\inf I = -\infty$. For $f : I \rightarrow \mathbb{R}$ and $\beta > 0$ we define the β -backward integral of f from a to b by

$$\int_a^b f(t) \nabla_{\beta} t = \int_{-\infty}^b f(t) \nabla_{\beta} t - \int_{-\infty}^a f(t) \nabla_{\beta} t,$$

where $\int_{-\infty}^x f(t) \nabla_{\beta} t = \beta \sum_{k=0}^{+\infty} f(x - k\beta)$, provided the series converges at $x = a$ and $x = b$. In that case, f is called β -backward integrable on $[a, b]$. We say that f is β -backward integrable over I if it is β -backward integrable for all $a, b \in I$.

The β -backward Nörlund sum has similar results and properties as the α -forward Nörlund sum. In particular, the β -backward integral is the inverse operator of ∇_{β} .

3 The α, β -symmetric Nörlund sum

We define the α, β -symmetric integral as a linear combination of the α -forward and the β -backward integrals.

Definition 3. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ and $a, b \in \mathbb{R}$, $a < b$. If f is α -forward and β -backward integrable on $[a, b]$, $\alpha, \beta \geq 0$ with $\alpha + \beta > 0$, then we define the α, β -symmetric integral of f from a to b by

$$\int_a^b f(t) d_{\alpha, \beta} t = \frac{\alpha}{\alpha + \beta} \int_a^b f(t) \Delta_{\alpha} t + \frac{\beta}{\alpha + \beta} \int_a^b f(t) \nabla_{\beta} t.$$

Function f is α, β -symmetric integrable if it is α, β -symmetric integrable for all $a, b \in \mathbb{R}$.

Remark 2. Note that if $\alpha \in \mathbb{R}^+$ and $\beta = 0$, then $\int_a^b f(t) d_{\alpha, \beta} t = \int_a^b f(t) \Delta_{\alpha} t$ and we do not need to assume in Definition 3 that f is β -backward integrable; if $\alpha = 0$ and $\beta \in \mathbb{R}^+$, then $\int_a^b f(t) d_{\alpha, \beta} t = \int_a^b f(t) \nabla_{\beta} t$ and we do not need to assume that f is α -forward integrable.

Example 1. Let $f(t) = 1/t^2$. Then $\int_1^3 \frac{1}{t^2} d_{2, 2} t = \frac{10}{9}$.

The α, β -symmetric integral has the following properties:

Theorem 6. Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be α, β -symmetric integrable on $[a, b]$. Let $c \in [a, b]$ and $k \in \mathbb{R}$. Then,

1. $\int_a^a f(t) d_{\alpha, \beta} t = 0$;

2. $\int_a^b f(t) d_{\alpha, \beta} t = \int_a^c f(t) d_{\alpha, \beta} t + \int_c^b f(t) d_{\alpha, \beta} t$, when the integrals exist;
3. $\int_a^b f(t) d_{\alpha, \beta} t = - \int_b^a f(t) d_{\alpha, \beta} t$;
4. kf is α, β -symmetric integrable on $[a, b]$ and $\int_a^b kf(t) d_{\alpha, \beta} t = k \int_a^b f(t) d_{\alpha, \beta} t$;
5. $f + g$ is α, β -symmetric integrable on $[a, b]$ and

$$\int_a^b (f + g)(t) d_{\alpha, \beta} t = \int_a^b f(t) d_{\alpha, \beta} t + \int_a^b g(t) d_{\alpha, \beta} t;$$

6. fg is α, β -symmetric integrable on $[a, b]$ provided g is a nonnegative function.

Proof. These results are easy consequences of the α -forward and β -backward integral properties.

The next result follows immediately from Theorem 3 and the corresponding β -backward version.

Theorem 7. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ and $p > 1$. If $|f|$ is symmetric α, β -integrable on $[a, b]$, then $|f|^p$ is also α, β -symmetric integrable on $[a, b]$.

Theorem 8. Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be α, β -symmetric integrable functions on $[a, b]$, $\mathcal{A} := \{a + k\alpha : k \in \mathbb{N}_0\}$ and $\mathcal{B} := \{b - k\beta : k \in \mathbb{N}_0\}$. For $b \in \mathcal{A}$ and $a \in \mathcal{B}$ one has:

1. if $|f(t)| \leq g(t)$ for all $t \in \mathcal{A} \cup \mathcal{B}$, then $\left| \int_a^b f(t) d_{\alpha, \beta} t \right| \leq \int_a^b g(t) d_{\alpha, \beta} t$;
2. if $f(t) \geq 0$ for all $t \in \mathcal{A} \cup \mathcal{B}$, then $\int_a^b f(t) d_{\alpha, \beta} t \geq 0$;
3. if $g(t) \geq f(t)$ for all $t \in \mathcal{A} \cup \mathcal{B}$, then $\int_a^b g(t) d_{\alpha, \beta} t \geq \int_a^b f(t) d_{\alpha, \beta} t$.

Proof. It follows from Theorem 4 and Corollary 1 and the corresponding β -backward versions.

In Theorem 9 we assume that $a, b \in \mathbb{R}$ with $b \in \mathcal{A} := \{a + k\alpha : k \in \mathbb{N}_0\}$ and $a \in \mathcal{B} := \{b - k\beta : k \in \mathbb{N}_0\}$, where $\alpha, \beta \in \mathbb{R}_0^+$, $\alpha + \beta \neq 0$.

Theorem 9 (Mean value theorem). Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be bounded and α, β -symmetric integrable on $[a, b]$ with g nonnegative. Let m and M be the infimum and the supremum, respectively, of function f . Then, there exists a real number K satisfying the inequalities $m \leq K \leq M$ such that $\int_a^b f(t) g(t) d_{\alpha, \beta} t = K \int_a^b g(t) d_{\alpha, \beta} t$.

Proof. Since $m \leq f(t) \leq M$ for all $t \in \mathbb{R}$ and $g(t) \geq 0$, then $mg(t) \leq f(t)g(t) \leq Mg(t)$ for all $t \in \mathcal{A} \cup \mathcal{B}$. All functions mg , $f g$ and Mg are α, β -symmetric integrable on $[a, b]$. By Theorems 6 and 8, $m \int_a^b g(t) d_{\alpha, \beta} t \leq \int_a^b f(t) g(t) d_{\alpha, \beta} t \leq M \int_a^b g(t) d_{\alpha, \beta} t$. If $\int_a^b g(t) d_{\alpha, \beta} t = 0$, then $\int_a^b f(t) g(t) d_{\alpha, \beta} t = 0$; if $\int_a^b g(t) d_{\alpha, \beta} t > 0$, then $m \leq \frac{\int_a^b f(t) g(t) d_{\alpha, \beta} t}{\int_a^b g(t) d_{\alpha, \beta} t} \leq M$. Therefore, the middle term of these inequalities is equal to a number K , which yields the intended result.

4 α, β -Symmetric Integral Inequalities

Inspired in the work by Agarwal et al. [1], we now present α, β -symmetric versions of Hölder, Cauchy–Schwarz and Minkowski inequalities. As before, we assume that $a, b \in \mathbb{R}$ with $b \in \mathcal{A} := \{a + k\alpha : k \in \mathbb{N}_0\}$ and $a \in \mathcal{B} := \{b - k\beta : k \in \mathbb{N}_0\}$, where $\alpha, \beta \in \mathbb{R}_0^+$, $\alpha + \beta \neq 0$.

Theorem 10 (Hölder’s inequality). *Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ and $a, b \in \mathbb{R}$ with $a < b$. If $|f|$ and $|g|$ are α, β -symmetric integrable on $[a, b]$, then*

$$\int_a^b |f(t)g(t)| d_{\alpha, \beta} t \leq \left(\int_a^b |f(t)|^p d_{\alpha, \beta} t \right)^{\frac{1}{p}} \left(\int_a^b |g(t)|^q d_{\alpha, \beta} t \right)^{\frac{1}{q}}, \quad (1)$$

where $p > 1$ and $q = p/(p-1)$.

Proof. For $\alpha, \beta \in \mathbb{R}_0^+$, $\alpha + \beta \neq 0$, the following inequality holds: $\alpha^{\frac{1}{p}} \beta^{\frac{1}{q}} \leq \frac{\alpha}{p} + \frac{\beta}{q}$.

Without loss of generality, suppose that $\left(\int_a^b |f(t)|^p d_{\alpha, \beta} t \right) \left(\int_a^b |g(t)|^q d_{\alpha, \beta} t \right) \neq 0$ (note that both integrals exist by Theorem 7). Set $\xi(t) = |f(t)|^p / \int_a^b |f(\tau)|^p d_{\alpha, \beta} \tau$ and $\gamma(t) = |g(t)|^q / \int_a^b |g(\tau)|^q d_{\alpha, \beta} \tau$. Since both functions α and β are symmetric α, β -integrable on $[a, b]$, then (1) holds:

$$\begin{aligned} \int_a^b \frac{|f(t)|}{\left(\int_a^b |f(\tau)|^p d_{\alpha, \beta} \tau \right)^{\frac{1}{p}}} \frac{|g(t)|}{\left(\int_a^b |g(\tau)|^q d_{\alpha, \beta} \tau \right)^{\frac{1}{q}}} d_{\alpha, \beta} t &= \int_a^b \xi(t)^{\frac{1}{p}} \gamma(t)^{\frac{1}{q}} d_{\alpha, \beta} t \\ &\leq \int_a^b \left(\frac{\xi(t)}{p} + \frac{\gamma(t)}{q} \right) d_{\alpha, \beta} t \\ &= \frac{1}{p} \int_a^b \left(\frac{|f(t)|^p}{\int_a^b |f(\tau)|^p d_{\alpha, \beta} \tau} \right) d_{\alpha, \beta} t + \frac{1}{q} \int_a^b \left(\frac{|g(t)|^q}{\int_a^b |g(\tau)|^q d_{\alpha, \beta} \tau} \right) d_{\alpha, \beta} t = 1. \end{aligned}$$

The particular case $p = q = 2$ of (1) gives the Cauchy–Schwarz inequality.

Corollary 3 (Cauchy–Schwarz’s inequality). *Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ and $a, b \in \mathbb{R}$ with $a < b$. If f and g are α, β -symmetric integrable on $[a, b]$, then*

$$\int_a^b |f(t)g(t)| d_{\alpha, \beta} t \leq \sqrt{\left(\int_a^b |f(t)|^2 d_{\alpha, \beta} t \right) \left(\int_a^b |g(t)|^2 d_{\alpha, \beta} t \right)}.$$

We prove the Minkowski inequality using Hölder’s inequality.

Theorem 11 (Minkowski’s inequality). *Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ and $a, b, p \in \mathbb{R}$ with $a < b$ and $p > 1$. If f and g are α, β -symmetric integrable on $[a, b]$, then*

$$\left(\int_a^b |f(t) + g(t)|^p d_{\alpha, \beta} t \right)^{\frac{1}{p}} \leq \left(\int_a^b |f(t)|^p d_{\alpha, \beta} t \right)^{\frac{1}{p}} + \left(\int_a^b |g(t)|^p d_{\alpha, \beta} t \right)^{\frac{1}{p}}.$$

Proof. One has

$$\begin{aligned} \int_a^b |f(t) + g(t)|^p d_{\alpha, \beta} t &= \int_a^b |f(t) + g(t)|^{p-1} |f(t) + g(t)| d_{\alpha, \beta} t \\ &\leq \int_a^b |f(t)| |f(t) + g(t)|^{p-1} d_{\alpha, \beta} t + \int_a^b |g(t)| |f(t) + g(t)|^{p-1} d_{\alpha, \beta} t. \end{aligned}$$

Applying Hölder's inequality (Theorem 10) with $q = p/(p-1)$, we obtain

$$\begin{aligned} \int_a^b |f(t) + g(t)|^p d_{\alpha, \beta} t &\leq \left(\int_a^b |f(t)|^p d_{\alpha, \beta} t \right)^{\frac{1}{p}} \left(\int_a^b |f(t) + g(t)|^{(p-1)q} d_{\alpha, \beta} t \right)^{\frac{1}{q}} \\ &\quad + \left(\int_a^b |g(t)|^p d_{\alpha, \beta} t \right)^{\frac{1}{p}} \left(\int_a^b |f(t) + g(t)|^{(p-1)q} d_{\alpha, \beta} t \right)^{\frac{1}{q}} \\ &= \left[\left(\int_a^b |f(t)|^p d_{\alpha, \beta} t \right)^{\frac{1}{p}} + \left(\int_a^b |g(t)|^p d_{\alpha, \beta} t \right)^{\frac{1}{p}} \right] \left(\int_a^b |f(t) + g(t)|^{(p-1)q} d_{\alpha, \beta} t \right)^{\frac{1}{q}}. \end{aligned}$$

Therefore,

$$\frac{\int_a^b |f(t) + g(t)|^p d_{\alpha, \beta} t}{\left(\int_a^b |f(t) + g(t)|^{(p-1)q} d_{\alpha, \beta} t \right)^{\frac{1}{q}}} \leq \left(\int_a^b |f(t)|^p d_{\alpha, \beta} t \right)^{\frac{1}{p}} + \left(\int_a^b |g(t)|^p d_{\alpha, \beta} t \right)^{\frac{1}{p}}.$$

Our α, β -symmetric calculus is more general than the standard h -calculus. In particular, all our results give, as corollaries, results in the classical quantum h -calculus by choosing $\alpha = h > 0$ and $\beta = 0$.

Acknowledgements Work supported by FEDER and Portuguese funds, COMPETE reference FCOMP-01-0124-FEDER-022690, and CIDMA and FCT, project PEst-C/MAT/UI4106/2011. Brito da Cruz is also supported by FCT through the Ph.D. fellowship SFRH/BD/33634/2009.

References

1. R. Agarwal, M. Bohner and A. Peterson, Inequalities on time scales: a survey, *Math. Inequal. Appl.* **4** (2001), no. 4, 535–557.
2. K. A. Aldwoah, *Generalized time scales and associated difference equations*, Ph.D. thesis, Cairo University, 2009.
3. A. M. C. Brito da Cruz, N. Martins and D. F. M. Torres, Higher-order Hahn's quantum variational calculus, *Nonlinear Anal.* **75** (2012), no. 3, 1147–1157. [arXiv:1101.3653](#)
4. V. Kac and P. Cheung, *Quantum calculus*, Springer, New York, 2002.
5. A. B. Malinowska and D. F. M. Torres, The Hahn quantum variational calculus, *J. Optim. Theory Appl.* **147** (2010), no. 3, 419–442. [arXiv:1006.3765](#)
6. B. S. Thomson, *Symmetric properties of real functions*, Dekker, New York, 1994.