

LENGTH-EXPANDING LIPSCHITZ MAPS ON TOTALLY REGULAR CONTINUA

VLADIMÍR ŠPITALSKÝ

ABSTRACT. The tent map is an elementary example of an interval map possessing many interesting properties, such as dense periodicity, exactness, Lipschitzness and a kind of length-expansiveness. It is often used in constructions of dynamical systems on the interval/trees/graphs. The purpose of the present paper is to construct, on totally regular continua (i.e. on topologically rectifiable curves), maps sharing some typical properties with the tent map. These maps will be called length-expanding Lipschitz maps, briefly LEL maps. We show that every totally regular continuum endowed with a suitable metric admits a LEL map. As an application we obtain that every totally regular continuum admits an exactly Devaney chaotic map with finite entropy and the specification property.

1. INTRODUCTION

The tent map is the piecewise linear map f on the interval $I = [0, 1]$ given by $x \mapsto 2 \min\{x, 1 - x\}$. The properties of this map, conjugate to the full logistic map $x \mapsto 4x(1 - x)$, include Lipschitzness, length-expansiveness (in a sense that it doubles the length of every subinterval J of I not containing $1/2$), exactness, specification, finite positive topological entropy and dense periodicity, just to name a few. This map, together with “generalized” tent maps, i.e. piecewise linear continuous maps $f_k : I \rightarrow I$ ($k \geq 3$) fixing 0 and mapping linearly every interval $[(i - 1)/k, i/k]$ onto I , are frequently used in dynamics. Usefulness of these maps lies in the fact that on one hand they are very simple (and so we have easy explicit formulae for iterates, periodic points, horseshoes, etc.) and on the other hand they are very “powerful”. They are often used in constructions of systems on the interval/trees/graphs with special properties. For example, it is known that to construct a transitive map on the unit interval with the smallest possible topological entropy, one can define $g : I \rightarrow I$ in such a way that $1/2$ is a fixed point, g maps linearly $I_0 = [0, 1/2]$ onto $I_1 = [1/2, 1]$ and $g|_{I_1} : I_1 \rightarrow I_0$ is “tent-like”. Analogously one can define a transitive map with the smallest possible entropy $(1/n) \log 2$ on any n -star S_n ($n \geq 3$), see [2]; the map fixes the branch point of S_n , maps cyclically each branch to the next one, all but one linearly and the remaining one in a “tent-like” way.

Unfortunately, when one wants to construct a map with given properties on curves more general than graphs, he/she faces the problem that no direct analogue of the tent map on such curves is known. Take e.g. the ω -star X , which is a very simple dendrite defined as an infinite wedge of arcs. A construction of a transitive finite entropy map on X is much more complicated than on n -stars and, as far as

2010 *Mathematics Subject Classification.* Primary 37B05, 37B20, 37B40; Secondary 54H20.

Key words and phrases. Lipschitz map, length-expanding map, tent map, totally regular continuum, rectifiable curve, exact Devaney chaos, specification property.

we know, no such construction has been available in literature. The only result in this direction known to us is the theorem of Agronsky and Ceder [1] stating that any finite-dimensional Peano continuum (hence also the ω -star) admits a transitive map; however, the proof does not say anything about the entropy of the map.

The purpose of the present paper is to construct, on continua more general than graphs, a family of maps sharing some typical properties with the tent map. Since the key property of these maps will deal with the *length* (Hausdorff one-dimensional measure) of subcontinua and their images, the natural class of spaces to consider is the class of *rectifiable curves*, i.e. continua of finite length. Topologically they coincide with the class of totally regular continua. Recall that a continuum X is *totally regular* if for every point $x \in X$ and every countable set $P \subseteq X$ there is a basis of neighborhoods of x with finite boundary not intersecting P . This notion was introduced in [20], but the class of these continua was studied a long time before, see e.g. [21, 8, 10, 9]. For more details on totally regular continua see Section 3.4.

Before stating the main results of the paper we need to introduce the notion of a length-expanding Lipschitz map. Let X be a non-degenerate totally regular continuum. We say that a family \mathcal{C} of non-degenerate subcontinua of X is *dense* if every nonempty open set in X contains a member of \mathcal{C} . Recall that a map $f : (X, d) \rightarrow (X', d')$ between metric spaces is *Lipschitz- L* if $d'(f(x), f(y)) \leq L \cdot d(x, y)$ for every $x, y \in X$. For a metric space (X, d) , the Hausdorff one-dimensional measure is denoted by \mathcal{H}_d^1 .

Definition A. Let $X = (X, d)$, $X' = (X', d')$ be non-degenerate (totally regular) continua of finite length and let $\mathcal{C}, \mathcal{C}'$ be dense systems of subcontinua of X, X' , respectively. We say that a continuous map $f : X \rightarrow X'$ is *length-expanding* with respect to $\mathcal{C}, \mathcal{C}'$ if there exists $\varrho > 1$ (called *length-expansivity constant* of f) such that, for every $C \in \mathcal{C}$, $f(C) \in \mathcal{C}'$ and

$$(1.1) \quad \text{if } f(C) \neq X' \quad \text{then } \mathcal{H}_{d'}^1(f(C)) \geq \varrho \cdot \mathcal{H}_d^1(C).$$

Moreover, if f is surjective and Lipschitz- L we say that $f : (X, d, \mathcal{C}) \rightarrow (X', d', \mathcal{C}')$ is *(ϱ, L) -length-expanding Lipschitz*. Sometimes we briefly say that f is *(ϱ, L) -LEL* or only *LEL*. On the other hand, when we wish to be more precise, we say that f is *$(\mathcal{C}, \mathcal{C}', \varrho, L)$ -LEL*.

A few comments are necessary. Assume that $f : (X, d, \mathcal{C}) \rightarrow (X', d', \mathcal{C}')$ is (ϱ, L) -LEL and denote by \mathcal{C}_X and $\mathcal{C}_{X'}$ the systems of *all* subcontinua of X and X' , respectively. Obviously, then also $f : (X, d, \mathcal{C}) \rightarrow (X', d', \mathcal{C}_{X'})$ is (ϱ, L) -LEL. However, one cannot claim that $f : (X, d, \mathcal{C}_X) \rightarrow (X', d', \mathcal{C}')$ is (ϱ, L) -LEL. In fact, for some spaces (X, d) , (X', d') there is no LEL map $f : (X, d, \mathcal{C}_X) \rightarrow (X', d', \mathcal{C}_{X'})$. For instance this is the case when X is the ω -star and $X' = I$. To show this, suppose that there is a (ϱ, L) -LEL map $f : (X, d, \mathcal{C}_X) \rightarrow (X', d', \mathcal{C}_{X'})$. Take $k \in \mathbb{N}$ such that $\varrho > L/k$ and find a k -star C in X such that every edge of C is mapped onto the same proper subinterval of X' . Then $\mathcal{H}_{d'}^1(f(C)) \leq (L/k) \cdot \mathcal{H}_d^1(C) < \varrho \cdot \mathcal{H}_d^1(C)$, a contradiction.

Our first result says that in the special case when $X = X'$ and $\mathcal{C} = \mathcal{C}'$, LEL maps have interesting dynamical properties. (For the definitions of the corresponding notions, see Section 3.)

Proposition B. *Let $f : (X, d, \mathcal{C}) \rightarrow (X, d, \mathcal{C})$ be a LEL map. Then f is exact and has finite positive entropy. Moreover, if f is the composition $\varphi \circ \psi$ of some maps*

$\psi : X \rightarrow I$ and $\varphi : I \rightarrow X$, then f has the specification property and so it is exactly Devaney chaotic.

The above mentioned tent-like maps $f_k : I \rightarrow I$ (where $k \geq 3$ and I is endowed with the Euclidean metric d_I) are $(\mathcal{C}_I, \mathcal{C}_I, k/2, k)$ -LEL, where \mathcal{C}_I is the system of all non-degenerate closed subintervals of I . Here $k \geq 3$ because the classical tent map f_2 is not $(\mathcal{C}_I, \mathcal{C}_I, \varrho, L)$ -LEL for any $\varrho > 1$ and any L . However, it becomes $(\mathcal{C}_I, \mathcal{C}_I, \varrho, L)$ -LEL (for some $\varrho > 1$ and L) after a slight change of the metric. One can easily construct examples of LEL maps between arbitrary graphs, even in the form of the composition $\varphi \circ \psi$ as in Proposition B; one can use e.g. the maps from [3, Lemma 3.6]. Further, for a given continuum (X, d) of finite length, one can often find $\mathcal{C}, \mathcal{C}'$ and construct LEL-maps $\varphi : (I, d_I, \mathcal{C}_I) \rightarrow (X, d, \mathcal{C})$ and $\psi : (X, d, \mathcal{C}') \rightarrow (I, d_I, \mathcal{C}_I)$. However, it is not so easy to obtain $\mathcal{C}' \supseteq \mathcal{C}$; this inclusion is desirable since then also the composition $\psi \circ \varphi$ is LEL (see Lemma 9).

Our main results, the proofs of which were inspired by [1] and [6], assert that such LEL maps can always be found provided we allow to change the metric on X (the new metric still being compatible with the topology). Recall that a metric d on X is *convex* if for every $x, y \in X$ there is $z \in X$ such that $d(x, z) = d(z, y) = d(x, y)/2$. For two points $a, b \in X$ of a continuum X , $\text{Cut}_X(a, b)$ denotes the set of points $x \in X$ such that a, b lie in different components of $X \setminus \{x\}$.

Theorem C. *For every non-degenerate totally regular continuum X and every $a, b \in X$ we can find a convex metric $d = d_{X,a,b}$ on X and Lipschitz surjections $\varphi_{X,a,b} : I \rightarrow X$, $\psi_{X,a,b} : X \rightarrow I$ with the following properties:*

- (a) $\mathcal{H}_d^1(X) = 1$;
- (b) the system $\mathcal{C} = \mathcal{C}_{X,a,b} = \{\varphi_{X,a,b}(J) : J \text{ is a closed subinterval of } I\}$ is a dense system of subcontinua of X ;
- (c) for every $\varrho > 1$ there are a constant L_ϱ (depending only on ϱ) and (ϱ, L_ϱ) -LEL maps

$$\varphi : (I, d_I, \mathcal{C}_I) \rightarrow (X, d, \mathcal{C}) \quad \text{and} \quad \psi : (X, d, \mathcal{C}) \rightarrow (I, d_I, \mathcal{C}_I)$$

with $\varphi(0) = a$, $\varphi(1) = b$, $\psi(a) = 0$ and such that $\varphi = \varphi_{X,a,b} \circ f_k$, $\psi = f_l \circ \psi_{X,a,b}$ for some $k, l \geq 3$.

Moreover, if $\text{Cut}_X(a, b)$ is uncountable, d, φ, ψ can be assumed to satisfy

- (d) $d(a, b) > 1/2$ and $\psi(b) = 1$.

Theorem D. *Keeping the notation from Theorem C, for every $\varrho > 1$, every non-degenerate totally regular continua X, X' and every points $a, b \in X$, $a', b' \in X'$ there are a constant L_ϱ (depending only on ϱ) and (ϱ, L_ϱ) -LEL map*

$$f : (X, d_{X,a,b}, \mathcal{C}_{X,a,b}) \rightarrow (X', d_{X',a',b'}, \mathcal{C}_{X',a',b'})$$

with $f(a) = a'$ and, provided $\text{Cut}_X(a, b)$ is uncountable, $f(b) = b'$. Moreover, f can be chosen to be the composition $\varphi \circ \psi$ of two LEL-maps $\psi : X \rightarrow I$ and $\varphi : I \rightarrow X'$.

In [1] it was shown that every non-degenerate finite-dimensional Peano continuum admits an exactly Devaney chaotic map and that every finite union of non-degenerate finite-dimensional Peano continua admits a Devaney chaotic map. Theorem D and Proposition B imply the following results which, on one hand, deal with smaller class of spaces, but on the other hand ensure finiteness of the entropy.

Corollary E. *Every non-degenerate totally regular continuum admits an exactly Devaney chaotic map with finite positive entropy and specification.*

Corollary F. *Every finite union of non-degenerate totally regular continua admits a Devaney chaotic map with finite positive entropy.*

In a subsequent paper we deal with the problem of determining the infima of entropies of transitive/exact/(exactly) Devaney chaotic maps on a given totally regular continuum and we show that under some conditions this infimum is zero. The constructions are heavily based on Theorems C and D. To illustrate usefulness of LEL maps let us sketch here an example which shows how easy is to construct a small entropy transitive system on the ω -star.

Example 1. Let X be the ω -star with the branch point a and edges A_i ($i = 1, 2, \dots$); i.e. $X = \bigcup_i A_i$ and $A_i \cap A_j = \{a\}$ for every $i \neq j$. Take arbitrarily large k , put $Y = \bigcup_{i \geq k} A_i$ and define a convex metric d on X in such a way that it coincides with $d_{Y,a,a}$ on Y and each of the sets A_1, \dots, A_{k-1} has length 1. Fix $\varrho > 1$. By Theorem C there are (ϱ, L_ϱ) -maps $f_{k-1} : A_{k-1} \rightarrow Y$, $f_k : Y \rightarrow A_1$ fixing a . Let $f_i : A_i \rightarrow A_{i+1}$ ($i = 1, \dots, k-2$) be isometries fixing a . Then it suffices to define $f : X \rightarrow X$ by $f|_{A_i} = f_i$ for $i < k$ and $f|_Y = f_k$. The map $f^k|_Y : Y \rightarrow Y$ is exact and has dense periodic points by Proposition B; moreover, it is Lipschitz- L_ϱ^2 . So f is Devaney chaotic with entropy $h(f) \leq (2/k) \log L_\varrho$, where L_ϱ does not depend on k .

The paper is organized as follows. In the next section we give an outline of the proofs of Theorems C and D. In Section 3 we recall all the needed definitions and facts. In Section 4 we prove some basic properties of LEL maps. The main part of the paper — Sections 5 and 6 — are devoted to the construction of LEL maps from the unit interval onto a given totally regular continuum and vice versa, see Proposition 25. Finally, in Section 7 we prove the main results of the paper, namely Theorems C, D and Corollaries E, F.

2. OUTLINE OF THE PROOFS OF THEOREMS C AND D

Since the proofs of Theorems C and D consist of a series of lemmas and propositions, for reader's convenience we decided to summarize here the main steps of them. To increase readability we skip some technical details, hence the outline is only “informal” view of the proofs.

In Section 5, for a given totally regular continuum X , we construct a convex metric d and two Lipschitz-1 surjections $g : [0, \alpha] \rightarrow (X, d)$ and $h : (X, d) \rightarrow [0, \beta]$ such that $\mathcal{H}_d^1(X) \leq 1$ and

$$\gamma \cdot |J| \leq \mathcal{H}_d^1(g(J)) \leq \Gamma \cdot |h \circ g(J)|$$

for every closed subinterval J of $[0, \alpha]$, where $0 < \gamma < \Gamma$ are constants not depending on J ; see Lemma 24. The metric d and maps g, h are defined as follows.

- By [6] we can realize X as the inverse limit

$$X = \varprojlim (X_n, f_n)$$

of graphs X_n with monotone surjective bonding maps $f_n : X_{n+1} \rightarrow X_n$ ($n = 1, 2, \dots$), see (5.6). We may assume that for every n there is exactly one point \tilde{x}_n of X_n having non-degenerate f_n -preimage $\tilde{X}_{n+1} = f_n^{-1}(\tilde{x}_n)$.

- The convex metric d is defined by

$$d(x, y) = \sup_{n \in \mathbb{N}} d_n(x_n, y_n) \quad \text{for } x = (x_n)_n, y = (y_n)_n \in X,$$

see (5.15). Here the metric d_1 on X_1 is defined by Lemma 12 and (5.7), and the metrics d_n on X_n ($n \geq 2$) are defined inductively in such a way that d_n “coincides” with d_{n-1} on $X_n \setminus \tilde{X}_n$ and the length of \tilde{X}_n is “very small” when compared to the length of any edge of X_{n-1} , see (5.8)–(5.10).

- In Lemmas 13–15 we prove that d is a convex metric on X compatible with the topology and that $\mathcal{H}_d^1(X) \leq 1$.
- In (5.18) we define $g : [0, \alpha] \rightarrow X$ as the inverse-limit map

$$g = \varprojlim g_n,$$

where $g_n : [0, \alpha_n] \rightarrow X_n$ are natural parametrizations of appropriately chosen paths in X_n ; see (5.11)–(5.14).

- The map $h : X \rightarrow [0, \beta]$ is defined in (5.19) simply by

$$h(x) = d(a, x) \quad \text{for } x \in X,$$

where $a \in X$ is a point fixed in advance.

- In Lemma 24 we summarize the properties of d , g and h .




FIGURE 1. The maps φ and ψ

In Section 6 we construct LEL maps $\varphi : I \rightarrow X$ and $\psi : X \rightarrow I$ basically by linear reparametrizations of g, h , see Proposition 25, Corollary 26 and, for an illustration, Figure 1.

Using the above described tools and results, Theorems C and D can already be easily proved (the proofs themselves can be found in Section 7).

3. PRELIMINARIES

Here we briefly recall all the notions and results which will be needed in the rest of the paper. The terminology is taken mainly from [16, 19, 17, 12].

If M is a set, its cardinality is denoted by $\#M$. The cardinality of infinite countable sets is denoted by \aleph_0 . If M is a singleton set we often identify it with its only point. We write \mathbb{N} for the set of positive integers $\{1, 2, 3, \dots\}$, \mathbb{R} for the set of reals and I for the unit interval $[0, 1]$. By an interval we mean any nonempty connected subset of \mathbb{R} (possibly degenerate to a point). For intervals J, J' we write $J \leq J'$ if $t \leq s$ for every $t \in J, s \in J'$.

By a *space* we mean any nonempty metric space. A space is called *degenerate* provided it has only one point; otherwise it is called *non-degenerate*. If E is a subset of a space $X = (X, d)$ we denote the closure, the interior and the boundary of E by \overline{E} , $\text{int}(E)$ and ∂E , respectively, and we write $d(E)$ for the diameter of E . We say that two sets $E, F \subseteq X$ are *non-overlapping* if they have disjoint interiors. For $x \in X$ and $r > 0$ we denote the closed ball with the center x and radius r by $B(x, r)$. If f is a map defined on X and \mathcal{C} is a system of subsets of X we denote the system $\{f(C) : C \in \mathcal{C}\}$ by $f(\mathcal{C})$.

A (*discrete*) *dynamical system* is a pair (X, f) where $X = (X, d)$ is a compact metric space and $f : X \rightarrow X$ is a continuous map. For $n \in \mathbb{N}$ we denote the composition $f \circ f \circ \dots \circ f$ (n -times) by f^n . A point $x \in X$ is a *periodic point* of f if $f^n(x) = x$ for some $n \in \mathbb{N}$. The *topological entropy* of a dynamical system (X, f) is denoted by $h(f)$. We say that (X, f) is (*topologically*) *transitive* if for every nonempty open sets $U, V \subseteq X$ there is $n \in \mathbb{N}$ such that $f^n(U) \cap V \neq \emptyset$. A system (X, f) is (*topologically*) *exact* or *locally eventually onto* if for every nonempty open subset U of X there is $n \in \mathbb{N}$ such that $f^n(U) = X$. Further, (X, f) is *Devaney chaotic* (*exactly Devaney chaotic*) provided X is infinite, f is transitive (exact) and has dense set of periodic points. Finally, a system (X, f) is said to satisfy the *specification property* if for every $\varepsilon > 0$ there is m such that for every $k \geq 2$, for every k points $x_1, \dots, x_k \in X$, for every integers $a_1 \leq b_1 < \dots < a_k \leq b_k$ with $a_i - b_{i-1} \geq m$ ($i = 2, \dots, k$) and for every integer $p \geq m + b_k - a_1$, there is a point $x \in X$ with $f^p(x) = x$ such that

$$d(f^n(x), f^n(x_i)) \leq \varepsilon \quad \text{for } a_i \leq n \leq b_i, 1 \leq i \leq k.$$

3.1. Continua. A *continuum* is a connected compact metric space. A *cut point* (or a *separating point*) of a continuum X is any point $x \in X$ such that $X \setminus \{x\}$ is disconnected. A point x of a continuum X is called a *local separating point* of X if there is a connected neighborhood U of x such that $U \setminus \{x\}$ is not connected. If a, b are points of X then any cut point of X such that a, b belong to different components of $X \setminus \{x\}$ is said to *separate* a, b . The set of all such points is denoted by $\text{Cut}(a, b)$ or $\text{Cut}_X(a, b)$. If $a = b$ then obviously $\text{Cut}(a, b) = \emptyset$.

Let X be a continuum, let $x \in X$ and let m be a cardinal number. We say that the *order* of x is at most m , written $\text{ord}_X(x) \leq m$, provided X has a local basis of open neighborhoods of x the boundary of which has cardinality at most m . If m is the least such cardinal we write $\text{ord}_X(x) = m$ with one exception: if $m = \aleph_0$ and x has a basis of neighborhoods with finite boundary, we write $\text{ord}_X(x) = \omega$. If $\text{ord}_X(x) = \omega$ or $\text{ord}_X(x)$ is finite we say that x has *finite order* and we write $\text{ord}_X(x) \leq \omega$. The points of order 1 are called *end points*, the points of order 2 are called *ordinary points* and the points of order at least 3 are called *branch points* of X ; the sets of all end, ordinary and branch points are denoted by $E(X)$, $O(X)$ and $B(X)$, respectively.

Tightly connected with the order of a point is the following notion, see e.g. [21]. A point x of a continuum X is said to be of *degree* m , written $\text{deg}_X(x) = m$, provided m is the least cardinal such that for every $\varepsilon > 0$ there exists an *uncountable* family of neighborhoods of x with diameters less than ε , each having the boundary of cardinality at most m and such that for any two neighborhoods U, V either $\overline{U} \subseteq V$ or $\overline{V} \subseteq U$. Again if $m = \aleph_0$ and the neighborhoods can be chosen with finite boundary we write $\text{deg}_X(x) = \omega$ instead of $\text{deg}_X(x) = \aleph_0$. We say that x has

finite degree and write $\deg_X(x) \leq \omega$ if the degree of x is either finite or ω . Trivially always $\text{ord}_X(x) \leq \deg_X(x)$ but there are examples when $\text{ord}_X(x) < \deg_X(x)$; e.g. if X is the Sierpiński triangle then the order of every point $x \in X$ is at most 4 and the degree is equal to the cardinality of the continuum [21].

Let X be a continuum. A metric d on X is said to be *convex* provided for every distinct $x, y \in X$ there is $z \in X$ such that $d(x, z) = d(z, y) = d(x, y)/2$. By [4, Theorem 8] every locally connected continuum admits a compatible convex metric.

3.2. Graphs. An *arc* A in X is any homeomorphic image of the unit interval I ; the end points of A are the images of the points 0, 1. A *simple closed curve* is any homeomorphic image of the unit circle S^1 .

By a *graph* we mean a continuum which can be written as the union of finitely many arcs which are either disjoint or intersect only at their end points. These arcs are called *edges* and their end points are called *vertices* of the graph. So we allow vertices of order 2 and thus the edges and vertices are not defined uniquely. Notice also that we do not allow simple closed curves to be edges of a graph. By a *subgraph* of a graph G we mean any non-degenerate subcontinuum H of G ; so the vertices/edges of H need not be vertices/edges of G .

Let $G = (G, d)$ be a graph with $\mathcal{H}_d^1(G) < \infty$ and let a, b be vertices of G . By a *path* in G from a to b we mean a sequence $\pi = a_0 E_1 a_1 E_2 \dots a_{k-1} E_k a_k$, where a_i ($i = 0, \dots, k$) are vertices of G such that $a_0 = a$, $a_k = b$ and E_j ($j = 1, \dots, k$) are edges of G with end points a_{j-1}, a_j ; the number k will be called the *length* of the path π . A *natural parametrization* of a path $\pi = a_0 E_1 a_1 E_2 \dots a_{k-1} E_k a_k$ is any continuous map $\kappa : J \rightarrow G$ defined on a compact interval $J = [s, t] \subseteq \mathbb{R}$ such that $\kappa(s) = a_0$, $\kappa(t) = a_k$ and we can write $J = J_1 \cup J_2 \cup \dots \cup J_k$ of non-overlapping closed subintervals such that $J_1 \leq J_2 \leq \dots \leq J_k$ and the restriction of $\kappa|_{J_j} : J_j \rightarrow E_j$ is an isometry for every $j = 1, \dots, k$.

3.3. Hausdorff one-dimensional measure and Lipschitz maps. For a Borel subset B of a metric space (X, d) the *one-dimensional Hausdorff measure* of B is defined by

$$\mathcal{H}_d^1(B) = \lim_{\delta \rightarrow 0} \mathcal{H}_{d,\delta}^1(B), \quad \mathcal{H}_{d,\delta}^1(B) = \inf \left\{ \sum_{i=1}^{\infty} d(E_i) : B \subseteq \bigcup_{i=1}^{\infty} E_i, d(E_i) < \delta \right\}.$$

We say that (X, d) has *finite length* if $\mathcal{H}_d^1(X) < \infty$. By e.g. [13, Proposition 4A],

$$(3.1) \quad \mathcal{H}_d^1(C) \geq d(C) \quad \text{whenever } C \text{ is a connected Borel subset of } X.$$

If $A \subseteq X$ is an arc then $\mathcal{H}_d^1(A)$ is equal to the length of A [12, Lemma 3.2]. In the case when (X, d) is the Euclidean real line \mathbb{R} and $J \subset \mathbb{R}$ is an interval $\mathcal{H}_d^1(J)$ is equal to the length of J and we denote it simply by $|J|$.

If (X, d) is a continuum of finite length endowed with a convex metric d , then it has the so-called *geodesic property* (see e.g. [13, Corollary 4E]): for every distinct $x, y \in X$ there is an arc A with end points x, y such that $d(x, y) = \mathcal{H}_d^1(A)$; any such arc A is called a *geodesic arc* or shortly a *geodesic*. Every subarc of a geodesic is again a geodesic. If x, y are the end points of a geodesic A and $z \in A$ then $d(x, y) = d(x, z) + d(z, y)$.

A map $f : (X, d) \rightarrow (Y, \varrho)$ between metric spaces is called *Lipschitz* with a Lipschitz constant $L \geq 0$, shortly *Lipschitz-L*, provided $\varrho(f(x), f(x')) \leq L \cdot d(x, x')$ for every $x, x' \in X$; the smallest such L is denoted by $\text{Lip}(f)$ and is called the

Lipschitz constant of f . If $f : X \rightarrow Y$ is Lipschitz- L then $\mathcal{H}_d^1(f(B)) \leq L \cdot \mathcal{H}_d^1(B)$ for every Borel set $B \subset X$ such that $f(B)$ is Borel-measurable [12, p. 10]. We omit the proof of the following lemma.

Lemma 2. *Let $X = (X, d)$ be a non-degenerate (totally regular) continuum of finite length and let $\varphi : I \rightarrow X$ be a Lipschitz surjection. Put*

$$\mathcal{C} = \varphi(\mathcal{C}_I) = \{\varphi(J) : J \text{ is a non-degenerate closed subinterval of } I\}.$$

Then the following hold:

- (1) \mathcal{C} is a dense system of subcontinua of X ;
- (2) for every $\varepsilon > 0$ the space X can be covered by some $C_1, \dots, C_k \in \mathcal{C}$ satisfying $\mathcal{H}_d^1(C_i) < \varepsilon$ for $i = 1, \dots, k$.

3.4. Totally regular continua. By e.g. [16, 20], a continuum X is called

- a *dendrite* if it is locally connected and contains no simple closed curve;
- a *local dendrite* if it is locally connected and contains at most finitely many simple closed curves;
- *completely regular* if it contains no non-degenerate nowhere dense subcontinuum;
- *totally regular* if for every $x \in X$ and every countable set $P \subseteq X$ there is a basis of neighborhoods of x with finite boundary not intersecting P ;
- *regular* if every $x \in X$ has a basis of neighborhoods with finite boundary, i.e. $\text{ord}_X(x) \leq \omega$ for every x ;
- *hereditarily locally connected* if every subcontinuum of X is locally connected;
- *rational* if every $x \in X$ has a basis of neighborhoods with countable boundary, i.e. $\text{ord}_X(x) \leq \aleph_0$ for every x ;
- a *curve* if it is one-dimensional.

Notice that (local) dendrites as well as completely regular continua are totally regular and (totally) regular continua are hereditarily locally connected, hence they are locally connected curves. Totally regular continua are also called *continua of finite degree* since they are just those continua X for which every point x has finite degree $\text{deg}_X(x) \leq \omega$ [8]. This and other conditions equivalent to total regularity are summarized in the following theorem.

Theorem 3. *For a continuum X the following are equivalent:*

- (1) X is totally regular;
- (2) X is of finite degree (i.e. $\text{deg}_X(x) \leq \omega$ for every x);
- (3) X has a (convex) metric d such that (X, d) has finite length;
- (4) X has a (convex) metric d such that (X, d) is a Lipschitz image of the unit interval;
- (5) X has a (convex) metric d such that for every $x \in X$ and for almost every $r > 0$ the boundary of the closed ball $B(x, r)$ is finite;
- (6) every non-degenerate subcontinuum of X contains uncountably many local separating points;
- (7) X is locally connected and for every disjoint closed sets $E, F \subseteq X$ there are disjoint perfect sets N_1, \dots, N_k such that every subcontinuum of X intersecting both E and F contains some N_i .

Proof. The equivalence of (1), (2), (3), (6) and (7) follows from [8], [21], [10], [15] and [9]. Immediately (4) implies (3) and (5) implies (2). By e.g. [14, Lemma 2A], (3) implies (4). Finally, the fact that (3) implies (5) follows from the following inequality (see e.g. [13, 1A(f)] or [18, Theorem 7.7]) applied to the map $f : X \rightarrow \mathbb{R}$, $f(x') = d(x, x')$. The inequality says that if $f : (X, d) \rightarrow (Y, \rho)$ is Lipschitz-1 then

$$\mathcal{H}_d^1(X) \geq \int_Y^* \#^* f^{-1}(y) d\mathcal{H}_\rho^1(y)$$

where $\#^*$ denotes the cardinality of a finite set and ∞ for infinite sets and $\int_Y^* h d\mu$ is the infimum of integrals $\int_Y g d\mu$ as g runs over μ -measurable functions from Y to $[0, \infty]$ such that $g \geq h$. Hence if (X, d) has finite length then $f^{-1}(y)$ is finite for \mathcal{H}_ρ^1 -almost every $y \in Y$. \square

By [9], if d is a metric on a totally regular continuum X with $\mathcal{H}_d^1(X) < \infty$ then there is a (unique) convex metric d^* on X such that $d^*(x, y) \geq d(x, y)$ for every $x, y \in X$ and $\mathcal{H}_{d^*}^1(B) = \mathcal{H}_d^1(B)$ for every Borel $B \subseteq X$; it is defined by $d^*(x, y) = \inf \{ \mathcal{H}_d^1(A) : A \text{ is an arc from } x \text{ to } y \}$.

3.5. Monotone inverse limits. An *inverse sequence* is a sequence $(X_n, f_n)_{n \in \mathbb{N}}$ where X_n is a compact metric space and $f_n : X_{n+1} \rightarrow X_n$ is a continuous map for every $n \in \mathbb{N}$. The *inverse limit* of an inverse sequence $(X_n, f_n)_{n \in \mathbb{N}}$ is the subspace $X_\infty = \varprojlim (X_n, f_n)$ of the product $\prod_{n=1}^\infty X_n$ given by

$$X_\infty = \varprojlim (X_n, f_n) = \left\{ (x_n)_{n=1}^\infty \in \prod_{n=1}^\infty X_n : f_n(x_{n+1}) = x_n \text{ for every } n \in \mathbb{N} \right\}.$$

The maps f_n are called *bonding maps*. For $n \in \mathbb{N}$ the projection from X_∞ onto the n -th coordinate will be denoted by $\pi_n : X_\infty \rightarrow X_n$. From now on we will assume that every f_n (and hence every π_n) is surjective.

A fundamental result states that the inverse limit of continua is a continuum [19, Theorem 2.1]. Moreover, if the dimension of every X_n is at most d then also $\dim X_\infty \leq d$ [11, Theorem 1.13.4]. Hence the inverse limit of curves is a curve.

The special case important for us is when the bonding maps are monotone. (Recall that a continuous map $f : X \rightarrow Y$ is *monotone* if every preimage $f^{-1}(y)$ is connected.) Then also every projection map π_n is monotone [17, Proposition 2.1.13]. The following theorem combines [17, Corollary 2.1.14], [20, Theorem 3.6] and [19, Theorem 10.36].

Theorem 4. *Let $X_\infty = \varprojlim (X_n, f_n)$ be the inverse limit of continua X_n with surjective monotone bonding maps. If every X_n is locally connected (totally regular, a dendrite) then also X_∞ is locally connected (totally regular, a dendrite).*

It is often the case that a continuum X is homeomorphic to the inverse limit of some ‘‘simpler’’ continua X_n . For example every continuum is the inverse limit of compact connected polyhedra [19, Theorem 2.15] and every curve is the inverse limit of graphs [11, Theorem 1.13.2]. Fundamental results for *monotone* inverse limits and locally connected curves are summarized below, see [20, Theorem 2.2], [6, Theorem 3] and e.g. the proof of [19, Theorem 10.32].

Theorem 5. *Every locally connected curve (totally regular continuum, dendrite) is the monotone inverse limit of regular continua (graphs, trees).*

Notice that for non-degenerate totally regular continua and for non-degenerate dendrites the bonding maps f_n in the previous theorem can be chosen such that $f_n^{-1}(x)$ is non-degenerate for exactly one point x .

The following theorem gives us a way to define the so-called *induced map* between inverse limits, see e.g. [17, Theorems 2.1.46–48].

Theorem 6. *Let $(X_n, f_n)_n$, $(X'_n, f'_n)_n$ be inverse sequences and let $g_n : X_n \rightarrow X'_n$ ($n \in \mathbb{N}$) be continuous maps such that for every n the left-hand side diagram commutes:*

$$\begin{array}{ccc} X_{n+1} & \xrightarrow{f_n} & X_n \\ g_{n+1} \downarrow & & \downarrow g_n \\ X'_{n+1} & \xrightarrow{f'_n} & X'_n \end{array} \qquad \begin{array}{ccc} X_\infty & \xrightarrow{\pi_n} & X_n \\ g_\infty \downarrow & & \downarrow g_n \\ X'_\infty & \xrightarrow{\pi'_n} & X'_n \end{array}$$

Then there is a unique continuous map $g_\infty = \varprojlim g_n : X_\infty \rightarrow X'_\infty$ such that for every n the right-hand side diagram commutes. The map g_∞ is given by

$$g_\infty(x_1, x_2, x_3, \dots) = (g_1(x_1), g_2(x_2), g_3(x_3), \dots).$$

Moreover, if every g_n is surjective (injective) then g_∞ is surjective (injective).

4. PROPERTIES OF LENGTH-EXPANDING LIPSCHITZ MAPS

Here we briefly state basic properties of the class of LEL maps. We start with the proof of Proposition B stated in the introduction.

Proposition B. *Let $f : (X, d, \mathcal{C}) \rightarrow (X, d, \mathcal{C})$ be a LEL map. Then f is exact and has finite positive entropy. Moreover, if f is the composition $\varphi \circ \psi$ of some maps $\psi : X \rightarrow I$ and $\varphi : I \rightarrow X$, then f has the specification property and so it is exactly Devaney chaotic.*

Proof. Let $f : (X, d, \mathcal{C}) \rightarrow (X, d, \mathcal{C})$ be a (ϱ, L) -LEL map. Take any nonempty open subset U of X and fix some $C \in \mathcal{C}$ contained in U . Then $f^n(C) \in \mathcal{C}$ for every n . If $f^n(C) \neq X$ for every n then $\mathcal{H}_d^1(f^n(C)) \geq \varrho^n \cdot \mathcal{H}_d^1(C) \rightarrow \infty$ for $n \rightarrow \infty$, which contradicts the fact that $X = (X, d)$ has finite length. So $f^n(U) \supseteq f^n(C) = X$ for some n , which proves the exactness of f .

Now assume that $f = \varphi \circ \psi$. Since f is exact, also the factor $f' = \psi \circ \varphi : I \rightarrow I$ of f is exact. Hence f' has the specification property by [5]. By [7, 21.4] also f , being a factor of f' , has the specification property. Finally, by [7, 21.3], f has dense periodic points. \square

Recall that d_I denotes the Euclidean metric on I and \mathcal{C}_I is the system of all non-degenerate closed subintervals of I . Note that the following lemma can be substantially generalized, but for our purposes this version is sufficient.

Lemma 7. *Let $k \geq 3$ and $f_k : I \rightarrow I$ be the piecewise linear map fixing 0 and mapping every $[(i-1)/k, i/k]$ onto I . Then $f_k : (I, d_I, \mathcal{C}_I) \rightarrow (I, d_I, \mathcal{C}_I)$ is $(k/2, k)$ -LEL.*

Proof. Only length-expansiveness needs a proof. Take any non-degenerate closed subinterval J of I . If there is i such that $J \supseteq [(i-1)/k, i/k]$ then $f(J) = I$. Otherwise there is i such that $J = J_0 \cup J_1$ where $J_0 \subseteq ((i-1)/k, i/k)$ and $J_1 \subseteq [i/k, (i+1)/k)$. Then $|f_k(J)| \geq k \cdot \max\{|J_0|, |J_1|\} \geq (k/2) \cdot |J|$. \square

Lemma 8. *Let $f : (X, d, \mathcal{C}) \rightarrow (X', d', \mathcal{C}')$ be a (ϱ, L) -LEL map. Then for every $\mathcal{D}' \supseteq \mathcal{C}'$, $1 < \varrho' \leq \varrho$ and $L' \geq L$, the map $f : (X, d, \mathcal{C}) \rightarrow (X', d', \mathcal{D}')$ is (ϱ', L') -LEL.*

Lemma 9. *Let $f : (X, d, \mathcal{C}) \rightarrow (X', d', \mathcal{C}')$ be (ϱ, L) -LEL and $f' : (X', d', \mathcal{C}') \rightarrow (X'', d'', \mathcal{C}'')$ be (ϱ', L') -LEL. Then $f' \circ f : (X, d, \mathcal{C}) \rightarrow (X'', d'', \mathcal{C}'')$ is $(\varrho\varrho', LL')$ -LEL.*

Proof. Put $g = f' \circ f$. Immediately $\text{Lip}(g) \leq LL'$. Take any $C \in \mathcal{C}$ and put $C' = f(C) \in \mathcal{C}'$. If $C' = X'$ then, by surjectivity of f' , $g(C) = X''$. Otherwise $\mathcal{H}_{\varrho'}^1(C') \geq \varrho\mathcal{H}_d^1(C)$ and, if $f'(C') \neq X''$, also $\mathcal{H}_{\varrho''}^1(f'(C')) \geq \varrho'\mathcal{H}_{d'}^1(C')$; hence $\mathcal{H}_{\varrho''}^1(g(C)) \geq \varrho\varrho'\mathcal{H}_d^1(C)$. \square

5. LIPSCHITZ-1 SURJECTIONS $g : [0, \alpha] \rightarrow X$, $h : X \rightarrow [0, \beta]$

In this section we show that for a totally regular continuum X there are a compatible convex metric d and two Lipschitz surjections $g : [0, \alpha] \rightarrow (X, d)$, $h : (X, d) \rightarrow [0, \beta]$ such that

$$\gamma \cdot |J| \leq \mathcal{H}_d^1(g(J)) \leq \Gamma \cdot |h \circ g(J)|$$

for every closed subinterval J of $[0, \alpha]$, where $0 < \gamma < \Gamma$ are constants not depending on J (see Lemma 24).

We start with a simple property of convex metrics on locally connected continua. For a metric space $X = (X, d)$ and a point $a \in X$ put

$$(5.1) \quad h_a : X \rightarrow \mathbb{R}, \quad h_a(x) = d(a, x) \quad \text{for } x \in X.$$

Lemma 10. *Let $X = (X, d)$ be a locally connected continuum endowed with a convex metric d and let $a \in X$. Then*

$$|h_a(A)| \geq \frac{1}{2} \cdot \mathcal{H}_d^1(A)$$

for any free arc A in X .

Proof. Let y, z be the end points of A . For distinct $u, v \in A$ we will denote by uv the subarc of A with end points u, v . Let α be the length of A and let $\kappa : [0, \alpha] \rightarrow A$ be the natural parametrization of A such that $\kappa(0) = y$ and $\kappa(\alpha) = z$. Put $y_t = \kappa(t)$ for $t \in [0, \alpha]$; hence $\mathcal{H}_d^1(y_t y_s) = |s - t|$ for every different $t, s \in [0, \alpha]$.

For every $t \in [0, \alpha]$ such that $y_t \neq a$ take a geodesic arc A_t from a to y_t . Assume first that a is not an interior point of A . Since A is a free arc, every arc (hence also every A_t) from a to a point of A must contain y or z . Take any $t \in [0, \alpha]$. If $y \in A_t$ then $d(a, y_t) = d(a, y) + d(y, y_t) = d(a, y) + t$ since A_t is geodesic. Analogously, if $z \in A_t$ then $d(a, y_t) = d(a, z) + d(z, y_t) = d(a, z) + (\alpha - t)$. So

$$h_a(y_t) = \min\{d(a, y) + t, d(a, z) + (\alpha - t)\}.$$

Hence immediately $|h_a(A)| \geq \alpha/2$.

Now assume that $a = y_s$ for some $s \in (0, \alpha)$; without loss of generality we may assume that $\mathcal{H}_d^1(ay) \leq \mathcal{H}_d^1(az)$. Then for every $t \in [0, \alpha]$, $t \neq s$ the geodesic arc A_t is either the subarc ay_t of A or an arc containing both y and z . Hence

$$h_a(y_t) = \min\{|t - s|, s + d(y, z) + (\alpha - t)\}.$$

So also in this case we easily have $|h_a(A)| \geq \alpha/2$. \square

5.1. Admissible maps on graphs. Let G be a graph with a metric d and let a, b be (not necessarily distinct) vertices of G . We say that a path $\pi = aE_{j_1}a_1 \dots a_{k-1}E_{j_k}b$ in G (from a to b) is *admissible* provided every edge of G is at least once but at most twice in π ; moreover, if a_i is a vertex of G of order 2 then $E_{j_i} \neq E_{j_{i+1}}$ (i.e. π “goes through” the ordinary vertices of G).

A continuous map κ from a compact interval $J = [\alpha, \beta]$ to G is called *fully-admissible* or, more precisely, *fully-admissible for (G, d) from a to b* , if it is the natural parametrization of some admissible path π from a to b . I.e. $\kappa(\alpha) = a$, $\kappa(\beta) = b$ and there is an admissible path $\pi_\kappa = aE_{j_1}a_1 \dots a_{k-1}E_{j_k}b$ and non-overlapping compact intervals $J_1 \leq J_2 \leq \dots \leq J_k$ such that $J = J_1 \cup \dots \cup J_k$ and every restriction $\kappa|_{J_i} : J_i \rightarrow E_{j_i}$ is an isometry.

A map is called *admissible* if it is a restriction of a fully-admissible map onto a compact interval. Notice that any admissible map is finite-to-one and outside of a finite set (the set of points mapped to the vertices of G) is at most two-to-one. Moreover, admissible maps are Lipschitz-1 provided the metric d is convex. The following lemma can be easily proved by induction on the number of edges of G .

Lemma 11. *Let G be a graph and let a, b be vertices of G . Then there is a fully-admissible map $\kappa : [\alpha, \beta] \rightarrow G$ for G from a to b .*

Lemma 12. *Let $0 < q < 1$ and let G be a graph. Then there is a convex metric d on G such that for every admissible map $\kappa : J \rightarrow G$ and every vertex a of G it holds that*

$$\mathcal{H}_d^1(\kappa(J)) \geq \frac{1}{2} \cdot |J| \quad \text{and} \quad |h_a \circ \kappa(J)| \geq \frac{1-q}{6} \cdot \mathcal{H}_d^1(\kappa(J)).$$

Moreover, $|h_a(G)| \geq \frac{1-q}{2} \cdot \mathcal{H}_d^1(G)$.

Proof. Fix any $0 < q < 1$. Let G be a graph and let E_0, \dots, E_k be the edges of G . Take a convex metric d on G such that $\mathcal{H}_d^1(G) < \infty$ and

$$(5.2) \quad \mathcal{H}_d^1(E_i) \leq q \cdot \mathcal{H}_d^1(E_{i-1}) \quad \text{for every } i \geq 1.$$

Such a metric can be constructed as follows: We may assume that G is a subset of \mathbb{R}^3 endowed with the Euclidean metric and that the (Euclidean) lengths of edges of G are finite and exponentially decreasing with quotient q . Then it suffices to take the convex metric on G generated by the Euclidean one.

Let a be a vertex of G and let $\kappa : J \rightarrow G$ be an admissible map for (G, d) ; put $Y = \kappa(J)$. Let $\pi = a_0E_{i_1}a_1E_{i_2} \dots a_{k-1}E_{i_k}a_k$ be the admissible path given by a fully-admissible extension of κ . Since π is admissible, we immediately have $\mathcal{H}_d^1(Y) \geq \frac{1}{2} \cdot |J|$.

Now we show the lower bound for the length of $h_a(Y)$. Realize that there are at most two indices j such that

$$(5.3) \quad Y \cap E_j \text{ is non-degenerate and } Y \not\subseteq E_j$$

(indeed, for any such j the edge E_j must contain the κ -image of an end point of J in its interior). For simplicity we will assume that there are exactly two j 's satisfying (5.3) — we denote them by j_1, j_2 — and that there is an index j such that $E_j \subseteq Y$; the other cases can be described analogously. Let j_0 be the smallest index j such

that $E_j \subseteq Y$. Then using (5.2) we have

$$(5.4) \quad \begin{aligned} \mathcal{H}_d^1(Y) &= \sum_j \mathcal{H}_d^1(E_j \cap Y) \leq \mathcal{H}_d^1(E_{j_1} \cap Y) + \mathcal{H}_d^1(E_{j_2} \cap Y) + \sum_{j \geq j_0} \mathcal{H}_d^1(E_j) \leq \\ &\leq \mathcal{H}_d^1(E_{j_1} \cap Y) + \mathcal{H}_d^1(E_{j_2} \cap Y) + \mathcal{H}_d^1(E_{j_0})/(1-q). \end{aligned}$$

On the other hand, Lemma 10 gives

$$\begin{aligned} |h_a(Y)| &\geq \max\{|h_a(E_{j_1} \cap Y)|, |h_a(E_{j_2} \cap Y)|, |h_a(E_{j_0})|\} \\ &\geq \frac{1}{2} \cdot \max\{\mathcal{H}_d^1(E_{j_1} \cap Y), \mathcal{H}_d^1(E_{j_2} \cap Y), \mathcal{H}_d^1(E_{j_0})\}. \end{aligned}$$

The simple fact that

$$(5.5) \quad \max_{i=1, \dots, p} c_i \leq \sum_{i=1}^p c_i \leq p \cdot \max_{i=1, \dots, p} c_i \quad \text{for any non-negative } c_1, \dots, c_p,$$

applied to (5.4) immediately implies $|h_a(Y)| \geq \mathcal{H}_d^1(Y) \cdot (1-q)/6$.

The final assertion of the lemma follows from the facts that $\mathcal{H}_d^1(G) \leq \mathcal{H}_d^1(E_0)/(1-q)$ and $|h_a(G)| \geq |h_a(E_0)| \geq \mathcal{H}_d^1(E_0)/2$ by Lemma 10. \square

5.2. The construction of d, g, h . Now we embark on the construction of a convex metric d on X and Lipschitz surjections $g : [0, \alpha] \rightarrow X$, $h : X \rightarrow [0, \beta]$ for a given totally regular continuum X (see Lemma 24).

Let $0 < q < 1$, let X be a non-degenerate totally regular continuum and let a, b be two points of X . By [6] there is an inverse sequence $(X_n, f_n)_{n \in \mathbb{N}}$ of graphs X_n with monotone surjective bonding maps $f_n : X_{n+1} \rightarrow X_n$ such that X is (homeomorphic to) the inverse limit

$$(5.6) \quad \varprojlim (X_n, f_n).$$

Without loss of generality we may assume that for every integer $n \geq 1$ the following hold:

- there is $\tilde{x}_n \in X_n$ such that $\tilde{X}_{n+1} = f_n^{-1}(\tilde{x}_n)$ is a non-degenerate subgraph of X_{n+1} ;
- $f_n^{-1}(x)$ is a singleton for every $x \neq \tilde{x}_n$;
- \tilde{x}_n is a vertex of X_n ;
- every vertex of \tilde{X}_{n+1} is a vertex of X_{n+1} ; moreover, every point of the boundary (in X_{n+1}) of \tilde{X}_{n+1} is a vertex of both \tilde{X}_{n+1} and X_{n+1} ; so an edge of \tilde{X}_{n+1} is also an edge of X_{n+1} ;
- the f_n -preimage of every vertex $x \neq \tilde{x}_n$ of X_n is a vertex of X_{n+1} ; so the f_n -image of any edge in X_{n+1} which is not an edge of \tilde{X}_{n+1} is a free arc contained in an edge of X_n .

Let $\pi_n : X \rightarrow X_n$ ($n \in \mathbb{N}$) be the natural projections; put $a_n = \pi_n(a)$, $b_n = \pi_n(b)$. We may assume that a_n, b_n are vertices of X_n and, if $a \neq b$, $a_1 \neq b_1$ (otherwise we remove finitely many of the first X_n 's). Then $a_n \neq b_n$ for every n provided $a \neq b$.

Let d_1 be a convex metric on X_1 obtained using Lemma 12 such that

$$(5.7) \quad \mathcal{H}_{d_1}^1(X_1) = 1 - q$$

and let $g_1 : I_1 \rightarrow X_1$ be a fully-admissible map for (X_1, d_1) from a_1 to b_1 . Assume that $n \geq 2$ and that for every $1 \leq m \leq n-1$ we have defined a metric d_m on X_m , maps $g_m : I_m \rightarrow X_m$ and, provided $m \geq 2$, a map $\varrho_{m-1} : I_m \rightarrow I_{m-1}$. Put

$$(5.8) \quad \mu_{n-1} = \min\{\mathcal{H}_{d_{n-1}}^1(E) : E \text{ is an edge of } X_{n-1}\}.$$

Let \tilde{d}_n be a convex metric on \tilde{X}_n obtained from Lemma 12 such that

$$(5.9) \quad \mathcal{H}_{\tilde{d}_n}^1(\tilde{X}_n) < \frac{q \cdot \mu_{n-1}}{2p} \quad \text{where } p = \#g_{n-1}^{-1}(\tilde{x}_{n-1}).$$

Denote by d_n the only convex metric on X_n such that for every edge E of X_n and every two points $x, y \in E$ the following holds:

$$(5.10) \quad d_n(x, y) = \begin{cases} \tilde{d}_n(x, y) & \text{if } E \subseteq \tilde{X}_n; \\ d_{n-1}(f_{n-1}(x), f_{n-1}(y)) & \text{otherwise.} \end{cases}$$

Let $s_1 < s_2 < \dots < s_p$ be the points of I_{n-1} mapped by g_{n-1} to \tilde{x}_{n-1} . Write I_{n-1} as the union $J'_0 \cup J'_1 \cup \dots \cup J'_p$ of non-overlapping compact subintervals such that $J'_0 \leq s_1 \leq J'_1 \leq s_2 \leq \dots \leq s_p \leq J'_p$ (here J'_0, J'_p can be degenerate). For every $i = 1, \dots, p$ let K'_i be an interval and $\kappa_i : K'_i \rightarrow (\tilde{X}_n, \tilde{d}_n)$ be a fully-admissible map (see Lemma 11); the images of end points of K'_i will be fixed later. Now let $I_n = [0, \alpha_n]$ be a compact interval of length $\alpha_n = |I_{n-1}| + \sum_{i=1}^p |K'_i|$ and define $g_n : I_n \rightarrow X_n$ by “concatenating” the maps

$$g_{n-1}|_{J'_0}, \kappa_1, g_{n-1}|_{J'_1}, \kappa_2, \dots, \kappa_p, g_{n-1}|_{J'_p}.$$

I.e. we write I_n as the union of non-overlapping compact intervals

$$(5.11) \quad I_n = J_0 \cup K_1 \cup J_1 \cup K_2 \cdots \cup K_p \cup J_p$$

such that $J_0 \leq K_1 \leq \dots \leq K_p \leq J_p$ and $|J_i| = |J'_i|$, $|K_j| = |K'_j|$ for every i, j ; then we define g_n such that

$$(5.12) \quad g_n|_{J_i} \approx g_{n-1}|_{J'_i} \quad \text{and} \quad g_n|_{K_j} \approx \kappa_j \quad \text{for every } i, j.$$

(Here we write $f \approx g$ for maps f, g defined on real intervals J, K if there is a constant s_0 such that $J = K + s_0$ and $f(s + s_0) = g(s)$ for every $s \in K$.) By an “appropriate” specification of κ_i -images of the end points of K'_i we obtain that g_n is continuous and that $g_n(0) = a_n$, $g_n(\alpha_n) = b_n$. Notice that

$$(5.13) \quad g_n : I_n \rightarrow (X_n, d_n) \quad \text{is a natural parametrization of some} \\ \text{(not necessarily admissible) path in } X_n \text{ from } a_n \text{ to } b_n.$$

Let $\varrho_{n-1} : I_n \rightarrow I_{n-1}$ be the piecewise linear continuous surjection with slopes 0 and 1 which collapses every K_i into a point. For $1 \leq k < n$ denote the composition $\varrho_k \circ \varrho_{k+1} \circ \dots \circ \varrho_{n-1}$ by $\varrho_{n,k} : I_n \rightarrow I_k$; for convenience put $\varrho_{n,n} = \text{id}_{I_n}$. Analogously define $f_{n,k} : X_n \rightarrow X_k$ for $1 \leq k \leq n$. Notice that the following diagram commutes for every $1 \leq k \leq n$:

$$(5.14) \quad \begin{array}{ccc} X_n & \xrightarrow{f_{n,k}} & X_k \\ g_n \uparrow & & \uparrow g_k \\ I_n & \xrightarrow{\varrho_{n,k}} & I_k \end{array}$$

After finishing the induction we obtain the metrics d_n on X_n and the maps $g_n : I_n \rightarrow X_n$. As in [6] define

$$(5.15) \quad d(x, y) = \sup_{n \in \mathbb{N}} d_n(x_n, y_n) \quad \text{for } x = (x_n)_n, y = (y_n)_n \in X.$$

(In Lemma 13 we will show that d is a convex metric on X .) Define also

$$I_\infty = \varprojlim (I_n, \varrho_n).$$

The corresponding projection map from I_∞ onto I_n ($n \in \mathbb{N}$) will be denoted by π'_n . It is easy to see that the map

$$\eta : I_\infty \rightarrow [0, \alpha], \quad (s_n)_{n \in \mathbb{N}} \mapsto t = \lim_{n \rightarrow \infty} s_n = \sup_{n \in \mathbb{N}} s_n, \quad \text{where } \alpha = \lim_{n \rightarrow \infty} \alpha_n,$$

defines a homeomorphism of I_∞ onto the interval $[0, \alpha]$, which is even isometry if we use the following metric d' on I_∞ (see Lemma 16):

$$(5.16) \quad d'(s, t) = \sup_n |s_n - t_n| \quad \text{for } s = (s_n)_n, t = (t_n)_n \in I_\infty.$$

Since the diagrams in (5.14) commute, the surjective maps $g_n : I_n \rightarrow X_n$ induce the continuous surjective map $g = \varprojlim \{g_n\} : I_\infty \rightarrow X$ between $I_\infty = \varprojlim (I_n, \varrho_n)$ and $X = \varprojlim (X_n, f_n)$ such that the following diagram commutes

$$(5.17) \quad \begin{array}{ccc} X & \xrightarrow{\pi_n} & X_n \\ g \uparrow & & \uparrow g_n \\ I_\infty & \xrightarrow{\pi'_n} & I_n \end{array}$$

(see Theorem 6); the map g is given by

$$(5.18) \quad g(s_1, s_2, s_3, \dots) = (g_1(s_1), g_2(s_2), g_3(s_3), \dots).$$

Finally define $h_n : X_n \rightarrow \mathbb{R}$, $h : X \rightarrow \mathbb{R}$ by

$$(5.19) \quad h_n(x_n) = d_n(x_n, a_n) \quad \text{for } x_n \in X_n, \quad h(x) = d(x, a) \quad \text{for } x \in X.$$

5.3. Properties of the metrics d_n , d . Notice that (5.9) and the fact that every \tilde{X}_n is non-degenerate immediately give

$$(5.20) \quad \mu_n < q \cdot \mu_{n-1} \quad \text{for every } n \geq 2.$$

Since $\mu_1 \leq 1 - q$ by (5.7) we have

$$(5.21) \quad \mu_n \leq q^{n-1} \cdot (1 - q) \quad \text{for every } n \in \mathbb{N}.$$

Lemma 13. *The map d is a convex metric on X compatible with the topology of X .*

Proof. (See [6].) Let $n \geq 2$. From the definition (5.10) of the metrics d_n we have that for every $x, y \in X_n$, $x' = f_{n-1}(x)$, $y' = f_{n-1}(y)$

$$(5.22) \quad d_{n-1}(x', y') \leq d_n(x, y) < d_{n-1}(x', y') + q \cdot \mu_{n-1}$$

and that, for every free arc A in $X_n \setminus \text{int}(\tilde{X}_n)$,

$$(5.23) \quad f_{n-1}|_A : A \rightarrow f_{n-1}(A) \quad \text{is a bijection and } \mathcal{H}_{d_n}^1(A) = \mathcal{H}_{d_{n-1}}^1(f_{n-1}(A)).$$

Combining (5.22) and (5.20) we obtain that for $m > n$

$$d_m(x_m, y_m) < d_n(x_n, y_n) + q \cdot (\mu_{m-1} + \dots + \mu_n) < d_n(x_n, y_n) + \frac{q}{1 - q} \cdot \mu_n,$$

so, since $\mu_n \leq q^{n-1}\mu_1 \leq q^{n-1}(1-q)$,

$$(5.24) \quad d_n(x_n, y_n) \leq d(x, y) \leq d_n(x_n, y_n) + q^n \quad \text{for every } n \in \mathbb{N}, x, y \in X.$$

Hence $d(x, y)$ is always finite. Since trivially d is symmetric, satisfies the triangle inequality and $d(x, y) = 0$ if and only if $x = y$, d defines a metric on X .

To prove that the metric d is compatible with the topology of X it suffices to show that any sequence $(x^{(k)})_k$ converges to x in (X, d) if and only if $(\pi_n x^{(k)})_k$ converges to $\pi_n x$ in (X_n, d_n) for every n . The implication from the left to the right is trivial. Assume that $\lim_k d_n(x_n^{(k)}, x_n) = 0$ for every n , where $x_n^{(k)} = \pi_n x^{(k)}$ and $x_n = \pi_n x$. By (5.24) we have for every n

$$\limsup_{k \rightarrow \infty} d(x^{(k)}, x) \leq \limsup_{k \rightarrow \infty} [d_n(x_n^{(k)}, x_n) + q^n] = q^n.$$

Since $q < 1$ we have that $\limsup_k d(x^{(k)}, x) = 0$.

Now it suffices to show that d is convex. Let $x = (x_n)_n, y = (y_n)_n \in X$. For every n the metric d_n is convex so there is a point $z_n \in X_n$ such that

$$d_n(x_n, z_n) = d_n(y_n, z_n) = \frac{1}{2} \cdot d(x_n, y_n).$$

Let $z^{(n)} \in X$ be such that $\pi_n(z^{(n)}) = z_n$. Compactness of X gives that $(z^{(n)})_n$ has a subsequence $(z^{(n_k)})_k$ converging to a point $z \in X$. Now (5.24) gives

$$\begin{aligned} d(x, z) &\leq d(x, z^{(n_k)}) + d(z^{(n_k)}, z) \leq d_{n_k}(x_{n_k}, z_{n_k}) + q^{n_k} + d(z^{(n_k)}, z) \\ &= \frac{1}{2} \cdot d_{n_k}(x_{n_k}, y_{n_k}) + q^{n_k} + d(z^{(n_k)}, z). \end{aligned}$$

Using (5.24) and taking the limit $k \rightarrow \infty$ and we obtain $d(x, z) \leq \frac{1}{2} \cdot d(x, y)$. Analogously, $d(y, z) \leq \frac{1}{2} \cdot d(x, y)$ and so $d(x, z) = d(y, z) = \frac{1}{2} \cdot d(x, y)$. Hence d is convex. \square

The inequalities (5.24) immediately imply that for the diameters of a subset B of X and its projections $B_n = \pi_n(B)$ ($n \in \mathbb{N}$) it holds that

$$(5.25) \quad d_n(B_n) \leq d(B) \leq d_n(B_n) + q^n.$$

In Lemma 15 we show a relation between the Hausdorff one-dimensional measure of a subset of X and of its projections. To this end we need the following refinement of (5.25) for the special case when B is a subcontinuum of X .

Lemma 14. *Let Y be a subcontinuum of X and let $n \in \mathbb{N}$. Put $Y_n = \pi_n(Y)$ and for every integer $k \geq n$ put $\tilde{y}_k = f_{k,n}(\tilde{x}_k) \in X_n$. Then*

$$d(Y) \leq d_n(Y_n) + q \cdot \sum_{k \geq n, \tilde{y}_k \in Y_n} \mu_k.$$

Proof. For $m > n$ let $N_{m,n}$ be the set of all integers $k \in \{n, n+1, \dots, m-1\}$ such that $\tilde{y}_k \in Y_n$. By the definition (5.15) of d it suffices to show that for every $m > n$

$$(5.26) \quad d_m(Y_m) \leq d_n(Y_n) + q \cdot \sum_{k \in N_{m,n}} \mu_k.$$

We prove this by induction through $m-n$. Assume first that $m-n=1$; then $N_{m,n}$ is either the singleton $\{n\}$ or an empty set according to whether $\tilde{y}_n = \tilde{x}_n$ either belongs to Y_n or not. In the latter case $\tilde{y}_n = \tilde{x}_n \notin Y_n$; since Y_{n+1} is a subgraph of X_{n+1} , (5.23) and convexness of d_{n+1} give that $d_{n+1}(Y_{n+1}) = d_n(Y_n)$. In the former

case (when $\tilde{x}_n \in Y_n$) we analogously have $d_{n+1}(Y_{n+1}) \leq d_n(Y_n) + d_{n+1}(\tilde{X}_{n+1}) \leq d_n(Y_n) + q \cdot \mu_n$. Hence (5.26) is true for any m, n such that $m - n = 1$.

Now assume that for some $p \geq 1$ (5.26) is true whenever $m - n \leq p$; let m, n be such that $m - n = p + 1$. By the induction hypothesis

$$\begin{aligned} d_m(Y_m) &\leq d_{n+1}(Y_{n+1}) + q \cdot \sum_{k \in N_{m, n+1}} \mu_k \\ d_{n+1}(Y_{n+1}) &\leq d_n(Y_n) + q \cdot \sum_{k \in N_{n+1, n}} \mu_k. \end{aligned}$$

Now (5.26) follows since $N_{m, n} = N_{m, n+1} \cup N_{n+1, n}$ (indeed, for $k \geq n + 1$ we have $f_{k, n+1}(\tilde{x}_k) \in Y_{n+1}$ if and only if $f_{k, n}(\tilde{x}_k) \in Y_n$). \square

Lemma 15. *Let $B \subseteq X$ be a closed set and let $B_n = \pi_n(B) \subseteq X_n$ for every $n \in \mathbb{N}$. Then*

$$\mathcal{H}_d^1(B) = \sup_{n \in \mathbb{N}} \mathcal{H}_{d_n}^1(B_n) = \lim_{n \rightarrow \infty} \mathcal{H}_{d_n}^1(B_n).$$

Moreover, $\mathcal{H}_d^1(X) \leq 1$.

Proof. First realize that for every closed (open) set $B \subseteq X$ the set $B_n = \pi_n(B)$ is a closed (open) subset of X_n , hence Borel measurable. Moreover,

$$(5.27) \quad \mathcal{H}_d^1(B) \geq \mathcal{H}_{d_n}^1(B_n) \quad \text{for every } n$$

since $\pi_n|_B : (B, d) \rightarrow (B_n, d_n)$ is Lipschitz-1. We need to show that

$$(5.28) \quad \mathcal{H}_d^1(B) \leq \sup_{n \in \mathbb{N}} \mathcal{H}_{d_n}^1(B_n).$$

We start with the case $B = X$. To this end take any $\delta > 0$ and arbitrary n such that $q^n < \delta/2$. Write X_n as the union $\bigcup_{i=1}^k Y^i$ of non-overlapping subgraphs Y^i such that for every i the diameter $d_n(Y^i)$ is less than $\delta/2$ and the boundary $\partial(Y^i)$ does not contain any $\tilde{y}_k = f_{k, n}(\tilde{x}_k)$ ($k \geq n$). Then $\mathcal{H}_{d_n}^1(X_n) = \sum_{i=1}^k \mathcal{H}_{d_n}^1(Y^i)$. For $1 \leq i \leq k$ put $Z^i = \pi_n^{-1}(Y^i)$; this is a subcontinuum of X since π_n is monotone. By (5.25), $d(Z^i) \leq d_n(Y^i) + q^n < \delta$; since $X = \bigcup_{i=1}^k Z^i$ we have

$$\mathcal{H}_{d, \delta}^1(X) \leq \sum_{i=1}^k d(Z^i).$$

Since for every $k \geq n$ there is just one i such that $\tilde{y}_k \in Y^i$, Lemma 14 gives

$$\sum_{i=1}^k d(Z^i) \leq \sum_{i=1}^k \left(d_n(Y^i) + q \cdot \sum_{k \geq n, \tilde{y}_k \in Y^i} \mu_k \right) \leq \sum_{i=1}^k d_n(Y^i) + q \cdot \sum_{k=n}^{\infty} \mu_k.$$

Thus, using (5.21),

$$\mathcal{H}_{d, \delta}^1(X) \leq \mathcal{H}_{d_n}^1(X_n) + q^n.$$

Since this is true for every sufficiently large n we have $\mathcal{H}_{d, \delta}^1(X) \leq \sup_n \mathcal{H}_{d_n}^1(X_n)$. So (5.28) is proved for $B = X$ (recall that $\mathcal{H}_d^1(X) = \lim_{\delta \rightarrow 0} \mathcal{H}_{d, \delta}^1(X)$).

Now let B be an arbitrary closed subset of X and let $\varepsilon > 0$. Take $n \in \mathbb{N}$ such that $\mathcal{H}_d^1(X) < \mathcal{H}_{d_n}^1(X_n) + \varepsilon$. Put $B_n = \pi_n(B)$ and $C = X \setminus \pi_n^{-1}(B_n) = \pi_n^{-1}(X_n \setminus B_n)$. Then $B \cap C = \emptyset$ and $\mathcal{H}_d^1(C) \geq \mathcal{H}_{d_n}^1(X_n \setminus B_n)$ by (5.27). So

$$\mathcal{H}_d^1(B) \leq \mathcal{H}_d^1(X) - \mathcal{H}_d^1(C) < (\mathcal{H}_{d_n}^1(X_n) + \varepsilon) - \mathcal{H}_{d_n}^1(X_n \setminus B_n) = \mathcal{H}_{d_n}^1(B_n) + \varepsilon.$$

Thus $\mathcal{H}_d^1(X) \leq \sup_n \mathcal{H}_{d_n}^1(B_n) + \varepsilon$. Since $\varepsilon > 0$ was arbitrary (5.28) follows.

Finally, $\mathcal{H}_d^1(X) \leq 1$ follows from (5.28), (5.7) and (5.9). \square

The following lemma describes the basic properties of the inverse limit space $(I_\infty, d') = \varprojlim (I_n, \varrho_n)$ (recall the definition of d' in (5.16)). The fact that I_∞ is an arc follows from general results. Indeed, I_∞ is a locally connected continuum since I_n 's are such and the bonding maps ϱ_n are monotone (see Theorem 4). Moreover, I_∞ is hereditarily unicoherent since every I_n is such [17, 2.1.26]. So I_∞ is a dendrite. Since I_∞ , being an arc-like continuum, does not contain a triod [17, 2.1.41], it must be an arc. In our simple case, however, we can easily prove this fact “from scratch”.

Lemma 16. *Put $\alpha = \sup_n \alpha_n = \lim_n \alpha_n$ (recall that α_n is the length of $I_n = [0, \alpha_n]$). Then $\alpha < \infty$ and the following hold:*

- (a) d' is a metric on I_∞ compatible with the topology;
- (b) for every $s = (s_n)_n, s' = (s'_n)_n \in I_\infty$ we have

$$d'(s, s') = \lim_{n \rightarrow \infty} |s_n - s'_n|$$

- (c) the projection maps $\pi'_n : (I_\infty, d') \rightarrow I_n$ ($n \in \mathbb{N}$) are Lipschitz-1;
- (d) the map

$$\eta : (I_\infty, d') \rightarrow [0, \alpha], \quad s = (s_n)_{n \in \mathbb{N}} \mapsto t = \lim_{n \rightarrow \infty} s_n = \sup_{n \in \mathbb{N}} s_n$$

is an isometry;

- (e) for every subcontinuum J of I_∞ it holds that

$$\mathcal{H}_{d'}^1(J) = \lim_{n \rightarrow \infty} |J_n| = \sup_{n \in \mathbb{N}} |J_n|, \quad \text{where } J_n = \pi'_n(J) \text{ for } n \in \mathbb{N}.$$

Proof. Since $\alpha_n \leq \alpha_{n-1} + 2p \cdot \mathcal{H}_{d_n}^1(\tilde{X}_n) < \alpha_{n-1} + q \cdot \mu_{n-1}$ by (5.9), the finiteness of α follows immediately from (5.20). The assertion (a) can be proved similarly as the fact that d is a compatible metric on X (see Lemma 13). The assertion (b) follows from (5.16) and the fact that every $\varrho_n : I_{n+1} \rightarrow I_n$ is Lipschitz-1 (indeed, if $0 \leq s' < s \leq \alpha_{n+1}$ then $0 \leq \varrho_n(s) - \varrho_n(s') \leq s - s'$). Since (c) is immediate from the definition (5.16) of d' and (e) follows from (d) (indeed, $\mathcal{H}_{d'}^1(J) = |\eta(J)|$), we only need to show (d). First realize that every $s = (s_n)_{n \in \mathbb{N}} \in I_\infty$ is a non-decreasing sequence (in fact, every ϱ_n is Lipschitz-1 and $\varrho_n(0) = 0$) bounded from above by α , hence $\eta(s)$ is well defined. Continuity of η is trivial. Since $\eta(0, 0, \dots) = 0$ and $\eta(\alpha_1, \alpha_2, \dots) = \alpha$, η is surjective. It remains to show that η is an isometry. But this is trivial by (b): for any $s = (s_n)_n, s' = (s'_n)_n \in I_\infty$ we have $d'(s, s') = \lim_n |s_n - s'_n| = |\lim_n s_n - \lim_n s'_n|$. The proof is finished. \square

5.4. Properties of the maps g_n, h_n . In what follows we prove that the lengths of g_n -images and $h_n \circ g_n$ images of any closed interval $J \subseteq I_n$ are bounded from below by some constant multiple of the length of J , where the constant does not depend on n, J . Till the end of this subsection fix $n \in \mathbb{N}$.

Lemma 17. *Let $J \subseteq I_n$ be a compact interval and let $Y = g_n(J)$, $L = h_n(Y)$. If Y contains at most one vertex of X_n then*

$$\mathcal{H}_{d_n}^1(Y) \geq \frac{1}{2} \cdot |J| \quad \text{and} \quad |L| \geq \frac{1}{4} \cdot \mathcal{H}_{d_n}^1(Y).$$

Proof. By the assumption we can write $J = J^{(0)} \cup J^{(1)}$ and $Y = Y^{(0)} \cup Y^{(1)}$ such that $J^{(0)}, J^{(1)}$ are non-overlapping compact intervals and, for $i = 0, 1$, $Y^{(i)} = g_n(J^{(i)})$

is a free arc and $g_n|_{J^{(i)}} : J^{(i)} \rightarrow Y^{(i)}$ is an isometry (see (5.13)). For $i = 0, 1$ put $L^{(i)} = h_n(Y^{(i)})$. Then the inequalities

$$\begin{aligned} \max|J^{(i)}| &\leq |J| \leq 2 \cdot \max|J^{(i)}| \\ \max \mathcal{H}_{d_n}^1(Y^{(i)}) &\leq \mathcal{H}_{d_n}^1(Y) \leq 2 \cdot \max \mathcal{H}_{d_n}^1(Y^{(i)}) \\ \max|L^{(i)}| &\leq |L| \end{aligned}$$

together with Lemma 10 and $\mathcal{H}_{d_n}^1(Y^{(i)}) = |J^{(i)}|$ ($i = 0, 1$) give the assertions of the lemma. \square

Lemma 18. *Let $J \subseteq I_n$ be a compact interval and let $Y = g_n(J)$, $L = h_n(Y)$. Assume that Y contains at least two vertices of X_n and that $Y' = f_{n-1}(Y)$ contains at most one vertex of X_{n-1} . Then*

$$\mathcal{H}_{d_n}^1(Y) \geq \frac{1}{2} \cdot |J| \quad \text{and} \quad |L| \geq \frac{1-q}{12} \cdot \mathcal{H}_{d_n}^1(Y).$$

Proof. Put $Y' = f_{n-1}(Y)$. If $Y \cap \tilde{X}_n$ is empty or degenerate we can proceed as in the proof of the previous lemma, since under this assumption $f_{n-1}|_Y : Y \rightarrow Y'$ is an isometry. So assume that $Y \cap \tilde{X}_n$ is non-degenerate. Put $J' = \varrho_{n-1}(J)$. Since $Y' = f_{n-1}(Y)$ contains at most one vertex of X_{n-1} , it is either degenerate or can be written as the union of two free arcs. Using the fact that just one point (namely the point \tilde{x}_{n-1}) has non-degenerate f_{n-1} -preimage, we have that J and Y can be written as the non-overlapping unions

$$J = J_0 \cup \tilde{J} \cup J_1 \quad \text{and} \quad Y = Y_0 \cup \tilde{Y} \cup Y_1,$$

where (for $i = 0, 1$)

- $J_0 \leq \tilde{J} \leq J_1$ are compact intervals (J_0, J_1 can be degenerate);
- Y_i is either a degenerate subset of X_n or a free arc in $X_n \setminus \text{int}(\tilde{X}_n)$;
- $g_n|_{J_i} : J_i \rightarrow Y_i$ is an isometry;
- $\tilde{Y} = g_n(\tilde{J}) \subseteq \tilde{X}_n$ is non-degenerate;
- $g_n|_{\tilde{J}} : \tilde{J} \rightarrow \tilde{Y}$ is admissible.

(Notice that it can happen that Y_i 's contain more than one vertex of X_n ; however, every vertex contained in the interior of Y_i has order 2, so g_n “goes-through” it, see the definition of an admissible path.)

For $i = 0, 1$ put $L_i = h_n(Y_i)$; we have

$$\mathcal{H}_d^1(Y_i) = |J_i| \quad \text{and} \quad |L_i| \geq \frac{1}{2} \cdot \mathcal{H}_d^1(Y_i).$$

For $\tilde{L} = h_n(\tilde{Y})$ Lemma 12 gives

$$\mathcal{H}_{d_n}^1(\tilde{Y}) \geq \frac{1}{2}|\tilde{J}| \quad \text{and} \quad |\tilde{L}| \geq \frac{1-q}{6} \mathcal{H}_{d_n}^1(\tilde{Y}).$$

Now the first inequality of Lemma 18 follows since

$$\mathcal{H}_{d_n}^1(Y) \geq \mathcal{H}_{d_n}^1(\tilde{Y}) + \max_i \mathcal{H}_{d_n}^1(Y_i) \geq \mathcal{H}_{d_n}^1(\tilde{Y}) + \frac{1}{2} \cdot (\mathcal{H}_{d_n}^1(Y_1) + \mathcal{H}_{d_n}^1(Y_2)).$$

To show the second inequality it suffices to use (5.5) and the fact that either one of J_0, J_1 is degenerate or $\tilde{Y} = \tilde{X}_n$ and, in the latter case, $|\tilde{L}| \geq \frac{1-q}{2} \cdot \mathcal{H}_{d_n}^1(\tilde{Y})$ by Lemma 12. \square

Lemma 19. *Let $J_n \subseteq I_n$ be a compact interval and let $Y_n = g_n(J_n)$, $L_n = h_n(Y_n)$. For $1 \leq m < n$ put*

$$J_m = \varrho_{n,m}(J_n), \quad Y_m = f_{n,m}(Y_n) = g_m(J_m) \quad \text{and} \quad L_m = h_m(Y_m).$$

Let m be such that Y_m contains at least two vertices of X_m . Then

$$|J_m| \leq |J_n| < \frac{1}{1-q} \cdot |J_m|, \quad \mathcal{H}_{d_m}^1(Y_m) \leq \mathcal{H}_{d_n}^1(Y_n) < \frac{1}{1-q} \cdot \mathcal{H}_{d_m}^1(Y_m)$$

and

$$\frac{1-3q}{1-q} \cdot |L_m| < |L_n|.$$

Proof. If $m = n$ there is nothing to prove; hence we may assume that $m \leq n-1$. The first and the third inequalities are immediate consequences of the fact that $\varrho_{n,m}$ and $f_{n,m}$ are Lipschitz-1. By the definition of ϱ_n we have that $|J_n| = |J_{n-1}| + \sum_{i=1}^p |J \cap K_i|$, where the intervals K_1, \dots, K_p are such that every restriction $g_n|_{K_i} : K_i \rightarrow \tilde{X}_n$ is a fully-admissible map, see (5.11) and (5.12). The simple estimate $|J_n \cap K_i| \leq |K_i| \leq 2 \cdot \mathcal{H}_{d_n}^1(\tilde{X}_n)$ together with (5.9) gives that $|J_n| < |J_{n-1}| + q \cdot \mu_{n-1}$. By (5.23) we have that $\mathcal{H}_{d_{n-1}}^1(Y_{n-1}) = \mathcal{H}_{d_n}^1(Y_n \setminus \text{int}(\tilde{X}_n))$; so, again by (5.9), $\mathcal{H}_{d_n}^1(Y_n) < \mathcal{H}_{d_{n-1}}^1(Y_{n-1}) + q \cdot \mu_{n-1}$. Repeating the previous arguments with n replaced by $n-1, n-2, \dots, m+1$ gives that $|J_n| < |J_m| + q \cdot (\mu_{n-1} + \mu_{n-2} + \dots + \mu_m)$ and $\mathcal{H}_{d_n}^1(Y_n) < \mathcal{H}_{d_m}^1(Y_m) + q \cdot (\mu_{n-1} + \mu_{n-2} + \dots + \mu_m)$. Now the second and the fourth inequality immediately follows from (5.20) and the fact that $\mu_m \leq \mathcal{H}_{d_m}^1(Y_m) \leq |J_m|$ since Y_m contains an edge of X_m .

It remains to show the fifth inequality. Let $x', y' \in Y_{n-1}$ be such that $|L_{n-1}| = d_{n-1}(a_{n-1}, x') - d_{n-1}(a_{n-1}, y')$. Take any $x \in Y_n \cap f_{n-1}^{-1}(x')$, $y \in Y_n \cap f_{n-1}^{-1}(y')$. Then, by (5.22),

$$|L_n| \geq d_n(a_n, x) - d_n(a_n, y) > d_{n-1}(a_{n-1}, x') - (d_{n-1}(a_{n-1}, y') + q \cdot \mu_{n-1})$$

so $|L_n| > |L_{n-1}| - q \cdot \mu_{n-1}$. Continue in this fashion to obtain

$$|L_n| > |L_m| - q \cdot (\mu_{n-1} + \mu_{n-2} + \dots + \mu_m) > |L_m| - \frac{q}{1-q} \cdot \mu_m.$$

Since Y_m contains an edge E of X_m , Lemma 10 gives that $|L_m| \geq |h_m(E)| \geq (1/2) \cdot \mathcal{H}_{d_m}^1(E)$, i.e. $|L_m| \geq (1/2) \cdot \mu_m$. So

$$|L_n| > |L_m| - \frac{2q}{1-q} \cdot |L_m| = \frac{1-3q}{1-q} \cdot |L_m|.$$

□

Combining Lemmas 17–19 gives the following estimates.

Lemma 20. *The map $g_n : I_n \rightarrow (X_n, d_n)$ is a Lipschitz-1 surjection. Moreover, for every compact interval $J \subseteq I_n$ and for $Y = g_n(J)$, $L = h_n(Y)$ it holds that*

$$\mathcal{H}_{d_n}^1(Y) \geq \frac{1-q}{2} \cdot |J| \quad \text{and} \quad |L| \geq \frac{1-4q}{12} \cdot \mathcal{H}_{d_n}^1(Y).$$

Proof. The fact that g_n is a Lipschitz-1 surjection is an immediate consequence of (5.13). To prove the second part of the assertion take any non-degenerate compact interval $J \subseteq I_n$ and put $Y = g_n(J)$, $L = h_n(Y)$. If Y contains at most one vertex of X_n we can use Lemma 17. So assume that Y contains at least two vertices of X_n . As in Lemma 19, for every $1 \leq m \leq n$ put $J_m = \varrho_{n,m}(J)$, $Y_m = f_{n,m}(Y_n) = g_m(J)$

and $J_m = h_m(Y)$. Let $m \geq 1$ be the smallest integer such that Y_m contains at least two vertices of X_m . Using Lemma 12 (if $m = 1$) or Lemma 18 (if $m \geq 2$) we obtain that $\mathcal{H}_{d_m}^1(Y_m) \geq \frac{1}{2} \cdot |J_m|$ and $|L_m| \geq \frac{1-q}{12} \cdot \mathcal{H}_{d_m}^1(Y_m)$. Now Lemma 19 gives

$$\frac{\mathcal{H}_{d_n}^1(Y)}{|J|} \geq (1-q) \cdot \frac{\mathcal{H}_{d_m}^1(Y_m)}{|J_m|} \geq \frac{1-q}{2}$$

and

$$\frac{|L|}{\mathcal{H}_{d_n}^1(Y)} \geq (1-3q) \cdot \frac{|L_m|}{\mathcal{H}_{d_m}^1(Y_m)} \geq (1-3q) \cdot \frac{1-q}{12}.$$

The desired inequalities follow. \square

5.5. Properties of the maps g, h . Recall that $g : I_\infty \rightarrow X$ is given by $g(s_1, s_2, \dots) = (g_1(s_1), g_2(s_2), \dots)$.

Lemma 21. *The map $g : (I_\infty, d') \rightarrow (X, d)$ is a Lipschitz-1 surjection.*

Proof. By the definitions of d, d' and the fact that the maps g_n are Lipschitz-1 we have that for every $s = (s_n)_n, t = (t_n)_n \in I_\infty$

$$d(g(s), g(t)) = \sup_n d_n(g_n(s_n), g_n(t_n)) \leq \sup_n |s_n - t_n| = d'(s, t).$$

So g is Lipschitz-1. The surjectivity of g follows from the surjectivity of the maps g_n by Theorem 6. \square

Lemma 22. *Let J be a subcontinuum of I_∞ and let $Y = g(J), L = h(Y)$. Then*

$$\mathcal{H}_d^1(Y) \geq \frac{1-q}{2} \cdot \mathcal{H}_{d'}^1(J) \quad \text{and} \quad |L| \geq \frac{1-4q}{12} \cdot \mathcal{H}_d^1(Y).$$

Proof. For every n put $J_n = \pi'_n(J), Y_n = \pi_n(Y)$ and $L_n = h_n(Y_n)$. By Lemmas 15 and 16 we have

$$\mathcal{H}_{d'}^1(J) = \lim_{n \rightarrow \infty} |J_n| \quad \text{and} \quad \mathcal{H}_d^1(Y) = \lim_{n \rightarrow \infty} \mathcal{H}_{d_n}^1(Y_n).$$

So the first inequality immediately follows from Lemma 20. If we prove that

$$(5.29) \quad |L| = \lim_{n \rightarrow \infty} |L_n|$$

then we analogously obtain the second inequality. (In fact, just an inequality in (5.29) is sufficient.) But (5.29) is an immediate consequence of the definitions of h and d . Indeed,

$$|L| = \sup_{y, y' \in Y} d(a, y) - d(a, y'), \quad |L_n| = \sup_{y, y' \in Y} d_n(a_n, \pi_n(y)) - d_n(a_n, \pi_n(y')),$$

so (5.24) immediately gives (5.29). \square

5.6. Summarization. Here we prove Lemma 24, which is the main result of Section 5. We start with some simple observations concerning cut points.

Lemma 23. *Let $X = \varprojlim (X_n, f_n)$ be the monotone inverse limit of continua and let $\pi_n : X \rightarrow X_n$ ($n \in \mathbb{N}$) be the natural projections. Take any two points $x, y \in X$ and put $x_n = \pi_n(x), y_n = \pi_n(y)$ ($n \in \mathbb{N}$). Then*

- (a) $\pi_n \text{Cut}_X(x, y) \subseteq \text{Cut}_{X_n}(x_n, y_n)$ for every $n \in \mathbb{N}$;
- (b) if $\text{Cut}_X(x, y)$ is uncountable then $\text{Cut}_{X_n}(x_n, y_n)$ is uncountable for every sufficiently large n ;
- (c) if $\text{Cut}_X(x, y)$ is countable and X is rational then every $\text{Cut}_{X_n}(x_n, y_n)$ is countable.

Proof. Put $C = \text{Cut}_X(x, y)$ and $C_n = \text{Cut}_{X_n}(x_n, y_n)$ for $n \in \mathbb{N}$.

(a) If $z \in X$ is such that $z_n = \pi_n z \notin C_n$ then there is a connected subset D_n of $X_n \setminus \{z_n\}$ containing both x_n, y_n . Then $D = \pi_n^{-1}(D_n)$ is a connected subset of $X \setminus \{z\}$ containing both x, y , hence $z \notin C$.

(b) Assume that C is uncountable. For every $z \neq z'$ from X there is $n_{z, z'} \in \mathbb{N}$ such that $\pi_n z \neq \pi_n z'$ for every $n \geq n_{z, z'}$. Since C is uncountable there is n_0 such that $n_{z, z'} = n_0$ for uncountably many pairs of distinct points z, z' from C . Hence, by (a), $C_n \supseteq \pi_n C$ is uncountable for every $n \geq n_0$.

(c) If C_n is uncountable for some n then $D = \pi_n^{-1}(C_n)$ has uncountably many components and every component of D separates x, y . Since X is rational, only countably many components of D are non-degenerate (see [16, Theorem 51.IV.5]) and so uncountably many components of D are singletons. Thus uncountably many points of D separate x, y , i.e. C is uncountable. \square

Lemma 24. *There are constants $0 < \gamma < \Gamma$ such that for any $\delta > 0$ the following hold: For any non-degenerate totally regular continuum X and any two points a, b of X there are a compatible convex metric d on X and maps $g : [0, \alpha] \rightarrow X$, $h : X \rightarrow [0, \beta]$ with the following properties:*

- (a) $g(0) = a$, $g(\alpha) = b$ and $h(a) = 0$;
- (b) g, h are Lipschitz-1 surjections;
- (c) $\gamma \cdot |J| \leq \mathcal{H}_d^1(g(J)) \leq \Gamma \cdot |h \circ g(J)|$ for every closed subinterval J of $[0, \alpha]$;
- (d) $\mathcal{H}_d^1(X) \in [1 - \delta, 1]$, $\mathcal{H}_d^1(X) \leq \alpha \leq 2 \cdot \mathcal{H}_d^1(X)$ and $(1/2 - \delta) \cdot \mathcal{H}_d^1(X) \leq \beta \leq \mathcal{H}_d^1(X)$.

Moreover, if $\text{Cut}(a, b)$ is uncountable then a metric d and maps g, h can be chosen such that also:

- (e) $h(b) = \beta$;
- (f) $d(a, b) > (1 - \delta) \cdot \mathcal{H}_d^1(X)$.

Proof. Let $0 < \gamma < \frac{1}{2}$, $\Gamma > 24$ and $0 < \delta \leq \frac{1}{2}$. Take $0 < q < 1$ such that

$$\frac{1 - q}{2} \geq \gamma, \quad \frac{1 - 4q}{12} \geq \frac{2}{\Gamma} \quad \text{and} \quad 2q < \delta.$$

For a non-degenerate totally regular continuum X and $a, b \in X$ construct a compatible convex metric d on X and maps $g : [0, \alpha] \rightarrow X$, $h : X \rightarrow [0, \beta]$ as in Section 5.2; particularly, $h(x) = d(a, x)$ for $x \in X$ and $\beta = \max_{x \in X} d(a, x)$. By Lemma 21, g is a Lipschitz-1 surjection; since h is such trivially, we have (b). The property (a) is also immediate: $h(a) = d(a, a) = 0$ and $g_n(0) = a_n$, $g_n(\alpha_n) = b_n$ for every n by (5.13), so $g(0) = a$ and $g(\alpha) = b$. The property (c) follows from Lemma 22 and the choice of q .

To finish the proof of the first part we have to show (d). The inequalities $1 - \delta \leq \mathcal{H}_d^1(X) \leq 1$ follow from (5.7) and Lemma 15. Since g, h are Lipschitz-1 surjections we immediately have $\beta \leq \mathcal{H}_d^1(X) \leq \alpha$. Lemma 12 applied to $J = [0, \alpha_n]$, $\kappa = g_n$ and $a = a_n$ gives that

$$\alpha_n \leq 2 \cdot \mathcal{H}_{d_n}^1(X_n) \quad \text{and} \quad |h_n(X_n)| \geq \frac{1 - q}{2} \cdot \mathcal{H}_{d_n}^1(X_n).$$

Thus, by Lemmas 16 and 15,

$$\alpha = \lim_{n \rightarrow \infty} \alpha_n \leq 2 \cdot \lim_{n \rightarrow \infty} \mathcal{H}_{d_n}^1(X_n) = 2 \cdot \mathcal{H}_d^1(X)$$

and

$$\beta = \lim_{n \rightarrow \infty} |h_n(X_n)| \geq \frac{1-q}{2} \cdot \lim_{n \rightarrow \infty} \mathcal{H}_{d_n}^1(X_n) = \frac{1-q}{2} \cdot \mathcal{H}_d^1(X) > \left(\frac{1}{2} - \delta\right) \cdot \mathcal{H}_d^1(X).$$

Now assume that a, b are such that the set $\text{Cut}(a, b)$ of points which separate them is uncountable (hence $a \neq b$). By Lemma 23 for every sufficiently large n the set $\text{Cut}_{X_n}(a_n, b_n)$ of points in X_n separating a_n, b_n is uncountable; without loss of generality we may assume that $\text{Cut}_{X_1}(a_1, b_1)$ is uncountable. Hence there is a free arc A in X_1 such that $X_1 \setminus \text{int}(A)$ has exactly two components, one of which contains a_1 and the other one contains b_1 . We may assume (adding two vertices of order 2 if necessary) that A is an edge of X_1 . We modify the construction of the metric d_1 on X_1 such that the longest edge is $E_0 = A$ (see the proof of Lemma 12). Then

$$d_1(a_1, b_1) \geq \mathcal{H}_{d_1}^1(A) \geq (1-q) \cdot \mathcal{H}_{d_1}^1(X_1).$$

Lemma 15 and (5.23), (5.9), (5.8) give

$$\begin{aligned} \mathcal{H}_d^1(X) &= \lim_{n \rightarrow \infty} \mathcal{H}_{d_n}^1(X_n) = \lim_{n \rightarrow \infty} \left[\mathcal{H}_{d_1}^1(X_1) + \mathcal{H}_{d_2}^1(\tilde{X}_2) + \cdots + \mathcal{H}_{d_n}^1(\tilde{X}_n) \right] \\ &\leq \frac{1}{1-q} \cdot \mathcal{H}_{d_1}^1(X_1). \end{aligned}$$

So, by (5.24),

$$d(a, b) \geq d_1(a_1, b_1) \geq (1-q)^2 \cdot \mathcal{H}_d^1(X) > (1-2q) \cdot \mathcal{H}_d^1(X)$$

and (f) is satisfied.

To obtain also (e) we must replace h by

$$\tilde{h} = \lambda \circ h : X \rightarrow [0, \tilde{\beta}] \quad \text{where } \tilde{\beta} = d(a, b) \text{ and } \lambda(s) = \begin{cases} s & \text{if } s \leq \tilde{\beta}; \\ 2\tilde{\beta} - s & \text{if } \tilde{\beta} < s \leq \beta. \end{cases}$$

Notice that $\tilde{\beta} = d(a, b) > (1-\delta) \cdot \mathcal{H}_d^1(X) \geq (1-\delta) \cdot d(X) \geq \frac{1}{2}\beta$, so $\lambda(s) \in [0, \tilde{\beta}]$ for every $s \in [0, \beta]$. Notice also that $\tilde{\beta} > (\frac{1}{2} - \delta) \cdot \mathcal{H}_d^1(X)$; thus to prove that the triple d, g, \tilde{h} satisfies (a)–(f) we only need to show that $\mathcal{H}_d^1(g(J)) \leq \Gamma \cdot |\tilde{h} \circ g(J)|$ for every closed subinterval J of $[0, \alpha]$, since the other properties are satisfied trivially. To this end fix a closed subinterval $J \subseteq [0, \alpha]$ and put $Y = g(J)$. Using

$$|\tilde{h}(Y)| \geq \max\{|h(Y) \cap [0, \tilde{\beta}]|, |h(Y) \cap [\tilde{\beta}, \beta]|\}$$

and Lemma 22 we immediately have

$$\Gamma \cdot |\tilde{h}(Y)| \geq \frac{\Gamma}{2} \cdot |h(Y)| \geq \frac{12}{1-4q} \cdot |h(Y)| \geq \mathcal{H}_d^1(Y).$$

Hence the proof is finished. \square

6. LENGTH-EXPANDING LIPSCHITZ MAPS FROM/TO THE INTERVAL

The following proposition provides the key tool for constructing LEL maps. Basically it is just a reformulation of Lemma 24.

Proposition 25. *There are constants $0 < \gamma < \Gamma$ and $L > 1$ such that the following hold: For every non-degenerate totally regular continuum X and every two points $a, b \in X$ there are a compatible convex metric d on X and maps $\varphi : I \rightarrow X$, $\psi : X \rightarrow I$ with the following properties:*

- (a) $\varphi(0) = a$, $\varphi(1) = b$ and $\psi(a) = 0$;

- (b) φ, ψ are Lipschitz- L surjections;
- (c) $\gamma \cdot |J| \leq \mathcal{H}_d^1(\varphi(J)) \leq \Gamma \cdot |\psi \circ \varphi(J)|$ for every closed subinterval J of I ;
- (d) $\mathcal{H}_d^1(X) = 1$.

Moreover, if $\text{Cut}(a, b)$ is uncountable then for any $\delta > 0$ a metric d and maps φ, ψ can be chosen such that it also holds:

- (e) $\psi(b) = 1$;
- (f) $d(a, b) > 1 - \delta$.

Proof. Take any $L > 2$ and let $0 < \gamma < \Gamma$ be constants from Lemma 24. Fix a non-degenerate totally regular continuum X , a pair $a, b \in X$ and a positive real δ ; we may assume that $2/(1 - 2\delta) < L$. We give the proof only in the case when $\text{Cut}(a, b)$ is uncountable; the other case can be described analogously.

Let \tilde{d} be a convex metric on X and $g : [0, \alpha] \rightarrow X$, $h : X \rightarrow [0, \beta]$ be maps satisfying (a)–(f) from Lemma 24. Now define $d : X \times X \rightarrow \mathbb{R}$, $\varphi : I \rightarrow X$ and $\psi : X \rightarrow I$ by

$$d(x, y) = \frac{1}{c} \cdot \tilde{d}(x, y), \quad \varphi(t) = g(\alpha t) \quad \text{and} \quad \psi(x) = \frac{1}{\beta} \cdot h(x),$$

where $c = \mathcal{H}_d^1(X)$. Then (a) and (d)–(f) are immediately satisfied. Since

$$\text{Lip}_d(\varphi) = \frac{\alpha}{c} \cdot \text{Lip}_{\tilde{d}}(g) \leq 2 < L \quad \text{and} \quad \text{Lip}_d(\psi) = \frac{c}{\beta} \cdot \text{Lip}_{\tilde{d}}(h) \leq \frac{2}{1 - 2\delta} < L,$$

also (b) is fulfilled. The property (c) follows from

$$\mathcal{H}_d^1(\varphi(J)) \geq \frac{\gamma\alpha}{c} \cdot |J| \quad \text{and} \quad \Gamma \cdot |\psi \circ \varphi(J)| \geq \frac{c}{\beta} \cdot \mathcal{H}_d^1(\varphi(J))$$

and from $\alpha \geq c \geq \beta$. □

Corollary 26. *Every non-degenerate totally regular continuum X , endowed with a suitable convex metric d and a dense systems \mathcal{C} of subcontinua of X , admits LEL-maps $\tilde{\varphi} : (I, d_I, \mathcal{C}_I) \rightarrow (X, d, \mathcal{C})$ and $\tilde{\psi} : (X, d, \mathcal{C}) \rightarrow (I, d_I, \mathcal{C}_I)$.*

Proof. Fix arbitrary $a, b \in X$; let d, φ, ψ be as in Proposition 25. Put $\mathcal{C} = \varphi(\mathcal{C}_I)$; this is a dense system by Lemma 2. Let f_k be the map from Lemma 7, where $k \geq 3$ is such that $\varrho = \gamma k/2 > 1$. Then the map $\tilde{\varphi} = \varphi \circ f_k : I \rightarrow X$ is (ϱ, kL) -LEL. Analogously, if $k' \geq 3$ is such that $\varrho' = k'/(2\Gamma) > 1$ then $\tilde{\psi} = f_{k'} \circ \psi : X \rightarrow I$ is $(\varrho', k'L)$ -LEL. □

Notice that from the proofs of Lemma 24 and Proposition 25 we can see that to fulfill only the conditions (a)–(d) we can find d, φ, ψ such that

$$\psi(x) = c \cdot d(a, x) \quad \text{for every } x \in X,$$

where c is a constant. One can also see that any constants $0 < \gamma < \frac{1}{2}$, $\Gamma > 24$ and $L > 2$ are suitable in Proposition 25. Derivation of the “best” values for γ, Γ and L is out of the scope of this paper. However, we can at least say that L and the ratio Γ/γ cannot be arbitrarily close to 1. In fact, if X is the 3-star then easy arguments show that we must have $\Gamma/\gamma \geq 3$. Further, if $X = (X, d)$ is a simple closed curve of length 1 then, for any $\psi : X \rightarrow I$ from Proposition 25, we can write X as the union $A \cup B$ of two non-overlapping arcs such that $\psi(A) = \psi(B) = I$; so $L \geq \text{Lip}(\psi) \geq 2$.

The following example shows that in the second part of Proposition 25 one cannot replace the assumption $\text{Cut}(a, b)$ is uncountable by $\text{Cut}(a, b)$ is infinite.

Example 27. Take an integer $p \geq 3$, put $a = (-1, 0)$, $b = (1, 0)$, $a_0 = (0, 0)$, $a_k = (1 - 2^{-k}, 0)$, $a_{-k} = -a_k$ ($k \in \mathbb{N}$) and define a continuum $X_p \subseteq \mathbb{R}^2$ by

$$X_p = \bigcup_{k \in \mathbb{Z}} G_k \cup \{a, b\}$$

where every G_k ($k \in \mathbb{Z}$) is a graph with exactly two vertices a_{k-1}, a_k , these vertices have order p (in G_k) and $G_k \cap G_l$ is empty for $l > k + 1$ and is equal to $\{a_k\}$ for $l = k + 1$; see Figure 2 for $p = 3$. In this case a, b are end points of X_p (so $\text{Cut}(a, b)$ is infinite), but neither (e) nor (f) can be fulfilled for small δ .




FIGURE 2. The continuum X_3

To show this realize that $\mathcal{H}_d^1(X_p) \geq p \cdot d(a, b)$ for any convex metric d on X_p ; indeed, X_p is the union of p arcs with ends a, b (so the length of any of them is greater than or equal to $d(a, b)$) and with countable intersections. So immediately (f) is not true for $\delta < 1 - \frac{1}{p}$. Moreover, since ψ is Lipschitz- L and $\psi(a) = 0$ we have that

$$\psi(b) = \psi(b) - \psi(a) \leq L \cdot d(a, b) \leq L \cdot \frac{\mathcal{H}_d^1(X_p)}{p} = \frac{L}{p}$$

which is smaller than 1 for $p > L$. So also (e) is not true. Notice that any metric d satisfying (a)–(d) must be such that the diameter of X_p is approximately p -times larger than the distance of a, b ; so for some k the shortest edge of G_k must be “very small” when compared to the longest one.

Remark 28. If we replace the metric d from Proposition 25 by $d' = c \cdot d$ (where $c > 0$ is a constant), the Lipschitz constants of φ, ψ change to $\text{Lip}_{d'}(\varphi) = c \cdot \text{Lip}_d(\varphi)$, $\text{Lip}_{d'}(\psi) = (1/c) \cdot \text{Lip}_d(\psi)$. So instead of the conditions (d) and (f) we can have

- (d') $\mathcal{H}_d^1(X) < 1/(1 - \delta)$;
- (f') $d(a, b) = 1$.

7. PROOFS OF THE MAIN RESULTS

Now we are ready to prove the main results of the paper stated in the introduction. For convenience we repeat the statements of them.

Theorem C. *For every non-degenerate totally regular continuum X and every $a, b \in X$ we can find a convex metric $d = d_{X,a,b}$ on X and Lipschitz surjections $\varphi_{X,a,b} : I \rightarrow X$, $\psi_{X,a,b} : X \rightarrow I$ with the following properties:*

- (a) $\mathcal{H}_d^1(X) = 1$;
- (b) *the system $\mathcal{C} = \mathcal{C}_{X,a,b} = \{\varphi_{X,a,b}(J) : J \text{ is a closed subinterval of } I\}$ is a dense system of subcontinua of X ;*

(c) for every $\varrho > 1$ there are a constant L_ϱ (depending only on ϱ) and (ϱ, L_ϱ) -LEL maps

$$\varphi : (I, d_I, \mathcal{C}_I) \rightarrow (X, d, \mathcal{C}) \quad \text{and} \quad \psi : (X, d, \mathcal{C}) \rightarrow (I, d_I, \mathcal{C}_I)$$

with $\varphi(0) = a$, $\varphi(1) = b$, $\psi(a) = 0$ and such that $\varphi = \varphi_{X,a,b} \circ f_k$, $\psi = f_l \circ \psi_{X,a,b}$ for some $k, l \geq 3$.

Moreover, if $\text{Cut}_X(a, b)$ is uncountable, d, φ, ψ can be assumed to satisfy

(d) $d(a, b) > 1/2$ and $\psi(b) = 1$.

Proof. Let γ, Γ and L be constants from Proposition 25. Let X be a non-degenerate totally regular continuum and a, b be two points of X . Put $\delta = 1/2$ and fix a metric $d = d_{X,a,b}$ on X and maps $\varphi_{X,a,b} : I \rightarrow X$, $\psi_{X,a,b} : X \rightarrow I$ satisfying (a)–(d) (or (a)–(f) if $\text{Cut}_X(a, b)$ is uncountable) from Proposition 25. Recall that $\mathcal{H}_d^1(X) = 1$ and, provided $\text{Cut}_X(a, b)$ is uncountable, $d(a, b) > 1/2$. By Lemma 2, $\mathcal{C}_{X,a,b} = \varphi_{X,a,b}(\mathcal{C}_I)$ is a dense system of subcontinua of X .

Let $k, l \geq 3$ be the smallest odd integers such that $\gamma k/2 \geq \varrho$ and $l/(2\Gamma) \geq \varrho$. Put $L_\varrho = 2L\varrho \cdot (1 + \max\{\Gamma, 1/\gamma\}) > 1$. As in the proof of Corollary 26, the maps $\varphi = \varphi_{X,a,b} \circ f_k : I \rightarrow X$ and $\psi = f_l \circ \psi_{X,a,b} : X \rightarrow I$ are (ϱ, L_ϱ) -LEL. Since k, l are odd we have $\varphi(0) = \varphi_{X,a,b}(0) = a$, $\varphi(1) = \varphi_{X,a,b}(1) = b$, $\psi(a) = \psi_{X,a,b}(a) = 0$ and, provided $\text{Cut}_X(a, b)$ is uncountable, $\psi(b) = \psi_{X,a,b}(b) = 1$. \square

Theorem D. *Keeping the notation from Theorem C, for every $\varrho > 1$, every non-degenerate totally regular continua X, X' and every points $a, b \in X$, $a', b' \in X'$ there are a constant L_ϱ (depending only on ϱ) and (ϱ, L_ϱ) -LEL map*

$$f : (X, d_{X,a,b}, \mathcal{C}_{X,a,b}) \rightarrow (X', d_{X',a',b'}, \mathcal{C}_{X',a',b'})$$

with $f(a) = a'$ and, provided $\text{Cut}_X(a, b)$ is uncountable, $f(b) = b'$. Moreover, f can be chosen to be the composition $\varphi \circ \psi$ of two LEL-maps $\psi : X \rightarrow I$ and $\varphi : I \rightarrow X'$.

Proof. The theorem follows from Theorem C and Lemma 9. \square

Corollary E. *Every non-degenerate totally regular continuum admits an exactly Devaney chaotic map with finite positive entropy and specification.*

Proof. This immediately follows from Theorem D and Proposition B. \square

Corollary F. *Every finite union of non-degenerate totally regular continua admits a Devaney chaotic map with finite positive entropy.*

Proof. Let $X = \bigsqcup_{i=1}^k X_i$, where X_i 's are non-degenerate totally regular continua. Fix $\varrho > 1$, $a_i \in X_i$ ($i = 1, \dots, k$) and put $d_i = d_{X_i, a_i, a_i}$, $\mathcal{C}_i = \mathcal{C}_{X_i, a_i, a_i}$. Let $f_i : X_i \rightarrow X_{i+1}$ ($i = 1, \dots, k-1$) and $f_k : X_k \rightarrow X_1$ be LEL maps from Theorem D. Finally, let d be the metric on X such that $d(x, y) = d_i(x, y)$ for any $x, y \in X_i$ ($i = 1, \dots, k$) and $d(x, y) = 2$ for $x \in X_i, y \in X_j$ ($i \neq j$). Since $d_i(X_i) \leq 1$, the metric d is compatible with the topology of X .

Define $f : X \rightarrow X$ by $f|_{X_i} = f_i$ for $i = 1, \dots, k$. For every i the restriction $f^k|_{X_i} : X_i \rightarrow X_i$ is LEL, hence it is exactly Devaney chaotic with positive finite entropy and specification by Proposition B. Since f permutes X_1, \dots, X_k , the assertion follows. \square

Acknowledgment. The author wishes to express his thanks to Lubomír Snoha for his help with the preparation of the paper. The author was supported by the Slovak

Research and Development Agency under the contract No. APVV-0134-10 and by the Slovak Grant Agency under the grants VEGA 1/0855/08 and VEGA 1/0978/11.

REFERENCES

- [1] S. Agronsky and J. G. Ceder, *Each Peano subspace of E^k is an ω -limit set*, Real Anal. Exchange **17** (1991/92), no. 1, 371–378.
- [2] L. Alsedà, S. Kolyada, J. Llibre and E. Snoha, *Entropy and periodic points for transitive maps*, Trans. Amer. Math. Soc. **351** (1999), no. 4, 1551–1573.
- [3] L. Alsedà, M. A. del Río and J. A. Rodríguez, *A splitting theorem for transitive maps*, J. Math. Anal. Appl. **232** (1999), no. 2, 359–375.
- [4] R. H. Bing, *Partitioning a set*, Bull. Amer. Math. Soc. **55** (1949), 1101–1110.
- [5] A. M. Blokh, *Decomposition of dynamical systems on an interval*, Russ. Math. Surv. **38** (1983), 133–134.
- [6] R. D. Buskirk, J. Nikiel and E. D. Tymchatyn, *Totally regular curves as inverse limits*, Houston J. Math. **18** (1992), no. 3, 319–327.
- [7] M. Denker, C. Grillenberger and K. Sigmund, *Ergodic theory on compact spaces*, Springer-Verlag, Berlin, 1976.
- [8] S. Eilenberg, *On continua of finite length*, Ann. Soc. Pol. Math. **17** (1938), 253–254.
- [9] S. Eilenberg, *Continua of finite linear measure II*, Amer. J. Math. **66** (1944), 425–427.
- [10] S. Eilenberg and O. G. Harrold, *Continua of finite linear measure I*, Amer. J. Math. **65** (1943), 137–146.
- [11] R. Engelking, *Dimension theory*, North-Holland and PWN—Polish Scientific Publishers, Warsaw, 1978.
- [12] K. J. Falconer, *The geometry of fractal sets*, Cambridge University Press, Cambridge, 1986.
- [13] D. H. Fremlin, *Spaces of finite length*, Proc. London Math. Soc. (3) **64** (1992), 449–486.
- [14] D. H. Fremlin, *Embedding spaces of finite length in \mathbb{R}^3* , J. London Math. Soc. (2) **49** (1994), 150–162.
- [15] O. G. Harrold, *The construction of a certain metric*, Duke Math. J. **11** (1944), 23–34.
- [16] K. Kuratowski, *Topology, vol. 2*, Academic Press and PWN, Warszawa, 1968.
- [17] S. Macías, *Topics on continua*, Chapman & Hall/CRC, Boca Raton, FL, 2005.
- [18] P. Mattila, *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability*, Cambridge University Press, Cambridge, 1995.
- [19] S. B. Nadler, *Continuum theory. An introduction*, Monographs and Textbooks in Pure and Applied Mathematics, 158, Marcel Dekker, Inc., New York, 1992.
- [20] J. Nikiel, *Locally connected curves viewed as inverse limits*, Fund. Math. **133** (1989), no. 2, 125–134.
- [21] G. T. Whyburn, *Concerning continua of finite degree and local separating points*, Amer. J. Math. **57** (1935), no. 1, 11–16.

DEPARTMENT OF MATHEMATICS, FACULTY OF NATURAL SCIENCES, MATEJ BEL UNIVERSITY,
TAJOVSKÉHO 40, 974 01 BANSKÁ BYSTRICA, SLOVAKIA
E-mail address: vladimir.spitalsky@umb.sk