Spectra originated from semi-B-Fredholm theory and commuting perturbations *

Qingping ZENG,[†] Qiaofen JIANG and Huaijie ZHONG

(School of Mathematics and Computer Science, Fujian Normal University, Fuzhou 350007, P.R. China)

Abstract: In [19], Burgos, Kaidi, Mbekhta and Oudghiri provided an affirmative answer to a question of Kaashoek and Lay and proved that an operator F is power finite rank if and only if $\sigma_{dsc}(T + F) = \sigma_{dsc}(T)$ for every operator T commuting with F. Later, several authors extended this result to the essential descent spectrum, the left Drazin spectrum and the left essentially Drazin spectrum. In this paper, using the theory of operator with eventual topological uniform descent and the technique used in [19], we generalize this result to various spectra originated from seni-B-Fredholm theory. As immediate consequences, we give affirmative answers to several questions posed by Berkani, Amouch and Zariouh. Besides, we provide a general framework which allows us to derive in a unify way commuting perturbational results of Weyl-Browder type theorems and properties (generalized or not). These commuting perturbational results, in particular, improve many recent results of [11, 14, 17, 18, 38] by removing certain extra assumptions.

2010 Mathematics Subject Classification: primary 47A10, 47A11; secondary 47A53, 47A55

Key words: Semi-B-Fredholm operators; eventual topological uniform descent; power finite rank; commuting perturbation.

1 Introduction

In 1972, Kaashoek and Lay have shown in [31] that the descent spectrum is invariant under commuting power finite rank perturbation F (that is, F^n is finite rank for some $n \in \mathbb{N}$). Also they have conjectured that this perturbation property characterizes such operators F. In 2006, Burgos, Kaidi, Mbekhta and Oudghiri provided in [19] an affirmative answer to this question and proved that an operator F is power finite rank if and only if $\sigma_{dsc}(T + F) = \sigma_{dsc}(T)$ for every operator T commuting with F. Later,

^{*}This work has been supported by National Natural Science Foundation of China (11171066), Specialized Research Fund for the Doctoral Program of Higher Education (2010350311001, 20113503120003), Natural Science Foundation of Fujian Province (2009J01005, 2011J05002) and Foundation of the Education Department of Fujian Province (JB10042, JA08036).

[†]Email address: zqpping2003@163.com.

Fredj generalized this result in [24] to the essential descent spectrum. Fredj, Burgos and Oudghiri extended this result in [25] to the left Drazin spectrum and the left essentially Drazin spectrum. The present paper is concern with commuting power finite rank perturbations of semi-B-Fredholm operators. As seen in Theorem 2.11 (i.e., main result), we generalize the previous results to various spectra originated from semi-B-Fredholm theory. The proof of our main result is mainly dependent upon the theory of operator with eventual topological uniform descent and the technique used in [19].

Spectra originated from semi-B-Fredholm theory include, in particular, the upper semi-B-Weyl spectrum σ_{USBW} (resp. the B-Weyl spectrum σ_{BW}) which is closely related to generalized a-Weyl's theorem, generalized a-Browder's theorem, property (gw) and property (gb) (resp. generalized Weyl's theorem, generalized Browder's theorem, property (gaw) and property (gab)). Concerning the upper semi-B-Weyl spectrum σ_{USBW} , Berkani and Amouch posed in [11] the following question:

Question 1.1. Let $T \in \mathcal{B}(X)$ and let $N \in \mathcal{B}(X)$ be a nilpotent operator commuting with T. Do we always have

$$\sigma_{USBW}(T+N) = \sigma_{USBW}(T) ?$$

Similarly, for the B-Weyl spectrum σ_{BW} , Berkani and Zariouh posed in [17] the following question:

Question 1.2. Let $T \in \mathcal{B}(X)$ and let $N \in \mathcal{B}(X)$ be a nilpotent operator commuting with T. Do we always have

$$\sigma_{BW}(T+N) = \sigma_{BW}(T) ?$$

Recently, Amouch, Zguitti, Berkani and Zariouh have given partial answers in [5, 7, 11, 14] to Question 1.1. As immediate consequences of our main result (see Theorem 2.11), we provide positive answers to Questions 1.1 and 1.2 and some other questions posed by Berkani and Zariouh (see Corollaries 3.1, 3.3 and 3.8). Besides, we provide a general framework which allows us to derive in a unify way commuting perturbational results of Weyl-Browder type theorems and properties (generalized or not). These commuting perturbational results, in particular, improve many recent results of [11, 14, 17, 18, 38] by removing certain extra assumptions (see Corollary 3.9 and Remark 3.10).

Throughout this paper, let $\mathcal{B}(X)$ denote the Banach algebra of all bounded linear operators acting on an infinite dimensional complex Banach space X, and $\mathcal{F}(X)$ denote its ideal of finite rank operators on X. For an operator $T \in \mathcal{B}(X)$, let T^* denote its dual, $\mathcal{N}(T)$ its kernel, $\alpha(T)$ its nullity, $\mathcal{R}(T)$ its range, $\beta(T)$ its defect, $\sigma(T)$ its spectrum and $\sigma_a(T)$ its approximate point spectrum. If the range $\mathcal{R}(T)$ is closed and $\alpha(T) < \infty$ (resp. $\beta(T) < \infty$), then T is said to be *upper semi-Fredholm* (resp. *lower semi-Fredholm*). If $T \in \mathcal{B}(X)$ is both upper and lower semi-Fredholm, then T is said to be *Fredholm*. If $T \in \mathcal{B}(X)$ is either upper or lower semi-Fredholm, then T is said to be *semi-Fredholm*, and its index is defined by $\operatorname{ind}(T) = \alpha(T) - \beta(T)$. For each $n \in \mathbb{N}$, we set $c_n(T) = \dim \mathcal{R}(T^n)/\mathcal{R}(T^{n+1})$ and $c'_n(T) = \dim \mathcal{N}(T^{n+1})/\mathcal{N}(T^n)$. It follows from [30, Lemmas 3.1 and 3.2] that, for every $n \in \mathbb{N}$,

$$c_n(T) = \dim X/(\mathcal{R}(T) + \mathcal{N}(T^n)), \quad c'_n(T) = \dim \mathcal{N}(T) \cap \mathcal{R}(T^n).$$

Hence, it is easy to see that the sequences $\{c_n(T)\}_{n=0}^{\infty}$ and $\{c'_n(T)\}_{n=0}^{\infty}$ are decreasing. Recall that the *descent* and the *ascent* of $T \in \mathcal{B}(X)$ are $dsc(T) = \inf\{n \in \mathbb{N} : \mathcal{R}(T^n) = \mathcal{R}(T^{n+1})\}$ and $asc(T) = \inf\{n \in \mathbb{N} : \mathcal{N}(T^n) = \mathcal{N}(T^{n+1})\}$, respectively (the infimum of an empty set is defined to be ∞). That is,

$$dsc(T) = \inf\{n \in \mathbb{N} : c_n(T) = 0\}$$

and

$$asc(T) = \inf\{n \in \mathbb{N} : c'_n(T) = 0\}.$$

Similarly, the esential descent and the esential ascent of $T \in \mathcal{B}(X)$ are

$$dsc_e(T) = \inf\{n \in \mathbb{N} : c_n(T) < \infty\}$$

and

$$asc_e(T) = \inf\{n \in \mathbb{N} : c'_n(T) < \infty\}.$$

If $asc(T) < \infty$ and $\mathcal{R}(T^{asc(T)+1})$ is closed, then T is said to be left Drazin invertible. If $dsc(T) < \infty$ and $\mathcal{R}(T^{dsc(T)})$ is closed, then T is said to be right Drazin invertible. If $asc(T) = dsc(T) < \infty$, then T is said to be Drazin invertible. Clearly, $T \in \mathcal{B}(X)$ is both left and right Drazin invertible if and only if T is Drazin invertible. If $asc_e(T) < \infty$ and $\mathcal{R}(T^{asc_e(T)+1})$ is closed, then T is said to be left essentially Drazin invertible. If $dsc_e(T) < \infty$ and $\mathcal{R}(T^{dsc_e(T)})$ is closed, then T is said to be right essentially Drazin invertible.

For $T \in \mathcal{B}(X)$, let us define the *left Drazin spectrum*, the *right Drazin spectrum*, the *Drazin spectrum*, the *left essentially Drazin spectrum*, and the *right essentially Drazin spectrum* of T as follows respectively:

 $\sigma_{LD}(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a left Drazin invertible operator}\};$

 $\sigma_{RD}(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a right Drazin invertible operator}\};$

 $\sigma_D(T) = \{ \lambda \in \mathbb{C} : T - \lambda I \text{ is not a Drazin invertible operator} \};$

 $\sigma_{LD}^e(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a left essentially Drazin invertible operator}\};$

 $\sigma_{RD}^e(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a right essentially Drazin invertible operator}\}.$

These spectra have been extensively studied by several authors, see e.g [2, 7, 8, 9, 22, 24, 25, 33].

Recall that an operator $T \in \mathcal{B}(X)$ is said to be *Browder* (resp. *upper semi-Browder*, *lower semi-Browder*) if T is Fredholm and $asc(T) = dsc(T) < \infty$ (resp. T is upper semi-Fredholm and $asc(T) < \infty$, T is lower semi-Fredholm and $dsc(T) < \infty$).

For each integer n, define T_n to be the restriction of T to $\mathcal{R}(T^n)$ viewed as the map from $\mathcal{R}(T^n)$ into $\mathcal{R}(T^n)$ (in particular $T_0 = T$). If there exists $n \in \mathbb{N}$ such that $\mathcal{R}(T^n)$ is closed and T_n is Fredholm (resp. upper semi-Fredholm, lower semi-Fredholm, Browder, upper semi-Browder, lower semi-Browder), then T is called *B*-Fredholm (resp. upper semi-B-Fredholm, lower semi-B-Fredholm, B-Browder, upper semi-B-Browder, lower semi-B-Browder). If $T \in \mathcal{B}(X)$ is upper or lower semi-B-Browder, then T is called semi-B-Browder. If $T \in \mathcal{B}(X)$ is upper or lower semi-B-Fredholm, then T is called semi-B-Fredholm. It follows from [13, Proposition 2.1] that if there exists $n \in \mathbb{N}$ such that $\mathcal{R}(T^n)$ is closed and T_n is semi-Fredholm, then $\mathcal{R}(T^m)$ is closed, T_m is semi-Fredholm and $\operatorname{ind}(T_m) = \operatorname{ind}(T_n)$ for all $m \geq n$. This enables us to define the index of a semi-B-Fredholm operator T as the index of the semi-Fredholm. An operator T_n , where n is an integer satisfying $\mathcal{R}(T^n)$ is closed and T_n is semi-Fredholm. An operator $T \in \mathcal{B}(X)$ is called B-Weyl (resp. upper semi-B-Weyl, lower semi-B-Weyl) if T is B-Fredholm and $\operatorname{ind}(T) = 0$ (resp. T is upper semi-B-Fredholm and $\operatorname{ind}(T) \leq 0$, T is lower semi-B-Fredholm and $\operatorname{ind}(T) \geq 0$). If $T \in \mathcal{B}(X)$ is upper or lower semi-B-Weyl, then T is called semi-B-Weyl.

For $T \in \mathcal{B}(X)$, let us define the upper semi-B-Fredholm spectrum, the lower semi-B-Fredholm spectrum, the semi-B-Fredholm spectrum, the B-Fredholm spectrum, the upper semi-B-Weyl spectrum, the lower semi-B-Weyl spectrum, the semi-B-Weyl spectrum, the B-Weyl spectrum, the upper semi-B-Browder spectrum, the lower semi-B-Browder spectrum, the semi-B-Browder spectrum, and the B-Browder spectrum of T as follows respectively:

$$\begin{split} \sigma_{USBF}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a upper semi-B-Fredholm operator}\};\\ \sigma_{LSBF}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a lower semi-B-Fredholm operator}\};\\ \sigma_{SBF}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Fredholm operator}\};\\ \sigma_{BF}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a B-Fredholm operator}\};\\ \sigma_{USBW}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a upper semi-B-Weyl operator}\};\\ \sigma_{LSBW}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a lower semi-B-Weyl operator}\};\\ \sigma_{SBW}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Weyl operator}\};\\ \sigma_{BW}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Weyl operator}\};\\ \sigma_{USBB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a B-Weyl operator}\};\\ \sigma_{USBB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a upper semi-B-Browder operator}\};\\ \sigma_{LSBB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a lower semi-B-Browder operator}\};\\ \sigma_{BB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Browder operator}\};\\ \sigma_{BB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Browder operator}\};\\ \sigma_{BB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Browder operator}\};\\ \sigma_{BB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Browder operator}\};\\ \sigma_{BB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Browder operator}\};\\ \sigma_{BB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Browder operator}\};\\ \sigma_{BB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Browder operator}\};\\ \sigma_{BB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Browder operator}\};\\ \sigma_{BB}(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a semi-B-Browder operator}\}.\\ \end{array}$$

These spectra originated from semi-B-Fredholm theory also have been extensively studied by several authors, see e.g [2, 7, 8, 10, 13, 17, 22].

For any $T \in \mathcal{B}(X)$, Berkani have found in [8, Theorem 3.6] the following elegant equalities:

$$\sigma_{LD}(T) = \sigma_{USBB}(T), \quad \sigma_{RD}(T) = \sigma_{LSBB}(T);$$

$$\sigma_{LD}^{e}(T) = \sigma_{USBF}(T), \quad \sigma_{RD}^{e}(T) = \sigma_{LSBF}(T);$$
$$\sigma_{D}(T) = \sigma_{BB}(T).$$

This paper is organized as follows. In Section 2, by using the theory of operator with eventual topological uniform descent and the technique used in [19], we characterize power finite rank operators via various spectra originated from seni-B-Fredholm theory. In Section 3, as some applications, we provide affirmative answers to some questions of Berkani, Amouch and Zariouh. Besides, we provide a general framework which allows us to derive in a unify way commuting perturbational results of Weyl-Browder type theorems and properties (generalized or not). These commuting perturbational results, in particular, improve many recent results of [11, 14, 17, 18, 38] by removing certain extra assumptions.

2 Main result

We begin with the following lemmas in order to give the proof of the main result in this paper.

Lemma 2.1. Let $F \in \mathcal{B}(X)$ with $F^n \in \mathcal{F}(X)$ for some $n \in \mathbb{N}$. If $T \in \mathcal{B}(X)$ is upper semi-B-Fredholm and commutes with F, then T + F is also upper semi-B-Fredholm.

Proof. Since T is upper semi-B-Fredholm, by [8, Theorem 3.6], T is left essentially Drazin invertible. Hence by [25, Proposition 3.1], T + F is left essentially Drazin invertible. By [8, Theorem 3.6] again, T is upper semi-B-Fredholm.

Lemma 2.2. Let $F \in \mathcal{B}(X)$ with $F^n \in \mathcal{F}(X)$ for some $n \in \mathbb{N}$. If $T \in \mathcal{B}(X)$ is lower semi-B-Fredholm and commutes with F, then T + F is also lower semi-B-Fredholm.

Proof. Since $F^n \in \mathcal{F}(X)$ for some $n \in \mathbb{N}$, $\mathcal{R}(F^n)$ is a closed and finite-dimensional subspace, and hence dim $\mathcal{R}(F^{*n}) = \dim \mathcal{N}(F^n)^{\perp} = \dim \mathcal{R}(F^n)$, thus $\mathcal{R}(F^{*n})$ is finitedimensional, this infers that $F^{*n} \in \mathcal{F}(X^*)$. It is obvious that T^* commutes with F^* . Since T is lower semi-B-Fredholm, by [8, Theorem 3.6], T is right essentially Drazin invertible. Then from the presentation before Section IV of [33], it follows that T^* is left essentially Drazin invertible. Hence by [25, Proposition 3.1], $(T + F)^* = T^* + F^*$ is left essentially Drazin invertible. From the presentation before Section IV of [33] again, it follows that T + F is right essentially Drazin invertible. Consequently, by [8, Theorem 3.6] again, T + F is lower semi-B-Fredholm.

It follows from [8, Corollary 3.7 and Theorem 3.6] that T is B-Fredholm if and only if T is both upper and lower semi-B-Fredholm.

Corollary 2.3. Let $T \in \mathcal{B}(X)$ and let $F \in \mathcal{B}(X)$ with $F^n \in \mathcal{F}(X)$ for some $n \in \mathbb{N}$. If T commutes with F, then

- (1) $\sigma_{USBF}(T+F) = \sigma_{USBF}(T);$
- (2) $\sigma_{LSBF}(T+F) = \sigma_{LSBF}(T);$
- (3) $\sigma_{SBF}(T+F) = \sigma_{SBF}(T);$
- (4) $\sigma_{BF}(T+F) = \sigma_{BF}(T).$

Proof. From Lemma 2.1, the first equation follows easily. The second equation follows immediately from Lemma 2.2. The third equation is true because $\sigma_{SBF}(T) = \sigma_{USBF}(T) \cap \sigma_{LSBF}(T)$, for every $T \in \mathcal{B}(X)$. The fourth equation is also true because $\sigma_{BF}(T) = \sigma_{USBF}(T) \cup \sigma_{LSBF}(T)$, for every $T \in \mathcal{B}(X)$.

To continue the discussion of this paper, we recall some classical definitions. Using the isomorphism $X/N(T^d) \approx R(T^d)$ and following [27], a topology on $R(T^d)$ is defined as follows.

Definition 2.4. Let $T \in \mathcal{B}(X)$. For every $d \in \mathbb{N}$, the operator range topological on $R(T^d)$ is defined by the norm $||\cdot||_{R(T^d)}$ such that for all $y \in R(T^d)$,

$$||y||_{R(T^d)} = \inf\{||x|| : x \in X, y = T^d x\}.$$

For a detailed discussion of operator ranges and their topologies, we refer the reader to [23] and [26]. If $T \in \mathcal{B}(X)$, for each $n \in \mathbb{N}$, T induces a linear transformation from the vector space $\mathcal{R}(T^n)/\mathcal{R}(T^{n+1})$ to the space $\mathcal{R}(T^{n+1})/\mathcal{R}(T^{n+2})$. We will let $k_n(T)$ be the dimension of the null space of the induced map. From [27, Lemma 2.3] it follows that, for every $n \in \mathbb{N}$,

$$k_n(T) = \dim(\mathcal{N}(T) \cap \mathcal{R}(T^n)) / (\mathcal{N}(T) \cap \mathcal{R}(T^{n+1}))$$

= dim($\mathcal{R}(T) + \mathcal{N}(T^{n+1})$)/($\mathcal{R}(T) + \mathcal{N}(T^n)$).

Definition 2.5. Let $T \in \mathcal{B}(X)$ and let $d \in \mathbb{N}$. Then T has uniform descent for $n \geq d$ if $k_n(T) = 0$ for all $n \geq d$. If in addition $R(T^n)$ is closed in the operator range topology of $R(T^d)$ for all $n \geq d$, then we say that T has eventual topological uniform descent, and, more precisely, that T has topological uniform descent for $n \geq d$.

Operators with eventual topological uniform descent are introduced by Grabiner in [27]. It includes all classes of operators introduced in the Introduction of this paper. It also includes many other classes of operators such as operators of Kato type, quasi-Fredholm operators, operators with finite descent and operators with finite essential descent, and so on. A very detailed and far-reaching account of these notations can be seen in [1, 8, 33]. Especially, operators which have topological uniform descent for $n \geq 0$ are precisely the *semi-regular* operators studied by Mbekhta in [32]. Discussions of operators with eventual topological uniform descent may be found in [12, 20, 27, 28, 29, 40].

An operator $T \in \mathcal{B}(X)$ is said to be *essentially semi-regular* if $\mathcal{R}(T)$ is closed and $k(T) := \sum_{n=0}^{\infty} k_n(T) < \infty$. From [27, Theorem 3.7] it follows that

$$k(T) = \dim \mathcal{N}(T) / (\mathcal{N}(T) \cap \mathcal{R}(T^{\infty})) = \dim (\mathcal{R}(T) + \mathcal{N}(T^{\infty})) / \mathcal{R}(T).$$

Hence, every essentially semi-regular operator $T \in \mathcal{B}(X)$ can be characterized by $\mathcal{R}(T)$ is closed and and there exists a finite dimensional subspace $F \subseteq X$ such that $\mathcal{N}(T) \subseteq \mathcal{R}(T^{\infty}) + F$. In addition, if T is essentially semi-regular, then T^n is essentially semiregular, and hence $R(T^n)$ is closed for all $n \in \mathbb{N}$ (see Theorem 1.51 of [1]). Hence it is easy to verify that if $T \in \mathcal{B}(X)$ is essentially semi-regular, then there exist $p \in \mathbb{N}$ such that T has topological uniform descent for $n \geq p$.

Also, an operator $T \in \mathcal{B}(X)$ is called Riesz if its essential spectrum $\sigma_e(T) := \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Fredholm}\} = \{0\}$. The hyperrange and hyperkernel of $T \in \mathcal{B}(X)$ are the subspaces of X defined by $\mathcal{R}(T^{\infty}) = \bigcap_{n=1}^{\infty} \mathcal{R}(T^n)$ and $\mathcal{N}(T^{\infty}) = \bigcup_{n=1}^{\infty} \mathcal{N}(T^n)$, respectively.

Lemma 2.6. Suppose that $T \in \mathcal{B}(X)$ has topological uniform descent for $m \ge d$. If $S \in \mathcal{B}(X)$ is a Riesz operator commuting with T and V = S + T has topological uniform descent for $n \ge l$, then:

(a) dim $(\mathcal{R}(T^{\infty}) + \mathcal{R}(V^{\infty}))/(\mathcal{R}(T^{\infty}) \cap \mathcal{R}(V^{\infty})) < \infty;$

(b) dim $(\overline{\mathcal{N}(T^{\infty})} + \overline{\mathcal{N}(V^{\infty})})/(\overline{\mathcal{N}(T^{\infty})} \cap \overline{\mathcal{N}(V^{\infty})}) < \infty;$

(c) dim $\mathcal{R}(V^n)/\mathcal{R}(V^{n+1})$ = dim $\mathcal{R}(T^m)/\mathcal{R}(T^{m+1})$ for sufficiently large m and n;

(d) $\dim \mathcal{N}(V^{n+1})/\mathcal{N}(V^n) = \dim \mathcal{N}(T^{m+1})/\mathcal{N}(T^m)$ for sufficiently large m and n.

Proof. Parts (c) and (d) follow directly from [40, Theorems 3.8 and 3.12 and Remark 4.5].

When $d \neq 0$ (that is, T is not semi-regular), parts (a) and (b) follow also directly from [40, Theorems 3.8 and 3.12 and Remark 4.5].

When d = 0 (that is, T is semi-regular), then by [40, Theorems 3.8] we have that V = T + S is essentially semi-regular. So, there exist $p \in \mathbb{N}$ such that V has topological uniform descent for $n \geq p$. If $p \neq 0$ (that is, V is not semi-regular), then parts (a) and (b) follow directly from [40, Theorem 3.12 and Remark 4.5]. If p = 0 (that is, V is semi-regular), noting that $(M + N)/N \approx M/(M \cap N)$ for any subspaces M and N of X (see [30, Lemma 2.2]), then parts (a) and (b) follow from [40, Theorem 3.12 and Remark 4.5].

Theorem 2.7. Let $F \in \mathcal{B}(X)$ with $F^n \in \mathcal{F}(X)$ for some $n \in \mathbb{N}$.

(1) If $T \in \mathcal{B}(X)$ is semi-B-Fredholm and commutes with F, then

(a) dim $(\mathcal{R}(T^{\infty}) + \mathcal{R}((T+F)^{\infty}))/(\mathcal{R}(T^{\infty}) \cap \mathcal{R}((T+F)^{\infty})) < \infty;$

(b) dim $(\overline{\mathcal{N}(T^{\infty})} + \overline{\mathcal{N}((T+F)^{\infty})})/(\overline{\mathcal{N}(T^{\infty})} \cap \overline{\mathcal{N}((T+F)^{\infty})}) < \infty$.

(2) If $T \in \mathcal{B}(X)$ is upper (resp. lower) semi-B-Fredholm and commutes with F, then T + F is also upper (resp. lower) semi-B-Fredholm and $\operatorname{ind}(T + F) = \operatorname{ind}(T)$.

Proof. Suppose that $F \in \mathcal{B}(X)$ with $F^n \in \mathcal{F}(X)$ for some $n \in \mathbb{N}$. Then F^n is Riesz, that is, $\sigma_e(F^n) = \{0\}$. By the spectral mapping theorem for the essential spectrum, we get that $\sigma_e(F) = \{0\}$, so F is Riesz.

(1) Since T is semi-B-Fredholm and commutes with F, by Lemmas 2.1 and 2.2, T + F is also semi-B-Fredholm. Since every semi-B-Fredholm operator is an operator of eventual topological uniform descent, by Lemma 2.6(a) and (b), parts (a) and (b) follow immediately.

(2) By Lemmas 2.1 and 2.2, it remains to prove that $\operatorname{ind}(T+F) = \operatorname{ind}(T)$. Since every semi-B-Fredholm operator is an operator of eventual topological uniform descent, by Lemma 2.6(c) and (d) and [13, Proposition 2.1], we have that $\operatorname{ind}(T+F) = \operatorname{ind}(T)$. \Box

Theorem 2.8. Let $T \in \mathcal{B}(X)$ and let $F \in \mathcal{B}(X)$ with $F^n \in \mathcal{F}(X)$ for some $n \in \mathbb{N}$. If T commutes with F, then

(1) $\sigma_{USBW}(T+F) = \sigma_{USBW}(T);$

- (2) $\sigma_{LSBW}(T+F) = \sigma_{LSBW}(T);$
- (3) $\sigma_{SBW}(T+F) = \sigma_{SBW}(T);$
- (4) $\sigma_{BW}(T+F) = \sigma_{BW}(T).$

Proof. It follows directly from Theorem 2.7(2).

Next, we turn to the discussion of characterizations of power finite rank operators via various spectra originated from seni-B-Fredholm theory. Before this, some notations are needed.

For $T \in \mathcal{B}(X)$, let us define the descent spectrum, the essential descent spectrum and the eventual topological uniform descent spectrum of T as follows respectively:

$$\sigma_{dsc}(T) = \{\lambda \in \mathbb{C} : dsc(T - \lambda I) = \infty\};\\ \sigma^{e}_{dsc}(T) = \{\lambda \in \mathbb{C} : dsc_{e}(T - \lambda I) = \infty\};$$

 $\sigma_{ud}(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ does not have eventual topological uniform descent}\}.$

In [29], Jiang, Zhong and Zhang obtained a classification of the components of eventual topological uniform descent resolvent set $\rho_{ud}(T) := \mathbb{C} \setminus \sigma_{ud}(T)$. As an application of the classification, they show that $\sigma_{ud}(T) = \emptyset$ precisely when T is algebraic.

Lemma 2.9. ([29, Corollary 4.5]) Let $T \in \mathcal{B}(X)$ and let $\sigma_* \in \{\sigma_{ud}, \sigma_{dsc}, \sigma_{dsc}^e, \sigma_{USBF} = \sigma_{LD}^e, \sigma_{USBB} = \sigma_{LD}, \sigma_{BB} = \sigma_D\}$. Then the following statements are equivalent: (1) $\sigma_*(T) = \emptyset$;

(2) T is algebraic (that is, there exists a non-zero complex polynomial p for which p(T) = 0).

Corollary 2.10. Let $T \in \mathcal{B}(X)$ and let $\sigma_* \in \{\sigma_{ud}, \sigma_{dsc}, \sigma_{dsc}^e, \sigma_{USBF} = \sigma_{LD}^e, \sigma_{LSBF} = \sigma_{RD}^e, \sigma_{SBF}, \sigma_{BF}, \sigma_{USBW}, \sigma_{LSBW}, \sigma_{SBW}, \sigma_{USBB} = \sigma_{LD}, \sigma_{LSBB} = \sigma_{RD}, \sigma_{SBB}, \sigma_{BB} = \sigma_{D}\}$. Then the following statements are equivalent:

- (1) $\sigma_*(T) = \emptyset;$
- (2) T is algebraic.

Proof. If $\sigma_* \in \{\sigma_{ud}, \sigma_{dsc}, \sigma_{dsc}^e, \sigma_{USBF} = \sigma_{LD}^e, \sigma_{USBB} = \sigma_{LD}, \sigma_{BB} = \sigma_D\}$, the conclusion is given by Lemma 2.9. Note that

$$\sigma_{ud}(\cdot) \subseteq \sigma_{SBF}(\cdot) \subseteq \{ \begin{smallmatrix} \sigma_{SBW}(\cdot) \\ \sigma_{USBF}(\cdot) = \sigma_{LD}^{e}(\cdot) \end{smallmatrix} \subseteq \sigma_{USBW}(\cdot) \subseteq \{ \begin{smallmatrix} \sigma_{BW}(\cdot) \\ \sigma_{USBB}(\cdot) = \sigma_{LD}(\cdot) \end{smallmatrix} \subseteq \sigma_{BB}(\cdot) = \sigma_{D}(\cdot) \end{smallmatrix}$$

and that

$$\sigma_{ud}(\cdot) \subseteq \sigma_{SBF}(\cdot) \subseteq \{ \substack{\sigma_{SBW}(\cdot) \\ \sigma_{LSBF}(\cdot) = \sigma_{RD}^{e}(\cdot)} \subseteq \sigma_{LSBW}(\cdot) \subseteq \{ \substack{\sigma_{BW}(\cdot) \\ \sigma_{LSBB}(\cdot) = \sigma_{RD}(\cdot)} \subseteq \sigma_{BB}(\cdot) = \sigma_{D}(\cdot) .$$

By Lemma 2.9, if $\sigma_* \in \{\sigma_{LSBF} = \sigma_{RD}^e, \sigma_{SBF}, \sigma_{USBW}, \sigma_{LSBW}, \sigma_{SBW}, \sigma_{BW}, \sigma_{LSBB} = \sigma_{RD}\}$, the conclusion follows easily. Note that

$$\sigma_{ud}(\cdot) \subseteq \sigma_{SBB}(\cdot) \subseteq \sigma_{BB}(\cdot) = \sigma_D(\cdot)$$

and that

$$\sigma_{ud}(\cdot) \subseteq \sigma_{BF}(\cdot) \subseteq \sigma_{BB}(\cdot) = \sigma_D(\cdot).$$

Again by Lemma 2.9, if $\sigma_* \in \{\sigma_{SBB}, \sigma_{BF}\}$, the conclusion follows easily.

In [25, Theorem 3.2], O. Bel Hadj Fredj et al. proved that $F \in \mathcal{B}(X)$ with $F_n \in \mathcal{F}(X)$ for some $n \in \mathbb{N}$ if and only if $\sigma_{LD}^e(T+F) = \sigma_{LD}^e(T)$ (equivalently, $\sigma_{LD}(T+F) = \sigma_{LD}(T)$) for every operator T in the commutant of F.

We are now in a position to give the proof of the following main result.

Theorem 2.11. Let $F \in \mathcal{B}(X)$ and $\sigma_* \in \{\sigma_{dsc}, \sigma_{dsc}^e, \sigma_{USBF} = \sigma_{LD}^e, \sigma_{LSBF} = \sigma_{RD}^e, \sigma_{SBF}, \sigma_{BF}, \sigma_{USBW}, \sigma_{LSBW}, \sigma_{SBW}, \sigma_{USBB} = \sigma_{LD}, \sigma_{LSBB} = \sigma_{RD}, \sigma_{SBB}, \sigma_{BB} = \sigma_{D}\}$. Then the following statements are equivalent:

(1) $F^n \in \mathcal{F}(X)$ for some $n \in \mathbb{N}$;

(2) $\sigma_*(T+F) = \sigma_*(T)$ for all $T \in \mathcal{B}(X)$ commuting with F.

Proof. For $\sigma_* \in \{\sigma_{dsc}, \sigma_{dsc}^e, \sigma_{USBB} = \sigma_{LD}, \sigma_{USBF} = \sigma_{LD}^e\}$, the conclusion can be found in [19, Theorem 3.1], [24, Theorem 3.1] and [25, Theorem 3.2]. In the following, we prove the conclusion for the others spectra.

(1) \Rightarrow (2) For $\sigma_* \in \{\sigma_{LSBF} = \sigma_{RD}^e, \sigma_{SBF}, \sigma_{BF}, \sigma_{USBW}, \sigma_{LSBW}, \sigma_{SBW}, \sigma_{BW}\}$, the conclusion follows directly from Corollary 2.3 and Theorem 2.8.

For $\sigma_* \in \{\sigma_{LSBB} = \sigma_{RD}\}$, suppose that $F \in \mathcal{B}(X)$ with $F^n \in \mathcal{F}(X)$ for some $n \in \mathbb{N}$ and that $T \in \mathcal{B}(X)$ commutes with F. It is clear that $F^* \in \mathcal{B}(X^*)$ with $F^{*n} \in \mathcal{F}(X^*)$ and that $T^* \in \mathcal{B}(X^*)$ commutes with F^* . From the presentation before this theorem, we get that $\sigma_{LD}(T^* + F^*) = \sigma_{LD}(T^*)$, hence dually, $\sigma_{RD}(T + F) = \sigma_{RD}(T)$.

For $\sigma_* \in {\sigma_{SBB}, \sigma_{BB} = \sigma_D}$, noting that $\sigma_{SBB}(\cdot) = \sigma_{USBB}(\cdot) \cap \sigma_{LSBB}(\cdot)$ and that $\sigma_{BB}(\cdot) = \sigma_{USBB}(\cdot) \cup \sigma_{LSBB}(\cdot)$, the conclusion follows.

(2) \Rightarrow (1) Conversely, suppose that $\sigma_*(T+F) = \sigma_*(T)$ for all $T \in \mathcal{B}(X)$ commuting with F, where $\sigma_* \in \{\sigma_{LSBF} = \sigma_{RD}^e, \sigma_{SBF}, \sigma_{BF}, \sigma_{USBW}, \sigma_{LSBW}, \sigma_{SBW}, \sigma_{BW}, \sigma_{LSBB} = \sigma_{RD}, \sigma_{SBB}, \sigma_{BB} = \sigma_D\}$. By considering T = 0, then $\sigma_*(F) = \sigma_*(0+F) = \sigma_*(0) = \emptyset$. By Corollary 2.10, we know that F is algebraic. Therefore

$$X = X_1 \oplus X_2 \oplus \cdots \oplus X_n,$$

where $\sigma(F) = \{\lambda_1, \lambda_2, \dots, \lambda_1\}$ and the restriction of $F - \lambda_i$ to X_i is nilpotent for every $1 \le i \le n$. We claim that if $\lambda_i \ne 0$, dim X_i is finite. Suppose to the contrary that $\lambda_i \ne 0$ and X_i is infinite dimensional. For every $1 \le j \le n$, let F_j be the restriction of F to X_j . Then with respect to the decomposition $X = X_1 \oplus X_2 \oplus \cdots \oplus X_n$,

$$F = F_1 \oplus F_2 \oplus \cdots \oplus F_n.$$

By [19, Proposition 3.3], there exists a non-algebraic operator S_i on X_i commuting with the restriction F_i of F. Let S denote the extension of S_i to X given by S = 0 on each X_j such that $j \neq i$. Obviously SF = FS, and so $\sigma_*(S + F) = \sigma_*(S)$ by hypothesis. On the other hand, since $F = F_1 \oplus F_2 \oplus \cdots \oplus F_n$ is algebraic, F_j is algebraic for every $1 \leq j \leq n$. In particular, F_j is algebraic for every $j \neq i$. Hence by Corollary 2.10, $\sigma_{BB}(F_j) = \emptyset$ for every $j \neq i$, it follows easily that $\sigma_*(S + F) = \sigma_*(S_i + F_i)$. Since $\sigma_*(S) = \sigma_*(S_i)$,

we obtain that $\sigma_*(S_i) = \sigma_*(S_i + F_i) = \sigma_*(S_i + \lambda_i)$ because $F_i - \lambda_i$ is nilpotent. Choose an arbitrary complex number $\alpha \in \sigma_*(S) \neq \emptyset$, it follows that $k\lambda_i + \alpha \in \sigma_*(S)$ for every positive integer k, which implies that $\lambda_i = 0$, the desired contradiction.

Remark 2.12. (1) The argument we have given for the implication $(2) \Rightarrow (1)$ in Theorem 2.11 is, in fact, discovered by following the trail marked out by Burgos, Kaidi, Mbekhta and Oudghiri [19].

(2) By [11, Lemma 2.3], [33, pp. 135-136] and a similar argument of [10, Proposition 3.3], we know that $\sigma_*(T + F) = \sigma_*(T)$ for all finite rank operator F not necessarily commuting with T, where $\sigma_* \in \{\sigma_{dsc}^e, \sigma_{USBF} = \sigma_{LD}^e, \sigma_{LSBF} = \sigma_{RD}^e, \sigma_{SBF}, \sigma_{BF}, \sigma_{USBW}, \sigma_{LSBW}, \sigma_{SBW}, \sigma_{BW}\}$. By [33, Observation 5 in p. 136], we know that σ_* is not stable under non-commuting finite rank perturbation, where $\sigma_* \in \{\sigma_{dsc}, \sigma_{USBB} = \sigma_{LD}, \sigma_{LSBB} = \sigma_{RD}, \sigma_{SBB}, \sigma_{BB} = \sigma_D\}$.

3 Some applications

Rashid claimed in [37, Theorem 3.15] that if $T \in \mathcal{B}(X)$ and Q is a quasi-nilpotent operator that commute with T, then (in [37], σ_{USBW} is denoted as $\sigma_{SBF_{-}}$)

$$\sigma_{USBW}(T+Q) = \sigma_{USBW}(T).$$

In [41, Example 2.13], the authors showed that this equality does not hold in general.

As an immediate consequence of Theorem 2.11 (that is, main result), we obtain the following corollary which, in particular, is a corrected version of [37, Theorem 3.15] and also provide positive answers to Questions 1.1 and 1.2.

Corollary 3.1. Let $T \in \mathcal{B}(X)$ and let $N \in \mathcal{B}(X)$ be a nilpotent operator commuting with T. Then

- (1) $\sigma_{USBW}(T+N) = \sigma_{USBW}(T);$
- (2) $\sigma_{LSBW}(T+N) = \sigma_{LSBW}(T);$
- (3) $\sigma_{SBW}(T+N) = \sigma_{SBW}(T);$
- (4) $\sigma_{BW}(T+N) = \sigma_{BW}(T).$

Besides Question 1.2, Berkani and Zariouh also posed in [17] the following question:

Question 3.2. Let $T \in \mathcal{B}(X)$ and let $N \in \mathcal{B}(X)$ be a nilpotent operator commuting with T. Under which conditions

$$\sigma_{BF}(T+N) = \sigma_{BF}(T) ?$$

As an immediate consequence of Theorem 2.11, we also obtain the following corollary which, in particular, provide a positive answer to Question 3.2.

Corollary 3.3. Let $T \in \mathcal{B}(X)$ and let $N \in \mathcal{B}(X)$ be a nilpotent operator commuting with T. Then

- (1) $\sigma_{USBF}(T+N) = \sigma_{USBF}(T);$
- (2) $\sigma_{LSBF}(T+N) = \sigma_{LSBF}(T);$

(3) $\sigma_{SBF}(T+N) = \sigma_{SBF}(T);$ (4) $\sigma_{BF}(T+N) = \sigma_{BF}(T).$

We say that $\lambda \in \sigma_a(T)$ is a left pole of T if $T - \lambda I$ is left Drazin invertible. Let $\Pi_a(T)$ denote the set of all left poles of T. An operator $T \in \mathcal{B}(X)$ is called *a-polaroid* if $iso\sigma_a(T) = \Pi_a(T)$. Here and henceforth, for $A \subseteq \mathbb{C}$, iso A is the set of isolated points of A. Besides Questions 1.2 and 3.2, Berkani and Zariouh also posed in [18] the following three questions:

Question 3.4. Let $T \in \mathcal{B}(X)$ and let $N \in \mathcal{B}(X)$ be a nilpotent operator commuting with T. Under which conditions

$$asc(T+N) < \infty \iff \sigma_{asc}(T) < \infty$$
?

Question 3.5. Let $T \in \mathcal{B}(X)$ and let $N \in \mathcal{B}(X)$ be a nilpotent operator commuting with T. Under which conditions, $\mathcal{R}((T+N)^m)$ is closed for m large enough if and only if $\mathcal{R}(T^m)$ is closed for m large enough?

Question 3.6. Let $T \in \mathcal{B}(X)$ and let $N \in \mathcal{B}(X)$ be a nilpotent operator commuting with T. Under which conditions

$$\Pi_a(T+N) = \Pi_a(T) ?$$

We mention that Question 3.4 is, in fact, an immediate consequence of an earlier result of Kaashoek and Lay [31, Theorem 2.2]. To Question 3.5, suppose that $T \in \mathcal{B}(X)$ and that $N \in \mathcal{B}(X)$ is a nilpotent operator commuting with T. As a direct consequence of Theorem 2.11 (that is, main result), we know that if there exists $n \in \mathbb{N}$ such that $c_n(T) < \infty$ or $c'_n(T) < \infty$, then $\mathcal{R}((T+N)^m)$ is closed for m large enough if and only if $\mathcal{R}(T^m)$ is closed for m large enough.

To Question 3.6, we first recall a classical result.

Lemma 3.7. ([33]) If $T \in \mathcal{B}(X)$ and $Q \in \mathcal{B}(X)$ is a quasi-nilpotent operator commuting with T, then

$$\sigma(T+Q) = \sigma(T) \text{ and } \sigma_a(T+Q) = \sigma_a(T). \tag{3.1}$$

As an immediate consequence of Theorem 2.11 and Lemma 3.7, we also obtain the following corollary which provide a positive answer to Question 3.6.

Corollary 3.8. Let $T \in \mathcal{B}(X)$ and let $N \in \mathcal{B}(X)$ be a nilpotent operator commuting with T. Then

$$\Pi_a(T+N) = \Pi_a(T). \tag{3.2}$$

Let $\Pi(T)$ denote the set of all poles of T. It is proved in [14, Lemma 2.2] that if $T \in \mathcal{B}(X)$ and $Q \in \mathcal{B}(X)$ is a nilpotent operator commuting with T, then

$$\Pi(T+N) = \Pi(T). \tag{3.3}$$

Let E(T) and $E_a(T)$ denote the set of all isolated eigenvalues of T and the set of all eigenvalues of T that are isolated in $\sigma_a(T)$, respectively. That is,

$$E(T) = \{\lambda \in iso\sigma(T) : 0 < \alpha(T - \lambda I)\}\$$

and

$$E_a(T) = \{\lambda \in iso\sigma_a(T) : 0 < \alpha(T - \lambda I)\}.$$

An operator $T \in \mathcal{B}(X)$ is called *a-isoloid* if $iso\sigma_a(T) = E_a(T)$.

We set $\Pi^0(T) = \{\lambda \in \Pi(T) : \alpha(T - \lambda I) < \infty\}, \ \Pi^0_a(T) = \{\lambda \in \Pi_a(T) : \alpha(T - \lambda I) < \infty\}, \ E^0(T) = \{\lambda \in E(T) : \alpha(T - \lambda I) < \infty\} \text{ and } E^0_a(T) = \{\lambda \in E_a(T) : \alpha(T - \lambda I) < \infty\}.$

Suppose that $T \in \mathcal{B}(X)$ and that $N \in \mathcal{B}(X)$ is a nilpotent operator commuting with T. Then from the proof of [17, Theorem 3.5], it follows that

$$0 < \alpha(T+N) \Longleftrightarrow 0 < \alpha(T)$$

and

$$\alpha(T+N) < \infty \Longleftrightarrow \alpha(T) < \infty.$$

Hence by Equation (3.1), we have the following equations:

$$E(T+N) = E(T); (3.4)$$

$$E_a(T+N) = E_a(T); (3.5)$$

$$E^{0}(T+N) = E^{0}(T); (3.6)$$

$$E_a^0(T+N) = E_a^0(T). (3.7)$$

An operator $T \in \mathcal{B}(X)$ is said to be *upper semi-Weyl* if T is upper semi-Fredholm and ind $(T) \leq 0$. An operator $T \in \mathcal{B}(X)$ is said to be *Weyl* if T is Fredholm and ind(T) = 0. For $T \in \mathcal{B}(X)$, let us define the *upper semi-Browder spectrum*, the *Browder spectrum*, the *upper semi-Weyl spectrum* and the *Weyl spectrum* of T as follows respectively:

 $\sigma_{USB}(T) = \{ \lambda \in \mathbb{C} : T - \lambda I \text{ is not a upper semi-Browder operator} \};$

- $\sigma_B(T) = \{\lambda \in \mathbb{C} : T \lambda I \text{ is not a Browder operator}\};$
- $\sigma_{USW}(T) = \{ \lambda \in \mathbb{C} : T \lambda I \text{ is not a upper semi-Weyl operator} \};$

 $\sigma_W(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a Weyl operator}\}.$

Suppose that $T \in \mathcal{B}(X)$ and that $R \in \mathcal{B}(X)$ is a Riesz operator commuting with T. Then it follows from [39, Proposition 5] and [36, Theorem 1] that

$$\sigma_{USW}(T+R) = \sigma_{USW}(T); \qquad (3.8)$$

$$\sigma_W(T+R) = \sigma_W(T); \tag{3.9}$$

$$\sigma_{USB}(T+R) = \sigma_{USB}(T); \qquad (3.10)$$

$$\sigma_B(T+R) = \sigma_B(T). \tag{3.11}$$

Suppose that $T \in \mathcal{B}(X)$ and that $Q \in \mathcal{B}(X)$ is a quasi-nilpotent operator commuting with T. Then, noting that $\Pi^0(T) = \sigma(T) \setminus \sigma_B(T)$ and $\Pi^0_a(T) = \sigma_a(T) \setminus \sigma_{USB}(T)$ for any $T \in \mathcal{B}(X)$, it follows from Equations (3.1), (3.10) and (3.11) that

$$\Pi^0(T+Q) = \Pi^0(T); \tag{3.12}$$

$$\Pi_a^0(T+Q) = \Pi_a^0(T).$$
(3.13)

In the following table, we use the abbreviations gaW, aW, gW, W, (gw), (w), (gaw)and (aw) to signify that an operator $T \in \mathcal{B}(X)$ obeys generalized a-Weyl's theorem, a-Weyl's theorem, generalized Weyl's theorem, Weyl's theorem, property (gw), property (w), property (gaw) and property (aw). For example, an operator $T \in \mathcal{B}(X)$ is said to obey generalized a-Weyl's theorem (in symbol $T \in gaW$), if $\sigma_a(T) \setminus \sigma_{USBW}(T) = E_a(T)$. Similarly, the abbreviations gaB, aB, gB, B, (gb), (b), (gab) and (ab) have analogous meaning with respect to Browder's theorem or the properties.

gaW	$\sigma_a(T) \backslash \sigma_{USBW}(T) = E_a(T)$	gaB	$\sigma_a(T) \backslash \sigma_{USBW}(T) = \Pi_a(T)$
aW	$\sigma_a(T) \backslash \sigma_{USW}(T) = E_a^0(T)$	aB	$\sigma_a(T) \backslash \sigma_{USW}(T) = \Pi_a^0(T)$
gW	$\sigma(T) \backslash \sigma_{BW}(T) = E(T)$	gB	$\sigma(T) \setminus \sigma_{BW}(T) = \Pi(T)$
W	$\sigma(T) \setminus \sigma_W(T) = E^0(T)$	В	$\sigma(T) \setminus \sigma_W(T) = \Pi^0(T)$
(gw)	$\sigma_a(T) \backslash \sigma_{USBW}(T) = E(T)$	(gb)	$\sigma_a(T) \backslash \sigma_{USBW}(T) = \Pi(T)$
(w)	$\sigma_a(T) \backslash \sigma_{USW}(T) = E^0(T)$	(b)	$\sigma_a(T) \backslash \sigma_{USW}(T) = \Pi^0(T)$
(gaw)	$\sigma(T) \setminus \sigma_{BW}(T) = E_a(T)$	(gab)	$\sigma(T) \setminus \sigma_{BW}(T) = \Pi_a(T)$
(aw)	$\sigma(T) \setminus \sigma_W(T) = E_a^0(T)$	(ab)	$\sigma(T) \setminus \sigma_W(T) = \Pi^0_a(T)$

Weyl-Browder type theorems and properties, in their classical and more recently in their generalized form, have been studied by a large of authors. Theorem 2.11 and Equations (3.1)—(3.13) give us an unifying framework for establishing commuting perturbational results of Weyl-Browder type theorems and properties (generalized or not).

Corollary 3.9. (1) If $T \in \mathcal{B}(X)$ obeys gaW (resp. aW, gW, W, (gw), (w), (gaw), (aw), (gb), (gab)) and $N \in \mathcal{B}(X)$ is a nilpotent operator commuting with T, then T + N also obeys gaW (resp. aW, gW, W, (gw), (w), (gaw), (aw), (gb), (gab)).

(2) If $T \in \mathcal{B}(X)$ obeys gaB (resp. aB, gB, B) and $R \in \mathcal{B}(X)$ is a Riesz operator commuting with T, then T + R also obeys gaB (resp. aB, gB, B).

(3) If $T \in \mathcal{B}(X)$ obeys (b) (resp. (ab)) and $Q \in \mathcal{B}(X)$ is a quasi-nilpotent operator commuting with T, then T + Q also obeys (b) (resp. (ab)).

Proof. (1) It follows directly from Theorem 2.11 and Equations (3.1)—(3.9).

(2) By [6], we know that T obeys gB (resp. gaB) if and only if T obeys B (resp. aB) for any $T \in \mathcal{B}(X)$. Note that T obeys B (resp. aB) if and only if $\sigma_W(T) = \sigma_B(T)$

(resp. $\sigma_{USW}(T) = \sigma_{USB}(T)$). Hence by Equations (3.8)—(3.11), the conclusion follows immediately.

(3) It follows directly from Equations (3.1), (3.8), (3.9), (3.12) and (3.13).

The commuting perturbational results established in Corollary 3.9, in particular, improve many recent results of [11, 14, 17, 18, 38] by removing certain extra assumptions.

Remark 3.10. (1) For generalized a-Weyl's theorem, part (1) of Corollary 3.9 improves [18, Theorem 3.3] by removing the extra assumption that $E_a(T) \subseteq iso\sigma(T)$ and extends [18, Theorem 3.2]. For property (gw), on one hand, part (1) of Corollary 3.9 improves [38, Theorem 2.16] (resp. [11, Theorem 3.6]) by removing the extra assumption that T is a-isoloid (resp. T is a-polaroid) and extends [17, Theorem 3.8]; on the other hand, our proof for it is a corrected proof of [38, Theorem 2.16]. For property (gab), part (1) of Corollary 3.9 improves [17, Theorem 3.2] by removing the extra assumption that T is a-polaroid and extends [17, Theorem 3.4].

For generalized Weyl's theorem (resp. property (w), property (gaw)), part (1) of Corollary 3.9 has been proved in [11, Theorem 3.4] (resp. [3, Theorem 3.8] and [11, Theorem 3.1], [17, Theorem 3.6]) by using a different method.

For a-Weyl's theorem, some other commuting perturbational theorems for it have been proved in [18, 21, 35].

For Weyl's theorem (resp. property (aw), property (gb)), part (1) of Corollary 3.9 has been proved in [34, Theorem 3] (resp. [17, Theorem 3.5], [41, Theorem 2.6]).

(2) It has been discovered in [4] that Browder's theorem and a-Browder's theorem are stable under commuting Riesz perturbations.

(3) For property (b) (resp. (ab)), part (3) of Corollary 3.9 extends [14, Theorem 2.1] (resp. [17, Theorem 3.1]) from commuting nilpotent perturbations to commuting quasi-nilpotent perturbations.

We conclude this paper by some examples to illustrate our commuting perturbational results of Weyl-Browder type theorems and properties (generalized or not).

The following simple example shows that gaW, aW, gW, W, (gw), (w), (gaw) and (aw) are not stable under commuting quasi-nilpotent perturbations.

Example 3.11. Let $Q: l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be a quasi-nilpotent operator defined by

$$Q(x_1, x_2, \dots) = (\frac{x_2}{2}, \frac{x_3}{3}, \dots)$$
 for all $(x_n) \in l_2(\mathbb{N})$.

Then Q is quasi-nilpotent, $\sigma(Q) = \sigma_a(Q) = \sigma_W(Q) = \sigma_{USW}(Q) = \sigma_{BW}(Q) = \sigma_{USBW}(Q)$ = {0} and $E_a(Q) = E_a^0(Q) = E(Q) = E^0(Q) = \{0\}$. Take T = 0. Clearly, T satisfies gaW (resp. aW, gW, W, (gw), (w), (gaw), (aw)), but T + Q = Q fails gaW (resp. aW, gW, W, (gw), (w), (gaw), (aw)).

The following example was given in [41, Example 2.14] to show that property (gb) is not stable under commuting quasi-nilpotent perturbations. Now, we use it to illustrate that property (gab) is also unstable under commuting quasi-nilpotent perturbations. **Example 3.12.** Let $U: l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be the unilateral right shift operator defined by

$$U(x_1, x_2, \dots) = (0, x_1, x_2, \dots)$$
 for all $(x_n) \in l_2(\mathbb{N})$.

Let $V: l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be a quasi-nilpotent operator defined by

$$V(x_1, x_2, \dots) = (0, x_1, 0, \frac{x_3}{3}, \frac{x_4}{4} \dots)$$
 for all $(x_n) \in l_2(\mathbb{N})$.

Let $N: l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be a quasi-nilpotent operator defined by

$$N(x_1, x_2, \dots) = (0, 0, 0, -\frac{x_3}{3}, -\frac{x_4}{4} \dots)$$
 for all $(x_n) \in l_2(\mathbb{N})$.

It is easy to verify that VN = NV. We consider the operators T and Q defined by $T = U \oplus V$ and $Q = 0 \oplus N$, respectively. Then Q is quasi-nilpotent and TQ = QT. Moreover,

$$\sigma(T) = \sigma(U) \cup \sigma(V) = \{\lambda \in \mathbb{C} : 0 \le |\lambda| \le 1\},\$$

$$\sigma_a(T) = \sigma_a(U) \cup \sigma_a(V) = \{\lambda \in \mathbb{C} : |\lambda| = 1\} \cup \{0\},\$$

$$\sigma(T+Q) = \sigma(U) \cup \sigma(V+N) = \{\lambda \in \mathbb{C} : 0 \le |\lambda| \le 1\}$$

and

$$\sigma_a(T+Q) = \sigma_a(U) \cup \sigma_a(V+N) = \{\lambda \in \mathbb{C} : |\lambda| = 1\} \cup \{0\}.$$

It follows that $\Pi_a(T) = \Pi(T) = \emptyset$ and $\{0\} = \Pi_a(T+Q) \neq \Pi(T+Q) = \emptyset$. Hence by [16, Corollary 2.7], T+Q does not satisfy property (gab). But since T has SVEP, T satisfies Browder's theorem or equivalently, by [6, Theorem 2.2], T satisfies generalized Browder's theorem. Therefore by [16, Corollary 2.7] again, T satisfies property (gb).

The following example was given in [41, Example 2.12] to show that property (gb) is not preserved under commuting finite rank perturbations. Now, we use it to illustrate that property (b) and (ab) are also unstable under commuting finite rank (hence compact) perturbations.

Example 3.13. Let $U: l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be the unilateral right shift operator defined by

$$U(x_1, x_2, \cdots) = (0, x_1, x_2, \cdots)$$
 for all $(x_n) \in l_2(\mathbb{N})$.

For fixed $0 < \varepsilon < 1$, let $F_{\varepsilon} : l_2(\mathbb{N}) \longrightarrow l_2(\mathbb{N})$ be a finite rank operator defined by

$$F_{\varepsilon}(x_1, x_2, \cdots) = (-\varepsilon x_1, 0, 0, \cdots) \text{ for all } (x_n) \in l_2(\mathbb{N}).$$

We consider the operators T and F defined by $T = U \oplus I$ and $F = 0 \oplus F_{\varepsilon}$, respectively. Then F is a finite rank operator and TF = FT. Moreover,

$$\sigma(T) = \sigma(U) \cup \sigma(I) = \{\lambda \in \mathbb{C} : 0 \le |\lambda| \le 1\},\$$
$$\sigma_a(T) = \sigma_a(U) \cup \sigma_a(I) = \{\lambda \in \mathbb{C} : |\lambda| = 1\},\$$
$$\sigma(T+F) = \sigma(U) \cup \sigma(I+F_{\varepsilon}) = \{\lambda \in \mathbb{C} : 0 \le |\lambda| \le 1\}$$

and

$$\sigma_a(T+F) = \sigma_a(U) \cup \sigma_a(I+F_{\varepsilon}) = \{\lambda \in \mathbb{C} : |\lambda| = 1\} \cup \{1-\varepsilon\}.$$

It follows that $\Pi_a^0(T) = \Pi^0(T) = \emptyset$ and $\{1 - \varepsilon\} = \Pi_a^0(T + F) \neq \Pi^0(T + F) = \emptyset$. Hence by [15, Corollary 2.7] (resp. [16, Corollary 2.6]), T + F does not satisfy property (b) (resp. (ab)). But since T has SVEP, T satisfies a-Browder's theorem (resp. Browder's theorem), therefore by [15, Corollary 2.7] (resp. [16, Corollary 2.6]) again, T satisfies property (b) (resp. (ab)).

References

- [1] *P. Aiena*, Fredholm and local spectral Theory, with application to Multipliers, Kluwer Academic Publishers, Dordrecht 2004.
- [2] P. Aiena, M. T. Biondi and C. Carpintero, On Drazin invertibility, Proc. Amer. Math. Soc. 136 2008, 2839–2848.
- [3] P. Aiena, M. T. Biondi and F. Villafañe, Property (w) and perturbations III, J. Math. Anal. Appl. 353 2009, 205–214.
- [4] P. Aiena, C. Carpintero and E. Rosas, Browder's theorem and the spectral mapping theorem, Divulg. Mat. 15 2007, 207–226.
- [5] M. Amouch, Polaroid operators with SVEP and perturbations of property (gw), Mediterr. J. Math. 6 2009, 461–470.
- [6] M. Amouch, H. Zguitti, On the equivalence of Browder's and generalized Browder's theorem, Glasgow Math. J. 48 2006, 179–185.
- [7] M. Amouch, H. Zguitti, B-Fredholm and Drazin invertible operators through localized SVEP, Math. Bohem. 136 2011, 39–49.
- [8] M. Berkani, Restriction of an operator to the range of its powers, Studia Math. 140 2000, 163–175.
- [9] M. Berkani, An Atkinson-type theorem for B-Fredholm operators, Studia Math. 148 2001, 251–257.
- [10] M. Berkani, Index of B-Fredholm operators and generalization of a Weyl theorem, Proc. Amer. Math. Soc. 130 2001, 1717–1723.
- [11] M. Berkani and M. Amouch, Preservation of property (gw) under perturbations, Acta Sci. Math. (Szeged) 74 2008, 769–781.
- [12] M. Berkani, N. Castro, and S. Djordjević, Single valued extension property and generalized Weyl's theorem, Math. Bohem. 131(1) 2006, 29–38.
- [13] M. Berkani and M. Sarih, On semi B-Fredholm operators, Glasgow Math. J. 43 2001, 457–465.

- [14] M. Berkani and H. Zariouh, Extended Weyl type theorems and perturbations, Math. Proc. R. Ir. Acad. 110A 2010, 73–81.
- [15] M. Berkani and H. Zariouh, Extended Weyl type theorems, Math. Bohem. 134(4) 2009, 369–378.
- [16] M. Berkani and H. Zariouh, New extended Weyl type theorems, Mat. Vesnik. 62(2) 2010, 145–154.
- [17] M. Berkani and H. Zariouh, Perturbation results for Weyl type theorems, Acta Math. Univ. Comenianae 80 2011, 119–132.
- [18] M. Berkani and H. Zariouh, Generalized a-Weyl's theorem and perturbations, Functional Analysis, Approximation and Computation 2 2010, 7–18.
- [19] M. Burgos, A. Kaidi, M. Mbekhta and M. Oudghiri, The descent spectrum and perturbations, J. Operator Theory 56 2006, 259–271.
- [20] X.H. Cao, Topological uniform descent and Weyl type theorem, Linear Algebra Appl. 420 2007, 175–182.
- [21] X.H. Cao, M.Z. Guo and B. Meng, Weyl spectra and Weyl's theorem, J. Math. Anal. Appl. 288 2003, 758–767.
- [22] C.R. Carpintero, O. García, E.R. Rosas, J.E. Sanabria, B-Browder spectra and localized SVEP, Rend. Circ. Mat. Palermo (2) 57 2008, 239–254.
- [23] P.A. Fillmore and J.P. Williams, On operator ranges. Advances in Math. 7(3) 1971, 254–281.
- [24] O. Bel Hadj Fredj, Essential descent spectrum and commuting compact perturbations, Extracta Math. **21** 2006, 261–271.
- [25] O. Bel Hadj Fredj, M. Burgos, and M. Oudghiri, Ascent spectrum and essential ascent spectrum, Studia Math. 187 2008, 59–73.
- [26] S. Grabiner, Ranges of products of operators. Canad. J. Math. XXVI(6) 1974, 1430–1441.
- [27] S. Grabiner, Uniform ascent and descent of bounded operators, J. Math. Soc. Japan 34 1982, 317–337.
- [28] Q.F. Jiang, H.J. Zhong and Q.P. Zeng, Topological uniform descent and localized SVEP, J. Math. Anal. Appl. 390 2012, 355–361.
- [29] Q.F. Jiang, H.J. Zhong and S.F. Zhang, Components of topological uniform descent resolvent set and local spectral theory, Sumitted.
- [30] M.A. Kaashoek, Ascent, descent, nullity and defect, a note on a paper by A.E. Taylor. Math. Ann. 172(2) 1967, 105–115.

- [31] M.A. Kaashoek and D.C. Lay, Ascent, descent, and commuting perturbations, Trans. Amer. Math. Soc. 169 1972, 35–47.
- [32] M. Mbekhta, Résolvant généralisé et théorie spectrale, J. Operator Theory. 21 1989, 69–105.
- [33] M. Mbekhta and V. Müller, On the axiomatic theory of spectrum II, Studia Math. 119 1996, 129–147.
- [34] K.K. Oberai, On the Weyl spectrum II, Illinois J. Math. 21 1977, 84–90.
- [35] M. Oudghiri, a-Weyl's theorem and perturbations, Studia Math. 173 2006, 193–201.
- [36] V. Rakocěvić, Semi-Browder operators and perturbations, Studia Math. 122 1997, 131–137.
- [37] *M.H.M. Rashid*, Property (*gb*) and perturbations, J. Math. Anal. Appl. **383** 2011, 82–94.
- [38] *M.H.M. Rashid*, Property (*gw*) and perturbations, Bull. Belg. Math. Soc. Simon Stevin **18** 2011, 635–654.
- [39] *H.-O. Tylli*, On the asymptotic behaviour of some quantities related to semi-Fredholm operators, J. London Math. Soc. **31** 1985, 340–348.
- [40] Q.P. Zeng, H.J. Zhong and Z.Y. Wu, Small essential spectral radius perturbations of operators with topological uniform descent, Bull. Aust. Math. Soc. 85 2012, 26–45.
- [41] Q.P. Zeng and H.J. Zhong, A note on property (gb) and perturbations, arXiv:1203.1134