

Gauge fixing in (2+1)-gravity with vanishing cosmological constant^{*}

C. Meusburger[†]

T. Schönfeld[‡]

Department Mathematik,
Friedrich-Alexander-Universität Erlangen-Nürnberg
Cauerstraße 11, 91058 Erlangen, Germany

March 23, 2012

Abstract

We apply Dirac's gauge fixing procedure to (2+1)-gravity with vanishing cosmological constant. For general gauge fixing conditions based on two point particles, this yields explicit expressions for the Dirac bracket. We explain how gauge fixing is related to the introduction of an observer into the theory and show that the Dirac bracket is determined by a classical dynamical r -matrix. Its two dynamical variables correspond to the mass and spin of a cone that describes the residual degrees of freedom of the spacetime. We show that different gauge fixing conditions and different choices of observers are related by dynamical Poincaré transformations. This allows us to locally classify all Dirac brackets resulting from the gauge fixing and to relate them to a set of particularly simple solutions associated with the centre-of-mass frame of the spacetime.

1 Introduction

(2+1)-dimensional gravity plays an important role in quantum gravity as a simple model for higher dimensions. It allows one to investigate important physics questions arising in the quantisation of the theory in a simplified framework, in which a full and rigorous quantisation of the theory is possible. This includes conceptual questions such as the role of time and observers in the quantum theory as well as mathematical questions surrounding quantisation, for an overview see [1].

In addition to its role in quantum gravity, (2+1)-gravity is of interest intrinsically due to its rich mathematical structure, which includes its close relation to Chern-Simons gauge theory and moduli spaces of flat connections [2, 3], aspects of three-dimensional geometry [4, 5] as well as knot theory [6], Reshetikhin-Turaev Invariants [7] and conformal field theory [8].

While the quantisation of the Euclidean case is rather well understood, the quantisation of (2+1)-gravity with Lorentzian signature proves more difficult. This is due to the fact that (2+1)-gravity can be viewed as a constrained system and the implementation of the constraints in the quantum theory is achieved via representation-theoretical methods. For (2+1)-gravity

^{*}This work was funded by the research grant ME 3425/1-1 in the Emmy-Noether programme of the German Research Foundation (DFG).

[†]catherine.meusburger@math.uni-erlangen.de

[‡]torsten.schoenfeld@math.uni-erlangen.de

with Lorentzian signature, the relevant Lie groups and quantum groups whose representations arise in the construction of the quantum theory are non-compact, which causes difficulties that are not present in the compact cases.

This is a strong motivation to investigate approaches for the quantisation of this theory which implement the constraints directly into the classical theory via gauge fixing and then quantise the resulting classical description. An independent motivation to study the effect of gauge fixing on the classical theory are indications that gauge fixing is related to the inclusion of an observer in the classical and quantum theory [9, 10]. The inclusion of observers into the quantum theory is an important conceptual question of quantum gravity, which can be studied in detail in this model.

A further motivation is the debate surrounding quantum group symmetries and non-commutative structures in quantum gravity such as κ -Poincaré symmetries or Drinfel'd doubles [11, 12, 13, 14, 15, 16, 17]. While various models in quantum gravity exhibit quantum group symmetries, these quantum group symmetries are often associated with certain extended or enlarged Hilbert spaces, from which the gauge-invariant Hilbert space is obtained via the imposition of constraints. It is therefore unclear how much of this quantum group symmetry survives constraint implementation and if these symmetries are generic features of quantum gravity or merely technical tools in the construction of the quantum theory. It seems plausible that a model containing an observer would be useful in providing an answer to this question.

In this article, we summarise our results [10, 18] on gauge fixing in (2+1)-dimensional gravity with vanishing cosmological constant and discuss their physical interpretation and their implications in quantum gravity. We consider a rather general set of gauge fixing conditions based on two point particles in the spacetime. These gauge fixing conditions have a direct physical interpretation as conditions that specify an observer in the spacetime. Via Dirac's gauge fixing procedure, we then derive an explicit description of the resulting Dirac bracket. We show that the underlying mathematical structures that define this bracket are classical dynamical r -matrices for the Lie algebra $\mathfrak{iso}(2, 1)$. These classical dynamical r -matrices depend on two variables which have a direct interpretation as energy and angular momentum of the spacetime as measured by the associated observer.

We then discuss how different gauge fixing conditions are related by dynamical Poincaré transformations, which generalise the usual gauge transformations of classical dynamical r -matrices in the literature [19]. We show that these transformations allow one to map each Dirac bracket and the associated classical dynamical r -matrices to two simple standard solutions for almost all values of the dynamical variables. These standard solutions have a direct interpretation as the centre-of-mass frame of the spacetime.

Our article is structured as follows. In Section 2, we summarise the combinatorial description of the phase space of (2+1)-gravity which plays an important role in the quantisation of the theory and serves as the starting point for our gauge fixing procedure. We give a detailed discussion of its geometrical interpretation and show how the variables that parametrise the phase space describe the construction of spacetimes from regions in Minkowski space.

In Section 3 we show how this description of the phase space can be interpreted as a constrained system. We discuss the gauge fixing conditions imposed to obtain the gauge-invariant phase space of the theory and describe how gauge fixing the theory amounts to specifying an observer. In Section 4, we determine the Dirac bracket associated with these gauge fixing conditions and show that it is given by classical dynamical r -matrices. We find that different choices of gauge fixing conditions are related by dynamical Poincaré transformations which depend on

the total energy and angular momentum of the universe, as measured by this observer. We then discuss how these transformations allow one to obtain a centre-of-mass frame description of the spacetime, in which the centre of mass appears as an effective particle at rest at the origin or a “tachyonic” particle with a spacelike worldline. Section 5 contains our outlook and conclusions.

2 The phase space of (2+1)-gravity

2.1 Notations and conventions

We denote by $\mathbf{e}_0 = (1, 0, 0)$, $\mathbf{e}_1 = (0, 1, 0)$, $\mathbf{e}_2 = (0, 0, 1)$ the standard basis of \mathbb{R}^3 and use Einstein’s summation convention. Unless stated otherwise, all indices run from 0 to 2 and are raised and lowered with the three-dimensional Minkowski metric $\eta = \text{diag}(1, -1, -1)$. We write ε_{abc} for the totally antisymmetric tensor in three dimensions with $\varepsilon_{012} = 1$. For three-vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^3$, we write $\eta(\mathbf{x}, \mathbf{y}) = \mathbf{x} \cdot \mathbf{y} = \eta_{ab}x^ay^b$, $\mathbf{x}^2 = \mathbf{x} \cdot \mathbf{x}$ and $\mathbf{x} \wedge \mathbf{y}$ for the three-vector with components $(\mathbf{x} \wedge \mathbf{y})^a = \varepsilon^{abc}x_by_c$.

The proper orthochronous Lorentz group in three dimensions is the group $\text{SO}_+(2, 1) \cong \text{PSL}(2, \mathbb{R})$ with Lie algebra $\mathfrak{so}(2, 1) \cong \mathfrak{sl}(2, \mathbb{R})$. In the following, we write Ad for its adjoint representation, which coincides with its representation by $\text{SO}_+(2, 1)$ matrices. The Poincaré group in three dimensions is the semidirect product of the proper orthochronous Lorentz group and \mathbb{R}^3 : $\text{ISO}(2, 1) \equiv \text{SO}_+(2, 1) \ltimes \mathbb{R}^3$. We parametrise elements of $\text{ISO}(2, 1)$ as

$$(u, \mathbf{a}) = (u, 0) \cdot (\mathbb{1}, -\mathbf{j}) = (u, -\text{Ad}(u)\mathbf{j}) \quad \text{with } u \in \text{SO}_+(2, 1), \mathbf{j}, \mathbf{a} \in \mathbb{R}^3,$$

and the group multiplication law then takes the form

$$(u_1, \mathbf{a}_1) \cdot (u_2, \mathbf{a}_2) = (u_1 \cdot u_2, \mathbf{a}_1 + \text{Ad}(u_1)\mathbf{a}_2).$$

We fix a basis $\{J_a, P_a\}_{a=0,1,2}$ of the Lie algebra $\mathfrak{iso}(2, 1)$, in which the Lie bracket takes the form

$$[J_a, J_b] = \varepsilon_{ab}{}^c J_c, \quad [J_a, P_b] = \varepsilon_{ab}{}^c J_c, \quad [P_a, P_b] = 0.$$

The basis elements J_a are the generators of the Lorentz transformations, the basis elements P_a the generators of the translations. An Ad -invariant symmetric bilinear form on $\mathfrak{iso}(2, 1)$, which defines the Chern-Simons formulation of (2+1)-gravity, is given by

$$\langle J_a, P_b \rangle = \eta_{ab}, \quad \langle J_a, J_b \rangle = \langle P_a, P_b \rangle = 0. \quad (1)$$

We denote by P_a^L, P_a^R the right- and left-invariant vector fields on $\text{ISO}(2, 1)$ associated with the translations and by J_a^L, J_a^R the ones associated with the Lorentz transformations. More generally, for any basis $\{T_a\}_{a=0,\dots,5}$ of $\mathfrak{iso}(2, 1)$, we define

$$L_af(h) = \left. \frac{d}{dt} \right|_{t=0} f(e^{-tT_a} \cdot h), \quad R_af(h) = \left. \frac{d}{dt} \right|_{t=0} f(h \cdot e^{tT_a}) \quad \forall f \in C^\infty(\text{ISO}(2, 1)), \quad (2)$$

where $e : \mathfrak{iso}(2, 1) \rightarrow \text{ISO}(2, 1)$, $\mathbf{x} \mapsto e^{\mathbf{x}}$ is the exponential map for $\text{ISO}(2, 1)$.




Figure 1: Generators of the fundamental group for an n -punctured genus g surface $S_{g,n}$.

2.2 Phase space and Poisson structure

In the following we investigate gauge fixing in (2+1)-gravity with vanishing cosmological constant in its formulation as a Chern-Simons gauge theory with gauge group $\text{ISO}(2,1)$ [3]. We consider spacetimes of topology $M \approx \mathbb{R} \times S_{g,n}$, where $S_{g,n}$ is an oriented surface of genus g with n punctures representing massive point particles with spin.

It is shown in [3] that the physical or gauge-invariant phase space \mathcal{P} of $\text{ISO}(2,1)$ -Chern-Simons theory on a manifold $M \approx \mathbb{R} \times S_{g,n}$ is the moduli space of flat $\text{ISO}(2,1)$ -connections on $S_{g,n}$. This phase space can be parametrised in terms of group homomorphisms $h : \pi_1(S_{g,n}) \rightarrow \text{ISO}(2,1)$ from the fundamental group $\pi_1(S_{g,n})$ into $\text{ISO}(2,1)$ that map the homotopy equivalence class of a loop m_i around the i -th puncture to a fixed $\text{ISO}(2,1)$ -conjugacy class \mathcal{C}_i determined by the mass $\mu_i \in [0, 2\pi)$ and spin $s_i \in \mathbb{R}$ of the associated particle [20]:

$$h(m_i) \in \mathcal{C}_i = \{h \cdot \exp(-\mu_i J_0 - s_i P_0) \cdot h^{-1} \mid h \in \text{ISO}(2,1)\}.$$

Two group homomorphisms $h : \pi_1(S_{g,n}) \rightarrow \text{ISO}(2,1)$ describe the same physical state if and only if they are related by conjugation with $\text{ISO}(2,1)$. This implies that the gauge-invariant phase space of the theory is given by

$$\begin{aligned} \mathcal{P} &= \text{Hom}_{\mathcal{C}_1, \dots, \mathcal{C}_n}(\pi_1(S_{g,n}), \text{ISO}(2,1)) / \text{ISO}(2,1) \\ &= \{h : \pi_1(S_{g,n}) \rightarrow \text{ISO}(2,1) \mid h(m_i) \in \mathcal{C}_i\} / \text{ISO}(2,1). \end{aligned} \quad (3)$$

A set of generators of the fundamental group $\pi_1(S_{g,n})$ is depicted in Figure 1. It consists of a loop m_i ($i = 1, \dots, n$) around each puncture and the a - and b -cycles a_j, b_j ($j = 1, \dots, g$) for each handle. These generators are subject to a single defining relation which states that the curve c in Figure 1 is contractible:

$$\pi_1(S_{g,n}) = \langle m_1, \dots, m_n, a_1, b_1, \dots, a_g, b_g \mid b_g a_g^{-1} b_g^{-1} a_g \cdots b_1 a_1^{-1} b_1^{-1} a_1 m_n \cdots m_1 = 1 \rangle.$$

By characterising group homomorphisms $h : \pi_1(S_{g,n}) \rightarrow \text{ISO}(2,1)$ in terms of the images $M_i = h(m_i)$, $A_j = h(a_j)$, $B_j = h(b_j)$ of the generators, one obtains a parametrisation of the phase space \mathcal{P} in terms of $\text{ISO}(2,1)$ -matrices:

$$\begin{aligned} \mathcal{P} &= \{(M_1, \dots, M_n, A_1, B_1, \dots, A_g, B_g) \in \text{ISO}(2,1)^{n+2g} \mid \\ &\quad M_i \in \mathcal{C}_i, [B_g, A_g^{-1}] \cdots [B_1, A_1^{-1}] \cdot M_n \cdots M_1 = 1\} / \text{ISO}(2,1). \end{aligned} \quad (4)$$

In the following, we refer to the images of elements $d \in \pi_1(S_{g,n})$ under these group homomorphisms as holonomies and denote them by the associated capital letter $D = h(d)$ for $d \in \pi_1(S_{g,n})$.

The moduli space \mathcal{P} of flat $\text{ISO}(2,1)$ -connections is equipped with a symplectic structure that is induced by the canonical symplectic structure associated with the Chern-Simons action [21]. A convenient description of this Poisson structure which serves as the starting point for its quantisation is derived in [22, 23]. In this description, the Poisson structure on the moduli space is characterised in terms of an auxiliary, non-canonical Poisson structure on an extended phase space $\mathcal{P}_{\text{ext}} = \text{ISO}(2,1)^{n+2g}$. This Poisson structure depends on the choice of a classical r -matrix $r = r^{\alpha\beta} T_\alpha \otimes T_\beta \in \mathfrak{iso}(2,1) \otimes \mathfrak{iso}(2,1)$ and is given in terms of a Poisson bivector:

$$\begin{aligned} \{F, G\} &= B_{\text{FR}}^r(dF \otimes dG) \quad \forall F, G \in C^\infty(\text{ISO}(2,1)^{n+2g}), \\ B_{\text{FR}}^r &= \frac{1}{2} r_{(a)}^{\alpha\beta} \left(\sum_{i=1}^n L_\alpha^{M_i} + R_\alpha^{M_i} + \sum_{j=1}^g L_\alpha^{A_j} + R_\alpha^{A_j} + L_\alpha^{B_j} + R_\alpha^{B_j} \right) \otimes \\ &\quad \left(\sum_{i=1}^n L_\beta^{M_i} + R_\beta^{M_i} + \sum_{j=1}^g L_\beta^{A_j} + R_\beta^{A_j} + L_\beta^{B_j} + R_\beta^{B_j} \right) \\ &\quad + \frac{1}{2} r_{(s)}^{\alpha\beta} \left(\sum_{i=1}^n R_\alpha^{M_i} \wedge R_\beta^{M_i} + \sum_{j=1}^g R_\alpha^{A_j} \wedge L_\beta^{A_j} + L_\alpha^{B_j} \wedge L_\beta^{B_j} \right. \\ &\quad \left. + \sum_{1 \leq i < j \leq n} \left(L_\alpha^{M_i} + R_\alpha^{M_i} \right) \wedge \left(L_\beta^{M_j} + R_\beta^{M_j} \right) \right. \\ &\quad \left. + \sum_{i=1}^n \sum_{j=1}^g \left(L_\alpha^{M_i} + R_\alpha^{M_i} \right) \wedge \left(L_\beta^{A_j} + R_\beta^{A_j} + L_\beta^{B_j} + R_\beta^{B_j} \right) \right. \\ &\quad \left. + \sum_{1 \leq i < j \leq g} \left(L_\alpha^{A_i} + R_\alpha^{A_i} + L_\alpha^{B_i} + R_\alpha^{B_i} \right) \wedge \left(L_\beta^{A_j} + R_\beta^{A_j} + L_\beta^{B_j} + R_\beta^{B_j} \right) \right), \quad (5) \end{aligned}$$

where $r_{(s)}^{\alpha\beta} = \frac{1}{2}(r^{\alpha\beta} + r^{\beta\alpha})$, $r_{(a)}^{\alpha\beta} = \frac{1}{2}(r^{\alpha\beta} - r^{\beta\alpha})$ denote the symmetric and antisymmetric components of r and L_α^X, R_α^X the right- and left-invariant vector fields (2) associated with a basis $\{T_\alpha\}_{\alpha=0,\dots,5}$ of $\mathfrak{iso}(2,1)$ and with the different copies of $\text{ISO}(2,1)$. It is shown in [22] that this bivector defines a Poisson structure on \mathcal{P}_{ext} if and only if r is a classical r -matrix for $\mathfrak{iso}(2,1)$, i.e. a solution of the classical Yang-Baxter equation

$$[r_{12}, r_{13}] + [r_{12}, r_{23}] + [r_{13}, r_{23}] = 0,$$

with $r_{12} = r^{\alpha\beta} T_\alpha \otimes T_\beta \otimes 1$, $r_{13} = r^{\alpha\beta} T_\alpha \otimes 1 \otimes T_\beta$, $r_{23} = r^{\alpha\beta} 1 \otimes T_\alpha \otimes T_\beta$. In this case, the resulting Poisson structure induces a symplectic structure on the moduli space \mathcal{P} , which agrees with the canonical symplectic structure induced by the Chern-Simons action if and only if the symmetric part of r is dual to the Ad-invariant symmetric form (1) in the Chern-Simons action:

$$r_S \equiv r_{(s)}^{\alpha\beta} T_\alpha \otimes T_\beta = \frac{1}{2}(P_a \otimes J^a + J^a \otimes P_a).$$

In the application to (2+1)-gravity with vanishing cosmological constant, a natural choice for the r -matrix which satisfies this condition is given by $r = P_a \otimes J^a$. A detailed discussion of the Poisson structure on $\text{ISO}(2,1)^{n+2g}$ obtained from this choice of r -matrix is given in [24], see also [17].

2.3 Geometrical interpretation

Although the relation between $\text{ISO}(2,1)$ -Chern Simons theory and (2+1)-gravity with vanishing cosmological constant is subtle [25, 26], the link between the two theories is close enough to




Figure 2: Conical spacetime associated with a point particle. The two horizontal black lines depict the identification of the boundary of the wedge according to $(t, r, 0) \sim (t + s, r, \mu)$.

provide us with a direct geometrical interpretation of the description of the gauge-invariant phase space \mathcal{P} in formula (4).

As the Ricci tensor of a three-dimensional manifold determines its curvature uniquely, vacuum solutions of the three-dimensional Einstein equations with vanishing cosmological constant are flat and locally isometric to Minkowski space. The theory has no local gravitational degrees of freedom and only a finite number of non-local degrees of freedom due to the presence of matter (point particles) and the topology of the spacetime. These non-local degrees of freedom are encoded in Poincaré transformations that describe the construction of spacetimes from open regions in Minkowski space and are closely related to the holonomies in the previous subsection.

The construction of spacetimes from regions in Minkowski space is illustrated by the simplest example, namely the spacetime associated with a single point particle in Minkowski space. The metric associated with a particle in three-dimensional Minkowski space that is at rest at the origin was derived in [27], see [28] for earlier results on the Euclidean case. It is shown there that in cylindrical coordinates (τ, ρ, ϕ) it takes the form

$$ds^2 = (d\tau + \frac{s}{2\pi} d\phi)^2 - \frac{1}{(1 - \frac{\mu}{2\pi})^2} d\rho^2 - \rho^2 d\phi^2, \quad (6)$$

where $\mu \in [0, 2\pi)$ is the mass and $s \in \mathbb{R}$ the spin of the particle. By introducing a new set of coordinates (t, r, φ) that is related to the cylindrical coordinates (τ, ρ, ϕ) via

$$t(\tau, \rho, \phi) = \tau + \frac{s}{2\pi} \phi, \quad r(\tau, \rho, \phi) = \frac{\rho}{1 - \frac{\mu}{2\pi}}, \quad \varphi(\tau, \rho, \phi) = (1 - \frac{\mu}{2\pi})\phi,$$

one finds that the metric (6) can be related to the Minkowski metric

$$ds^2 = dt^2 - dr^2 - r^2 d\varphi^2.$$

However, this metric is only locally, not globally, isometric to the Minkowski metric. This is apparent from the range of the variable φ , which is no longer $[0, 2\pi)$ but $[0, 2\pi - \mu)$. The metric thus describes a conical spacetime obtained by cutting out a wedge of Minkowski space and identifying its boundary according to $(t, r, 0) \sim (t + s, r, \mu)$ as shown in Figure 2. This identification is given by the action of the Poincaré transformation $M = (\exp(-\mu J_0), -se_0)$.

By applying a Poincaré transformation $(v, \mathbf{x}) \in \text{ISO}(2, 1)$, this description of the point particle metric is easily generalised to particles whose worldline is a general timelike geodesic in Minkowski space. Such a geodesic can be parametrised as $g(T) = T\hat{\mathbf{p}} + \mathbf{x}$ with $\mathbf{x} \cdot \mathbf{p} = 0$,




Figure 3: Domain in Minkowski space for a spacetime of genus $g = 2$ with $n = 4$ particles.

$\hat{\mathbf{p}}^2 = 1$, where T is the eigentime of the particle, $\hat{\mathbf{p}}$ its unit momentum three-vector and \mathbf{x} the position of the particle at $T = 0$. In this case, the identification of the boundaries of the wedge is given by a Poincaré transformation $M \in \text{ISO}(2, 1)$ which takes the form

$$M = (u, -\text{Ad}(u)\mathbf{j}) = (v, \mathbf{x}) \cdot (\exp(-\mu J_0), -s e_0) \cdot (v, \mathbf{x})^{-1}, \quad u, v \in \text{SO}_+(2, 1), \mathbf{j}, \mathbf{x} \in \mathbb{R}^3.$$

The relation between the variables u, \mathbf{j} and v, \mathbf{x} is given by

$$u = v \cdot \exp(-\mu J_0) \cdot v^{-1} = \exp(-\mu \hat{\mathbf{p}}^c J_c), \quad \mathbf{j} = s\hat{\mathbf{p}} + (\mathbb{1} - \text{Ad}(u^{-1}))\mathbf{x}. \quad (7)$$

The quantity $\mathbf{p} = \mu\hat{\mathbf{p}} = \mu \text{Ad}(v)e_0$ has the interpretation of a momentum three-vector of the particle, whose 0-component describes its energy and whose 1- and 2-component its momentum. The quantity \mathbf{j} can be viewed as a generalised angular momentum three-vector. Its 0-component describes the angular momentum of the particle, its 2- and 3-component are related to Lorentz boosts. From the formula above it is apparent that the angular momentum three-vector is composed of two parts. The first one is parallel to its momentum three-vector and given by the particle's spin. The second one is orthogonal to the momentum three-vector and encodes the angular momentum of the particle due to its motion. In the limit $\mu \rightarrow 0$, one has $\mathbf{j} \rightarrow s\hat{\mathbf{p}} + \hat{\mathbf{p}} \wedge \mathbf{x}$ and hence recovers the usual expression for the angular momentum.

The general case of a spacetime $M \approx \mathbb{R} \times S_{g,n}$ of genus g with n point particles is more involved, but the construction of spacetimes by identifying the boundary of an open region in Minkowski space is similar. The main difference is that the associated region in Minkowski space is no longer obtained by cutting out a wedge but takes a more complicated form depicted schematically in Figure 3. Its boundary is given by $2(n + 2g)$ plane segments in Minkowski space, which are identified pairwise by certain Poincaré transformations. These Poincaré transformations correspond to the images of the generators of the fundamental group $\pi_1(S_{g,n})$ under the group homomorphisms $h : \pi_1(S_{g,n}) \rightarrow \text{ISO}(2, 1)$ in formula (4). The associated spacetime is obtained by gluing these sides pairwise as shown in Figures 3, 4.

The condition that the holonomies for the loops m_i around the punctures lie in fixed conjugacy classes \mathcal{C}_i is needed to ensure that the identification of the sides corresponding to point particles gives rise to a cone with the correct opening angle and time shift defined by the




Figure 4: Gluing of a spacetime of genus $g = 0$ with $n = 4$ point particles.

mass and spin of the particle. The condition that arises from the defining relation of the fundamental group ensures that all black vertices in Figure 3 are mapped to a single point in the spacetime and guarantees that the gluing procedure gives rise to a flat Lorentzian manifold with conical singularities. The holonomies from the previous subsection thus have a direct interpretation as gluing data that describes the construction of spacetimes from regions in Minkowski space.

Given a region in three-dimensional Minkowski space whose sides are identified by the holonomies $h(m_i), h(a_j), h(b_j)$, one obtains another such region by applying a Poincaré transformation $g \in \text{ISO}(2, 1)$ to it. The sides of the image are then identified by the images of the holonomies under the diagonal action of $\text{ISO}(2, 1)$ on $\mathcal{C}_1 \times \dots \times \mathcal{C}_n \times \text{ISO}(2, 1)^{2g}$: $h(m_i) \mapsto g \cdot h(m_i) \cdot g^{-1}$, $h(a_j) \mapsto g \cdot h(a_j) \cdot g^{-1}$, $h(b_j) \mapsto g \cdot h(b_j) \cdot g^{-1}$. As Poincaré transformations are isometries of Minkowski space, the spacetimes obtained by gluing these two regions are isometric and hence describe the same physical state. This motivates the appearance of the quotient by the action of $\text{ISO}(2, 1)$ in formulas (3), (4). Poincaré transformations which act diagonally on all holonomies have the status of gauge transformations: they are transitions between two different parametrisations of the same physical state.

3 Gauge fixing and observers

3.1 The phase space of (2+1)-gravity as a constrained system

Formula (4) implies that the description of the phase space of (2+1)-gravity in terms of the auxiliary Poisson structure (5) can be viewed as a constrained system in the sense of Dirac. From this viewpoint, one has an extended phase space $\mathcal{P}_{\text{ext}} = \text{ISO}(2, 1)^{n+2g}$, equipped with the Poisson structure (5). The physical (or gauge-invariant) phase space is obtained from this extended phase space by imposing a set of $2n$ constraints which restrict the holonomies of the particles to the conjugacy classes \mathcal{C}_i and an $\text{ISO}(2, 1)$ -valued constraint which arises from the defining relation of the fundamental group $\pi_1(S_{g,n})$. The former can be expressed as

$$\text{Tr}(u_{M_i}) - \cos \mu_i \approx 0, \quad \text{Tr}(j_{M_i}^a J_a \cdot u_{M_i}) - s_i \sin \mu_i \approx 0. \quad (8)$$

vanishes on the constraint surface $\bigcap_{i=1}^k \phi_i^{-1}(0) \subset M$. Gauge fixing consists in imposing an additional set of constraints $\{\chi_j\}_{j=1,\dots,k} \subset C^\infty(\mathcal{P}_{\text{ext}})$, the gauge fixing conditions, such that the following two requirements are met:

1. It is possible to map any point $p \in \bigcap_{i=1}^k \phi_i^{-1}(0)$ on the constraint surface to one that satisfies the gauge fixing conditions via the flows on M generated by the constraint functions ϕ_i .
2. The gauge fixing conditions must break the gauge symmetries completely. In other words, the matrix $C = (\{\phi_i, \chi_j\})_{i,j=1,\dots,k}$ must be invertible at least on the gauge-fixed constraint surface $\Sigma := \bigcap_{i=1}^k \phi_i^{-1}(0) \cap \chi_i^{-1}(0)$.

If these conditions are satisfied, the Dirac matrix $D = (\{C_i, C_j\})_{i,j=1,\dots,2k}$ obtained by combining the constraints ϕ_i and the gauge fixing conditions χ_j into a set of $2k$ constraints $\{C_i\}_{i=1,\dots,2k}$ is invertible on Σ . The Dirac bracket $\{, \}_D$ is given in terms of its inverse as

$$\{F, G\}_D := \{F, G\} - \sum_{i,j=1}^{2k} \{F, C_i\} (D^{-1})_{ij} \{C_j, G\} \quad \forall F, G \in C^\infty(M).$$

and defines a Poisson structure on Σ for which the constraints ϕ_i and gauge fixing conditions χ_i are Casimir functions. This allows one to strongly impose these constraints in the classical theory.

In the application to the phase space of (2+1)-gravity, the Poisson manifold $(M, \{, \})$ is the manifold $\mathcal{P}_{\text{ext}} = \text{ISO}(2, 1)^{n+2g}$ with the Poisson bracket (5) and the constraint functions ϕ_i are given by (9). The associated gauge transformations these constraints generate via the Poisson bracket correspond to the diagonal action of $\text{ISO}(2, 1)$ on $\text{ISO}(2, 1)^{n+2g}$ and encode the gauge freedom of applying a Poincaré transformation to the regions in Minkowski space from which spacetimes are obtained by gluing.

This gauge freedom is linked to the absence of a preferred observer in general relativity. Physical measurements of quantities such as the distance between certain particles or quantities associated with the geometry of the handles on $S_{g,n}$ depend on the choice of the observer. If one restricts attention to observers in free fall, each observer corresponds to a timelike geodesic in Minkowski space and the Poincaré transformations relating different domains in Minkowski space describe the transitions between different observers. Eliminating this gauge freedom via gauge fixing conditions thus corresponds to specifying an observer.

In the absence of a preferred reference frame, the only physically meaningful way of specifying an observer is with respect to the geometry of the spacetime itself, for instance with respect to the geometry of a handle or two point particles contained in the spacetime. A detailed discussion of this issue is given in [10] and in [9] for the case of vacuum spacetimes.

In the following, we will restrict attention to gauge fixing conditions based on the motion of two point particles. As permutations of the particles correspond to the action of the braid groups on $S_{g,n}$ and the associated surface $S_{g,n} \setminus D$ with a disc removed and these braid groups act by Poisson isomorphisms [24], we can suppose without restriction of generality that these point particles are the ones associated with the holonomies M_1, M_2 .

A set of particularly simple gauge fixing condition of this type is investigated in [10]. These gauge fixing conditions impose that the first particle is at rest at the origin and that the second particle moves in the direction of the x_1 -axis in such a way that its distance to the

Theorem 4.5. The role of the rescaling in Theorem 4.5 is to eliminate the freedom in defining ψ, α as functions of the Lorentzian and translational component of the holonomy $M_2 \cdot M_1$ in (12) and defines the mass and spin of the associated cone as a function of y_1 and y_2 .

From the discussion in the previous sections it then follows that the timelike solution in Theorem 4.5 corresponds to an observer in the centre-of-mass frame of the universe, to whom the centre of mass appears as a particle that is at rest at the origin. The interpretation of the spacelike solution in Theorem 4.5 is less direct. It arises for those values of the parameter ψ for which the gauge-fixed holonomies M_1, M_2 form a Gott pair [31] and can be viewed as a “tachyonic” particle whose worldline is identified with the x_1 -axis.

This amounts to a complete *local* classification of all possible Dirac brackets for those values of the parameter ψ for which $\mathbf{q}_\psi^2(\psi), \mathbf{q}_\alpha^2(\psi) \neq 0$. Theorem 4.5 states that every such solution can be identified with one of the two standard solutions above for all values of the parameter ψ for which $\mathbf{q}_\psi^2(\psi), \mathbf{q}_\alpha^2(\psi) \neq 0$. The possible outcomes of the gauge fixing procedure are thus equivalent to the ones associated with the centre-of-mass frames, where the centre of mass of the residual system appears as a particle at rest at the origin or a “tachyonic” particle associated with the x_1 -axis.

However, it is shown in [10, 18] that for generic Dirac brackets resulting from our gauge fixing conditions the signature of $\mathbf{q}_\psi, \mathbf{q}_\alpha$ varies with ψ . In particular, the solutions described there exhibit values of the variable ψ for which these vectors become lightlike. These lightlike solutions appear as transition points between the time- and spacelike cases discussed above and do not correspond to a fixed Cartan subalgebra of $\mathfrak{iso}(2, 1)$. The solutions of the CDYBE resulting from gauge fixing in (2+1)-gravity are therefore not simply given by standard classical dynamical r -matrices as in [19] but connect in a non-trivial way non-equivalent standard classical dynamical r -matrices associated with non-conjugate Cartan subalgebras of $\mathfrak{iso}(2, 1)$.

5 Outlook and conclusions

In this article we investigated gauge fixing in (2+1)-dimensional gravity with vanishing cosmological constant. We discussed how gauge fixing this theory amounts to the introduction of an observer and determined the resulting Dirac bracket for a rather general set of gauge fixing conditions. We showed that these Dirac brackets are determined by solutions of the classical dynamical Yang-Baxter equation, whose two dynamical variables have a direct interpretation as the total energy and total angular momentum of the spacetime measured by this observer. We showed how dynamical Poincaré transformations allow one to relate, for almost all values of the dynamical variables, any Dirac bracket obtained from this gauge fixing to two particularly simple standard solutions which correspond to the centre-of-mass frame of the spacetime.

That these statements hold for a very large class of gauge fixing conditions and that classical dynamical r -matrix symmetries were also obtained in a different and independent approach based on the regularisation of punctures coupled to Chern-Simons theory [32, 33, 34] suggest that the appearance of classical dynamical r -matrices in gauge-fixed (2+1)-gravity is generic and related to the introduction of observers into the theory.

From the physics perspective, this result suggests that the relevant quantum groups for the quantisation of (2+1)-gravity should become dynamical quantum groups when an observer is implemented in the quantum theory. As the dynamical variables in the classical theory are

the total energy and angular momentum measured by this observer, the dynamical quantum group symmetries, unlike the quantum groups acting on extended, non-gauge-invariant Hilbert spaces, should manifest themselves in the measurements of observers.

It is also instructive to compare the classical dynamical r -matrices arising in this description to the Hopf algebra symmetries associated with other models such as κ -Poincaré symmetries. By comparing the r -matrices appearing in these descriptions, one finds that the classical dynamical r -matrices in Theorem 4.5 are rather similar to the ones occurring in (2+1)-dimensional κ -Poincaré symmetries but do not coincide with them. The difference is that the classical dynamical r -matrices in this article also contain a term which involves the tensor product of two generators of translations, which is not present in the classical r -matrices of κ -Poincaré symmetries. It would be interesting to investigate if κ -Poincaré symmetries can be obtained from the classical dynamical r -matrices in Theorem 4.5 via a suitable limiting procedure in which the angular momentum of the cone tends to zero.

Finally, it would be desirable to quantise the gauge-fixed Poisson structure derived in this article. As the Dirac brackets are of a form very similar to the Poisson structure in [22, 23] which serves as the starting point of the combinatorial quantisation formalism, this formalism could also be applied to quantise the gauge-fixed theory. In that case the difficulties associated with the implementation of the constraints would no longer be present and the quantisation of the theory would reduce to the construction of the dynamical quantum groups that correspond to the classical dynamical r -matrices in the classical description. In particular, it would be interesting to see if this construction gives rise to quantisation conditions on the total mass and angular momentum of the spacetime which are encoded in its dynamical variables.

References

- [1] S. Carlip. *Quantum Gravity in 2+1 Dimensions*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2003. ISBN 0521545889.
- [2] A. Achúcarro and P. Townsend. “A Chern-Simons action for three-dimensional anti-de Sitter supergravity theories.” *Physics Letters B*, **180** (1986) 89–92. DOI:10.1016/0370-2693(86)90140-1.
- [3] E. Witten. “2 + 1 dimensional gravity as an exactly soluble system.” *Nuclear Physics B*, **311** (1988) 46–78. DOI:10.1016/0550-3213(88)90143-5.
- [4] G. Mess. “Lorentz spacetimes of constant curvature.” *Geometriae Dedicata*, **126** (2007) 3–45. DOI:10.1007/s10711-007-9155-7.
- [5] R. Benedetti and F. Bonsante. “Canonical Wick rotations in 3-dimensional gravity.” (2005) 201. arXiv:math/0508485.
- [6] E. Witten. “Quantum field theory and the Jones polynomial.” *Communications in Mathematical Physics*, **121** (1989) 351–399. DOI:10.1007/BF01217730.
- [7] N. Y. Reshetikhin and V. G. Turaev. “Invariants of 3-manifolds via link polynomials and quantum groups.” *Inventiones Mathematicae*, **103** (1991) 547–597. DOI:10.1007/BF01239527.

- [8] E. Witten. “Topology-changing amplitudes in $2 + 1$ dimensional gravity.” *Nuclear Physics B*, **323** (1989) 113–140. DOI:10.1016/0550-3213(89)90591-9.
- [9] C. Meusburger. “Cosmological measurements, time and observables in $(2+1)$ -dimensional gravity.” *Classical and Quantum Gravity*, **26** (2009) 55006. DOI:10.1088/0264-9381/26/5/055006. arXiv:0811.4155.
- [10] C. Meusburger and T. Schönfeld. “Gauge fixing in $(2+1)$ -gravity: Dirac bracket and spacetime geometry.” *Classical and Quantum Gravity*, **28** (2011) 125008. DOI:10.1088/0264-9381/28/12/125008. arXiv:1012.1835.
- [11] J. Lukierski, H. Ruegg, A. Nowicki, and V. N. Tolstoy. “q-deformation of Poincaré algebra.” *Physics Letters B*, **264** (1991) 331–338. DOI:10.1016/0370-2693(91)90358-W.
- [12] J. Lukierski, A. Nowicki, and H. Ruegg. “New quantum Poincaré algebra and κ -deformed field theory.” *Physics Letters B*, **293** (1992) 344–352. DOI:10.1016/0370-2693(92)90894-A.
- [13] S. Majid and H. Ruegg. “Bicrossproduct structure of κ -Poincare group and non-commutative geometry.” *Physics Letters B*, **334** (1994) 348–354. DOI:10.1016/0370-2693(94)90699-8. arXiv:hep-th/9405107.
- [14] J. Kowalski-Glikman and S. Nowak. “Doubly special relativity theories as different bases of κ -Poincaré algebra.” *Physics Letters B*, **539** (2002) 126–132. DOI:10.1016/S0370-2693(02)02063-4. arXiv:hep-th/0203040.
- [15] L. Freidel, J. Kowalski-Glikman, and L. Smolin. “ $2+1$ gravity and doubly special relativity.” *Physical Review D*, **69** (2004) 044001. DOI:10.1103/PhysRevD.69.044001. arXiv:hep-th/0307085.
- [16] C. Meusburger and B. J. Schroers. “Generalised Chern-Simons actions for 3d gravity and κ -Poincaré symmetry.” *Nuclear Physics B*, **806** (2009) 462–488. DOI:10.1016/j.nuclphysb.2008.06.023. arXiv:0805.3318.
- [17] C. Meusburger and K. Noui. “The Hilbert space of 3d gravity: quantum group symmetries and observables.” *Adv. Theor. Math. Phys.*, **14** (2010) 1651–1716. arXiv:0809.2875.
- [18] C. Meusburger and T. Schönfeld. “Gauge fixing and classical dynamical r-matrices in ISO(2,1)-Chern-Simons theory.” (2012). arXiv:1203.5609.
- [19] P. Etingof and A. Varchenko. “Geometry and Classification of Solutions of the Classical Dynamical Yang-Baxter Equation.” *Communications in Mathematical Physics*, **192** (1998) 77–120. DOI:10.1007/s002200050292. arXiv:q-alg/9703040.
- [20] P. de Sousa Gerbert. “On spin and (quantum) gravity in $2 + 1$ dimensions.” *Nuclear Physics B*, **346** (1990) 440–472. DOI:10.1016/0550-3213(90)90288-O.
- [21] M. Atiyah and R. Bott. “The Yang-Mills equations over Riemann surfaces.” *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, **54** (1983) 523–615. DOI:10.1098/rsta.1983.0017.
- [22] V. V. Fock and A. A. Rosly. “Poisson structure on moduli of flat connections on Riemann surfaces and r -matrix.” *Am. Math. Soc. Transl.*, **191** (1999) 67–86. arXiv:math/9802054.

- [23] A. Y. Alekseev and A. Z. Malkin. “Symplectic structure of the moduli space of flat connection on a Riemann surface.” *Communications in Mathematical Physics*, **169** (1995) 99–119. DOI:10.1007/BF02101598. arXiv:hep-th/9312004v1.
- [24] C. Meusburger and B. J. Schroers. “Poisson structure and symmetry in the Chern-Simons formulation of (2+1)-dimensional gravity.” *Classical and Quantum Gravity*, **20** (2003) 2193–2233. DOI:10.1088/0264-9381/20/11/318. arXiv:gr-qc/0301108.
- [25] H.-J. Matschull. “On the relation between 2+1 Einstein gravity and Chern-Simons theory.” *Classical and Quantum Gravity*, **16** (1999) 2599–2609. DOI:10.1088/0264-9381/16/8/303. arXiv:gr-qc/9903040.
- [26] E. Witten. “Three-Dimensional Gravity Revisited.” (2007). arXiv:0706.3359.
- [27] S. Deser, R. Jackiw, and G. ’t Hooft. “Three-dimensional Einstein gravity: Dynamics of flat space.” *Annals of Physics*, **152** (1984) 220–235. DOI:10.1016/0003-4916(84)90085-X.
- [28] A. Staruszkiewicz. “Gravitation theory in three-dimensional space.” *Acta Phys. Polon.*, **24** (1963) 735–740.
- [29] E. Buffenoir, K. Noui, and P. Roche. “Hamiltonian quantization of Chern Simons theory with $SL(2, \mathbb{C})$ group.” *Classical and Quantum Gravity*, **19** (2002) 4953–5015. DOI:10.1088/0264-9381/19/19/313. arXiv:hep-th/0202121.
- [30] C. Meusburger and K. Noui. “Combinatorial quantisation of the Euclidean torus universe.” *Nuclear Physics B*, **841** (2010) 463–505. DOI:10.1016/j.nuclphysb.2010.08.014. arXiv:1007.4615.
- [31] J. Gott. “Closed timelike curves produced by pairs of moving cosmic strings: Exact solutions.” *Physical Review Letters*, **66** (1991) 1126–1129. DOI:10.1103/PhysRevLett.66.1126.
- [32] E. Buffenoir and K. Noui. “Unfashionable observations about three dimensional gravity.” (2003). arXiv:gr-qc/0305079.
- [33] E. Buffenoir and P. Roche. “Chern-Simons Theory with Sources and Dynamical Quantum Groups I: Canonical Analysis and Algebraic Structures.” (2005). arXiv:hep-th/0505239.
- [34] K. Noui and A. Perez. “Three-dimensional loop quantum gravity: coupling to point particles.” *Classical and Quantum Gravity*, **22** (2005) 4489–4513. DOI:10.1088/0264-9381/22/21/005. arXiv:gr-qc/0402111.