

Neutral 3-3-1 Higgs Boson at LHC

J. E. Cieza Montalvo¹

¹*Instituto de Física, Universidade do Estado do Rio de Janeiro,
Rua São Francisco Xavier 524, 20559-900 Rio de Janeiro, RJ, Brazil*

R. J. Gil Ramírez, G. H. Ramírez Ulloa, A. I. Rivasplata Mendoza²

²*Universidad Nacional de Trujillo, Departamento de Física,
Av. Juan Pablo II S/N; Ciudad Universitaria*

M. D. Tonasse^{3*}

³*Instituto de Física Teórica, Universidade Estadual Paulista,
Rua Dr. Bento Teobaldo Ferraz 271, 01140-070 São Paulo, SP, Brazil*

(Dated: September 1, 2020)

Abstract

We present an analysis of production and signature of neutral Higgs bosons on the version of the 3-3-1 model containing heavy leptons at the Large Hadron Collider (LHC). The production rate is found to be significant and the signal is clear, showing that these scalars can be detected in this accelerator. We also studied the possibility to identify them using their respective branching ratios. Cross section are given for two collider energies, $\sqrt{s} = 8$ TeV and 14 TeV. Event rates and significances are discussed for two possible values of integrated luminosity, 10 fb^{-1} and 300 fb^{-1} .

PACS numbers:

11.15.Ex: Spontaneous breaking of gauge symmetries,

12.60.Fr: Extensions of electroweak Higgs sector,

14.80.Cp: Non-standard-model Higgs bosons.

* Permanent address: Universidade Estadual Paulista, *Campus* Experimental de Registro, Rua Nelson Brihi Badur 430, 11900-000 Registro, SP, Brazil

I. INTRODUCTION

The way to the understanding of the symmetry breaking mainly go through the scalars, although there are many other models that not contain elementary scalar fields, such as Nambu-Jona-Lasinio mechanism, technicolour theories, the strongly interacting gauge systems [1]. These scalars protect the renormalizability of the theory by moderating the cross section growth. But so far, despite many experimental and theoretical efforts in order to understanding the scalar sector, the Higgs mechanism remains still unintelligible. Nowadays, the major goal of the experimentalists in particle physics at the Large Hadron Collider (LHC), is to unravel the nature of electroweak symmetry breaking. The Standard Model (SM) is the prototype of a gauge theory with spontaneous symmetry breaking. This had great success in explaining the most of the experimental data. However, recent results from neutrino oscillation experiments makes clear that the SM is not complete, then the neutrino oscillation implies that at least two neutrino flavors are massive. Moreover, there are others crucial problems in particle physics that not get response in SM. For instance, it offers not solution to the dark matter problem, dark energy and the asymmetry of matter-antimatter in the Universe. Therefore, there is a consensus among the particle physicists that the SM must be extended.

In the SM appear only one elementary scalar, which arises through the breaking of electroweak symmetry and this is the Higgs boson. The Higgs Boson is an important prediction of several quantum field theories and is so crucial to our understanding of the Universe. The Higgs boson is the one missing piece, is a critical ingredient to complete our understanding of the SM. Different types of Higgs bosons, if they exist, may lead us into new realms of physics beyond the SM. So, the observation of any kind of Higgs particle must be an important step forward in the understanding of physics in the electroweak sector or beyond the SM.

Since the SM leaves many questions open, there are several extensions. For example, if the Grand Unified Theory (GUT) contains the SM at high energies, then the Higgs bosons associated with GUT symmetry breaking must have masses of order $M_X \sim \mathcal{O}(10^{15})$ GeV. Supersymmetry [2] provides a solution to hierarchy problem through the cancellation of the quadratic divergences via fermionic and bosonic loops contributions [3]. Moreover, the Minimal Supersymmetric extension of the SM (MSSM) can be derived as an effective theory

of supersymmetric Grand Unified Theories [4].

Among these extensions of the SM there are also other class of models based on $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ gauge symmetry (3-3-1 model) [5–7], where the anomaly cancellation mechanisms occur when the three basic fermion families are considered and not family by family as in the SM. This mechanism is peculiar because it requires that the number of families is an integer multiple of the number of colors. This feature combined together with the asymptotic freedom, which is a property of quantum chromodynamics (QCD), requires that the number of colors is less than five, and therefore as a consequence of this, the number of family of fermions must be exactly equal to three. Moreover, according to these models, the Weinberg angle is restricted to the value $s_W^2 = \sin^2 \theta_W < 1/4$ in the version of heavy-leptons [5], but when evolves to higher values, it shows that the model loses its perturbative character when it reaches to mass scale of about 4 TeV [8]. Then, the 3-3-1 model is one of the most interesting extensions of the SM and is phenomenologically well motivated to be probed at the LHC and other accelerators.

In this work we study the production and signatures of two neutral Higgs bosons, predicted by the 3-3-1 model, which incorporates the charged heavy leptons [5, 9]. One of these neutral Higgs is the standard one. We can show that the neutral Higgs boson signatures can be significant at LHC. Clear signal of these new particles can be obtained by studying the different decay modes. With respect to both mechanisms, that is the Drell-Yan and gluon-gluon fusion, we consider the Z , Z' , H_1^0 and H_2^0 as propagators. Therefore in Sec. II we present the relevant features of the model. In Sec. III we compute the total cross sections of the process and in Sec. IV we summarize our results and conclusions.

II. RELEVANT FEATURES OF THE MODEL

We are working here with the version of the 3-3-1 model that contains heavy leptons [5]. The model is based on the semi simple symmetry group $SU(3)_C \otimes SU(2)_L \otimes U(1)_N$. The electric charge operator is given by

$$\frac{Q}{e} = \left(T_3 - \sqrt{3} T_8 \right) + N, \quad (1)$$

where T_3 and T_8 are the generators of $SU(3)$ and e is the elementary electric charge. So, we can build three triplets of quarks of $SU(3)_L$ such that

$$Q_{1L} = \begin{pmatrix} u'_1 \\ d'_1 \\ J_1 \end{pmatrix}_L \sim \left(\mathbf{3}, \frac{2}{3} \right), \quad Q_{\alpha L} = \begin{pmatrix} J'_\alpha \\ u'_\alpha \\ d'_\alpha \end{pmatrix}_L \sim \left(\mathbf{3}^*, -\frac{1}{3} \right). \quad (2)$$

where the new quark J_1 carries $5/3$ units of electric charge while J_α ($\alpha = 2, 3$) carry $-4/3$ each. We must also introduce the right-handed fermionic fields $U_R \sim (\mathbf{1}, 2/3)$, $D_R \sim (\mathbf{1}, -1/3)$, $J_{1R} \sim (\mathbf{1}, 5/3)$ and $J'_{\alpha R} \sim (\mathbf{1}, -4/3)$. We have defined $U = (u' \ c' \ t')$ and $D = (d' \ s' \ b')$.

The spontaneous symmetry breaking is accomplished *via* three $SU(3)$ scalar triplets, which are,

$$\eta = \begin{pmatrix} \eta^0 \\ \eta_1^- \\ \eta_2^+ \end{pmatrix} \sim (\mathbf{3}, 0), \quad \rho = \begin{pmatrix} \rho^+ \\ \rho^0 \\ \rho^{++} \end{pmatrix} \sim (\mathbf{3}, 1), \quad \chi = \begin{pmatrix} \chi^- \\ \chi^{--} \\ \chi^0 \end{pmatrix} \sim (\mathbf{3}, -1). \quad (3)$$

For sake of simplicity, we will assume here that the model respects the $B+L$ symmetry, where B is the baryon number and L is the lepton number. Then, the more general renormalizable Higgs potential is given by

$$\begin{aligned} V(\eta, \rho, \chi) = & \mu_1^2 \eta^\dagger \eta + \mu_2^2 \rho^\dagger \rho + \mu_3^2 \chi^\dagger \chi + \lambda_1 (\eta^\dagger \eta)^2 + \lambda_2 (\rho^\dagger \rho)^2 + \lambda_3 (\chi^\dagger \chi)^2 + \\ & + \eta^\dagger \eta [\lambda_4 (\rho^\dagger \rho) + \lambda_5 (\chi^\dagger \chi)] + \lambda_6 (\rho^\dagger \rho) (\chi^\dagger \chi) + \lambda_7 (\rho^\dagger \eta) (\eta^\dagger \rho) + \\ & + \lambda_8 (\chi^\dagger \eta) (\eta^\dagger \chi) + \lambda_9 (\rho^\dagger \chi) (\chi^\dagger \rho) + \frac{1}{2} (f \varepsilon^{ijk} \eta_i \rho_j \chi_k + \text{c. H.}), \end{aligned} \quad (4)$$

where μ_i ($i = 1, 2, 3$) and f are constants with mass dimension and λ_j ($j = 1, \dots, 9$) are dimensionless constants [9]. The potential (4) is bounded from below when the neutral Higgs fields develops the vacuum expectation values (VEVs) $\langle \eta^0 \rangle = v_\eta$, $\langle \rho^0 \rangle = v_\rho$ and $\langle \chi^0 \rangle = v_\chi$, with $v_\eta^2 + v_\rho^2 = v_W^2 = 246^2 \text{ GeV}^2$. The scalar χ^0 is supposedly heavy and it is responsible for the spontaneous symmetry breaking of $SU(3)_L \otimes U(1)_N$ to $SU(2)_L \otimes U(1)_Y$ of the standard model. Meanwhile, η^0 and ρ^0 are lightweight and are responsible for the breaking of $SU(2)_L \otimes U(1)_Y$ to $U(1)_Q$, of the electromagnetism. Therefore, it is reasonable to expect

$$v_\chi \gg v_\eta, v_\rho. \quad (5)$$

The potential (4) provides the masses of neutral Higgs as

$$m_{H_1^0}^2 \approx 4 \frac{\lambda_2 v_\rho^4 - \lambda_1 v_\eta^4}{v_\eta^2 - v_\rho^2}, \quad m_{H_2^0}^2 \approx \frac{v_W^2 v_\chi^2}{2v_\eta v_\rho}, \quad (6a)$$

$$m_{H_3^0}^2 \approx -\lambda_3 v_\chi, \quad m_h^2 = -\frac{f v_\chi}{v_\eta v_\rho} \left[v_W^2 + \left(\frac{v_\eta v_\rho}{v_\chi} \right)^2 \right] \quad (6b)$$

with the corresponding eigenstates

$$\begin{pmatrix} \xi_\eta \\ \xi_\rho \end{pmatrix} \approx \begin{pmatrix} c_w & s_w \\ s_w & c_w \end{pmatrix} \begin{pmatrix} H_1^0 \\ H_2^0 \end{pmatrix}, \quad \xi_\chi \approx H_3^0, \quad \zeta_\chi \approx ih, \quad (7)$$

where the mixing parameters are $c_w = \cos w = v_\eta / \sqrt{v_\eta^2 + v_\rho^2}$ and $s_w = \sin w$ [9]. In Eqs. (6) and (7) we used the approximation (5) and, for not to introduce the new mass scale in the model, we assume $f \approx -v_\chi$. We can then note that H_3^0 is a typical 3-3-1 Higgs boson. The scalar H_1^0 is one that can be identified with the SM Higgs, since its mass and eigenstate do not depend on v_χ .

Now, we can write the Yukawa interactions for the ordinary quarks, i.e.

$$\mathcal{L}_q^Y = \sum_\alpha \left[\bar{Q}_{1L} \left(G_{1\alpha} U'_{\alpha R} \eta + \tilde{G}_{1\alpha} D'_{\alpha R} \rho \right) + \sum_i \bar{Q}_{iL} \left(F_{i\alpha} U'_{\alpha R} \rho^* + \tilde{F}_{i\alpha} D'_{\alpha R} \eta^* \right) \right], \quad (8)$$

where G_{ab} and G'_{ab} (a and b are generation indexes) are coupling constants.

The interaction eigenstates (2) and their right-handed counterparts can rotate about their respective physical eigenstates as

$$U'_{aL(R)} = \mathcal{U}_{ab}^{L(R)} U_{bL(R)}, \quad (9a)$$

$$D'_{aL(R)} = \mathcal{D}_{ab}^{L(R)} U_{bL(R)}, \quad J'_{aL(R)} = \mathcal{J}_{ab}^{L(R)} J_{bL(R)}, \quad (9b)$$

Since the cross sections involving the sum over the flavors and rotation matrices are unitary, then the mixing parameters have no major effect on the calculations. In terms of physical fields the Yukawa Lagrangian for the neutral Higgs can be written as

$$\begin{aligned} -\mathcal{L}_Q &= \frac{1}{2} \left\{ \bar{U} (1 + \gamma_5) \left[1 + \left[\frac{s_w}{v_\rho} + \left(\frac{c_w}{v_\eta} + \frac{s_w}{v_\rho} \right) \mathcal{V}^U \right] H_1^0 + \right. \right. \\ &\quad \left. \left. + \left[\frac{-c_w}{v_\rho} + \left(\frac{s_w}{v_\eta} - \frac{c_w}{v_\rho} \right) \mathcal{V}^U \right] H_2^0 \right] M^U U + \right. \end{aligned}$$

$$\begin{aligned}
& + \bar{D} (1 + \gamma_5) \left[1 + \left[\frac{c_w}{v_\eta} + \left(\frac{s_w}{v_\rho} - \frac{c_w}{v_\eta} \right) \mathcal{V}^D \right] H_1^0 + \right. \\
& \left. + \left[\frac{s_w}{v_\eta} - \left(\frac{c_w}{v_\rho} + \frac{s_w}{v_\eta} \right) \mathcal{V}^D \right] H_2^0 \right] M^D D \Big\} + \text{H. c.}, \tag{10}
\end{aligned}$$

where $V_L^U V_L^D = V_{\text{CKM}}$ is the Cabibbo-Kobayashi-Maskawa matrix, \mathcal{V}^U and \mathcal{V}^D are arbitrary mixing matrices and $M^U = \text{diag} (m_u \ m_c \ m_t)$ and $M^D = \text{diag} (m_d \ m_s \ m_b)$ are matrices which carrying the masses of the quarks.

In the gauge sector, beyond the standard particles γ , Z , and W^\pm the model predicts: one neutral (Z'), two single-charged (V^\pm), and two doubly-charged ($U^{\pm\pm}$) gauge bosons. The gauge interactions with Higgs bosons are given by

$$\mathcal{L}_{GH} = \sum_{\varphi} (\mathcal{D}_\mu \varphi)^\dagger (\mathcal{D}_\mu \varphi), \tag{11}$$

where the covariant derivatives are

$$\mathcal{D}_\mu \varphi_i = \partial_\mu \varphi_i - ig \left(W_\mu \cdot \frac{T}{2} \right)_i^j \varphi_j - ig' N_\varphi \varphi_i B_\mu, \tag{12}$$

where $\varphi = \eta, \rho, \chi$ ($N_\eta = 0, N_\rho = 1, N_\chi = -1$) are the Higgs triplets, W_μ and B_μ are the SU(2) and U(1) field tensors, g and g' are the U(1) and SU(2) coupling constants, respectively. Diagonalization of the Lagrangean (11), after symmetry breaking, gives masses for the neutral weak gauge bosons, i.e.,

$$m_Z \approx \frac{|e|}{2s_W c_W} v_W, \quad m_{Z'}^2 \approx \frac{1}{3(1 - 4s_W^2)} \left(\frac{|e| c_W v_\chi}{s_W} \right)^2, \tag{13}$$

where $s_W = \sin \theta_W$, with θ_W being the Weinberg angle, and $c_W^2 = 1 - s_W^2$. Then the eigenstates are

$$W_\mu^3 \approx s_W A_\mu - c_W Z_\mu \tag{14a}$$

$$W_\mu^8 \approx -\sqrt{3} s_W \left(A_\mu - \frac{s_W}{c_W} Z_\mu \right) + \frac{\sqrt{1 - 4s_W^2}}{c_W} Z'_\mu \tag{14b}$$

$$B_\mu \approx \frac{s_W}{\sqrt{1 - 4s_W^2}} A_\mu + \frac{s_W}{c_W} \left(Z_\mu + \sqrt{3} Z'_\mu \right). \tag{14c}$$

In Eqs. (13) and (14) we have used the approximation (5). Finally the weak neutral current in the sector of u and d quarks reads

$$-\mathcal{L}_Z = \frac{|e|}{2s_W c_W} \bar{q} \gamma^\mu [v(q) + a(q) \gamma_5] q Z_\mu \tag{15a}$$

$$-\mathcal{L}_{Z'} = \frac{|e|}{2s_W c_W} \bar{q} \gamma^\mu [v'(q) + a'(q) \gamma_5] q Z'_\mu \tag{15b}$$

whose coefficients are

$$v(u) = 1 - \frac{s_W^2}{8}, \quad a(u) = -a(d) = -1, \quad v(d) = -1 + \frac{4}{3}s_W^2, \quad (16a)$$

$$v'(u) = \sqrt{1 + 4s_W^2}, \quad a'(u) = \sqrt{\frac{1 - 4s_W^2}{3}}, \quad v'(d) = \frac{2s_W^2 - 1}{\sqrt{3}}, \quad (16b)$$

$$a'(d) = -v'(d). \quad (16c)$$

In this work we study the production of a neutral Higgs bosons at pp colliders. With respect to both mechanisms, that is the Drell-Yan and gluon-gluon fusion, we consider the Z , Z' , H_1^0 and H_2^0 as propagators.

III. CROSS SECTION PRODUCTION

The mechanisms for the production of neutral Higgs particles in pp collisions occurs in association with the boson Z , Z' , H_1^0 and H_2^0 , see Fig. 1 and Fig. 2. Contrarily to the standard model, where the gluon-gluon fusion dominates over Drell-Yan when the Higgs boson is heavier than 100 GeV [10], in 3-3-1 Model does not occur, the mechanism of Drell-Yan dominates over gluon-gluon fusion at leading order (LO) for H_1^0 and H_2^0 production at $\sqrt{s} = 8(14)$ TeV. The process $pp \rightarrow H_i^0 Z$ ($i = 1, 2$) takes place in the s channel. So using the interaction Lagrangian [5, 12], we obtain the differential cross section in the first place for Drell-Yan for H_1^0 and H_2^0

$$\frac{d\hat{\sigma}_{H_1^0}}{d\Omega} = \frac{1}{64\pi^2\hat{s}} \left(\overline{|A_Z|^2} + \overline{|A_{Z'}|^2} + \overline{|A_{H_1^0}|^2} + \overline{|A_{H_2^0}|^2} + 2\text{Re}\overline{A_Z^* A_{Z'}} + 2\text{Re}\overline{A_{H_1^0}^* A_{H_2^0}} \right), \quad (17)$$

$$\frac{d\hat{\sigma}_{H_2^0}}{d\Omega} = \frac{1}{64\pi^2\hat{s}} \left(\overline{|A_{Z'}|^2} + \overline{|A_{H_2^0}|^2} + \overline{|A_{H_1^0}|^2} + 2\text{Re}\overline{A_{H_1^0}^* A_{H_2^0}} \right). \quad (18)$$

where the A_i is the matrix element of each particle.

In the cross section for H_i^0 production, where $i = 1, 2$, the interference term between the $Z(Z')$ and H_i^0 should be absent because it gives an imaginary value, and in the cross section for H_2^0 production the term involving the boson Z is absent, because there is no coupling

between the Z and H_2^0 Z , then we write separately the differential cross section for H_1^0 and H_2^0 production

$$\begin{aligned}
\frac{d\hat{\sigma}_{H_1^0}}{d\cos\theta} = & \frac{\beta_{H_1^0} g^2}{192\pi \cos^2\theta_W s} \left\{ \frac{m_W^2 g^2}{4\cos^4\theta_W (s - m_Z^2 + im_Z\Gamma_Z)^2} \left((m_Z^2 + \frac{tu}{m_Z^2} - t - u + s)(g_{V^q}^2 + g_{A^q}^2) \right) \right. \\
& + \frac{g^4 \Lambda_{ZZ'H_1^0}^2}{4(s - m_{Z'}^2 + im_{Z'}\Gamma_{Z'})^2} \left((m_Z^2 + \frac{tu}{m_Z^2} - t - u + s)(g_{V^q}^2 + g_{A^q}^2) \right) \\
& + \frac{g^3 m_W \Lambda_{ZZ'H_1^0}}{2 \cos^2\theta_W (s - m_Z^2 + im_Z\Gamma_Z)(s - m_{Z'}^2 + im_{Z'}\Gamma_{Z'})} \left((m_Z^2 + \frac{tu}{m_Z^2} - t - u + s) \right. \\
& (g_V^q g_{V'}^q + g_A^q g_{A'}^q) \left. \right) + \left(\frac{m_q^2 (v_\rho^2 - v_\eta^2)^2}{32 v_W^6} |\chi^{(1)}(\hat{s})|^2 + \frac{v_\eta^2 v_\rho^2}{2 v_W^6} \left(m_u \frac{v_\eta}{v_\rho} - m_d \frac{v_\rho}{v_\eta} \right)^2 |\chi^{(2)}(\hat{s})|^2 \right. \\
& \left. + \frac{m_q \left(m_u \frac{v_\eta}{v_\rho} - m_d \frac{v_\rho}{v_\eta} \right) v_\eta v_\rho (v_\rho^2 - v_\eta^2)}{4 v_W^6} |\chi^{(1)}(\hat{s})| |\chi^{(2)}(\hat{s})| \right) \\
& \left(\frac{\hat{s}}{m_Z^2} (\hat{s}^2 - 2m_Z^2 \hat{s} + m_Z^4) - \frac{m_q^2}{m_Z^2} (2\hat{s}^2 - 4\hat{s} m_Z^2 + 2m_Z^4) + \right. \\
& \left. - \frac{m_{H_1^0}^2}{m_Z^2} (2\hat{s}^2 + 2\hat{s}m_q^2 - 4m_q^2\hat{s} - 4m_q^2m_Z^2) + \frac{m_{H_1^0}^4 \hat{s}}{m_Z^2} - \frac{2m_{H_1^0}^4 m_q^2}{m_Z^2} \right) \left. \right\}, \tag{19}
\end{aligned}$$

$$\begin{aligned}
\frac{d\hat{\sigma}_{H_2^0}}{d\cos\theta} = & \frac{\beta_{H_2^0} g^2}{192\pi \cos^2\theta_W s} \left\{ \frac{g^4 \Lambda_{ZZ'H_2^0}^2}{4(s - m_{Z'}^2 + im_{Z'}\Gamma_{Z'})^2} \left((m_Z^2 + \frac{tu}{m_Z^2} - t - u + s)(g_{V^q}^2 + g_{A^q}^2) \right) \right. \\
& + \left(\frac{v_\eta^2 v_\rho^2 m_q^2}{2 v_W^6} |\chi^{(1)}(\hat{s})|^2 + \frac{\left(m_u \frac{v_\eta}{v_\rho} - m_d \frac{v_\rho}{v_\eta} \right)^2 (v_\rho^2 - v_\eta^2)^2}{32 v_W^6} |\chi^{(2)}(\hat{s})|^2 \right. \\
& \left. + \frac{m_q \left(m_u \frac{v_\eta}{v_\rho} - m_d \frac{v_\rho}{v_\eta} \right) v_\eta v_\rho (v_\rho^2 - v_\eta^2)}{4 v_W^6} |\chi^{(1)}(\hat{s})| |\chi^{(2)}(\hat{s})| \right) \\
& \left(\frac{\hat{s}}{m_Z^2} (\hat{s}^2 - 2m_Z^2 \hat{s} + m_Z^4) - \frac{m_q^2}{m_Z^2} (2\hat{s}^2 - 4\hat{s} m_Z^2 + 2m_Z^4) + \right. \\
& \left. - \frac{m_{H_2^0}^2}{m_Z^2} (2\hat{s}^2 + 2\hat{s}m_q^2 - 4m_q^2\hat{s} - 4m_q^2m_Z^2) + \frac{m_{H_2^0}^4 \hat{s}}{m_Z^2} - \frac{2m_{H_2^0}^4 m_q^2}{m_Z^2} \right) \left. \right\}, \tag{20}
\end{aligned}$$

here g is the coupling constant of the weak interaction, the $\beta_{H_i^0}$ ($i = 1, 2$) is the Higgs velocity in the c.m. of the subprocess which is equal to

$$\beta_{H_i^0} = \frac{\left[\left(1 - \frac{(m_Z + m_{H_i^0})^2}{\hat{s}} \right) \left(1 - \frac{(m_Z - m_{H_i^0})^2}{\hat{s}} \right) \right]^{1/2}}{1 - \frac{m_Z^2 - m_{H_i^0}^2}{\hat{s}}},$$

and we have also defined

$$\chi^i(\hat{s}) = \frac{1}{\hat{s} - m_{H_i^0}^2 + im_{H_i^0}\Gamma_{H_i^0}},$$

with $\Gamma_{H_i^0}$ being the Higgs boson total width, the primes ($'$) are for the case when we take a Z' boson, Γ_Z and $\Gamma_{Z'}$ [11, 12], are the total width of the Z and Z' boson, $g_{V,A}^q$ are the standard quark coupling constants, m_q , where $q = u, d$ are the masses of the quark, $g_{V',A'}^q$ are the 3-3-1 quark coupling constants, $\sqrt{\hat{s}}$ is the center of mass energy of the $q\bar{q}$ system, $g = \sqrt{4\pi\alpha}/\sin\theta_W$ and α is the fine structure constant, which we take equal to $\alpha = 1/128$. For the Z' boson we take $M_{Z'} = (0.5 - 3)$ TeV, since $M_{Z'}$ is proportional to the VEV v_χ [6, 7]. For the standard model parameters we assume PDG values, *i. e.*, $M_Z = 91.19$ GeV, $\sin^2\theta_W = 0.2315$, and $M_W = 80.33$ GeV [13], t and u are the kinematic invariants. We have also defined the $\Lambda_{ZZH_i^0}$ ($\Lambda_{ZZ'H_i^0}$) as the coupling constants of the Z (Z') boson to Z boson and Higgs H_i^0 where i stands for H_1^0, H_2^0 , the $\Lambda_{H_i^0 H_i^0 Z}$ are the couplings constants of the H_1^0 boson to H_1^0 and Z boson, of the H_2^0 boson to H_2^0 and Z boson and of the H_1^0 boson to H_2^0 and Z boson, these three coupling constants should be multiplied by $p^\mu - q^\mu$ to get a $\Lambda_{ZZH_i^0}^\mu = \Lambda_{ZZH_i^0}(p^\mu - q^\mu)$ with p and q being the momentum four-vectors of the H_i and Z boson where $i = 1, 2$, and the $\Lambda_{q\bar{q}H_i^0}$ are the coupling constants of the H_1^0 (H_2^0) to $q\bar{q}$, the $v(q)$ $a(q)$ $v'(q)$ and $a'(q)$ are given in [11]. It can be noticed then that the coupling $H_1^0 Z_\mu Z_\nu$ is the same as the SM, as one would expect, while $H_2^0 Z_\mu Z_\nu$ does not exist. We remark still that in 3-3-1 model, the states H_1^0 and H_2^0 are mixed.

$$t = m_{q_i}^2 + m_Z^2 - \frac{s}{2} \left\{ \left(1 + \frac{m_Z^2 - m_H^2}{s} \right) - \cos\theta \left[\left(1 - \frac{(m_Z + m_H)^2}{s} \right) \left(1 - \frac{(m_Z - m_H)^2}{s} \right) \right]^{1/2} \right\},$$

$$u = m_{q_i}^2 + m_H^2 - \frac{s}{2} \left\{ \left(1 - \frac{m_Z^2 - m_H^2}{s} \right) + \cos\theta \left[\left(1 - \frac{(m_Z + m_H)^2}{s} \right) \left(1 - \frac{(m_Z - m_H)^2}{s} \right) \right]^{1/2} \right\},$$

$$(\Lambda_{q\bar{q}Z})_\mu \approx i \frac{|e|}{2s_W c_W} \gamma_\mu [v(q) + a(q) \gamma_5], \quad (21a)$$

$$(\Lambda_{q\bar{q}Z'})_\mu \approx i \frac{|e|}{2s_W c_W} \gamma_\mu [v'(q) + a'(q) \gamma_5], \quad (21b)$$

$$\Lambda_{q\bar{q}H_1^0} \approx -i \frac{m_q}{2v_W} (1 + \gamma_5), \quad (21c)$$

$$\Lambda_{q\bar{q}H_2^0} \approx \frac{i}{2v_W} \left(-m_u \frac{v_\eta}{2v_\rho} + m_d \frac{v_\rho}{v_\eta} \right) (1 + \gamma_5) \quad (21d)$$

$$\left(\Lambda_{ZZH_1^0} \right)_{\mu\nu} \approx -i \frac{g^2 v_W}{2} \left(\frac{m_Z}{m_W} \right)^2 g_{\mu\nu}, \quad (21e)$$

$$\left(\Lambda_{ZZ'H_1^0} \right)_{\mu\nu} \approx -\frac{g^2}{2\sqrt{3}v_W} \frac{m_Z (1 + 2s_W^2) v_\eta^2 - (1 - 4s_W^2) v_\rho^2}{m_W c_W \sqrt{1 - 4s_W^2}} g_{\mu\nu}, \quad (21f)$$

$$\left(\Lambda_{ZZ'H_2^0} \right)_{\mu\nu} \approx \frac{g^2}{\sqrt{3}(1 - 4s_W^2)} \frac{v_\eta v_\rho}{v_W} g_{\mu\nu}, \quad (21g)$$

$$\left(\Lambda_{H_1^0 H_1^0 Z} \right)_\mu \approx \frac{g}{2} \frac{m_Z}{m_W} \frac{(v_\rho^2 - v_\eta^2)}{v_W^2} (p - q)_\mu, \quad (21h)$$

$$\left(\Lambda_{H_2^0 H_2^0 Z} \right)_\mu \approx -\frac{g}{2} \frac{m_Z}{m_W} \frac{(v_\rho^2 - v_\eta^2)}{v_W^2} (p - q)_\mu, \quad (21i)$$

$$\left(\Lambda_{H_1^0 H_2^0 Z} \right)_\mu \approx -2g \frac{m_Z}{m_W} \frac{v_\rho v_\eta}{v_W^2} (p - q)_\mu, \quad (21j)$$

where θ is the angle between the Higgs and the incident quark in the CM frame.

The total cross section for the process $pp \rightarrow qq \rightarrow ZH_i$ is related to the subprocess $qq \rightarrow ZH_i$ total cross section $\hat{\sigma}$, through

$$\sigma = \int_{\tau_{min}}^1 \int_{\ln \sqrt{\tau_{min}}}^{-\ln \sqrt{\tau_{min}}} d\tau dy q(\sqrt{\tau}e^y, Q^2) q(\sqrt{\tau}e^{-y}, Q^2) \hat{\sigma}(\tau, s), \quad (22)$$

where $\tau_{min} = (m_Z + m_{H_i})^2/s$ ($\tau = \hat{s}/s$) and $q(x, Q^2)$ is the quark structure function.

Another form to produce a neutral Higgs is *via* the gluon-gluon fusion, namely through the reaction of the type $pp \rightarrow gg \rightarrow ZH_i$. Since the final state is neutral, the s channel involves the exchange of the boson Z , Z' , H_1^0 and H_2^0 . The exchange of a photon is not allowed by C conservation (Furry's theorem), which also indicates that only the axial-vector couplings of the bosons Z and Z' , contribute to this process. Therefore, the differential cross section for production of H_1^0 and H_2^0 we make separately in order to do explicit the different contributions:

$$\begin{aligned} \frac{d\hat{\sigma}_{H_1^0}}{d\Omega} = & \frac{1}{64\pi^2\hat{s}} \left(\overline{|A_Z|^2} + \overline{|A_{Z'}|^2} + \overline{|A_{H_1^0}|^2} + \overline{|A_{H_2^0}|^2} + 2\text{Re}\overline{A_Z^* A_{Z'}} + 2\text{Re}\overline{A_{Z(Z')} A_{H_1^0}^*} \right. \\ & \left. + 2\text{Re}\overline{A_{Z(Z')} A_{H_2^0}^*} + 2\text{Re}\overline{A_{H_1^0}^* A_{H_2^0}} \right), \end{aligned} \quad (23)$$

$$\begin{aligned} \frac{d\hat{\sigma}_{H_2^0}}{d\Omega} = & \frac{1}{64\pi^2\hat{s}} \left(\overline{|A_{Z'}|^2} + \overline{|A_{H_1^0}|^2} + \overline{|A_{H_2^0}|^2} + 2\text{Re}\overline{A_{Z'} A_{H_1^0}^*} + 2\text{Re}\overline{A_{Z'} A_{H_2^0}^*} \right. \\ & \left. + 2\text{Re}\overline{A_{H_1^0}^* A_{H_2^0}} \right). \end{aligned} \quad (24)$$

where the A_i is the matrix element of each particle.

It is important to emphasize that for production of H_1^0 the interference term between the $Z(Z')$, which are antisymmetric in the gluon polarizations, and the H_1^0 diagrams, we only consider the antisymmetric term of H_1^0 , because the other part is symmetric and therefore vanishes, and in the production of H_2^0 we take the interference between the Z' diagrams and the antisymmetric term of H_2^0 , then we write explicitly the Z, Z', H_1^0 and H_2^0 contributions to the elementary cross section

$$\left(\frac{d\hat{\sigma}}{d\cos\theta} \right)_{pp \rightarrow ZH_i^0}^{Z(Z')} = \frac{g^4(g^6) \alpha_s^2 (\Lambda_{Z(Z')ZH_i^0})^2 \Delta}{8192 \pi^3 \hat{s} \cos_W^2 M_{Z(Z')}^4} \beta_{H_i^0} \left| \sum_{q=u,d} T_3^q(q') (1 + 2\delta_q I_q) \right|^2, \quad (25)$$

$$\begin{aligned} \left(\frac{d\hat{\sigma}}{d\cos\theta} \right)_{pp \rightarrow ZH_i^0}^{H_1^0} = & \frac{g^2 \alpha_s^2 (v_\rho^2 - v_\eta^2)^2 \hat{s} \Omega \beta_{H_i^0}}{131072 \pi^3 v_W^6 \cos_{\theta_W}^2} |\chi^{(1)}(\hat{s})|^2 \left| \sum_{q=u,d} [2\delta_q + \delta_q(4\delta_q - 1)I_q] \right|^2 + \\ & + \frac{g^2 \alpha_s^2 (v_\rho^2 - v_\eta^2)^2 \Omega \beta_{H_i^0}}{65536 \pi^3 \hat{s} v_W^6 \cos_W^2} |\chi^{(1)}(\hat{s})|^2 \left| \sum_{q=u,d} m_q^2 I_q \right|^2, \end{aligned} \quad (26)$$

$$\begin{aligned} \left(\frac{d\hat{\sigma}}{d\cos\theta} \right)_{pp \rightarrow ZH_i^0}^{H_2^0} = & \frac{g^2 \alpha_s^2 v_\eta^2 v_\rho^2 \hat{s} \Omega \beta_{H_i^0}}{8192\pi^3 v_W^6 \cos_{\theta_W}^2} |\chi^{(2)}(\hat{s})|^2 \left| \sum_{q=u,d} \frac{(m_u \frac{v_\eta}{v_\rho} - m_d \frac{v_\rho}{v_\eta})}{m_q} [2\delta_q + \delta_q(4\delta_q - 1)I_q] \right|^2 + \\ & + \frac{g^2 \alpha_s^2 v_\eta^2 v_\rho^2 \Omega \beta_{H_i^0}}{4096 \pi^3 \hat{s} v_W^6 \cos_W^2} |\chi^{(2)}(\hat{s})|^2 \left| \sum_{q=u,d} m_q \left(m_u \frac{v_\eta}{v_\rho} - m_d \frac{v_\rho}{v_\eta} \right) I_q \right|^2, \end{aligned} \quad (27)$$

$$\begin{aligned} \left(\frac{d\hat{\sigma}}{d\cos\theta} \right)_{pp \rightarrow ZH_1^0}^{Z-Z'} = & \frac{g^5 \alpha_s^2 \Lambda_{ZZ'H_1^0} \Delta}{4096 \pi^3 \hat{s} \cos_{\theta_W}^2 m_Z^2 m_{Z'}^2} \beta_{H_1^0} \\ & \text{Re} \left[\sum_{q=u,d} T_3^q (1 + 2\delta_q I_q^*) \sum_{q=u,d} T_3^{q'} (1 + 2\delta_{q'} I_{q'}) \right], \end{aligned} \quad (28)$$

$$\left(\frac{d\hat{\sigma}}{d\cos\theta}\right)_{pp\rightarrow ZH_1^0}^{Z(Z')-H_1^0} = -\frac{g^3(g^4)\alpha_s^2\Lambda_{Z(Z')ZH_1^0}(v_\rho^2-v_\eta^2)\Pi\beta_{H_i^0}}{4096\pi^3\hat{s}\cos_{\theta_W}^2v_W^3}Re\left[\frac{\chi^{(1)}(\hat{s})}{(s-m_{Z(Z')}^2+im_{Z(Z')}\Gamma_{Z(Z')})}\right. \\ \left.\sum_{q=u,d}m_q^2T_3^q(1+2\delta_qI_q)\sum_{q=u,d}I_q^*\right], \quad (29)$$

$$\left(\frac{d\hat{\sigma}}{d\cos\theta}\right)_{pp\rightarrow ZH_1^0(H_2^0)}^{H_1^0-H_2^0} = \mp\frac{g^2\alpha_s^2(v_\rho^2-v_\eta^2)v_\rho v_\eta\Omega\beta_{H_i^0}}{8192\pi^3\hat{s}\cos_{\theta_W}^2v_W^6}Re\chi^{(1)}(\hat{s})\chi^{(2)}(\hat{s}) \\ \sum_{q=u,d}m_q^3\left(m_u\frac{v_\eta}{v_\rho}-m_d\frac{v_\rho}{v_\eta}\right)I_q\sum_{q=u,d}I_q^*\mp\frac{g^2\alpha_s^2(v_\rho^2-v_\eta^2)v_\rho v_\eta s\Omega\beta_{H_i^0}}{16384\pi^3\cos_{\theta_W}^2v_W^6} \\ Re\chi^{(1)}(\hat{s})\chi^{(2)}(\hat{s})\sum_{q=u,d}\frac{\left(m_u\frac{v_\eta}{v_\rho}-m_d\frac{v_\rho}{v_\eta}\right)}{m_q}I_q\sum_{q=u,d}I_q^*, \quad (30)$$

$$\left(\frac{d\hat{\sigma}}{d\cos\theta}\right)_{pp\rightarrow ZH_1^0}^{Z(Z')-H_2^0} = -\frac{g^3(g^4)\alpha_s^2\Lambda_{Z(Z')ZH_2^0}v_\eta v_\rho\Pi\beta_{H_i^0}}{1024\pi^3\hat{s}^2\cos_{\theta_W}^2v_W^3}Re\left[\frac{\chi^{(2)}(\hat{s})}{(s-m_{Z'}^2+im_{Z'}\Gamma_{Z'})}\right. \\ \left.\sum_{q=u,d}m_q\left(m_u\frac{v_\eta}{v_\rho}-m_d\frac{v_\rho}{v_\eta}\right)T_3^q(1+2\delta_qI_q)\sum_{q=u,d}I_q^*\right], \quad (31)$$

$$\left(\frac{d\hat{\sigma}}{d\cos\theta}\right)_{pp\rightarrow ZH_2^0}^{H_1^0} = \frac{g^2\alpha_s^2v_\eta^2v_\rho^2\hat{s}\Omega\beta_{H_i^0}}{8192\pi^3v_W^6\cos_{\theta_W}^2}|\chi^{(1)}(\hat{s})|^2\left|\sum_{q=u,d}2\delta_q+\delta_q(4\delta_q-1)I_q\right|^2+ \\ +\frac{g^2\alpha_s^2v_\eta^2v_\rho^2\Omega\beta_{H_i^0}}{4096\pi^3\hat{s}\cos_W^2v_W^6}|\chi^{(1)}(\hat{s})|^2\left|\sum_{q=u,d}m_q^2I_q\right|^2, \quad (32)$$

$$\left(\frac{d\hat{\sigma}}{d\cos\theta}\right)_{pp\rightarrow ZH_2^0}^{H_2^0} = \frac{g^2\alpha_s^2(v_\rho^2-v_\eta^2)^2\hat{s}\Omega\beta_{H_i^0}}{131072\pi^3v_W^6\cos_{\theta_W}^2}|\chi^{(2)}(\hat{s})|^2\left|\sum_{q=u,d}\frac{\left(m_u\frac{v_\eta}{v_\rho}-m_d\frac{v_\rho}{v_\eta}\right)}{m_q}[2\delta_q+\delta_q(4\delta_q-1)I_q]\right|^2 \\ +\frac{g^2\alpha_s^2(v_\rho^2-v_\eta^2)^2\Omega\beta_{H_i^0}}{65536\pi^3\hat{s}v_W^6\cos_W^2}|\chi^{(2)}(\hat{s})|^2\left|\sum_{q=u,d}m_q\left(m_u\frac{v_\eta}{v_\rho}-m_d\frac{v_\rho}{v_\eta}\right)I_q\right|^2, \quad (33)$$

$$\left(\frac{d\hat{\sigma}}{d\cos\theta}\right)_{pp\rightarrow ZH_2^0}^{Z'-H_1^0} = -\frac{g^4\alpha_s^2\Lambda_{ZZ'H_2^0}v_\eta v_\rho\Pi\beta_{H_i^0}}{1024\pi^3\hat{s}^2\cos_{\theta_W}^2v_W^3}Re\left[\frac{\chi^{(1)}(\hat{s})}{(s-m_{Z'}^2+im_{Z'}\Gamma_{Z'})}\right. \\ \left.\sum_{q=u,d}m_q^2T_3^q(1+2\delta_qI_q)\sum_{q=u,d}I_q^*\right], \quad (34)$$

$$\left(\frac{d\hat{\sigma}}{d\cos\theta}\right)_{pp\rightarrow ZH_2^0}^{Z'-H_2^0} = -\frac{g^4 \alpha_s^2 \Lambda_{ZZ'H_2^0} (v_\rho^2 - v_\eta^2) \Pi \beta_{H_i^0}}{4096 \pi^3 \hat{s}^2 \cos_{\theta_W}^2 v_W^3} \text{Re} \left[\frac{\chi^{(1)}(\hat{s})}{(s - m_{Z'}^2 + im_{Z'}\Gamma_{Z'})} \right. \\ \left. \sum_{q=u,d} m_q \left(m_u \frac{v_\eta}{v_\rho} - m_d \frac{v_\rho}{v_\eta} \right) T_3^q (1 + 2\delta_q I_q) \sum_{q=u,d} I_q^* \right], \quad (35)$$

where in Eq. (25) are considered the contribution of Z and Z' bosons, in Eq. (26) the contribution of H_1^0 , in Eq. (27) the contribution of H_2^0 , in Eq. (28) the contribution of interference of ZZ' , in Eq. (29) the contribution of interference of ZH_1^0 and $Z'H_1^0$, in Eq. (30) the contribution of interference of $H_1^0 H_2^0$ and in Eq. (31) the contribution of interference of ZH_2^0 and $Z'H_2^0$, all these contributions are to produce the ZH_1^0 . All other equations (30, 32, 33, 34, 35) are to produce the ZH_2^0 , we point out that the term $\left(\frac{d\hat{\sigma}}{d\cos\theta}\right)_{pp\rightarrow ZH_2^0}^{H_1^0-H_2^0}$ is similar to $\left(\frac{d\hat{\sigma}}{d\cos\theta}\right)_{pp\rightarrow ZH_1^0}^{H_1^0-H_2^0}$, because the coupling constants $\Lambda_{H_2^0 H_2^0 Z} = -\Lambda_{H_1^0 H_1^0 Z}$ are equal. The sum runs over all generations, T_3^q is the quark weak isospin [$T_3^{u(d)} = +(-)1/2$], Re stands for the real part of the expression. The loop function $I_i \equiv I(\delta_i = m_i^2/\hat{s})$, is defined by

$$I_i \equiv I_i(\delta_i) = \int_0^1 \frac{dx}{x} \ln \left[1 - \frac{(1-x)x}{\delta_i} \right] = \begin{cases} -2 \left[\sin^{-1} \left(\frac{1}{2\sqrt{\delta_i}} \right) \right]^2, & \delta_i > \frac{1}{4} \\ \frac{1}{2} \ln^2 \left(\frac{r_+}{r_-} \right) - \frac{\pi^2}{2} + i\pi \ln \left(\frac{r_+}{r_-} \right), & \delta_i < \frac{1}{4}, \end{cases}$$

with, $r_\pm = 1 \pm (1 - 4\delta_i)^{1/2}$ and $\delta_i = m_i^2/\hat{s}$. Here, $i = q$ stands for the particle (quark) running in the loop.

We have also defined Δ , Ω and Π which are equal to:

$$\Delta = 4\hat{s} - \frac{\hat{u}^2}{2m_Z^2} + \frac{\hat{t} \hat{u}}{m_Z^2} - \frac{\hat{t}^2}{2m_{Z'}^2}$$

$$\Omega = \frac{\hat{s}^2}{4m_Z^2} - \frac{\hat{s}}{2} + \frac{m_Z^2}{4} - \frac{\hat{s} m_{H_1^0}^2}{2m_Z^2} - \frac{m_{H_1^0}^2}{2} + \frac{m_{H_1^0}^4}{4m_Z^2}$$

$$\Pi = -\frac{\hat{s}^4}{8 m_Z^4} + \frac{\hat{s}^3}{4 m_Z^2} - \frac{\hat{s}^2 \hat{u}}{8 m_Z^2} - \frac{\hat{s}^2 \hat{t}}{8 m_Z^2} + \frac{\hat{s}^2}{8} + \frac{3 \hat{s} \hat{u}}{8} + \frac{3 \hat{s} \hat{t}}{8} - \frac{\hat{s} m_Z^2}{4} + \frac{\hat{s}^3 m_{H_i^0}^2}{4 m_Z^4} \\ + \frac{\hat{s}^2 m_{H_i^0}^2}{4 m_Z^2} + \frac{\hat{s} \hat{u} m_{H_i^0}^2}{8 m_Z^2} + \frac{\hat{s} \hat{t} m_{H_i^0}^2}{8 m_Z^2} - \frac{3 \hat{s} m_{H_i^0}^2}{4} - \frac{\hat{s}^2 m_{H_i^0}^4}{4 m_Z^4},$$

The total cross section for the process $pp \rightarrow gg \rightarrow ZH_i$ is related to the subprocess $gg \rightarrow ZH_i$ total cross section $\hat{\sigma}$ through

$$\sigma = \int_{\tau_{min}}^1 \int_{\ln \sqrt{\tau_{min}}}^{-\ln \sqrt{\tau_{min}}} d\tau dy G(\sqrt{\tau}e^y, Q^2) G(\sqrt{\tau}e^{-y}, Q^2) \hat{\sigma}(\tau, s), \quad (36)$$

where $G(x, Q^2)$ is the gluon structure function and τ_{min} is given above.

IV. RESULTS AND CONCLUSIONS

In the SM at the Tevatron, Higgs-boson production in association with W or Z bosons, $pp \rightarrow W H_1^0/Z H_1^0 + X$ is the most promising discovery channel for a SM Higgs H_1^0 masses below about 130 GeV, where $H_1^0 \rightarrow b\bar{b}$ decays are dominant [14], while Higgsstrahlung is only marginal at the LHC.

We consider Higgs masses spanning in the range $115 \text{ GeV} \leq m_{H_1^0} \leq 800 \text{ GeV}$. Our talk will be subdivided into two distinct classes, if $m_{H_1^0}$ is less than 145 GeV or greater than 466 GeV, then ATLAS excludes Higgs boson masses above 145 GeV till 466 GeV, [15, 16]. The best chance of discovering a SM Higgs at the LHC appears to be given by the following processes:

- $gg \rightarrow H_1^0 \rightarrow \gamma\gamma$, where the highest rates for this channel occur in the region 80-150 GeV and the combination of a rising branching ratio and a falling cross section [17], yields rates which are constant over the above range.
- $q\bar{q}' \rightarrow WH_1^0 \rightarrow \ell\nu_\ell\gamma\gamma$, these channels are different with respect to the case of the direct $gg \rightarrow H_1^0 \rightarrow \gamma\gamma$ production and decay. On the one hand owing to a larger number of reducible ($W\gamma j$ with the pions in the jet giving hard photons and $Z\gamma$ with a ℓ^\pm from the Z decay faking a photon) and irreducible ($W\gamma\gamma$) backgrounds. On the other hand owing to the isolated hard lepton from the W decays that allows for a strong reduction of the backgrounds [18]. Must be here noticed that unlike the signals from leptonic decay modes, those from hadronic decay channels undergo mostly QCD background processes.
- $H \rightarrow b\bar{b}$; $q\bar{q}' \rightarrow WH \rightarrow \ell\bar{\nu}_\ell b\bar{b}$; $gg, q\bar{q}' \rightarrow t\bar{t}H \rightarrow q\bar{q}q\bar{q}WW \rightarrow q\bar{q}q\bar{q}\ell\bar{\nu}_\ell X$, where the first decay mode is the dominant in the range $80 \text{ GeV} \leq m_{H_1^0} \leq 130 \text{ GeV}$. All these decay modes use the techniques of flavor identification of b-jets, thereby reducing the huge QCD background from light quarks and gluon jets. The chances to tag the Higgs boson in association with a W or a $t\bar{t}$ pair are $W \rightarrow \ell\bar{\nu}_\ell$ and $t\bar{t} \rightarrow \ell\bar{\nu}_\ell X$. The lepton is usually at

high p_T and isolated. The expected signatures would then be $\ell b\bar{b}X$ from WH_1^0 and $b\bar{b}b\bar{b}\ell X$ from $t\bar{t}H_1^0$. Higgs signal in the $b\bar{b}$ channel would appear as a peak in the invariant mass distribution of b-quark pairs.

The backgrounds to the channel $WH_1^0 \rightarrow \ell b\bar{b}X$ that we have considered are: $Wb\bar{b}$, $Zb\bar{b}$, WZ , ZZ , $q\bar{q}' \rightarrow t\bar{b}, t\bar{t}$, $qg \rightarrow t\bar{b}q'$, Wjb and Wjj . The dominant irreducible backgrounds are $Wb\bar{b}$, WZ with $Z \rightarrow b\bar{b}$. The dominant reducible background comes from Wjj production, through Wgg , Wgq and $Wq\bar{q}$ in which the two jets are faked as b-quarks and Wjb for one jet only. The irreducible background in $t\bar{t}Z$ events is small with respect to the signal $t\bar{t}H_1^0$.

- $H \rightarrow Z Z \rightarrow 4 \ell$ this channel with $\ell = e$ or μ nicknamed as the ‘gold-plated’ has recived much attention in the literature because of the opportunity for a fully reconstructed Higgs signal. This channel provides the best chances for Higgs detection over a large interval of the Higgs mass range, between $\simeq 120$ and 800 GeV. Its signature is relatively clean, especially if compared to the difficulties encountered for the $H_1^0 \rightarrow \gamma\gamma$ and $H_1^0 \rightarrow b\bar{b}$ cases, however, this process occurs with a small branching ratio, $B = \frac{1}{3} \times (0.034)^2 = 3.9 \times 10^{-4}$, that the predicted events rates are small. In the mass region $m_{H_1^0} \leq 2m_Z$ the main backgrounds to the four lepton signal come from $t\bar{t} \rightarrow b\bar{b}W^+W^-$, $Zb\bar{b}$ and ZZ production, where the last is irreducible. In the first two backgrounds two of the leptons come from semi-leptonic b-decays, while the remaining two from the decays of the massive vector bosons, that is $W^+W^- \rightarrow \ell^+\ell^-X$ and $Z \rightarrow \ell^+\ell^-$. In the case of the high mass region, when the Higgs decay $H_1^0 \rightarrow ZZ$ can occur, the only significant background, after applying the invariant mass cut $m_{\ell^+\ell^-} \simeq m_Z$ on both the lepton-antilepton pair, is the continuum ZZ production [17]. The Higgs signal would appear as a resonance, which has a maximum at $m_{4\ell} = 2m_Z$.

- $H_1^0 \rightarrow ZZ \rightarrow \ell\bar{\ell}\nu\bar{\nu}$, this channel should also be considered as it is enhanced relative to the gold-plated decay mode by a factor of six [19], owing to its BR which is six times larger. This channel can give good chances for Higgs detection for high values of $m_{H_1^0} \geq 700$ GeV. While this process may at first seem an unlikely channel since the second Z is not detected, the signature reveals a Z with two high p_T leptons and high missing transverse energy from the other Z. The main backgrounds to this channel are the continuum production of ZZ and WZ. Other possible reducible backgrounds are $q\bar{q} \rightarrow ZW \rightarrow \ell^+\ell^-\ell'\nu$ where the charged leptons $\ell' = e, \mu, \tau$ escapes detection. Another reducible backgrounds arises from Z+ jets, where enough of the jets escape detection to leave large missing p_T and the production of $t\bar{t}$, but to suppress this background the additional cut $|m_Z - m_{\ell^+\ell^-}| \leq 6$ GeV must be applied.

In this work we have calculated contributions regarding to the Drell-Yan and gluon-gluon fusion in 3-3-1 model. We present the cross section for the process $pp \rightarrow ZH_i$ involving the Drell-Yan mechanism and the gluon-gluon fusion, to produce such Higgs bosons for the LHC. In all calculations was taken for the parameters, the VEV and masses the following representative values, [9, 20] : $\lambda_1 = -0.20$, $\lambda_2 = 1.02$, $\lambda_3 = -\lambda_6 = -1$, $\lambda_4 = 2.98$ $\lambda_5 = -1.57$, $\lambda_7 = -2$, $\lambda_8 = -0.42$, $v_\eta = 195$ GeV, $\lambda_9 = -0.9(-0.76)$, for $v_\chi = 1000(1500)$ GeV, these parameters are used to estimate the values for the particles masses which are given in Table I, it is to notice that the value of λ_9 was chosen this way in order to guarantee the approximation $-f \simeq v_\chi$ [9, 20].

TABLE I: Values for the particle masses used in this work. All the values in this Table are given in GeV.

f	v_χ	m_E	m_M	m_T	$m_{H^{\pm\pm}}$	$m_{H_3^0}$	m_{h^0}	$m_{H_1^0}$	$m_{H_2^0}$	$m_{H_2^\pm}$	m_V	m_U
-1008.3	1000	148.9	875	2000	500	2000	1454.6	126	1017.2	183	467.5	464
-1499.7	1500	223.3	1312.5	3000	500	3000	2164.3	126	1525.8	285.2	694.1	691.7

	$m_{Z'}$	m_{J_1}	m_{J_2}	m_{J_3}
	1707.6	1000	1410	1410
	2561.3	1500	2115	2115

A. The Higgs H_1^0

We may have very strong hints of the standard lightweight Higgs ($m_{H_1^0} \propto 125$ GeV), with the b-tagging results from CDF and DZero of the TEVATRON and with the two-photon and four-lepton results from ATLAS and CMS of the LHC, at the level of 5 sigma, in the mass region around 125-126 GeV [21, 22]. We interpret this to be due to the production of a previously unobserved particle with a mass of around 125 GeV. At the end of 2012, CMS and ATLAS updated all the Higgs decay channels for 13 fb^{-1} and 8 TeV, except that CMS choose not to publish the diphoton result because it was smaller than expected. The hope is that they will get full results at 20 fb^{-1} and 8 TeV and they will enable to elucidate further

the nature of this newly observed particle. In addition to the cross-sections we can hope for an update to the tests of spin parity on the Higgs boson. This will be the final step to declare that the Higgs-Very-Like-Boson is indeed the standard Higgs-Boson. Even so, until we can not measure a signal of a Higgs in a completely convincing way, we have a right to make the phenomenology analysis on the intermediate mass Higgs.

So in Fig. 3 and 4, we show the cross section $pp \rightarrow ZH_1^0$ at $\sqrt{s} = 8$ (14) TeV, these processes will be studied in two cases, the one is for the vacuum expectation value $v_\chi = 1000$ GeV and the other is for $v_\chi = 1500$. Considering that the expected integrated luminosity for the LHC collider that will be reach is of order of 20 fb^{-1} (300 fb^{-1}) then the statistics for $v_\chi = 1000$ gives a total of $\simeq 2.6 \times 10^4$ (2.0×10^4) (2.1×10^6) (1.7×10^6) events per year for Drell-Yan and $\simeq 360$ (328) (2.9×10^5) (2.6×10^5) events per year for gluon-gluon fusion, if we take the mass of the Higgs boson $m_{H_1^0} = 650$ (800) GeV respectively ($\Gamma_{H_1^0} = 214, 551.7$ GeV). Here we must clarify that the first two number of events $\simeq 2.6 \times 10^4$ (2.0×10^4) are relative to Drell-Yan and the other two $\simeq 360$ (328) to gluon-gluon fusion and correspond to 8 TeV and the other pair $\simeq 2.1 \times 10^6$ (1.7×10^6) are relative to Drell-Yan and $\simeq 2.9 \times 10^5$ (2.6×10^5) to gluon-gluon fusion and correspond to 14 TeV for the LHC, respectively.

Next, we multiply the production cross sections by the respective branching ratios to obtain event rates for various channels. Considering that the signal for ZH_1^0 production for $m_{H_1^0} = 650$ (800) GeV will be $Z \rightarrow \ell^+\ell^-$ and $H_1^0 \rightarrow ZZ \rightarrow \ell^+\ell^-\ell^+\ell^-$, then taking into account that the branching ratios for both particles would be $B(Z \rightarrow \ell^+\ell^-) = 3.4 \%$ and $B(H_1^0 \rightarrow ZZ) = 29.0$ (30.9) %, see Fig. 5, we would have approximately $\simeq 0$ (0) (24) (21) events per year. In respect to gluon-gluon fusion we will have $\simeq 0$ (0) (3) (4) events per year to produce the same particles. We will remember that the first two number of events $\simeq 0$ (0) refer to Drell-Yan and the other two $\simeq 0$ (0) to gluon-gluon fusion and correspond to 8 TeV and the other pair $\simeq 24$ (21) refer to Drell-Yan and $\simeq 3$ (4) to gluon-gluon fusion and correspond to 14 TeV for the LHC, respectively.

Regarding the vacuum expectation value $v_\chi = 1500$ GeV for the same masses of $m_{H_1^0} = 650$ (800) ($\Gamma_{H_1^0} = 212.2, 551.4$ GeV) it will give a total of $\simeq 4.0 \times 10^3$ (3.6×10^3) (7.1×10^5) (6.6×10^5) events per year to produce H_1^0 for Drell-Yan. In respect to gluon-gluon fusion we will have $\simeq 2$ (2) (9.0×10^3) (8.9×10^3) events per year to produce the same particles. Taking into account the same signal as above, that is $B(Z \rightarrow \ell^+\ell^-) = 3.4 \%$ and $B(H_1^0 \rightarrow ZZ) = 29.2$ (30.9) %, see Fig. 6, we would have approximately $\simeq 0$ (0) (8) (8) events per year for

Drell-Yan and $\simeq 0(0)(0(0))$ for gluon-gluon fusion. Comparing these signatures with the standard model background, like $pp \rightarrow ZZZ$ and using COMPHEP [23], we have that a cross section at LO is $3.8 \times 10^{-3}(9.6 \times 10^{-3})$ pb, where the first value is for $\sqrt{s} = 8$ and the second for $\sqrt{s} = 14$ TeV, respectively. Considering now the same signature as above we have $\simeq 0(0)$ events for the background. So we will have a total of $\simeq 27$ events per year for the signal for $m_{H_1^0} = 650$ GeV and $\simeq 25$ events per year for $m_{H_1^0} = 800$ GeV, for $v_\chi = 1000$ GeV and for $\sqrt{s} = 14$ TeV and $\simeq 8(8)$ events per year for $m_{H_1^0} = 650(800)$ for $v_\chi = 1500$ GeV and for $\sqrt{s} = 14$ TeV. To extract the leptonic signal from the background we impose the Z window cut where the invariant mass of opposite-charge leptons must be far from the Z mass: $|m_{\ell^+\ell^-} - m_Z| > 10$ GeV, this removes events where the leptons come from Z decay [24]. Then the Higgs signal would appear as a broad Breit-Wigner resonance, which has a maximum at $m_{4\ell} = 2m_Z$.

Taking another signal such as $H_1^0 Z \rightarrow W^+W^-Z$, and taking into account that the branching ratios for these particles would be $B(H_1^0 \rightarrow W^+W^-) = 59.1(62.3)$ %, see Fig. 5 and $B(Z \rightarrow \ell^+\ell^-) = 3.4$ %, for the mass of the Higgs boson $m_{H_1^0} = 650(800)$ GeV, $v_\chi = 1000$ GeV, and that the particles W^\pm decay into $e^\pm\nu$ and $e^\mp\bar{\nu}$, whose branching ratios for these particles would be $B(W^\pm \rightarrow e^\pm\nu) = 10, 75$ %, then we would have approximately $\simeq 6(4)(487(414))$ events per year for Drell-Yan. In respect to gluon-gluon fusion we will have $\simeq 0(0)(67(65))$ events per year to produce the same particles. Regarding the vacuum expectation value $v_\chi = 1500$ GeV and considering that the branching ratios for H_1^0 would be $B(H_1^0 \rightarrow W^+W^-) = 59.4(62.4)$ %, see Fig. 6 and taking the same parameters and branching ratios for the same particles given above, then we would have for $m_{H_1^0} = 650(800)$ a total of $\simeq 0(0)(166(161))$ events of H_1^0 produced per year for Drell-Yan and in respect to gluon-gluon fusion the number of events per year will be $\simeq 0(0)(2(2))$.

Taking the largest standard model background, like $pp \rightarrow W^+W^-Z$ and using CompHep [23] we have that this yields a cross section of $3.8 \times 10^{-2}(9.8 \times 10^{-2})$ pb at LO for $\sqrt{s} = 8(14)$ TeV respectively. Taking into account the $\ell^+\ell^-e^+e^-X$ backgrounds we have a total of $\simeq 0(12)$ events, so we will have a total of $\simeq 554$ events per year for the signal for $m_{H_1^0} = 650$ GeV and $\simeq 479$ events per year for $m_{H_1^0} = 800$ GeV, both for $v_\chi = 1000$ GeV and for $\sqrt{s} = 14$ TeV and $\simeq 168(163)$ events per year for $m_{H_1^0} = 650(800)$ for $v_\chi = 1500$ GeV and for $\sqrt{s} = 14$ TeV and $\simeq 12$ events for the background. To make clear these signals we must isolate the hard lepton from the W with $p_T^\ell > 20$ GeV and adopt the cut on the missing transverse

momentum $p_T' > 20$ GeV.

The $H_1^0 Z$ will also decay into $H^{\pm\pm} H^{\mp\mp} Z$, and taking into account that the branching ratios for these particles would be $B(H_1^0 \rightarrow H^{\pm\pm} H^{\mp\mp}) = 6.6(2.8)$ %, see Fig. 5 and $B(Z \rightarrow b\bar{b}) = 15.2$ % for the mass of the Higgs boson $m_{H_1^0} = 650(800)$ GeV and $v_\chi = 1000$ GeV, and that the particles $H^{\pm\pm}$ decay into $e^\pm E^\pm$, whose branching ratios for these particles would be $BR(H^{\pm\pm} \rightarrow e^\pm E^\pm) = 44.0(31.1)$ % [25], these branching ratios are in good agreement with the upper cinematic limit of Higgs mass [20], which in this case are $m_{H^{\pm\pm}} = 325(400)$ GeV, then we would have approximately $\simeq 50(8)(4.1 \times 10^3(700))$ events per year for Drell-Yan. Must be taken into account that E is the new heavy lepton and we assume that $(P = E, M, T)$ [9, 26]. In respect to gluon-gluon fusion we will have $\simeq 0(0)(5.6 \times 10^2(107))$ events per year to produce the same particles.

With respect to $v_\chi = 1500$ and doing all the analysis given above for the mass of Higgs boson $m_{H_1^0} = 650(800)$, considering that $B(H_1^0 \rightarrow H^{\pm\pm} H^{\mp\mp}) = 6.1(2.6)$ %, see Fig. 6 and $B(Z \rightarrow b\bar{b}) = 15.2$ % and $BR(H^{\pm\pm} \rightarrow e^\pm E^\pm) = 50.0(31.1)$ % [25], then it will take a total of $\simeq 10(2)(1.6 \times 10^3(252))$ events per year for Drell-Yan and $\simeq 0(0)(21(3))$ events for gluon-gluon fusion. In this way we have as signal four leptons and Z would appear in the invariant mass distribution of b-quark pairs, therefore, if we observe this signal, additionally to H_1^0 we observe also the doubly charged Higgs boson and heavy leptons, consequently it is a very striking and important signal. So we will have for $\sqrt{s} = 8$ TeV a total of $\simeq 50(8)$ events for signal $b\bar{b}e^+E^-e^-E^+$, for $m_{H_1^0} = 650(800)$ and for $v_\chi = 1000$ GeV and $\simeq 10(2)$ events for $v_\chi = 1500$ GeV and for the same parameters cited above. It is important to be noticed if the standard lightweight Higgs ($m_{H_1^0} \propto 125$ GeV) has not been discovered, this channel of decay cited above is the most important. For $\sqrt{s} = 14$ TeV we will have a total of $\simeq 4.6 \times 10^3(807)$ events per year for $m_{H_1^0} = 650(800)$, for $v_\chi = 1000$ GeV and $\simeq 1.6 \times 10^3(255)$ events per year for $m_{H_1^0} = 650(800)$ and for $v_\chi = 1500$ GeV to produce the same signal.

The main background to this signature is the ZZZ production which gives a cross section at LO, using COMPHEP [23], of $3.8 \times 10^{-3}(9.6 \times 10^{-3})$ pb, for $\sqrt{s} = 8(14)$ TeV respectively. We consider the irreducible background, that is $ZZZ \rightarrow b\bar{b}\ell^+\ell^-\ell^+\ell^-$, then we have $\simeq 0(0)$ events, therefore to observe these signals with the Z which decay into $b\bar{b}$ channel, first we must select the $b\bar{b}$ channel out of the huge QCD backgrounds of quark and gluon jets by using the b-tagging capabilities of vertex detectors, later we apply the Z window cut $|m_{\ell+\ell^-} - m_Z| > 10$ GeV, which removes events where the leptons come from Z decay [24]. Therefore the

Higgs signal would appear as a resonance, which has a maximum at $m_{4\ell} = 2m_{H_{\pm\mp}}$.

Taking another channel of decay such as $H_1^0 Z \rightarrow t\bar{t} \ell^+ \ell^-$, and considering that the branching ratios for these particles would be $B(H_1^0 \rightarrow t\bar{t}) = 5.1(3.9) \%$, see Fig. 5 and $B(Z \rightarrow \ell^+ \ell^-) = 3.4 \%$ for the mass of the Higgs boson $m_{H_1^0} = 650(800) \text{ GeV}$, $v_\chi = 1000 \text{ GeV}$, and that the $t\bar{t}$ particles decay into $b\bar{b}W^+W^-$, whose branching ratios for these particles would be $B(t \rightarrow bW) = 99.8 \%$ followed by leptonic decay of the boson W, that is $B(W \rightarrow e\nu) = 10.75 \%$, then our signal will be $b\bar{b}e^+e^-\ell^+\ell^-X$ and we would have approximately $\simeq 0(0)(42(26))$ events per year for Drell-Yan for the parameters listed above.

Regarding to gluon-gluon fusion we will have $\simeq 0(0)(6(4))$ events per year to produce the same particles. Considering the vacuum expectation value $v_\chi = 1500 \text{ GeV}$ and the branching ratios $B(H_1^0 \rightarrow t\bar{t}) = 5.2(3.9) \%$, see Fig. 6 and taking the same parameters and branching ratios for the same particles given above, then we would have for $m_{H_1^0} = 650(800)$ a total of $\simeq 0(0)(14(10))$ events of H_1^0 produced per year and in respect to gluon-gluon fusion the number of events per year will be $\simeq 0(0)(0(0))$. In this way we have a total of $\simeq 0(0)(48(30))$ events per year for $\sqrt{s} = 8(14) \text{ TeV}$, for $m_{H_1^0} = 650(800) \text{ GeV}$, for $v_\chi = 1000 \text{ GeV}$ and $\simeq 0(0)(14(10))$ events per year for $\sqrt{s} = 8(14) \text{ TeV}$, for $m_{H_1^0} = 650(800) \text{ GeV}$ and for $v_\chi = 1500 \text{ GeV}$. Taking the irreducible background $t\bar{t}Z \rightarrow b\bar{b}e^+e^-\ell^+\ell^-X$, and using CompHep we have that a cross section at LO is $\simeq 2.3 \times 10^{-1}(1) \text{ pb}$ for $\sqrt{s} = 8(14) \text{ TeV}$. Then the number of events will be $\simeq 1(117)$, in this case we have that the number of backgrounds is greater than the number of signals and consequently the statistical significance is $\simeq 4.4(2.8)\sigma$ for $m_{H_1^0} = 6500(800) \text{ GeV}$, for $v_\chi = 1000 \text{ GeV}$, for $\sqrt{s} = 14 \text{ TeV}$, and $\simeq 1.3(0.9)\sigma$ for $m_{H_1^0} = 650(800) \text{ GeV}$, for $v_\chi = 1500 \text{ GeV}$ and for $\sqrt{s} = 14$. Must be imposed a set of kinematic cuts on all the missing transverse momentum and the lepton momenta to improve the statistical significance of a signal, must be isolated a hard lepton from the W decay with $p_T^\ell > 20 \text{ GeV}$, put the cut on the missing transverse momentum $\cancel{p}_T > 20 \text{ GeV}$ and apply the Z window cut $|m_{\ell^+\ell^-} - m_Z| > 10 \text{ GeV}$, which removes events where the leptons come from Z decay [24].

B. The Higgs H_2^0

The Higgs H_2^0 in 3-3-1 model is not coupled to a pair of standard bosons, it couples to quarks, leptons, Z Z', Z'Z' gauge bosons, $H_1^- H_1^+$, $H_2^- H_2^+$, $h^0 h^0$, $H_1^0 H_3^0$ higgs bosons, $V^- V^+$

charged bosons, $U^{--}U^{++}$ double charged bosons, H_1^0Z , H_1^0Z' bosons and $H^{--}H^{++}$ double charged Higgs bosons [12]. The Higgs H_2^0 can be much heavier than 1018 GeV for $v_\chi = 1000$ GeV, and 1526 for $v_\chi = 1500$ GeV, so the Higgs H_2^0 is a heavy particle. The coupling of the H_2^0 with H_1^0 contributes to the enhancement of the total cross section *via* the Drell-Yan and gluon-gluon fusion.

In Fig. 7 and 8, we show the cross section $pp \rightarrow ZH_2^0$, these processes will also be studied in two cases, for the vacuum expectation value $v_\chi = 1000$ GeV and for $v_\chi = 1500$. Considering the expected integrated luminosity for the LHC collider given above, then the analysis for $v_\chi = 1000$ gives a total of $\simeq 1.7 \times 10^3(7.5 \times 10^2)(2.7 \times 10^5(1.4 \times 10^5))$ events per year for Drell-Yan and $\simeq 57(45)(9.5 \times 10^4(7.7 \times 10^4))$ events per year for gluon-gluon fusion, if we take the mass of the Higgs boson $m_{H_2^0} = 1100(1300)$ GeV ($\Gamma_{H_2^0} = 442.9, 715$ GeV). These values are in accord with the Table I. It must be noticed that must take care with large Higgs masses, as the width approaches the value of the mass itself for a very heavy Higgs and one loses the concept of resonance. Remember that the first two number of events $\simeq 1.7 \times 10^3(7.5 \times 10^2)$ are relative to Drell-Yan and the other two $\simeq 57(45)$ to gluon-gluon fusion and correspond to 8 TeV and the other pair $\simeq (2.7 \times 10^5(1.4 \times 10^5))$ are relative to Drell-Yan and $\simeq (9.5 \times 10^4(7.7 \times 10^4))$ to gluon-gluon fusion and correspond to 14 TeV for the LHC, respectively.

To obtain event rates we multiply the production cross sections by the respective branching ratios. Considering that the signal for H_2^0Z production for $m_{H_2^0} = 1100(1300)$ GeV and $v_\chi = 1000$ GeV will be $H_2^0Z \rightarrow ZH_1^0Z$, and taking into account that the branching ratios for these particles would be $B(H_2^0 \rightarrow ZH_1^0) = 46.2(47.3)$ % and $B(Z \rightarrow b\bar{b}) = 15.2$ %, see Fig. 9, and that the particles H_1^0 decay into W^+W^- , and taking into account that the branching ratios for these particles would be $B(H_1^0 \rightarrow W^+W^-) = 64.6(63.2)$ %, see Fig. 9, and then the W^+ decay into $\ell^+\nu$ and W^- into $\ell^-\bar{\nu}$ whose branching ratios for these particles would be $BR(W \rightarrow \ell\nu) = 10.8$ %, consequently we would have approximately $\simeq 0(0)(22(12))$ events per year for Drell-Yan and $\simeq 0(0)(8(6))$ for gluon-gluon fusion for the signal $b\bar{b}b\bar{b}\ell^+\ell^-X$.

With respect to vacuum expectation value $v_\chi = 1500$ GeV for the masses of $m_{H_2^0} = 1600(1800)$ ($\Gamma_{H_2^0} = 1303, 1838$ GeV) it will give a total of $\simeq 228(121)(6.8 \times 10^4(4.4 \times 10^4))$ events per year to produce H_2^0 for Drell-Yan and in respect to gluon-gluon fusion we will have $\simeq 0(0)(2.9 \times 10^3(2.5 \times 10^3))$ events per year to produce the same particles. Taking into

account the same signal as above, that is $b\bar{b}b\bar{b}\ell^+\ell^-X$ and considering that the branching ratios would be $B(H_2^0 \rightarrow ZH_1^0) = 48.3(48.7) \%$, $B(H_1^0 \rightarrow W^+W^-) = 65.7(65.1) \%$, $B(Z \rightarrow b\bar{b}) = 15.2 \%$ and $BR(W \rightarrow \ell\nu) = 10.8 \%$ we would have approximately $\simeq 0(0)(6(4))$ events per year for Drell-Yan and $\simeq 0(0)(0(0))$ for gluon-gluon fusion. The main background to this signal is $t\bar{t}Z \rightarrow b\bar{b}b\bar{b}\ell^+\ell^-X$, which cross section at LO is $\simeq 1$ pb for $\sqrt{s} = 14$ TeV. Then we have a total of $\simeq 530$ events for the background and $\simeq 30(18)$ events for the signal for $m_{H_2^0} = 1100(1300)$ GeV, for $\sqrt{s} = 14$ TeV and for $v_\chi = 1000$, by other side for $v_\chi = 1500$ the number of events for the signal is insignificant.

Therefore we have that the statistical significance is $\simeq 1.3(0.8)\sigma$ for $m_{H_2^0} = 1100(1300)$ GeV, that is a low probability to detect the signals. The improvement will be significant if we consider a luminosity $\simeq 10$ times higher than original LHC design, that is what we are awaiting to happen for 2025, then we will have $\simeq 300(180)$ events for the signals for $m_{H_2^0} = 1100(1300)$ GeV, for $\sqrt{s} = 14$ TeV and for $v_\chi = 1000$, which corresponds to have a 5σ discovery in the $b\bar{b}b\bar{b}\ell^+\ell^-X$ final state, by other side for $v_\chi = 1500$ we have $60(40)$ events for $m_{H_2^0} = 1600(1800)$ GeV, for $\sqrt{s} = 14$ TeV and which corresponds to $2.6(1.7)\sigma$, for this last scenario the signal is too small to be observed even with $3000 fb^{-1}$. To extract the signal from the background we must select the $b\bar{b}$ channel using the techniques of b-flavour identification, thus reducing the huge QCD backgrounds of quark and gluon jets, later the Z which coming together with the H_2^0 and the other Z coming from the decay of H_2^0 would appear as a peak in the invariant mass distribution of b-quark pairs. The charged lepton track from the W decay and the cut on the missing transverse momentum $\cancel{p}_T > 20$ GeV allows for a very strong reduction of the backgrounds.

The H_2^0Z will also decay into $t\bar{t}\ell^+\ell^-$, and consider that the branching ratios for these particles would be $B(H_2^0 \rightarrow t\bar{t}) = 6.0(4.5) \%$, see Fig. 9 and $B(Z \rightarrow \ell^+\ell^-) = 3.4 \%$ for the mass of the Higgs boson $m_{H_2^0} = 1100(1300)$ GeV, $v_\chi = 1000$ GeV, and that the particles $t\bar{t}$ decay into $b\bar{b}W^+W^-$, whose branching ratios for these particles would be $B(t \rightarrow bW) = 99.8 \%$, followed by leptonic decay of the boson W, that is $B(W \rightarrow e\nu) = 10.75 \%$, then we would have approximately $\simeq 0(0)(6(2))$ events per year. Regarding to gluon-gluon fusion we will have $\simeq 0(0)(2(1))$ events per year to produce the same particles. Considering the vacuum expectation value $v_\chi = 1500$ GeV and the branching ratios $B(H_2^0 \rightarrow t\bar{t}) = 3.1(2.5) \%$, see Fig. 10 and taking the same parameters and branching ratios for the same particles given above, then we would have for $m_{H_2^0} = 1600(1800)$ a total of $\simeq 0(0)(2(1))$ events of H_2^0

produced per year and in respect to gluon-gluon fusion the number of events per year will be $\simeq 0(0)(0(0))$. Taking again the irreducible background $t\bar{t}Z \rightarrow b\bar{b}e^+e^-\ell^+\ell^-X$, and using CompHep we have that a cross section at LO is $\simeq 1$ pb for $\sqrt{s} = 14$ TeV, which gives $\simeq 117$ events. We consider only the events for the signal for $v_\chi = 1000$, which gives $\simeq 8(2)$, for $v_\chi = 1500$, the number of events is insignificant.

Then we have that the statistical significance is $\simeq 0.7(0.3)\sigma$ for $m_{H_2^0} = 1100(1300)$ GeV, for $v_\chi = 1000$ GeV, for $\sqrt{s} = 14$ TeV. For this scenario the signal significance is smaller than 1σ and discovery can not be accomplished unless the luminosity will be improved. So, if we enhance the integrated luminosity up to $3000 fb^{-1}$, then we will have $\simeq 60(30)$ events for the signals for $m_{H_2^0} = 1100(1300)$ GeV, for $\sqrt{s} = 14$ TeV and for $v_\chi = 1000$, which corresponds to have a $5.6(2.8)\sigma$ discovery in the $b\bar{b}e^+e^-\ell^+\ell^-X$ final state and for $v_\chi = 1500$ the signal will be not visible in this channel. We impose the following cuts to improve the statistical significance of a signal, i. e. we isolate a hard lepton from the W decay with $p_T^\ell > 20$ GeV, put the cut on the missing transverse momentum $\cancel{p}'_T > 20$ GeV and apply the Z window cut $|m_{\ell^+\ell^-} - m_Z| > 10$ GeV, which removes events where the leptons come from Z decay [24]. However, all this scenarios can only be cleared by a careful Monte Carlo work to determine the size of the signal and background.

In summary, we showed in this work that in the context of the 3-3-1 model the signatures for neutral Higgs bosons can be significant in LHC collider. Our study indicates the possibility of obtaining a clear signal of these new particles through its different modes of decay. If this model is realizable in the nature, certainly new particles will appear such as $H_1^0, H_2^0, Z', P^\pm, H^{\pm\pm}$ in the context of this study.

Acknowledgments

MDT is grateful to the Instituto de Física Teórica for hospitality UNESP, the Brazilian agencies CNPq for a research grant, and FAPESP for financial support (Processo No. 2009/02272-2).

[1] Damiano Anselmi, Eur. Phys. J. C **65**, 523 (2010); P. Binétruy, S. Chadha, P. Sikivie, Nucl. Phys. **B207**, 505 (1982); S. Dimopoulos, L. Susskind, Nucl. Phys. **B155**, 237 (1979).

- [2] J. Wess and B. Zumino, Nucl. Phys. **B70** (1974) 39.
- [3] J. Wess and B. Zumino, Phys. Lett. **49B** (1974) 52; J. Iliopoulos and B. Zumino, Nucl. Phys. **B76** (1974) 310; S. Ferrara, J. Iliopoulos and B. Zumino, Nucl. Phys. **B77** (1974) 413; E. Witten, Nucl. Phys. **B188** (1981) 513.
- [4] S. Dimopoulos and H. Georgi, Nucl.Phys. **B193** (1985) 150; S. Dimopoulos, S. Raby, and F. Wilczek, Phys. Rev. D **24** (1981) 1681; L. Ibañez and G. G. Ross, Phys. Lett. **105B** (1981) 439.
- [5] V. Pleitez and M. D. Tonasse, Phys. Rev. D **48**, 2353 (1993).
- [6] F. Pisano and V. Pleitez, Phys. Rev. D **46**, 410 (1992);
- [7] P. H. Frampton, Phys. Rev. Lett. **69**, 2889 (1992).
- [8] A. G. Dias, Phys. Rev. D **71**, 015009 (2005).
- [9] M. D. Tonasse, Phys. Lett. B **381**, 191 (1996).
- [10] V. Barger, E. W. N. Glover, K. Hikasa, W. Y. Keung, M. G. Olsson, C. J. Suchyta and X. R. Tata, Phys. Rev. Lett. **57**, 1672 (1986); Duane A. Dicus and Chung Kao, Phys. Rev. D **38**, 1008 (1988).
- [11] J. E. Cieza Montalvo and M. D. Tonasse, Nucl. Phys. **B623** (2002) 325.
- [12] J. E. Cieza Montalvo and M. D. Tonasse, Phys. Rev. D **71**, 095015 (2005).
- [13] K. Nakamura *et al.*, (Particle Data Group), J. Phys. G **37**, 075021 (2010).
- [14] M. Carena *et al.*, Report of the Tevatron Higgs Working Group, hep-ph/0010338; A. Stange, W. J. Marciano and S. Willenbrock, Phys. Rev. D **49**, 1354(1994) and Phys. Rev. D **50**, 4491(1994).
- [15] G. Aad *et al.*, Phys. Lett. B **710**, 49 (2012).
- [16] ATLAS Technical Proposal, CERN/LHC/94-43 LHCC/P2 (December 1994); CMS Technical Proposal, CERN/LHC/94-43 LHCC/P1 (December 1994).
- [17] Z. Kunszt, S. Moretti and W. J. Stirling, Zeit. Phys. **C 74** (1997)479; ATLAS Technical Proposal, CERN/LHC/94-43 LHCC/P2 (December 1994); CMS Technical Proposal, CERN/LHC/94-43 LHCC/P1 (December 1994).
- [18] S. Moretti, hep-ph/9612310.
- [19] R. N. Cahn and M. S. Chanowitz, Phys. Rev. Lett. **56**, (1986)1327.
- [20] J. E. Cieza Montalvo, Nelson V. Cortez and M. D. Tonasse, Phys. Rev. D **76**, 117703 (2007).
- [21] CMS collaboration (2012). “Observation of a new boson with a mass near 125 GeV”. Cms-

- [22] ATLAS collaboration (2012). “Observation of an Excess of Events in the Search for the Standard Model Higgs boson with the ATLAS detector at the LHC”. Atlas-Conf-2012-093.
- [23] A. Pukhov *et. al.*, hep-ph/9908288.
- [24] F. del Aguila, J. A. Aguilar-Saavedra, Nucl. Phys. B 813 (2009) 22; A. G. Akeroyd, Cheng-Wei Chiang, Naveen Gaur, JHEP 1011 (2010) 005.
- [25] J. E. Cieza Montlavo, G. H. Ramirez Ulloa and Mauro D. Tonasse, Eur.Phys.J.C72, 2210 (2012).
- [26] J. E. Cieza Montalvo, Nelson V. Cortez, J. Sá Borges and M. D. Tonasse, Nucl. Phys. **B756** (2006) 1-15.

FIGURE CAPTIONS

Figure 1: Feynman diagrams for production of neutral Higgs *via* Drell-Yan process.

Figure 2: Feynman diagrams for production of neutral Higgs *via* gluon-gluon fusion.

figure 3: Total cross section for the process $p p \rightarrow ZH_1^0$ as a function of $m_{H_1^0}$ for a $v_\chi = 1.0(1.5)$ TeV at $\sqrt{s} = 8$ TeV for Drell-Yan- solid line(dot-dash line) and Gluon-Gluon fusion-dash line(dot-dot dash line);

figure 4: Total cross section for the process $p p \rightarrow ZH_1^0$ as a function of $m_{H_1^0}$ for a $v_\chi = 1.0(1.5)$ TeV at $\sqrt{s} = 14$ TeV for Drell-Yan- solid line(dot-dash line) and Gluon-Gluon fusion-dash line(dot-dot dash line);

figure 5: BRs for the H_1^0 decays as functions of $m_{H_1^0}$ for $v_\chi = 1.0$ TeV.

figure 6: BRs for the H_1^0 decays as functions of $m_{H_1^0}$ for $v_\chi = 1.5$ TeV.

figure 7: Total cross section for the process $p p \rightarrow ZH_1^0$ as a function of $m_{H_2^0}$ for a $v_\chi = 1.0(1.5)$ TeV at $\sqrt{s} = 8$ TeV for Drell-Yan- solid line(dot-dash line) and Gluon-Gluon fusion-dash line(dot-dot dash line);

figure 8: Total cross section for the process $p p \rightarrow ZH_1^0$ as a function of $m_{H_2^0}$ for a $v_\chi = 1.0(1.5)$ TeV at $\sqrt{s} = 14$ TeV for Drell-Yan- solid line(dot-dash line) and Gluon-Gluon fusion-dash line(dot-dot dash line);

figure 9: BRs for the H_2^0 decays as functions of $m_{H_2^0}$ for $v_\chi = 1.0$ TeV.

figure 10: BRs for the H_2^0 decays as functions of $m_{H_2^0}$ for $v_\chi = 1.5$ TeV.



















