Possible High-Tc Superconductivity in Z = 7.33 Materials from Combinatorial Synthesis of Z = n (n=4.67, 10, 12.67) Families

O. P. Isikaku-Ironkwe^{1, 2}

¹The Center for Superconductivity Technologies (TCST)

Department of Physics,

Michael Okpara University of Agriculture, Umudike (MOUAU),

Umuahia, Abia State, Nigeria

and

²RTS Technologies, San Diego, CA 92122

Abstract

The search for MgB₂-like superconductivity has before now been limited to binary and ternary systems. However systems with more-than-three-elements that meet the MgB₂ model have yet to be investigated either computationally or experimentally. Here we show that through computational synthesis of two binary or ternary materials from the Z=4.67, 10 and 12.67 families, we can get Z=7.33 systems with more-than-three elements. We have identified over 300 of such possibilities for further studies. We propose that the resultant combinatorial synthesis of such Z=7.33 systems with more-than-three-elements and that meet the MgB₂ model will be superconducting with Tcs higher than 39K.

Keywords: more-than-three-elements (mtte) MgB₂-like superconductors, mtte-MgB₂-like superconductors, Material specific characterization dataset (MCSD)

Introduction

The search for novel high-temperature superconductors has been described as "high-risk, high-payoff" project [1] that requires "passionate optimism" [2] and lots of serendipity [3, 4]. The goal is to move the field from discovery by serendipity to discovery by rational design [3, 4]. Achieving this requires understanding the chemical building blocks of superconductivity or the material specificity [5] of superconductivity. The discovery of magnesium diboride as a superconductor [6] shattered long believed search paradigms [7, 8] and introduced a new paradigm [5] of superconductivity search. It was not immediately obvious to many researchers [11, 12] that structural and electronic similarities were not sufficient conditions for basing a search for MgB_2 -like superconductivity. This led to predictions and experimental results [9 – 12] far below the anticipated high-Tc of MgB_2 . Turning the search to material

specific parameters in the Periodic Table such as electronegativity, valence electrons, atomic number and formula weight [5], we have been able to show that novel binary and ternary MgB₂-like superconductors with high-Tc do exist [13 - 21], awaiting experimental confirmation. We observe that all predicted MgB₂-like superconductors have been binary or ternary systems. In this paper, we extend the search to potential quaternary systems that still meet the MgB₂-like model. These new class of more-than-three-elements (mtte) materials is composed entirely of light elements and supersede the binaries and ternaries with similar properties proposed in [22]. These new mtte-MgB₂-like compounds are formed by the appropriate combinations of the Z =4.67 family [20] with either the Z =10 family [19] or the Z =12.67 family [21]. In this paper, we show the computational combinatorial synthesis process and some of the resultant compounds and their MSCDs. We also provide computational predictions of their Tcs using our Tc estimation formula [5] and superconductivity search rules.

Binary and Ternary Z=7.33 Superconductors

The Z=7.33 family represents materials whose average atomic number is 7.33. MgB₂ is one of them. Many binary and ternary Z=7.33 materials have been computationally proposed that show MgB₂-like superconductivity [13 - 21]. Table 1 shows some of them and their MSCDs. Their Tcs have been estimated using the new material specific Tc equation derived in [5], given by:

$$\mathbf{T_c} = \mathcal{X} \frac{\mathrm{Ne}}{\sqrt{Z}} \, \mathbf{K_o} \tag{1}$$

where Ko = n(Fw/Z) and n can be estimated empirically. For magnesium diboride, n = 3.65. Fw is formula weight and Z is average atomic number. The value of n may be less for other Z = 7.33 materials.

Combinatorial Synthesis of Novel Z=7.33 Materials

There have been no reports of the existence of more-than-three-elements (mtte)-MgB₂-like superconductors. Such materials would be required to have the same valence electron count

and atomic number as MgB₂. We propose that mtte-MgB₂-like superconductors could result from the combinatorial synthesis of two families (Fz): where Z=4.67, 10 or 12.67 materials, with Ne=2.67 in each case, according to the equations:

$$F_{7=4.67} + F_{7=10} = F_{7=7.33}$$
 (2)

$$2F_{z=4.67} + F_{z=12.67} = F_{z=7.33}$$
 (3)

These combinations are illustrated in figures 1 and 2. Tables 2, 3 and 4 show some of the potential materials in the three families under consideration. Equation (2) states that 1:1 combinatorial synthesis of the $Fz_{=4.67}$ and $Fz_{=10}$ families will yield a new family $Fz_{=7.33}$. From equation (3), the combinatorial synthesis of two parts of a $Fz_{=4.67}$ family member with one part of a $Fz_{=12.67}$ family member, will be a new compound belonging to the family $Fz_{=7.33}$. The expected MSCD [5] of the new compound will be:

MSCD of compound =
$$\langle \mathcal{X}, \text{Ne}, \text{Z}, \text{Ne}/\sqrt{Z} \rangle$$
 (4)
= $\langle \mathcal{X}, 2.667, 7.333, 0.9847, \text{Fw}, \text{Fw/Z} \rangle$

where Ne is the valence electron count of each family = 2.667, the average atomic number Z =7.333 and the average final electronegativity, \mathcal{X} . Fw is the total formula weight of the new Z=7.33 material. In [5] we showed that a high-Tc should be expected when:

$$0.8 < Ne/\sqrt{Z} < 1.0$$
 (5)

Thus from equation (4) and (5) the resultant combinatorial synthesis of the families as defined in equations (2) and (3), should lead to MgB₂-like materials with high Tc.

Combinatorial Results

Tables 5 and 6 show some of the results of implementing the combinatorial synthesis of two families ($F_{z=4.67}$ and $F_{z=10}$) of materials with the same Ne =2.67 in the 1:1 ratio indicated. In each case we end up with MgB₂-like materials with Z=7.333 and Ne=2.67 as predicted from equation (4) above. The number of possible compounds is the product of the numbers of compounds in each family, in this case 170. In Table 6, we show just 12 out of the 170 MSCDs possibilities. As observed, some will be 3-, 4-, 5- or 6-elements MgB₂-like materials with Z=7.33 while the electronegativities vary. From Table 6, except for the 3-atom case with Fw/Z

=6.26, the 6-atom cases have average Fw/Z of 12.52. Tables 7 and 8 show some of the results of computational combinatorial synthesis of two parts of a member of family $F_{\mathbf{Z}=4.67}$ with one part of a member of the $F_{\mathbf{Z}=12.67}$ family. With 10 members of $F_{\mathbf{Z}=4.67}$ and 16 members in $F_{\mathbf{Z}=12.67}$, we have 160 minimum possible combinations. Again, some will be 3-, 4-, 5- or 6-elements MgB_2 -like materials with Z=7.33 while the electronegativity varies. Only 6 possible results are listed in Table 7 with their MSCDs in Table 8. From Table 8 we note that we get 9-atom systems with 4 or 5 elements. The average Fw/z is 18.77. We summarize the variation of Fw/Z with number of atoms in Table 9 and plot Fw/z vs computed Tc in Figure 3.

Analysis of Combinatorial Results

Figures 1 and 2 shows the combinatorial process which results in the creation of Z=7.33 family, noted for hosting MgB_2 . A significant aspect of this process is the creation of 4-, 5-, 6-element compounds that meet the model of MgB_2 . If they are found to exist and are stable, we may have for the first time, materials with Tcs higher than MgB_2 but meet its specifications. This will be because Fw/z is higher as explained in [5] with the derived equation for Tc [5] in terms of electronegativity, x, valence electron count, Ne, atomic number, Z, and formula weight, Fw.

Discussion

Rosner et al. [9] proposed that hole-doping LiBC, a Z=4.67 material to Li_{0.5}BC might lead to higher Tc than found in MgB₂. Though their concept was right, the approach was not. Working with the paradigm that such doping should not change the valence electron count and the average atomic number, we have been able to "dope" Z=4.67 materials with Z = 10.0 and Z=12.67 materials resulting in Z=7.33 materials with Fw/Z much higher than that of MgB₂. Looking at the elements involved we see that they are mostly combinations of the 12 lightest elements with Z<20 in such a manner as to give an average Z= 7.333 and Ne=2.667. The electronegativity rarely exceeds 1.733. For Fw/Z of 12.52 the computed Tc is 63K for the 6-

atom systems with Z=7.33 MSCD and electronegativity same as MgB₂. The highest Fw/Z for these compounds was 18.77 which we have computed should correspond to a Tc of 75K.

Conclusions

This research revealed over 300 potential new high-Tc MgB₂-like superconductors in the hitherto unexplored chemical world of 6- and 9- atom mtte-compounds with average atomic number, Z, of 7.333 and average valence electron count, Ne, of 2.667. These novel 6-atom and 9-atom MgB₂-like materials with Fw/Z as high as 18.77 strongly suggest that they may have Tcs as high as 75K (table 9 and figure 3). Experimental confirmation of these predictions presents a strong challenge to experimentalists and computational material scientists and opens a whole new world of mtte-MgB₂-like superconductors, too early to evaluate.

Acknowledgements

The author acknowledges useful discussions with Professor M. Brain Maple at U.C. San Diego. This research was executed with grants from Dr. M. J. Schaffer, formerly at General Atomics, San Diego. Research literature from Dr. Jim O'Brien facilitated my work.

References

- DOE-BES, "Basic Research Needs for Superconductivity": Report of the Basic Energy Sciences Workshop on Superconductivity (2006). pp 99, http://science.energy.gov/~/media/bes/pdf/reports/files/sc_rpt.pdf
- 2. D. Larbalestier and P.C. Canfield, "Superconductivity at 100---where we've been and where we're going", MRS bulletin 36, 590 -593, August 2011
- DOE-BES, "Basic Research Needs for Superconductivity": Report of the Basic Energy Sciences Workshop on Superconductivity 2006. pp 21 - 23 http://science.energy.gov/~/media/bes/pdf/reports/files/sc_rpt.pdf
- 4. L.H. Greene, "Taming Serendipity", Physics World 24, 41 43, 2011
- 5. O.P. Isikaku-Ironkwe, "Transition temperatures of superconductors estimated from Periodic Table properties" Arxiv: 1204.0233 (2012) and references therein.
- 6. J. Nagamatsu, N. Nakagawa, T. Muranaka, Y. Zenitani and J. Akimitsu, "Superconductivity at 39K in Magnesium Diboride," Nature 410, 63 (2001)

- 7. B.T. Matthias, "The search for high-temperature superconductors", Physics Today, August 1971, 23 -27
- 8. A.S. Cooper, E. Corenzwit, L.D. Longinotti, B.T. Matthias and W.H. Zachariassen, "Superconductivity: the transition temperature peak below four electrons per atom", Proc. Nat. Acad. Sci., 67, 313 -319 (1970).
- 9. H. Rosner, A. Kitaigorodsky and W.E. Pickett, "Prediction of High Tc superconductivity in hole-doped LiBC", Phys. Rev. Lett. 88, 127001 (2002)
- 10. A. Bharathi, S. Jemima Balaselvi, M. Premila, T.N. Sairam, G.L.N. Reddy, C.S. Sundar, Y. Hariharan, "Synthesis and search for superconductivity in LiBC", Solid State Comm., (2002), 124, 423
- 11. R.L. Meng, B. Lorenz, J. Cmaidalka, Y.S. Wang, Y.Y. Sun, J. Lenzi, J.K. Meen, Y.Y. Xue and C.W. Chu, "Study of intermetallic compounds iso-structural to MgB_{2,}" IEEE Trans. Applied Superconductivity, Vol. 13, 3042- 3046 (2002).
- 12. R.L. Meng, B. Lorenz, Y.S. Wang, J. Cmaidalka, Y.Y. Xue, J.K. Meen. C.W. Chu "Study of binary and pseudo-binary intermetallic compounds with AlB₂ structure." Physica C: 382 113–116(2002).
- 13. O. P. Isikaku-Ironkwe, "Possible High-Tc Superconductivity in LiMgN: A MgB₂-like Material", Arxiv: 1204.5389 (2012) and references therein.
- 14. O. P. Isikaku-Ironkwe, "Prospects for superconductivity in LiBSi", Arxiv: 1205. 2407 (2012)
- 15. O. P. Isikaku-Ironkwe, "Is Lithium Sulfide a MgB₂-like Superconductor?", Arxiv: 1205.4051
- 16. O. P. Isikaku-Ironkwe, "Beryllium Silicide Clusters, Be_nSi_n , $Be_{2n}Si_n$ (n = 1 4) and possible MgB₂-like superconductivity in some of them", Arxiv: 1205.5931
- 17. O. P. Isikaku-Ironkwe, "MgBeC: A potential MgB₂-like Superconductor", Arxiv:1205.6237
- 18. O. P. Isikaku-Ironkwe, "Lithium Beryllium Phosphide (LiBeP): A possible MgB₂-like Superconductor", Arxiv:1206.1915
- 19. O. P. Isikaku-Ironkwe, "Searching for superconductivity in Z=10 Family of Materials", Arxiv:1206.3006
- 20. O. P. Isikaku-Ironkwe, "Searching for Superconductivity in the Z=4.67 Family", Arxiv:1206.4016
- 21. O. P. Isikaku-Ironkwe, "Searching for Superconductivity in the Z=12.67 Family", Arxiv:1206.4791
- 22. A. Hermann, N.W. Ashcroft and R. Hoffman, "Light element ternary compounds—searching for new superconductors in the 'upper left corner' " http://meetings.aps.org/BAPS.2012.MAR.W21.10

Tables

Ma	terial	x	Ne	Z	Ne/ \sqrt{Z}	Fw	Fw/Z
1	MgB ₂	1.733	2.6667	7.333	0.9847	45.93	6.263
2	Li ₂ S	1.5	2.6667	7.333	0.9847	45.95	6.266
3	Be₂Si	1.6	2.6667	7.333	0.9847	46.11	6.290
4	LiB eP	1.533	2.6667	7.333	0.9847	46.92	6.398
5	LiMgN	1.733	2.6667	7.333	0.9847	45.26	5.999
6	MgBeC	1.733	2.6667	7.333	0.9847	45.33	6.181
7	LIBSi	1.6	2.6667	7.333	0.9847	45.84	6.251

Table 1 MSCD of some 7 binary and ternary Z =7.333 materials, with Ne=2.667. \mathcal{X} = electronegativity, Ne = valence electron count, Z= average atomic number and Fw = formula weight Reference [5] describes how to compute MSCD.

Ma	terial	x	Ne	Z	Ne/ \sqrt{Z}	Fw	Fw/Z
1	LiBC	1.8333	2.6667	4.6667	1.2344	29.76	6.377
2	BeB ₂	1.8333	2.6667	4.6667	1.2344	30.63	6.564
3	Be ₂ C	1.8333	2.6667	4.6667	1.2344	30.03	6.435
4	Li₂O	1.8333	2.6667	4.6667	1.2344	29.88	6.403
5	LiBeN	1.8333	2.6667	4.6667	1.2344	29.96	6.420
6	LiB ₅	1.8333	2.6667	4.6667	1.2344	60.99	13.07
7	Li ₃ BN ₂	1.8333	2.6667	4.6667	1.2344	59.65	12.78
8	NaAlH ₄	1.8	1.3333	4.6667	0.6172	54.01	11.574
9	KBH ₄	1.8667	1.3333	4.6667	0.6172	53.95	11.56
10	MgH ₂	1.8	1.3333	4.6667	0.6172	26.33	5.642

Table 2: 10 MSCDs of some Z =4.67 materials with Ne=2.67 and 1.33

Ma	terial	x	Ne	Z	Ne/√Z	Fw	Fw/Z
1	MgBeSi	1.833	2.667	10.0	0.8433	61.41	6.141
2	Na₂O	1.767	2.667	10.0	0.8433	61.98	6.198
3	KBeN	1.767	2.667	10.0	0.8433	62.12	6.212
4	NaMgN	1.7	2.667	10.0	0.8433	61.31	6.131
5	LiCaN	1.667	2.667	10.0	0.8433	61.03	6.103
6	CaB ₂	1.667	2.667	10.0	0.8433	61.70	6.170
7	CaBeC	1.667	2.667	10.0	0.8433	61.10	6.110
8	Mg ₂ C	1.633	2.667	10.0	0.8433	60.63	6.063
9	NaAIC	1.633	2.667	10.0	0.8433	61.98	6.198
10	BeBSc	1.6	2.667	10.0	0.8433	64.78	6.478
11	NaBSi	1.567	2.667	10.0	0.8433	61.89	6.189
12	BeAl ₂	1.5	2.667	10.0	0.8433	62.97	6.297
13	Be₂Ti	1.5	2.667	10.0	0.8433	65.90	6.590
14	NaBeP	1.5	2.667	10.0	0.8433	62.97	6.297
15	LiNaS	1.467	2.667	10.0	0.8433	62.00	6.200
16	LiMgP	1.433	2.667	10.0	0.8433	62.22	6.222
17	LiAlSi	1.433	2.667	10.0	0.8433	62.01	6.201

Table 3: 17 MSCDs of Z =10.0 materials.

Ma	terial	X	Ne	Z	Ne/ \sqrt{Z}	Fw	Fw/Z
1	CaBeSi	1.433	2.667	12.667	0.7493	77.18	6.09
2	Na₂S	1.433	2.667	12.667	0.7493	78.05	6.16
3	LiKS	1.433	2.667	12.667	0.7493	78.11	6.17
4	NaAlSi	1.4	2.667	12.667	0.7493	78.06	6.16
5	Mg₂Si	1.4	2.667	12.667	0.7493	76.71	6.06
6	NaMgP	1.4	2.667	12.667	0.7493	78.27	6.18
7	KAIC	1.6	2.667	12.667	0.7493	78.09	6.16
8	NaCaN	1.633	2.667	12.667	0.7493	77.08	6.09
9	LiBGe _{0.89} Si _{0.11}	1.6	2.667	12.667	0.7493	85.46	6.75
10	KBSi	1.533	2.667	12.667	0.7493	78.0	6.16
11	KBeP	1.467	2.667	12.667	0.7493	79.09	6.24
12	MgAl ₂	1.4	2.667	12.667	0.7493	78.27	6.18
13	LiAlTi	1.333	2.667	12.667	0.7493	81.8	6.46
14	LiCaP	1.367	2.667	12.667	0.7493	77.99	6.16
15	Na ₃ AlP ₂	1.4	2.667	12.667	0.7493	157.89	12.465
16	K ₃ BN ₂	1.7333	2.667	12.667	0.7493	156.13	12.326

Table 4: 16 MSCDs of Z =12.667 materials.

	Fz=4.67	Fz=10	12 of the combinatorial results: Fz=4.67 + Fz=10 for 170 combinatorial possibilities.			
1	BeB ₂	Na ₂ O	Na ₂ BeB ₂ O	LiNaBeBCP		
2	Li ₂ O	Mg ₂ C	LiNa ₂ BCO	LiNa ₂ B ₅ O		
3	Be ₂ C	NaBeP	Li ₃ MgPO	Li ₂ NaAlCO		
4	MgH ₂	KBeN	Na ₂ Be ₂ CO	Li ₂ Mg ₂ CO		
5	LiB ₅	LiMgP	MgBeC	NaBe ₂ AlC ₂		
6	LiBC	NaAlC	LiMg ₂ BC ₂	LiNaBAIC ₂		

Table 5: 12 of the combinatorial results for 170 combinatorial possibilities from the two families of materials (Fz=4.67 + Fz=10).

Ma	terial	x	Ne	Z	Ne/ \sqrt{Z}	Fw	Fw/Z
1	Na ₂ BeB ₂ O	1.8	2.6667	7.3333	0.9847	92.61	12.63
2	LiNa₂BCO	1.8	2.6667	7.3333	0.9847	91.74	12.51
3	Li₃MgPO	1.6333	2.6667	7.3333	0.9847	92.10	12.56
4	Na ₂ Be ₂ CO	1.8	2.6667	7.3333	0.9847	92.01	12.55
5	MgBeC	1.7333	2.6667	7.3333	0.9847	45.93	6.26
6	LiMg ₂ BC ₂	1.7333	2.6667	7.3333	0.9847	90.39	12.34
7	Li ₂ Mg ₂ CO	1.7333	2.6667	7.3333	0.9847	90.51	12.34
8	Li ₂ NaAlCO	1.7333	2.6667	7.3333	0.9847	91.86	12.53
9	LiNaBeBCP	1.6667	2.6667	7.3333	0.9847	92.73	12.64
10	Mg ₃ H ₂ C	1.7167	2. 0	7.3333	0.7386	86.96	11.86
11	NaBe ₂ AIC ₂	1.7333	2.6667	7.3333	0.9847	92.01	12.55
12	LiNaBAlC ₂	1.7333	2.6667	7.3333	0.9847	91.74	12.51

Table 6: MSCDs of 12 out of 170 combinatorial materials possibilities from (Fz=4.67 + Fz=10). The combinations result in 6-atom compounds with 3, 4, 5 and 6 different elements.

	Fz=4.67	Fz=12.667	6 of the combinatorial results:		
	12	11 11:007	2Fz=4.67 + Fz=12.67 for 160		
			combinatorial possibilities.		
1	Li ₂ O	Mg ₂ Si	Li ₄ Mg ₂ SiO ₂		
2	Be ₂ C	Na ₂ S	Na ₂ Be ₄ C ₂ S		
3	LiBC	CaBeSi	CaBe ₅ SiC ₂		
4	LiBeN	KBSi	KBe ₄ BSiC ₂		
5	LiB ₅	LiAlTi	LiBe₄AlTiC₂		
6	Li ₃ BN ₂	LiCaP	Li ₂ Na ₂ B ₂ C ₂ S		

Table 7: 6 of the combinatorial results for 160 combinatorial possibilities in (2Fz=4.67 + Fz=12.67).

	Material	x	Ne	Z	Ne/ \sqrt{Z}	Fw	Fw/Z
1	Li ₄ Mg ₂ SiO ₂	1.689	2.6667	7.3333	0.9847	136.47	18.61
2	Na ₂ Be ₄ C ₂ S	1.7	2.6667	7.3333	0.9847	138.11	18.33
3	CaBe₅SiC₂	1.7	2.6667	7.3333	0.9847	137.24	18.72
4	KBe ₄ BSiC ₂	1.733	2.6667	7.3333	0.9847	138.06	18.83
5	LiBe ₄ AlTiC ₂	1.667	2.6667	7.3333	0.9847	141.86	19.35
6	Li ₂ Na ₂ B ₂ C ₂ S	1.7	2.6667	7.3333	0.9847	137.57	18.76

Table 8: MSCD of 6 out of 160 combinatorial materials possibilities from (Fz=4.67 + Fz=10) of data from tables 2, 4 and 7. The combinations result in 9-atom compounds with 4 and 5 elements.

#Atoms (An)	Fw/Z(Ave)	n	Ko=n(Fw/Z)	Tc(K)	Example	Reference
3	6.26	3.65	22.85	39	MgB ₂	Figure 2 of [5]
6	12.52	2.95	36.91	63	NaBe ₂ AlC ₂	Table 6 of this paper + Figure 2 of [5]
9	18.77	2.34	43.94	75	KBe ₄ BSiC ₂	Table 8 of this paper + Figure 2 of [5]

Table 9: Data on variation of Fw/Z with number of atoms for the Z=7.33 family of materials of which MgB₂ belongs. All these examples have same \mathcal{X} , Ne, and Z , but different Fw/Z. Using the material specific Tc formula: $\mathbf{T_c} = \mathcal{X} \frac{\mathrm{Ne}}{\sqrt{Z}} \, \mathrm{K_o}$ derived in reference [5].

Figures




Figure 1: Combining materials from Fz=4.67 and Fz=10.0 yields a new family Fz=7.33 material with Ne=2.67 and Z=7.33. The electronegativity will be an average of the electronegativities of Fz=4.67 and Fz=10.0




Figure 2: Combining 2x a material from Fz=4.67 with one material from Fz=12.67 yields a new family Fz=7.33 material with Ne=2.67 and Z=7.33. The electronegativity will be an average of the electronegativities of 2(Fz=4.67) and Fz=12.67.




Figure 3: Fw/Z versus Tc. From Reference[5] we know that Tc increases with Fw/Z. The Z=7.33 materials with the same MSCD as MgB2 should have the same Tc but since they have higher Fw/Z, we should expect a higher Tc by equation (1) in this paper and figure 2 of Ref [5].