

Stretched Magnetic Moments

L. Zamick and Y.Y. Sharon

Department of Physics and Astronomy, Rutgers University, Piscataway, New Jersey 08854 U.S.A.

We here wish to compile some simple expressions for nuclear magnetic moments. The results are not new but it may be useful to have things all in one place.

To make things concrete we use as an example a paper by G. Ilie et al. [1]. In that work they measured the g factor of the lowest 7^- state of ^{126}Sn . They note that the leading configuration is $[h_{11/2} d_{3/2}]J=7$ and they correctly give the value of the g factor of this configuration as -0.11.

Note that the above configuration is stretched. That is to say $J=7$ is the largest possible value one can have. For a stretched configuration the magnetic moments that add. We now give more details of this not unknown result.

By convention the magnetic moment of a system of particles with total angular momentum J is defined as the expectation value of the z component of the magnetic moment operator in a state with $M=J$. For a system of two nucleons with configuration $[j_1, j_2]$ with $J=j_1+j_2$ (stretched case) the wave function factorizes.

$$\Psi(j_1, j_2, J, M=J) = \psi(j_1, m=j_1) \psi(j_2, m=j_2)$$

Note that for each of the 2 wave functions on the right we have $m=j$ which is just what you want for getting the magnetic moment. From this it obviously follows that $\mu(J) = \mu(j_1) + \mu(j_2)$ for this stretched case. For the above configuration the values for the 2 terms are -1.913 (free neutron) and -3/5 (-1.913) so that the total is 2/5 (-1.913).

More generally for any J value the relation satisfied by the g factor is a bit more complicated:

$$g = (g_1 + g_2)/2 + (g_1 - g_2) / (2 J(J+1)) * [j_1(j_1+1) - j_2(j_2+1)]$$

For the above configuration one obtains $g = -2/35 * \mu(\text{free neutron}) = -0.109$. This is close to the experimental value -0.098(9).

In some cases the formula for g simplifies. If j_1 is equal to j_2 we have

$$g = (g_1 + g_2)/2$$

We also have this result if $g_1 = g_2$. This has the consequence that all g factors for identical nucleons in a single j-shell (e.g. only neutrons) are the same. We also get this result for a system with an equal number of protons and neutrons in a single j shell; also equal number of protons and neutron- holes or neutrons and proton- holes.

[1] G. Ilie et al., Phys.Lett. B 687 (2010)305