

TDGL Equations for Two-Band Superconductor

V.L.OSTROVSKI

ORT Braude college,Karmiel,21982, Israel

Abstract

Green's function techniques for studying nonequilibrium processes in two-band superconductor are discussed. Perturbation expansions and Green's function equations are developed. A time dependent modification of the Ginsburg-Landau equation is given.

Introduction.

Time dependent Ginzburg-Landau (TDGL) equation has proven to be very successful in the theory of superconductivity [1-11]. The usual classification of superconductors characterizes materials by one-component order parameter. However the properties of some compounds (e.g. MgB_2 , Nb_2Se , V_3Si) are described by two-band model [12] with two- component order parameter. The two-gap GL model (the two gaps arise from the existence of two bands) has been proposed [13, 14] at the dawn of superconductivity theory. The interest to the two-gap model has recently been renewed after the discovery the superconductivity in magnesium diboride[15-42]. The studies below are based on a set of Ginzburg-Landau equations for the order parameters of a two-band system. We focus our attention mainly on the overlapping of energy bands on the Fermi surface and on anisotropy due to different dimensions of the energy bands under study.

E-mail: vostrovski@gmail.com

1. Green's functions in the Gor'kov representation.

The two-band pairing Hamiltonian in the BCS theory in the absence of external field is given by

$$\hat{H} = \hat{H}_0 + \hat{H}_{\text{int}} \quad (1.1)$$

Where

$$\hat{H}_0 = \int d^3\vec{r} \sum_{n\sigma} \hat{\psi}_{n\sigma}^+(\vec{r}) \hat{\varepsilon}_n(\hat{\vec{p}}) \hat{\psi}_{n\sigma}(\vec{r}) \quad (1.2)$$

is the Hamiltonian of noninteracting quasiparticles, $\hat{\varepsilon}_n(\hat{\vec{p}}) = \zeta_n + \frac{\hat{\vec{p}}^2}{2m_n} - \mu$, μ is the chemical potential, $n=1, 2$ are the band indexes, $\varepsilon_n(\vec{p}) = \zeta_n + \frac{\vec{p}^2}{2m_n} - \mu$ is the dispersion relation of quasiparticles in the n -th band, m_n is the effective mass, and $\hat{\psi}_{n\sigma}^+, \hat{\psi}_{n\sigma}$ are the creation and annihilation field operators in the Heisenberg representation for quasiparticles in band n with spin $\sigma = \uparrow$ or \downarrow .

$$\begin{aligned} \hat{H}_{\text{int}} &= - \sum_{nm} V_{nm} \int d^3\vec{r} \sum_{\sigma\sigma'} \hat{\psi}_{n\sigma}^+(\vec{r}) \hat{\psi}_{n\sigma'}^+(\vec{r}) \hat{\psi}_{m\sigma'}(\vec{r}) \hat{\psi}_{m\sigma}(\vec{r}) (1 - \delta_{\sigma\sigma'}) = \\ &= - \sum_{nm} V_{nm} \int d^3\vec{r} [\hat{\psi}_{n\uparrow}^+(\vec{r}) \hat{\psi}_{n\downarrow}^+(\vec{r}) \hat{\psi}_{m\downarrow}(\vec{r}) \hat{\psi}_{m\uparrow}(\vec{r}) + \\ &\quad + \hat{\psi}_{n\downarrow}^+(\vec{r}) \hat{\psi}_{n\uparrow}^+(\vec{r}) \hat{\psi}_{m\uparrow}(\vec{r}) \hat{\psi}_{m\downarrow}(\vec{r})] \end{aligned} \quad (1.3)$$

is the Hamiltonian of interaction, V_{11}, V_{22} are the constants of intraband interaction, V_{12}, V_{21} are the constants of interband

interaction ($V_{nm} = V_{mn}$). The model assumes the formation of Cooper pairs in each band and the tunneling of these pairs as a whole from one band into another. The equation of motion for Heisenberg operators $\hat{\psi}_n(\vec{r}, t), \hat{\psi}_n^+(\vec{r}, t)$ have the form.

The formulas (1.11) are equivalent to

$$\begin{aligned} i \frac{\partial \hat{\psi}_{n\sigma}(\vec{r}, t)}{\partial t} &= \hat{\varepsilon}_n \hat{\psi}_{n\sigma}(\vec{r}, t) - \sum_{m\sigma'} V_{nm} \hat{\psi}_{m\sigma'}^+(\vec{r}, t) \hat{\psi}_{m\sigma'}(\vec{r}, t) \hat{\psi}_{n\sigma}(\vec{r}, t) (1 - \delta_{\sigma\sigma'}), \\ i \frac{\partial \hat{\psi}_{n\sigma}^+(\vec{r}, t)}{\partial t} &= -\hat{\varepsilon}_n \hat{\psi}_{n\sigma'}^+(\vec{r}, t) + \sum_{m\sigma'} V_{nm} \hat{\psi}_{n\sigma}^+(\vec{r}, t) \hat{\psi}_{m\sigma'}^+(\vec{r}, t) \hat{\psi}_{m\sigma'}(\vec{r}, t) (1 - \delta_{\sigma\sigma'}), \end{aligned} \quad (1.4)$$

Inserting $\hat{\varepsilon}_n = \frac{\hat{\vec{p}}^2}{2m_n} - \mu + \zeta_n$ in the (1.4) we obtain

$$\begin{aligned} i \frac{\partial \hat{\psi}_{n\sigma}(\vec{r}, t)}{\partial t} &= \left(\frac{\hat{\vec{p}}^2}{2m_n} - \mu + \zeta_n \right) \hat{\psi}_{n\sigma}(\vec{r}, t) - \sum_{m\sigma'} V_{nm} \hat{\psi}_{m\sigma'}^+(\vec{r}, t) \hat{\psi}_{m\sigma'}(\vec{r}, t) \hat{\psi}_{n\sigma}(\vec{r}, t) (1 - \delta_{\sigma\sigma'}), \\ i \frac{\partial \hat{\psi}_{n\sigma}^+(\vec{r}, t)}{\partial t} &= -\left(\frac{\hat{\vec{p}}^2}{2m_n} - \mu + \zeta_n \right) \hat{\psi}_{n\sigma'}^+(\vec{r}, t) + \sum_{m\sigma'} V_{nm} \hat{\psi}_{n\sigma}^+(\vec{r}, t) \hat{\psi}_{m\sigma'}^+(\vec{r}, t) \hat{\psi}_{m\sigma'}(\vec{r}, t) (1 - \delta_{\sigma\sigma'}) \end{aligned} \quad (1.5)$$

The Green's function is defined by

$$G_{n\sigma, m\sigma'}(\vec{r}_1, t_1; \vec{r}_2, t_2) = -i \left\langle T_t \left(\hat{\psi}_{n\sigma}(\vec{r}_1, t_1) \hat{\psi}_{m\sigma'}^+(\vec{r}_2, t_2) \right) \right\rangle_{N,N} \quad (1.6)$$

where $\langle \dots \rangle_{N,N'}$ denotes the statistical averaging of matrix element between states with N and N' particles.

For systems having no net spin polarization

$$G_{n\sigma,m\sigma'}(\vec{r}_1,t_1;\vec{r}_2,t_2) = \delta_{\sigma\sigma'} G_{nn}(\vec{r}_1,t_1;\vec{r}_2,t_2)$$

Then

$$\begin{aligned} i \frac{\partial}{\partial t_1} G_{n\sigma,m\sigma'}(\vec{r}_1,t_1;\vec{r}_2,t_2) &= -i \left\langle T_t \left(i \frac{\partial}{\partial t_1} \hat{\psi}_{n\sigma}^+(\vec{r}_1,t_1) \hat{\psi}_{m\sigma'}^+(\vec{r}_2,t_2) \right) \right\rangle_{N,N} = \\ &= -i \left\langle \left[T_t \left(\frac{\hat{\vec{p}}_1^2}{2m_n} - \mu + \zeta_n \right) \hat{\psi}_{n\sigma}^+(\vec{r}_1,t_1) - \sum_{l\sigma''} V_{nl} \hat{\psi}_{l\sigma''}^+(\vec{r}_1,t_1) \hat{\psi}_{n\sigma}^-(\vec{r}_1,t_1) (1 - \delta_{\sigma\sigma''}) \right] \hat{\psi}_{m\sigma'}^+(\vec{r}_2,t_2) \right\rangle_{N,N} = \\ &= \left(\frac{\hat{\vec{p}}_1^2}{2m_n} - \mu + \zeta_n \right) (-i) \left\langle T_t \left(\hat{\psi}_{n\sigma}^+(\vec{r}_2,t_2) \hat{\psi}_{m\sigma'}^+(\vec{r}_1,t_1) \right) \right\rangle_{N,N} - \\ &\quad - \sum_{l\sigma''} V_{nl} (-i) \left\langle T_t \left(\hat{\psi}_{l\sigma''}^+(\vec{r}_1,t_1) \hat{\psi}_{n\sigma}^-(\vec{r}_1,t_1) \hat{\psi}_{m\sigma'}^+(\vec{r}_2,t_2) (1 - \delta_{\sigma\sigma''}) \right) \right\rangle_{N,N} - \\ &\quad - \delta(\vec{r}_1 - \vec{r}_2) \delta(t_1 - t_2) \delta_{\sigma\sigma'} = \\ &= \left(\frac{\hat{\vec{p}}_1^2}{2m_n} - \mu + \zeta_n \right) G_{n\sigma,m\sigma'}(\vec{r}_1,t_1;\vec{r}_2,t_2) - i \sum_{l\sigma''} V_{nl} (-i) \left\langle T_t \left(\hat{\psi}_{l\sigma''}^+(\vec{r}_1,t_1) \hat{\psi}_{n\sigma}^-(\vec{r}_1,t_1) \right) \right\rangle_{N,N+2} \times \\ &\quad \times (-i) \left\langle T_t \left(\hat{\psi}_{l\sigma''}^+(\vec{r}_1,t_1) \hat{\psi}_{m\sigma'}^+(\vec{r}_2,t_2) \right) \right\rangle_{N+2,N} (1 - \delta_{\sigma\sigma''}) - \delta(\vec{r}_1 - \vec{r}_2) \delta(t_1 - t_2) \delta_{\sigma\sigma'} = \\ &= \left(\frac{\hat{\vec{p}}_1^2}{2m_n} - \mu + \zeta_n \right) G_{n\sigma,m\sigma'}(\vec{r}_1,t_1;\vec{r}_2,t_2) - i \sum_{l\sigma''} V_{nl} F_{l\sigma'',n\sigma}(\vec{r}_1,t_1;\vec{r}_1,t_1) F_{l\sigma'',m\sigma'}^+(\vec{r}_1,t_1;\vec{r}_2,t_2) (1 - \delta_{\sigma\sigma''}) - \\ &\quad - \delta(\vec{r}_1 - \vec{r}_2) \delta(t_1 - t_2) \delta_{\sigma\sigma'} \end{aligned}$$

$$\text{where } F_{n\sigma,m\sigma'}^+(\vec{r}_1,t_1;\vec{r}_2,t_2) = -i \left\langle T_t \left(\hat{\psi}_{n\sigma}^+(\vec{r}_1,t_1) \hat{\psi}_{m\sigma'}^+(\vec{r}_2,t_2) \right) \right\rangle_{N+2,N}$$

$$\text{and } F_{n\sigma,m\sigma'}(\vec{r}_1,t_1;\vec{r}_2,t_2) = -i \left\langle T_t \left(\hat{\psi}_{n\sigma}^-(\vec{r}_1,t_1) \hat{\psi}_{m\sigma'}^-(\vec{r}_2,t_2) \right) \right\rangle_{N,N+2}$$

are the anomalous Green's functions which satisfy the relations

$$\begin{aligned} F_{n\sigma,m\sigma'}(\vec{r}_1, t_1; \vec{r}_2, t_2) &= g_{\sigma\sigma'} F_{nm}(\vec{r}_1, t_1; \vec{r}_2, t_2) \\ F^+_{n\sigma,m\sigma'}(\vec{r}_1, t_1; \vec{r}_2, t_2) &= g_{\sigma\sigma'} F^+_{nm}(\vec{r}_1, t_1; \vec{r}_2, t_2) \\ g_{\sigma\sigma'} &= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \end{aligned}$$

For the function $G_{nm}(\vec{r}_1, t_1; \vec{r}_2, t_2)$ we obtain the following equation

$$\begin{aligned} &\left[i \frac{\partial}{\partial t_1} - \frac{\hat{\vec{p}}_1^2}{2m_n} + \mu - \zeta_n \right] G_{nm}(\vec{r}_1, t_1; \vec{r}_2, t_2) + i \sum_l V_{nl} F_{ln}(\vec{r}_1, t_1; \vec{r}_1, t_1) F^+_{lm}(\vec{r}_1, t_1; \vec{r}_2, t_2) = \\ &= \delta(\vec{r}_1 - \vec{r}_2) \delta(t_1 - t_2) \end{aligned} \quad (1.7)$$

The function $F^+_{n\sigma,m\sigma'}(\vec{r}_1, t_1; \vec{r}_2, t_2)$ satisfies the equation

$$\begin{aligned} i \frac{\partial}{\partial t_1} F^+_{n\sigma,m\sigma'}(\vec{r}_1, t_1; \vec{r}_2, t_2) &= -i \left\langle T_t \left(i \frac{\partial}{\partial t_1} \hat{\psi}_{n\sigma}^+(\vec{r}_1, t_1) \hat{\psi}_{m\sigma'}^+(\vec{r}_2, t_2) \right) \right\rangle_{N+2,N} = \\ &= -i \left\langle T_t \left(\left[-\left(\frac{\hat{\vec{p}}^2}{2m_n} - \mu + \zeta_n \right) \hat{\psi}_{n\sigma}^+(\vec{r}_1, t_1) + \sum_{l\sigma''} V_{nl} \hat{\psi}_{n\sigma}^+(\vec{r}_1, t_1) \hat{\psi}_{l\sigma''}^+(\vec{r}_1, t_1) \hat{\psi}_{l\sigma''}^-(\vec{r}_1, t_1) (1 - \delta_{\sigma\sigma''}) \right] \hat{\psi}_{m\sigma'}^+(\vec{r}_2, t_2) \right) \right\rangle_{N+2,N} = \\ &= -\left(\frac{\hat{\vec{p}}_1^2}{2m_n} - \mu + \zeta_n \right) \left[-i \left\langle T_t \left(\hat{\psi}_{n\sigma}^+(\vec{r}_1, t_1) \hat{\psi}_{m\sigma'}^+(\vec{r}_2, t_2) \right) \right\rangle_{N+2,N} \right] + \\ &+ \sum_{l\sigma''} V_{nl} \left[-i \left\langle T_t \left(\hat{\psi}_{n\sigma}^+(\vec{r}_1, t_1) \hat{\psi}_{l\sigma''}^+(\vec{r}_1, t_1) \hat{\psi}_{l\sigma''}^-(\vec{r}_1, t_1) \hat{\psi}_{m\sigma'}^+(\vec{r}_2, t_2) \right) \right\rangle_{N+2,N} (1 - \delta_{\sigma\sigma''}) \right] = \\ &= -\left(\frac{\hat{\vec{p}}_1^2}{2m_n} - \mu + \zeta_n \right) F^+_{n\sigma,m\sigma'}(\vec{r}_1, t_1; \vec{r}_2, t_2) + i \sum_{l\sigma''} V_{nl} (-i) \left\langle T_t \left(\hat{\psi}_{n\sigma}^+(\vec{r}_1, t_1) \right) \hat{\psi}_{l\sigma''}^+(\vec{r}_1, t_1) \right\rangle_{N+2,N} \times \\ &\times (-i) \left\langle T_t \left(\hat{\psi}_{l\sigma''}^-(\vec{r}_1, t_1) \hat{\psi}_{m\sigma'}^+(\vec{r}_2, t_2) \right) \right\rangle_{N,N} (1 - \delta_{\sigma\sigma''}) = -\left(\frac{\hat{\vec{p}}_1^2}{2m_n} - \mu + \zeta_n \right) F^+_{n\sigma,m\sigma'}(\vec{r}_1, t_1; \vec{r}_2, t_2) + \\ &+ i \sum_{l\sigma''} V_{nl} F^+_{n\sigma,l\sigma''}(\vec{r}_1, t_1; \vec{r}_1, t_1) G_{l\sigma'',m\sigma'}(\vec{r}_1, t_1; \vec{r}_2, t_2) (1 - \delta_{\sigma\sigma''}) \end{aligned}$$

or

$$\left[i \frac{\partial}{\partial t_1} + \frac{\hat{p}_1^2}{2m_n} - \mu + \zeta_n \right] F_{nm}^+(\vec{r}_1, t_1; \vec{r}_2, t_2) - i \sum_l V_{nl} F_{nl}^+(\vec{r}_1, t_1; \vec{r}_1, t_1) G_{lm}(\vec{r}_1, t_1; \vec{r}_2, t_2) = 0 \quad (1.8)$$

Since the nondiagonal functions are supposed neglected

$$G_{12}, G_{21} \ll G_{11}, G_{22},$$

$$F_{12}, F_{21} \ll F_{11}, F_{22}$$

the system (1.7),(1.8) takes the form

$$\begin{aligned} & \left[i \frac{\partial}{\partial t_1} - \frac{\hat{p}_1^2}{2m_n} + \mu - \zeta_n \right] G_{nn}(\vec{r}_1, t_1; \vec{r}_2, t_2) + i \sum_l V_{nl} F_{ll}(\vec{r}_1, t_1; \vec{r}_1, t_1) F_{nn}^+(\vec{r}_1, t_1; \vec{r}_2, t_2) = \\ & = \delta(\vec{r}_1 - \vec{r}_2) \delta(t_1 - t_2) \end{aligned} \quad (1.9)$$

$$\left[i \frac{\partial}{\partial t_1} + \frac{\hat{p}_1^2}{2m_n} - \mu + \zeta_n \right] F_{nn}^+(\vec{r}_1, t_1; \vec{r}_2, t_2) - i \sum_l V_{nl} F_{ll}^+(\vec{r}_1, t_1; \vec{r}_1, t_1) G_{nn}(\vec{r}_1, t_1; \vec{r}_2, t_2) = 0$$

Let's define the order parameter $\Delta_n(\vec{r}, t)$ as follows

$$\begin{aligned} \Delta_n(\vec{r}, t) &= i \sum_l V_{nl} F_{ll}(\vec{r}, t; \vec{r}, t), \\ \Delta_n^*(\vec{r}, t) &= -i \sum_l V_{nl} F_{ll}^+(\vec{r}, t; \vec{r}, t) \end{aligned} \quad (1.10)$$

Then (1.8), (1.7) may be rewritten in the form

$$\begin{aligned} & \left[i \frac{\partial}{\partial t_1} - \frac{\hat{p}_1^2}{2m_n} + \mu - \zeta_n \right] G_{nn}(\vec{r}_1, t_1; \vec{r}_2, t_2) + \Delta_n(\vec{r}_1, t_1) F_{nn}^+(\vec{r}_1, t_1; \vec{r}_2, t_2) = \\ & = \delta(\vec{r}_1 - \vec{r}_2) \delta(t_1 - t_2) \end{aligned} \quad (1.11)$$

$$\left[i \frac{\partial}{\partial t_1} + \frac{\hat{p}_1^2}{2m_n} - \mu + \zeta_n \right] F_{nn}^+(\vec{r}_1, t_1; \vec{r}_2, t_2) + \Delta_n^*(\vec{r}_1, t_1) G_{nn}(\vec{r}_1, t_1; \vec{r}_2, t_2) = 0 \quad (1.12)$$

The pair of equations (1.11),(1.12) are Gor'kov equations for two-band superconductor.

The equation (1.10) is the self-consistency condition.

2. A time dependent Ginzburg-Landau equation

In the limit case $V_{nm} = 0$ we have $\Delta_n = 0$ and the system (1.11), (1.12) obtains the form

$$\left[i \frac{\partial}{\partial t_1} - \frac{\hat{p}_1^2}{2m_n} + \mu - \zeta_n \right] G^{(0)}_{mn}(\vec{r}_1, t_1; \vec{r}_2, t_2) = \delta(\vec{r}_1 - \vec{r}_2) \delta(t_1 - t_2) \quad (2.1)$$

$$F^{+(0)}_{mn}(\vec{r}_1, t_1; \vec{r}_2, t_2) = 0 \quad (2.2)$$

The system (1.10), (1.11) we may rewrite as integral equations

$$G_{nn}(\vec{r}_1, t_1; \vec{r}_2, t_2) + \iint d\vec{r}_3 dt_3 g^{(0)}_n(\vec{r}_1, t_1; \vec{r}_3, t_3) \Delta_n(\vec{r}_3, t_3) F_{nn}^+(\vec{r}_3, t_3; \vec{r}_2, t_2) = \\ = g^{(0)}_n(\vec{r}_1, t_1; \vec{r}_2, t_2) \quad (2.3)$$

$$F_{nn}^+(\vec{r}_1, t_1; \vec{r}_2, t_2) + \iint d\vec{r}_3 dt_3 \tilde{g}_n^{(0)}(\vec{r}_1, t_1; \vec{r}_3, t_3) \Delta_n^*(\vec{r}_3, t_3) G_{nn}(\vec{r}_3, t_3; \vec{r}_2, t_2) = 0 \quad (2.4)$$

$$\left[i \frac{\partial}{\partial t_1} + \frac{\hat{p}_1^2}{2m_n} - \mu + \zeta_n \right] \tilde{g}_n^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2) = \delta(\vec{r}_1 - \vec{r}_2) \delta(t_1 - t_2) \quad (2.5)$$

$$\left[i \frac{\partial}{\partial t_1} - \frac{\hat{p}_1^2}{2m_n} + \mu - \zeta_n \right] g^{(0)}_n(\vec{r}_1, t_1; \vec{r}_2, t_2) = \delta(\vec{r}_1 - \vec{r}_2) \delta(t_1 - t_2)$$

The functions $g_n^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2)$ and $\tilde{g}_n^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2)$ are functions of the differences $\vec{r}_1 - \vec{r}_2, t_1 - t_2$:

$$\begin{aligned} g_n^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2) &= g_n^{(0)}(\vec{r}_1 - \vec{r}_2, t_1 - t_2), \\ \tilde{g}_n^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2) &= \tilde{g}_n^{(0)}(\vec{r}_1 - \vec{r}_2, t_1 - t_2) \end{aligned}$$

Then

$$\begin{aligned} g_n^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2) &= \iint g_n^{(0)}(\omega, \vec{p}) e^{i\vec{p}(\vec{r}_1 - \vec{r}_2) - i\omega(t_1 - t_2)} \frac{d\omega d^3 p}{(2\pi)^4} \\ [\omega - \varepsilon_n(\vec{p})] g_n^{(0)}(\omega, \vec{p}) &= 1 \end{aligned} \quad (2.6)$$

$$\begin{aligned} \tilde{g}_n^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2) &= \iint \tilde{g}_n^{(0)}(\omega, \vec{p}) e^{i\vec{p}(\vec{r}_1 - \vec{r}_2) - i\omega(t_1 - t_2)} \frac{d\omega d^3 p}{(2\pi)^4}, \\ [\omega + \varepsilon_n(\vec{p})] \tilde{g}_n^{(0)}(\omega, \vec{p}) &= 1, \\ \tilde{g}_n^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2) &= -g_n^{(0)}(\vec{r}_2, t_2; \vec{r}_1, t_1) \end{aligned} \quad (2.7)$$

With (2.7) the equation (2.4) takes the form

$$F_{nm}^+(\vec{r}_1, t_1; \vec{r}_2, t_2) = - \iint d\vec{r}_3 dt_3 g_n^{(0)}(\vec{r}_3, t_3; \vec{r}_1, t_1) \Delta_n^*(\vec{r}_3, t_3) G_{nm}(\vec{r}_3, t_3; \vec{r}_2, t_2) \quad (2.8)$$

To solve the system (2.6),(2.11) iteratively we write Δ_{nl} as $\lambda \Delta_{nl}$ and expand $G_{nm}(\vec{r}_1, t_1; \vec{r}_2, t_2)$, $F_{nm}^+(\vec{r}_1, t_1; \vec{r}_2, t_2)$ in the form

$$G_{nm}(\vec{r}_1, t_1; \vec{r}_2, t_2) = G_{nm}^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2) + \lambda^2 G_{nm}^{(2)}(\vec{r}_1, t_1; \vec{r}_2, t_2) + \dots$$

$$F_{nm}^+(\vec{r}_1, t_1; \vec{r}_2, t_2) = \lambda F_{nm}^{+(1)}(\vec{r}_1, t_1; \vec{r}_2, t_2) + \lambda^3 F_{nm}^{+(3)}(\vec{r}_1, t_1; \vec{r}_2, t_2) \dots$$

Substituting these forms into (1.21),(1.22) and associating equal powers of λ we have

$$F^{+(1)}_{nm}(\vec{r}_1, t_1; \vec{r}_2, t_2) = - \iint d\vec{r}_3 dt_3 g^{(0)}_n(\vec{r}_3, t_3; \vec{r}_1, t_1) \Delta_n^*(\vec{r}_3, t_3) g^{(0)}_n(\vec{r}_3, t_3; \vec{r}_2, t_2) \quad (2.9)$$

$$\begin{aligned} G^{(2)}_{nm}(\vec{r}_1, t_1; \vec{r}_2, t_2) &= - \iint d\vec{r}_3 dt_3 g^{(0)}_n(\vec{r}_1, t_1; \vec{r}_3, t_3) \Delta_n(\vec{r}_3, t_3) F^{+(1)}_{nn}(\vec{r}_3, t_3; \vec{r}_2, t_2) = \\ &= \iint d\vec{r}_3 dt_3 g^{(0)}_n(\vec{r}_1, t_1; \vec{r}_3, t_3) \Delta_n(\vec{r}_3, t_3) \iint d\vec{r}_4 dt_4 g^{(0)}_n(\vec{r}_4, t_4; \vec{r}_3, t_3) \Delta_n^*(\vec{r}_4, t_4) g^{(0)}_n(\vec{r}_4, t_4; \vec{r}_2, t_2) = \quad (2.10) \\ &= \iint d\vec{r}_3 dt_3 \iint d\vec{r}_4 dt_4 g^{(0)}_n(\vec{r}_1, t_1; \vec{r}_3, t_3) \Delta_n(\vec{r}_3, t_3) g^{(0)}_n(\vec{r}_4, t_4; \vec{r}_3, t_3) \Delta_n^*(\vec{r}_4, t_4) g^{(0)}_n(\vec{r}_4, t_4; \vec{r}_2, t_2) \end{aligned}$$

$$\begin{aligned} F^{+(3)}_{mm}(\vec{r}_1, t_1; \vec{r}_2, t_2) &= \iint d\vec{r}_3 dt_3 g^{(0)}_n(\vec{r}_3, t_3; \vec{r}_1, t_1) \Delta_n^*(\vec{r}_3, t_3) G^{(2)}_{mn}(\vec{r}_3, t_3; \vec{r}_2, t_2) = \\ &= \iint d\vec{r}_3 dt_3 \iint d\vec{r}_4 dt_4 \iint d\vec{r}_5 dt_5 g^{(0)}_n(\vec{r}_3, t_3; \vec{r}_1, t_1) \Delta_n^*(\vec{r}_3, t_3) g^{(0)}_n(\vec{r}_3, t_3; \vec{r}_4, t_4) \times \quad (2.11) \\ &\quad \times \Delta_n(\vec{r}_4, t_4) g^{(0)}_n(\vec{r}_5, t_5; \vec{r}_4, t_4) \Delta_n^*(\vec{r}_5, t_5) g^{(0)}_n(\vec{r}_5, t_5; \vec{r}_2, t_2) \end{aligned}$$

Adding up the results (1.10), (2.9), (2.11) we obtain

$$\begin{aligned} \Delta_n^*(\vec{r}, t) &= -i \sum_m V_{nm} F^+_{mm}(\vec{r}, t; \vec{r}, t) = -i \sum_m V_{nm} \left[F^{+(1)}_{mm}(\vec{r}, t; \vec{r}, t) + F^{+(3)}_{mm}(\vec{r}, t; \vec{r}, t) \right] = \\ &= -i \sum_m V_{nm} \iint d\vec{r}_1 dt_1 g^{(0)}_m(\vec{r}_1, t_1; \vec{r}, t) \Delta_m^*(\vec{r}_1, t_1) g^{(0)}_m(\vec{r}_1, t_1; \vec{r}, t) + \\ &\quad + i \sum_m V_{nm} \iint d\vec{r}_1 dt_1 \iint d\vec{r}_2 dt_2 \iint d\vec{r}_3 dt_3 g^{(0)}_m(\vec{r}_1, t_1; \vec{r}, t) \Delta_m^*(\vec{r}_1, t_1) g^{(0)}_m(\vec{r}_1, t_1; \vec{r}_2, t_2) \times \quad (2.12) \\ &\quad \times \Delta_m(\vec{r}_2, t_2) g^{(0)}_m(\vec{r}_3, t_3; \vec{r}_2, t_2) \Delta_m^*(\vec{r}_3, t_3) g^{(0)}_m(\vec{r}_3, t_3; \vec{r}, t) \end{aligned}$$

Let us denote w_{mn} elements of inverse matrix for (V_{nm}) then (2.12) obtains the form

$$\begin{aligned}
\sum_n W_{mn} \Delta_n^*(\vec{r}, t) = & -i \iint d\vec{r}_1 dt_1 g^{(0)}_m(\vec{r}_1, t_1; \vec{r}, t) \Delta_m^*(\vec{r}_1, t_1) g^{(0)}_m(\vec{r}_1, t_1; \vec{r}, t) + \\
& + i \iint d\vec{r}_1 dt_1 \iint d\vec{r}_2 dt_2 \iint d\vec{r}_3 dt_3 g^{(0)}_m(\vec{r}_1, t_1; \vec{r}, t) \Delta_m^*(\vec{r}_1, t_1) g^{(0)}_m(\vec{r}_1, t_1; \vec{r}_2, t_2) \times \\
& \times \Delta_m(\vec{r}_2, t_2) g^{(0)}_m(\vec{r}_3, t_3; \vec{r}_2, t_2) \Delta_m^*(\vec{r}_3, t_3) g^{(0)}_m(\vec{r}_3, t_3; \vec{r}, t)
\end{aligned} \tag{2.13}$$

We now expand $\Delta_m(\vec{r}_1, t_1)$ in the first integral of the r.h.s. of eq. (2.13) as follows

$$\begin{aligned}
\Delta_m(\vec{r}_1, t_1) = & \Delta_m(\vec{r}, t) + \frac{\partial \Delta_m(\vec{r}, t)}{\partial x_\alpha} (x_{1\alpha} - x_\alpha) + \frac{1}{2} \frac{\partial^2 \Delta_m(\vec{r}, t)}{\partial x_\alpha \partial x_\beta} (x_{1\alpha} - x_\alpha)(x_{1\beta} - x_\beta) + \\
& + \frac{\partial \Delta_m(\vec{r}, t)}{\partial t} (t_1 - t)
\end{aligned} \tag{2.14}$$

where $(x_1, x_2, x_3) = \vec{r}$.

In the second integral we take the $\Delta_m(\vec{r}_1, t_1), \Delta_m(\vec{r}_2, t_2), \Delta_m(\vec{r}_3, t_3)$ just equal to $\Delta_m(\vec{r}, t)$.

Using (2.14), we put eq. (2.13) in the following form

$$\sum_n W_{mn} \Delta_n(\vec{r}, t) = A_m \Delta_m(\vec{r}, t) - \Gamma_m \frac{\partial \Delta_m(\vec{r}, t)}{\partial t} + D_m^{\alpha\beta} \frac{\partial^2 \Delta_m(\vec{r}, t)}{\partial x_\alpha \partial x_\beta} - B_m \Delta_m(\vec{r}, t) |\Delta_m(\vec{r}, t)|^2 \tag{2.15}$$

$(m=1, 2)$

Where

$$A_m = -i \iint d\vec{r}_1 dt_1 \left[g^{(0)}_m(\vec{r}_1, t_1; \vec{r}, t) \right]^2 \tag{2.16}$$

$$\Gamma_m = -i \iint d\vec{r}_1 dt_1 (t_1 - t) \left[g^{(0)}_m(\vec{r}_1, t_1; \vec{r}, t) \right]^2 \tag{2.17}$$

$$D_m^{\alpha\beta} = -\frac{i}{2} \iint d\vec{r}_1 dt_1 (x_{1\alpha} - x_\alpha)(x_{1\beta} - x_\beta) \left[g^{(0)}_m(\vec{r}_1, t_1; \vec{r}, t) \right]^2 \tag{2.18}$$

$$B_m = -i \iint d\vec{r}_1 dt_1 \iint d\vec{r}_2 dt_2 \iint d\vec{r}_3 dt_3 g^{(0)}_m(\vec{r}_1, t_1; \vec{r}, t) g^{(0)}_m(\vec{r}_1, t_1; \vec{r}_2, t_2) \times \\ \times g^{(0)}_m(\vec{r}_3, t_3; \vec{r}_2, t_2) g^{(0)}_m(\vec{r}_3, t_3; \vec{r}, t) \quad (2.19)$$

The eq.(2.15) is the Ginzburg-Landau equation for two-band superconductor

$$\Gamma_m \frac{\partial \Delta_m(\vec{r}, t)}{\partial t} = \sum_{n=1,2} [A_m \delta_{nm} - W_{mn}] \Delta_n(\vec{r}, t) + D_m^{\alpha\beta} \frac{\partial^2 \Delta_m(\vec{r}, t)}{\partial x_\alpha \partial x_\beta} - B_m \Delta_m(\vec{r}, t) |\Delta_m(\vec{r}, t)|^2 \quad (2.20)$$

(m=1,2)

or

$$\Gamma_1 \frac{\partial \Delta_1(\vec{r}, t)}{\partial t} = (A_1 - W_{11}) \Delta_1(\vec{r}, t) - W_{12} \Delta_2(\vec{r}, t) + D_1^{\alpha\beta} \frac{\partial^2 \Delta_1(\vec{r}, t)}{\partial x_\alpha \partial x_\beta} - B_1 \Delta_1(\vec{r}, t) |\Delta_1(\vec{r}, t)|^2 \quad (2.21)$$

$$\Gamma_2 \frac{\partial \Delta_2(\vec{r}, t)}{\partial t} = -W_{21} \Delta_1(\vec{r}, t) + (A_2 - W_{22}) \Delta_2(\vec{r}, t) + D_2^{\alpha\beta} \frac{\partial^2 \Delta_2(\vec{r}, t)}{\partial x_\alpha \partial x_\beta} - B_2 \Delta_2(\vec{r}, t) |\Delta_2(\vec{r}, t)|^2 \quad (2.22)$$

Equations (2.21), (2.22) can be rewritten as

$$\Gamma_1 \frac{\partial \Delta_1}{\partial t} = -\frac{\delta F}{\delta \Delta_1^*},$$

$$\Gamma_2 \frac{\partial \Delta_2}{\partial t} = -\frac{\delta F}{\delta \Delta_2^*}$$

where

$$F = \int d\vec{r} \left[\frac{1}{2} (W_{11} - A_1) |\Delta_1|^2 + \frac{1}{2} (W_{22} - A_2) |\Delta_2|^2 + W_{12} \Delta_1^* \Delta_2 + W_{21} \Delta_2^* \Delta_1 + \right. \\ \left. + \frac{1}{2} D_1^{\alpha\beta} \frac{\partial \Delta_1}{\partial x_\alpha} \frac{\partial \Delta_1^*}{\partial x_\beta} + \frac{1}{2} D_2^{\alpha\beta} \frac{\partial \Delta_2}{\partial x_\alpha} \frac{\partial \Delta_2^*}{\partial x_\beta} + \frac{1}{4} B_1 |\Delta_1|^4 + \frac{1}{4} B_2 |\Delta_2|^4 \right]$$

is the GL functional.

3. The coefficients of the model.

At present section we derive the formulas for evaluation of the coefficients $A_m, \Gamma_m, D_m^{\alpha\beta}, B_m$. Since the functions $g_n^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2)$ are functions of the differences $\vec{r}_1 - \vec{r}_2, t_1 - t_2$:

$$g_n^{(0)}(\vec{r}_1, t_1; \vec{r}_2, t_2) = g_n^{(0)}(\vec{r}_1 - \vec{r}_2, t_1 - t_2),$$

We obtain

$$A_m = -i \iint d\vec{r}_1 dt_1 \left[g_m^{(0)}(\vec{r}_1, t_1; \vec{r}, t) \right]^2 = -i \iint d\vec{r}_1 dt_1 \left[g_m^{(0)}(\vec{r}_1 - \vec{r}, t_1 - t) \right]^2 = \\ = -i \int d\vec{r}_1 \int_{-\infty}^t dt_1 \left[g_m^{(0)}(\vec{r}_1 - \vec{r}, t_1 - t) \right]^2 - i \int d\vec{r}_1 \int_t^\infty dt_1 \left[g_m^{(0)}(\vec{r}_1 - \vec{r}, t_1 - t) \right]^2 = \\ = -i \int d\vec{r}_1 \int_{-\infty}^0 d\tau \left[g_m^<(\vec{r}_1 - \vec{r}, \tau) \right]^2 - i \int d\vec{r}_1 \int_0^\infty d\tau \left[g_m^>(\vec{r}_1 - \vec{r}, \tau) \right]^2 = \\ = -i \int d\vec{r} \int_{-\infty}^0 d\tau \left[g_m^<(\vec{r}, \tau) \right]^2 - i \int d\vec{r} \int_0^\infty d\tau \left[g_m^>(\vec{r}, \tau) \right]^2 \quad (3.1)$$

where $g_m^<(\vec{r}, t)$, $g_m^>(\vec{r}, t)$ are the correlation functions which Fourier transform is

$$g_m^>(\vec{p}, t) = -i \frac{e^{-i\varepsilon_m(\vec{p})t}}{e^{-\beta\varepsilon_m(\vec{p})} + 1}, \\ g_m^<(\vec{p}, t) = i \frac{e^{-i\varepsilon_m(\vec{p})t}}{e^{\beta\varepsilon_m(\vec{p})} + 1}, \quad (3.2)$$

$$\beta = \frac{1}{k_B T}$$

and

$$\begin{cases} f(\vec{r}) = \int f(\vec{p}) e^{i\vec{p}\cdot\vec{r}} \frac{d\vec{p}}{(2\pi)^3} \\ f(\vec{p}) = \int f(\vec{r}) e^{-i\vec{p}\cdot\vec{r}} d\vec{r} \end{cases}$$

Then

$$\begin{aligned} A_m &= \frac{1}{2(2\pi)^3} \int \frac{\tanh(\frac{1}{2}\beta\varepsilon_m(\vec{p}))}{\varepsilon_m(\vec{p})} d\vec{p} = \\ &= \frac{\hbar^3}{2} \int_0^{\hbar\omega_D} \frac{\tanh(\frac{1}{2}\beta\varepsilon_m)}{\varepsilon_m} \cdot N_m(\varepsilon) d\varepsilon_m = \\ &= \frac{\hbar^3 N_m(0)}{2} \int_0^{\hbar\omega_D} \frac{d\varepsilon}{\varepsilon} \tanh(\frac{1}{2}\beta\varepsilon) = \frac{\hbar^3 N_m(0)}{2} \int_0^{\frac{\beta\hbar\omega_D}{2}} \frac{\tanh x}{x} dx \end{aligned} \quad (3.3)$$

where $N_m(0)$ is the density of states at the Fermi surface of m -tm band, ω_D is the Debye frequency. For most superconductors $\frac{\hbar\omega_D}{2k_B T_c} \gg 1$, where T_c is the critical temperature.

The integral (3.3) can be evaluated using

$$\int_0^a \frac{\tanh x}{x} dx = \tanh a \cdot \ln a - \int_0^a \frac{\ln x}{\cosh^2 x} dx$$

For $a \gg 1$ we obtain

$$\int_0^a \frac{\tanh x}{x} dx \approx \ln a - \int_0^\infty \frac{\ln x}{\cosh^2 x} dx = \ln a - \ln\left(\frac{\pi}{4e^C}\right) = \ln\left(\frac{4e^C a}{\pi}\right)$$

with $C \approx 0.577$ being the Euler constant[43].

Therefore

$$A_m(T) = \frac{N_m(0)\hbar^3}{4\pi} \ln\left(\frac{2e^c\hbar\omega_D}{\pi k_B T}\right) \quad (3.4)$$

The critical temperature of a two-band superconductor is determined by the condition[15,17,18,19,20,21,27,33]

$$\det\begin{pmatrix} A_1(T_c) - W_{11} & -W_{12} \\ -W_{21} & A_2(T_c) - W_{22} \end{pmatrix} = 0 \quad (3.5)$$

The damping constants are

$$\begin{aligned} \Gamma_m &= -i \iint d\vec{r}_1 dt_1 (t_1 - t) [g_m^{(0)}(\vec{r}_1, t_1; \vec{r}, t)]^2 = \\ &= -i \iint d\vec{r}_1 dt_1 (t_1 - t) [g_m^{(0)}(\vec{r}_1 - \vec{r}, t_1 - t)]^2 = \\ &= -i \int d\vec{r}_1 \int_{-\infty}^t dt_1 (t_1 - t) [g_m^{(0)}(\vec{r}_1 - \vec{r}, t_1 - t)]^2 - i \int d\vec{r}_1 \int_t^\infty dt_1 (t_1 - t) [g_m^{(0)}(\vec{r}_1 - \vec{r}, t_1 - t)]^2 = \\ &= -i \int d\vec{r}_1 \int_{-\infty}^0 d\tau \tau [g_m^<(\vec{r}_1 - \vec{r}, \tau)]^2 - i \int d\vec{r}_1 \int_0^\infty d\tau \tau [g_m^>(\vec{r}_1 - \vec{r}, \tau)]^2 = \\ &= -i \int d\vec{r} \int_{-\infty}^0 d\tau \tau [g_m^<(\vec{r}, \tau)]^2 - i \int d\vec{r} \int_0^\infty d\tau \tau [g_m^>(\vec{r}, \tau)]^2 = \\ &= -i \int \frac{d\vec{p}}{(2\pi)^3} \int_{-\infty}^0 d\tau \tau [g_m^<(\vec{p}, \tau)]^2 - i \int \frac{d\vec{p}}{(2\pi)^3} \int_0^\infty d\tau \tau [g_m^>(\vec{p}, \tau)]^2 \end{aligned}$$

Using (3.2) we obtain

$$\begin{aligned} \Gamma_m &= -\frac{i}{4(2\pi)^3} \int \frac{\tanh\left(\frac{1}{2}\beta\varepsilon_m(\vec{p})\right)}{(\varepsilon_m(\vec{p}) + i0)^2} d\vec{p} = -\frac{i\hbar^3}{4} \int N_m(\varepsilon) \frac{\tanh\left(\frac{1}{2}\beta\varepsilon_m\right)}{(\varepsilon_m + i0)^2} d\varepsilon_m = \\ &= -\frac{i\hbar^3\beta N_m(0)}{8} \int \frac{\tanh x}{(x + i0)^2} dx = \frac{\pi\hbar^3 N_m(0)}{16k_B T_c} \end{aligned} \quad (3.6)$$

The diffusion constants are

$$D_n^{\alpha\beta} = -\frac{i}{2} \iint d\vec{r} dt x_\alpha x_\beta \left[g_n^{(0)}(\vec{r}, t) \right]^2 = -\frac{\beta}{4(2\pi)^3} \int d\vec{p} \frac{\partial^2}{\partial p_\alpha \partial p_\beta} \left[\frac{\tanh(\beta\epsilon_n(\vec{p}))}{\beta\epsilon_n(\vec{p})} \right]$$

The integral can be evaluated using

$$\frac{\tanh x}{x} = \sum_{n=-\infty}^{\infty} \frac{1}{x^2 + \left[\pi \left(n + \frac{1}{2} \right) \right]^2}$$

After the substitution we find

$$\begin{aligned} D_n^{\alpha\beta} &= -\frac{\beta}{4(2\pi)^3} \sum_{n=-\infty}^{\infty} \int d\vec{p} \frac{\partial^2}{\partial p_\alpha \partial p_\beta} \left[\frac{1}{\beta^2 \epsilon_n^2(\vec{p}) + \pi^2 \left(n + \frac{1}{2} \right)^2} \right] \approx \\ &\approx \frac{\beta^3}{2(2\pi)^3} \sum_{n=-\infty}^{\infty} \int d\vec{p} \frac{\frac{\partial \epsilon_n(\vec{p})}{\partial p_\alpha} \cdot \frac{\partial \epsilon_n(\vec{p})}{\partial p_\beta}}{\left[\beta^2 \epsilon_n^2(\vec{p}) + \pi^2 \left(n + \frac{1}{2} \right)^2 \right]^2} \approx \frac{\beta^3}{2(2\pi)^3} \sum_{n=-\infty}^{\infty} \int d\vec{p} \frac{v_{F\alpha} \cdot v_{F\beta}}{\left[\beta^2 \epsilon_n^2(\vec{p}) + \pi^2 \left(n + \frac{1}{2} \right)^2 \right]^2} = \\ &= \frac{\beta^3 v_F^2 \delta_{\alpha\beta}}{6(2\pi)^3} \sum_{n=-\infty}^{\infty} \int d\vec{p} \frac{1}{\left[\beta^2 \epsilon_n^2(\vec{p}) + \pi^2 \left(n + \frac{1}{2} \right)^2 \right]^2} \end{aligned}$$

where $\vec{v}_F = \frac{\vec{p}_F}{m_n}$ is the Fermi velocity. Inserting $d\vec{p} = (2\pi\hbar)^3 N_n(\epsilon) d\epsilon$ [44] we obtain

$$\begin{aligned}
D_n^{\alpha\beta} &= \frac{\beta^3 v_F^2 \delta_{\alpha\beta} N_n(0) \hbar^3}{6} \sum_{n=-\infty}^{\infty} \int_0^{\infty} \frac{d\varepsilon}{\left[\beta^2 \varepsilon^2 + \pi^2 \left(n + \frac{1}{2} \right)^2 \right]^2} = \\
&= \frac{\beta^2 v_F^2 \delta_{\alpha\beta} N_n(0) \hbar^3}{6} \sum_{n=-\infty}^{\infty} \int_0^{\infty} \frac{dx}{\left[x^2 + \pi^2 \left(n + \frac{1}{2} \right)^2 \right]^2} = \frac{\beta^2 v_F^2 \delta_{\alpha\beta} N_n(0) \hbar^3}{6} \sum_{n=-\infty}^{\infty} \frac{\pi}{4 \left(n + \frac{1}{2} \right)^3} = \\
&= \frac{7\pi\beta^2 v_F^2 \delta_{\alpha\beta} N_n(0) \hbar^3}{24} \zeta(3) \equiv D_n \delta_{\alpha\beta}
\end{aligned}$$

Here we use

$$\sum_{n=0}^{\infty} \frac{1}{\left(n + \frac{1}{2} \right)^z} = (2^z - 1) \zeta(z)$$

Where $\sum_{n=1}^{\infty} \frac{1}{n^z} = \zeta(z)$ is the Riemann ζ function.

The Green's function satisfies the anti-periodic boundary condition

$g_m^{(0)}(\vec{r}_1, t_1; \vec{r}, t) \Big|_{t_1=0} = -g_m^{(0)}(\vec{r}_1, t_1; \vec{r}, t) \Big|_{t_1=-i\beta}$ we express $g_m^{(0)}$ as Fourier series [45]

$$\begin{aligned}
g_m^{(0)}(\vec{p}, t-t') &= \frac{i}{\beta} \sum_{\nu} \frac{e^{-iz_{\nu}(t-t')}}{z_{\nu} - \varepsilon_m(\vec{p})}, \\
z_{\nu} &= \frac{\pi i \nu}{\beta}
\end{aligned} \tag{3.7}$$

where the sum is taken to run over all odd integers $(\nu = 2n-1, n \in \mathbb{Z})$. If we then substitute (3.7) into the (2.19) we obtain

$$\begin{aligned}
B_m &= \frac{1}{\pi\beta} \sum_{\nu} \int d\vec{p} \frac{1}{\left[\varepsilon_m^2(\vec{p}) + \frac{\pi^2 \nu^2}{\beta^2} \right]^2} = \frac{8\pi^2 N_m(0) \hbar^3}{\beta} \sum_{\nu} \int_0^{\infty} \frac{d\varepsilon}{\left(\varepsilon^2 + \frac{\pi^2 \nu^2}{\beta^2} \right)^2} = \\
&= 2N_m(0) \hbar^3 \beta^2 \sum_{n \in \mathbb{Z}} \frac{1}{(2n-1)^3} = \frac{7N_m(0) \hbar^3 \zeta(3)}{4(k_B T)^2}
\end{aligned}$$

4. The linearized model.

The equations (2.21), (2.22) are nonlinear differential equations of the diffusion type to describe the space-time variation of the order parameters near the critical temperature. The self consistent superconducting interaction drives the order parameters towards its equilibrium values $\bar{\Delta}_1, \bar{\Delta}_2$ which are the static uniform solution

$$0 = (A_1 - W_{11})\bar{\Delta}_1 - W_{12}\bar{\Delta}_2 - B_1\bar{\Delta}_1^3$$

$$0 = -W_{21}\bar{\Delta}_1 + (A_2 - W_{22})\bar{\Delta}_2 - B_2\bar{\Delta}_2^3$$

For $T > T_c$ we have $\bar{\Delta}_1 = \bar{\Delta}_2 = 0$ (symmetrical phase) while for $T < T_c$ we have $\bar{\Delta}_1, \bar{\Delta}_2 \propto \sqrt{\tau}$ (unsymmetrical phase) where $\tau = 1 - \frac{T}{T_c}$. In this section we linearize our model about equilibrium: $\Delta_n(\vec{r}, t) = \bar{\Delta}_n + \eta_n(\vec{r}, t)$ where $\eta_n(\vec{r}, t)$ is the deviation of the order parameters from its real equilibrium value. In the linear approximation $\eta_n(\vec{r}, t)$ satisfy the set of equations

$$\begin{aligned} \Gamma_1 \frac{\partial \eta_1}{\partial t} &= (A_1 - W_{11} - 3B_1\bar{\Delta}_1^2)\eta_1 - W_{12}\eta_2 + D_1\Pi^2\eta_1 \\ \Gamma_2 \frac{\partial \eta_2}{\partial t} &= -W_{21}\eta_1 + (A_2 - W_{22} - 3B_2\bar{\Delta}_2^2)\eta_2 + D_2\Pi^2\eta_2 \end{aligned} \quad (4.1)$$

where Π^2 is the Laplace operator. In a spatially homogeneous system the eq.(4.1) takes the form

$$\frac{d\vec{\eta}(t)}{dt} = -L(\tau)\vec{\eta}(t) \quad (4.2)$$

where $\vec{\eta}(t) = \begin{pmatrix} \eta_1(t) \\ \eta_2(t) \end{pmatrix}$, $L(\tau) = \begin{pmatrix} \frac{3B_1\bar{\Delta}_1^2 + W_{11} - A_1}{\Gamma_1} & \frac{W_{12}}{\Gamma_1} \\ \frac{W_{21}}{\Gamma_2} & \frac{3B_2\bar{\Delta}_2^2 + W_{22} - A_2}{\Gamma_2} \end{pmatrix}$. The general

solution of eq. (4.2) is

$$\vec{\eta}(t) = C_1 \vec{u}^{(1)} e^{\lambda_1(\tau)t} + C_2 \vec{u}^{(2)} e^{\lambda_2(\tau)t} \quad (4.3)$$

where C_1, C_2 are arbitrary constants, $\lambda_1(\tau), \lambda_2(\tau)$ are eigenvalues of $L(\tau)$ and $\vec{u}^{(1)}, \vec{u}^{(2)}$ are the corresponding eigenvectors. The system has two relaxation times : $\vartheta_1 = -\frac{1}{\lambda_1}, \vartheta_2 = -\frac{1}{\lambda_2}, (\vartheta_1 > \vartheta_2)$. The eq. (3.5) leads to

$$\lim_{\tau \rightarrow 0} \lambda_1(\tau) = 0$$

$$\lim_{\tau \rightarrow 0} \lambda_2(\tau) = -\mu < 0$$

Thus, for $t \gg \vartheta_2 = \mu^{-1}$ the second term in (4.3) is negligible. We have

$$\vec{\eta}(t) \underset{t \gg \mu^{-1}}{\cong} C_1 \vec{u}_1 e^{\lambda_1(\tau)t} \quad (4.4)$$

where $\vec{u}^{(1)} = \begin{pmatrix} u_1^{(1)} \\ u_2^{(1)} \end{pmatrix}$ is a constant vector. It is readily checked that

$$\frac{\eta_1(t)}{\eta_2(t)} \underset{t \gg \mu^{-1}}{=} \frac{u_1^{(1)}}{u_2^{(1)}} = \text{const}$$

So the ratio $\frac{\eta_1(t)}{\eta_2(t)}$ comes out to be time independent. Order parameters of a two-band superconductor are strongly correlated in the limit $t \rightarrow \infty$. Both order parameters vary on the same time scale. This conclusion is in agreement with [33]. The motion of pairs in one band causes a change in another one. The first subsystem follows adiabatically the second. We shall call this regime the “adiabatic regime”.

If $\lambda_1(\tau)$ is expanded in powers of τ , within the GL accuracy we can put $\lambda_1(\tau) = \text{const} \cdot \tau$. The largest relaxation time of the system is

$$\vartheta_1 = -\frac{1}{\lambda_1(\tau)} \propto \frac{1}{\tau}. \text{ Thus, in this approximation } \frac{\partial}{\partial t} \propto \tau, \text{ and } \frac{\partial \Delta_1}{\partial t} \propto \tau^{\frac{3}{2}}, \frac{\partial \Delta_2}{\partial t} \propto \tau^{\frac{3}{2}}.$$

5. The nonlinear model in adiabatic regime.

Our goal is to obtain the set of decoupled equations for Δ_1, Δ_2 . We now recall [33] that in the one-band GL equation

$$\gamma \frac{\partial \Delta}{\partial t} = a\Delta - b\Delta |\Delta|^2 + D\Pi^2\Delta$$

all terms are of the same order with respect to

$$\tau : a \propto \tau^1, b \propto \tau^0, \Delta \propto \tau^{1/2}, \frac{\partial}{\partial t} \propto \tau^1, \Pi^2 \propto \xi^{-2} \propto \tau \quad \text{where } \xi \text{ is the coherence length.}$$

Following [33] we express Δ_1 from eq.(2.21) and substitute the result in eq. (2.22), keeping only terms up to order $\tau^{3/2}$:

$$\gamma \frac{\partial \Delta_1}{\partial t} = a\Delta_1 - b_1\Delta_1 |\Delta_1|^2 + K\Pi^2\Delta_1 \quad (5.1)$$

with

$$\begin{aligned} \gamma &= \Gamma_1(A_2 - W_{22}) + \Gamma_2(A_1 - W_{11}), \quad a = (A_1 - W_{11})(A_2 - W_{22}) - W_{12}W_{21}, \\ K &= D_1(A_2 - W_{22}) + D_2(A_1 - W_{11}), \quad b_1 = B_1(A_2 - W_{22}) + \frac{B_2(A_1 - W_{11})^3}{W_{12}^2} \end{aligned}$$

The term with $\frac{\partial^2 \Delta_1}{\partial t^2}$ is neglected.

Similarly, one obtains an equation for Δ_2 :

$$\gamma \frac{\partial \Delta_2}{\partial t} = a\Delta_2 - b_2\Delta_2 |\Delta_2|^2 + K\Pi^2\Delta_2 \quad (5.2)$$

with $b_2 = B_2(A_1 - W_{11}) + \frac{B_1(A_2 - W_{22})^3}{W_{21}^2}$. Within the GL accuracy $a = k\tau, k > 0$.

We obtain upon dividing by a of both equations (5.1),(5.2) :

$$\begin{aligned} g \frac{\partial \Delta_1}{\partial t} &= \Delta_1 \left(1 - \frac{|\Delta_1|^2}{|\bar{\Delta}_1|^2} \right) + \xi^2 \Pi^2 \Delta_1, \\ g \frac{\partial \Delta_2}{\partial t} &= \Delta_2 \left(1 - \frac{|\Delta_2|^2}{|\bar{\Delta}_2|^2} \right) + \xi^2 \Pi^2 \Delta_2 \end{aligned} \quad (5.3)$$

where $\vartheta = \frac{\gamma}{a} = \frac{\gamma}{k\tau}$ is the largest relaxation time of the system,

$$\xi = \sqrt{\frac{K}{a}} = \sqrt{\frac{K}{k\tau}}, \bar{\Delta}_1 = \sqrt{\frac{a}{b_1}}, \bar{\Delta}_2 = \sqrt{\frac{a}{b_2}}.$$

The ratio $\frac{\bar{\Delta}_1}{\bar{\Delta}_2} = \frac{\Delta_1}{\Delta_2} = \Psi$ satisfies the equation

$$\vartheta \frac{\partial \Psi}{\partial t} = \Psi(1 - |\Psi|^2) + \xi^2 \Pi^2 \Psi \quad (5.4)$$

The two TDGL equations for the two-band case are reduced to a single equation for the normalized order parameter $\Psi(\vec{r}, t)$.

Acknowledgments

We are indebted to J.Berger, V. Zhuravlev for useful discussions and help.

References

- [1] A.Schmid, Phys. Kondens.Mater. **5**, 302 (1966).
- [2] E.Abrahms, T.Tsuneto, Phys.Rev. **152**, 416 (1966).
- [3] C.Caroli, K.Maki, Phys.Rev. **159**, 306 (1967), Phys.Rev. **159**, 316 (1967).
- [4] L.P.Gor'kov, G.M.Eliashberg, Sov.Phys.,JETP **27**,328 (1968).
- [5] J.W.F.Woo, E.Abrahms, Phys.Rev. **169**, 407 (1968).
- [6] G.M.Eliashberg, Sov.Phys.,JETP **29**,1298 (1969).

- [7] A.Houghton, K.Maki, Phys.Rev. B **3**, 1625 (1971).
- [8] C.R.Hu, R.S.Thompson, Phys.Rev. B **6**, 110 (1972).
- [9] M.Cyrot, Rep.Prog.Phys. **36**, 103 (1973).
- [10] L.Kramer, R.J.Watts-Tobin, Phys.Rev.Lett. **40**, 1041 (1978).
- [11] A.Levchenko, A.Kamenev, arXiv:1101.4599v1.
- [12] X X Xi, Rep.Prog.Phys. **71**, 1 (2008).
- [13] V.A.Moskalenko, Fiz.Met.Metalloved, **8**,503 (1959).
- [14] H. Suhl, B.T. Matthias, L.R. Walker, Phys. Rev. Lett. **3**, 552 (1959).
- [15] A.E.Koshelev, A.A.Golubov, arXiv:cond-mat/0211388.
- [16] A.Gurevich, V.M.Vinokur, arXiv: cond-mat/0207559.
- [17] A.Gurevich, arXiv: cond-mat/0212129.
- [18] A.A.Golubov, A.E.Koshelev, arXiv:cond-mat/0303237.
- [19] A.E.Koshelev, A.A.Golubov, arXiv:cond-mat/0401339.
- [20] M.E.Zhitomirsky, V.-H.Dao, Phys.Rev. B **69**, 054508 (2004).
- [21] A.E.Koshelev,A.A.Varlamov, V.M.Vinokur, Phys.Rev. B **72**, 064523 (2005).
- [22] A.Gurevich, arXiv: cond-mat/0701281.
- [23] K.Tanaka, M.Eschrig,D.F.Agterberg, arXiv: cond-mat/0702270v2.

- [24] A.Levchenko, A.Kamenev, arXiv: 0706.2884.
- [25] K.Tanaka,M.Eschrig, arXiv:0809.3989.
- [26] O.V.Dolgov, A.A.Golubov, arXiv:0803.1103.
- [27] O.V.Dolgov,I.I.Mazin,D.Parker, A.A.Golubov, arXiv:0810.1476.
- [28] V.G.Kogan, C.Martin, R.Prozorov, arXiv: 0905.0029.
- [29] J.Carlstrom, E.Babaev, M.Speight, arXiv: 1009.2196.
- [30] E.Babaev, J.Carlstrom, arXiv: 1007.1965.
- [31] E.Babaev, J.Carlstrom, M.Speight, arXiv: 0910.1607.
- [32] M.Silaev, E.Babaev, arXiv:1102.5734.
- [33] V.G.Kogan, J.Schmalian, Phys.Rev. B **83**, 054515 (2011).
- [34] J.Carlstrom,J.Garaud, E.Babaev, arXiv:1101.4599.
- [35] M.E.Palistrant, arXiv:cond-mat/0305496.
- [36] V.A.Moskalenko, M.E.Palistrant, V.M.Vakalyuk , arXiv:cond-mat/0309671.
- [37] M.E.Palistrant, arXiv:cond-mat/0312302.
- [38] M.E.Palistrant, V.A.Ursu, arXiv:cond-mat/0602551.
- [39] M.E.Palistrant, V.A.Ursu, arXiv:cond-mat/0611275.

- [40] M.E.Palistrant,I.D.Cebotari, V.A.Ursu, arXiv:0811.0897.
- [41] M.E.Palistrant, arXiv:1008.4833.
- [42] Shi-Zeng Lin,Xiao Hu, arXiv:1007.1940.
- [43] I. S. Gradshteyn , I. M. Ryzhik, Table of Integrals, Series, and Products, Academic Press, New York (1980).
- [44] N.Kopnin, Theory of nonequilibrium superconductivity, Oxford University Press , Oxford (2001).
- [45] L.P.Kadanoff, G.Baym, Quantum Statistical Mechanics, W.A.Benjamin, New York (1962).