

# Clausius inequality and optimality of quasi static transformations for nonequilibrium stationary states

L. Bertini,<sup>1</sup> D. Gabrielli,<sup>2</sup> G. Jona-Lasinio,<sup>3</sup> and C. Landim<sup>4</sup>

<sup>1</sup>*Dipartimento di Matematica, Università di Roma La Sapienza, Piazza A. Moro 2, 00185 Roma, Italy*

<sup>2</sup>*Dipartimento di Matematica, Università dell'Aquila, 67100 Coppito, L'Aquila, Italy*

<sup>3</sup>*Dipartimento di Fisica and INFN, Università di Roma La Sapienza, Piazza A. Moro 2, 00185 Roma, Italy*

<sup>4</sup>*IMPA, Estrada Dona Castorina 110, J. Botânico, 22460 Rio de Janeiro, Brazil,  
and CNRS UMR 6085, Université de Rouen, F76801 Saint-Étienne-du-Rouvray, France*

Nonequilibrium stationary states of thermodynamic systems dissipate a positive amount of energy per unit of time. If we consider transformations of such states that are realized by letting the driving depend on time, the amount of energy dissipated in an unbounded time window becomes then infinite. Following the general proposal by Oono and Paniconi and using results of the macroscopic fluctuation theory, we give a natural definition of a renormalized work performed along any given transformation. We then show that the renormalized work satisfies a Clausius inequality and prove that equality is achieved for very slow transformations, that is in the quasi static limit. We finally connect the renormalized work to the quasi potential of the macroscopic fluctuation theory, that gives the probability of fluctuations in the stationary nonequilibrium ensemble.

PACS numbers: 05.70.Ln, 05.20.-y, 05.40.-a, 05.60.-k

A main goal of nonequilibrium thermodynamics is to construct analogues of thermodynamic potentials for nonequilibrium stationary states. These potentials should describe the typical macroscopic behavior of the system as well as the asymptotic probability of fluctuations. As it has been shown in [1], this program can be implemented without the explicit knowledge of the stationary ensemble and requires as input the macroscopic dynamical behavior of systems which can be characterized by the transport coefficients. This theory, now known as *macroscopic fluctuation theory*, is based on an extension of Einstein equilibrium fluctuation theory to stationary nonequilibrium states combined with a dynamical point of view. It has been very powerful in studying concrete microscopic models but can be used also as a phenomenological theory. It has led to several new interesting predictions [2–7].

From a thermodynamic viewpoint, the analysis of transformations from one state to another one is most relevant. This issue has been addressed by several authors in different contexts. For instance, following the basic papers [8–10], the case of Hamiltonian systems with finitely many degrees of freedom has been recently discussed in [11, 12] while the case of Langevin dynamics is considered in [13].

We here consider thermodynamic transformations for driven diffusive systems in the framework of the macroscopic fluctuation theory. With respect to the authors mentioned above, the main difference is that we deal with systems with infinitely many degrees of freedom and the spatial structure becomes relevant. For simplicity of notation, we restrict to the case of a single conservation law, e.g., the conservation of the mass. We thus consider an open system in contact with boundary reservoirs, characterized by their chemical potential  $\lambda$ , and under the

action of an external field  $E$ . We denote by  $\Lambda \subset \mathbb{R}^d$  the bounded region occupied by the system, by  $x$  the macroscopic space coordinates and by  $t$  the macroscopic time. With respect to our previous work [1, 3, 4], we here consider the case in which  $\lambda$  and  $E$  depend explicitly on the time  $t$ , driving the system from a nonequilibrium state to another one. The macroscopic dynamics is given by the hydrodynamic equation for the density which satisfies the following general assumptions based on the notion of local equilibrium. For stochastic lattice gases these assumptions can be proven rigorously and the macroscopic transport coefficient can be characterized in terms of the underlying microscopic dynamics [14].

The macroscopic state is completely described by the local density  $u(t, x)$  and the associated current  $j(t, x)$ . In the sequel we drop the dependence on the space coordinate  $x$  from the notation. The macroscopic evolution is given by the continuity equation

$$\partial_t u(t) + \nabla \cdot j(t) = 0, \quad (1)$$

together with the constitutive equation  $j(t) = J(t, u(t))$  expressing the local current in function of the local density. For driven diffusive systems the constitutive equation takes the form

$$J(t, \rho) = -D(\rho) \nabla \rho + \chi(\rho) E(t) \quad (2)$$

where the *diffusion coefficient*  $D(\rho)$  and the *mobility*  $\chi(\rho)$  are positive matrices. In the case of time independent driving the right hand side does not depend explicitly on time and we denote the current simply by  $J(\rho)$ . The transport coefficients in (2) satisfy the local Einstein relation  $D(\rho) = \chi(\rho) f''(\rho)$ , where  $f$  is the equilibrium free energy per unit of volume. The interaction with the external reservoirs specify the boundary conditions for the evolution defined by (1)–(2). Recalling that  $\lambda(t)$  is the

chemical potential of the reservoirs, this boundary condition reads  $f'(u(t, x)) = \lambda(t, x)$ ,  $x \in \partial\Lambda$ . We shall also assume that when  $\lambda$  and  $E$  do not depend on time there is a unique and globally attractive stationary solution for the flow defined by (1)–(2) that is denoted by  $\bar{\rho} = \bar{\rho}_{\lambda, E}$ . In particular,  $\bar{\rho}_{\lambda, E}$  is the typical density profile in the stationary nonequilibrium state corresponding to time independent chemical potential  $\lambda$  and external field  $E$ .

Fix time dependent paths  $\lambda(t)$  of the chemical potential and  $E(t)$  of the driving field. Given a density profile  $\rho$ , denote by  $(u(t), j(t))$ ,  $t \geq 0$ , the solution of (1)–(2) with initial condition  $\rho$ . Let  $W_{[0, T]} = W_{[0, T]}(\lambda, E, \rho)$  be the energy exchanged between the system and the external driving in the time interval  $[0, T]$ , that is

$$W_{[0, T]} = \int_0^T dt \left\{ \int_{\Lambda} dx j(t) \cdot E(t) - \int_{\partial\Lambda} d\sigma \lambda(t) j(t) \cdot \hat{n} \right\}, \quad (3)$$

where  $\hat{n}$  is the outer normal to  $\partial\Lambda$  and  $d\sigma$  is the surface measure on  $\partial\Lambda$ . The first term on the right hand side is the energy provided by the external field while the second is the energy provided by the reservoirs.

In view of the boundary conditions and the Einstein relation, by using the divergence theorem in (3), we deduce that

$$W_{[0, T]} = F(u(T)) - F(\rho) + \int_0^T dt \int_{\Lambda} dx j(t) \cdot \chi(u(t))^{-1} j(t), \quad (4)$$

where  $F$  is the equilibrium free energy functional,

$$F(\rho) = \int_{\Lambda} dx f(\rho(x)). \quad (5)$$

Consider two stationary states corresponding to (time-independent)  $(\lambda_0, E_0)$  and  $(\lambda_1, E_1)$  and denote by  $\bar{\rho}_0 = \bar{\rho}_{\lambda_0, E_0}$  and  $\bar{\rho}_1 = \bar{\rho}_{\lambda_1, E_1}$  the associated density profiles. Such states can be either equilibrium or nonequilibrium states. We can drive the system from the initial state  $\bar{\rho}_0$  at time  $t = 0$  to the final state  $\bar{\rho}_1$  at time  $t = +\infty$  by considering a time dependent forcing  $(\lambda(t), E(t))$  satisfying  $(\lambda(0), E(0)) = (\lambda_0, E_0)$  and  $(\lambda(+\infty), E(+\infty)) = (\lambda_1, E_1)$ . As the second term on the right hand side of (4) is positive, by letting  $W = W_{[0, +\infty]}$  be the total energy exchanged in the transformation, we deduce the Clausius inequality

$$W \geq \Delta F = F(\bar{\rho}_1) - F(\bar{\rho}_0). \quad (6)$$

When the initial and final states are equilibrium states, e.g., the external field  $E$  vanishes and the chemical potential  $\lambda$  is constant, the inequality (6) is a standard formulation of the second law of thermodynamics. Moreover, by considering a sequence of transformations in which the variation of the driving becomes very slow, it is not difficult to show that equality in (6) is achieved in the quasi static limit, we refer to [15] for the details. On the

other hand, for nonequilibrium states the inequality (6) does not carry any information. Indeed, as nonequilibrium states support a non vanishing current, in the limit  $T \rightarrow +\infty$  the second term on the right hand side of (4) becomes infinite so that the left hand side of (6) is infinite while the right hand side is bounded. By interpreting the ideas in [16], further developed in [10, 17], we next define a renormalized work for which a significant Clausius inequality can be obtained also for nonequilibrium stationary states.

To this aim, we recall the quasi potential, which is the key notion of the macroscopic fluctuation theory. Consider a system with time independent driving and let  $(\hat{u}(t), \hat{j}(t))$ ,  $t \in [T_1, T_2]$  be a pair density–current satisfying the continuity equation  $\partial_t \hat{u} + \nabla \cdot \hat{j} = 0$ . According to the basic principles of the macroscopic fluctuation theory [1, 3, 4], the probability of observing this path is given, up to a prefactor, by  $\exp \left\{ -\varepsilon^{-d} \beta I_{[T_1, T_2]}(\hat{u}, \hat{j}) \right\}$  where  $\varepsilon$  is the scaling parameter, i.e., the ratio between the microscopic length scale (say the typical intermolecular distance) and the macroscopic one,  $\beta = 1/\kappa T$  (here  $T$  is the temperature and  $\kappa$  is Boltzmann's constant), and the action functional  $I$  has the form

$$I_{[T_1, T_2]}(\hat{u}, \hat{j}) = \frac{1}{4} \int_{T_1}^{T_2} dt \int_{\Lambda} dx [\hat{j}(t) - J(\hat{u}(t))] \cdot \chi(\hat{u}(t))^{-1} [\hat{j}(t) - J(\hat{u}(t))]. \quad (7)$$

In particular, if  $(\hat{u}, \hat{j})$  solves (1)–(2) then  $I_{[T_1, T_2]}(\hat{u}, \hat{j}) = 0$ . The above statement therefore implies that the typical behavior of the system is described by the hydrodynamic equations. The quasi potential is the functional on the set of density profiles defined by the variational problem

$$V(\rho) = \inf \left\{ I_{(-\infty, 0]}(\hat{u}, \hat{j}), \hat{u}(0) = \rho \right\} \quad (8)$$

where the infimum is carried out over all the trajectories satisfying the prescribed boundary condition. Namely,  $V(\rho)$  is the minimal action to bring the system from the typical density profile  $\bar{\rho}$  to the fluctuation  $\rho$ . The probability of a density profile  $\rho$  in the stationary nonequilibrium ensemble is then given, up to a prefactor, by  $\exp \left\{ -\varepsilon^{-d} \beta V(\rho) \right\}$ . In particular, the minimizer of  $V$  is the typical density profile  $\bar{\rho}$ . For equilibrium states it can be shown [4] that  $V$  coincides, apart an affine transformation, with the free energy functional (5). Moreover, as shown in [1], the functional  $V$  solves the stationary Hamilton-Jacobi equation

$$\int_{\Lambda} dx \nabla \frac{\delta V}{\delta \rho} \cdot \chi(\rho) \nabla \frac{\delta V}{\delta \rho} - \int_{\Lambda} dx \frac{\delta V}{\delta \rho} \nabla \cdot J(\rho) = 0 \quad (9)$$

where  $\delta V/\delta \rho$  vanishes at the boundary  $\partial\Lambda$  and  $\rho$  satisfies the boundary condition  $f'(\rho(x)) = \lambda(x)$ ,  $x \in \partial\Lambda$ .

In the case of time independent driving, the current  $J(\rho)$  in (2) can be decomposed as  $J(\rho) = J_S(\rho) + J_A(\rho)$ , where  $J_S(\rho) = -\chi(\rho) \nabla \frac{\delta V}{\delta \rho}$  and  $J_A(\rho) = J(\rho) - J_S(\rho)$ . In

view of the stationary Hamilton-Jacobi equation (9), the above decomposition is orthogonal in the sense that for each density profile  $\rho$

$$\int_{\Lambda} dx J_S(\rho) \cdot \chi(\rho)^{-1} J_A(\rho) = 0. \quad (10)$$

We shall refer to  $J_S(\rho)$  as the *symmetric* current and to  $J_A(\rho)$  as the *antisymmetric* current. This terminology refers to symmetric and antisymmetric part of the microscopic dynamics [1, 3]. We remark that  $J_S$  is proportional to the thermodynamic force and the above decomposition depends on the external driving.

Since the quasi potential  $V$  is minimal in the stationary profile, the above definitions imply that  $J_S(\bar{\rho}) = 0$ ; namely, the stationary current is purely antisymmetric. In particular,  $J_A(\bar{\rho})$  is the typical current in the stationary nonequilibrium ensemble associated and it is therefore experimentally accessible. In view of the general formula (4) for the total work, the amount of energy per unit of time needed to maintain the system in the stationary profile  $\bar{\rho}$  is

$$\int_{\Lambda} dx J_A(\bar{\rho}) \cdot \chi(\bar{\rho})^{-1} J_A(\bar{\rho}). \quad (11)$$

We shall next consider time dependent driving and define a renormalized work by subtracting from the total work the energy needed to maintain the system out of equilibrium. Fix, therefore,  $T > 0$ , a density profile  $\rho$ , and space-time dependent chemical potentials  $\lambda(t)$  and external field  $E(t)$ ,  $t \in [0, T]$ . Let  $(u(t), j(t))$  be the corresponding solution of (1)–(2) with initial condition  $\rho$ . Recalling (11), we define the renormalized work  $W_{[0,T]}^{\text{ren}} = W_{[0,T]}^{\text{ren}}(\lambda, E, \rho)$  performed by the reservoirs and the external field in the time interval  $[0, T]$  as

$$W_{[0,T]}^{\text{ren}} = W_{[0,T]} - \int_0^T dt \int_{\Lambda} dx J_A(t, u(t)) \cdot \chi(u(t))^{-1} J_A(t, u(t)). \quad (12)$$

In this formula  $W_{[0,T]} = W_{[0,T]}(\lambda, E, \rho)$  is given in (3),

$$J(t, u) = J_S(t, u) + J_A(t, u),$$

$$J_S(t, u) = -\chi(u) \nabla \frac{\delta V_{\lambda(t), E(t)}(u)}{\delta u}$$

in which  $u$  is a generic density profile,  $J(t, u)$  is given by (2), and  $V_{\lambda(t), E(t)}$  is the quasi potential relative to the state  $(\lambda(t), E(t))$  with frozen  $t$ . Observe that the definition of the renormalized work involves the antisymmetric current  $J_A(t)$  computed not at density profile  $\bar{\rho}_{\lambda(t), E(t)}$  but at the solution  $u(t)$  of the time dependent hydrodynamic equation. That is, at time  $t$  we subtract the power the system would have dissipated if its actual state  $u(t)$  had been the stationary profile corresponding to  $(\lambda(t), E(t))$ . This choice, which is certainly reasonable

for slow transformations, leads to a Clausius inequality. Indeed, by using (4) and the orthogonality between the symmetric and the antisymmetric part of the current,

$$W_{[0,T]}^{\text{ren}}(\lambda, E, \rho) = F(u(T)) - F(\rho) + \int_0^T dt \int_{\Lambda} dx J_S(t, u(t)) \cdot \chi(u(t))^{-1} J_S(t, u(t)).$$

Consider a density profile  $\rho$  and a space-time dependent chemical potential and external field  $(\lambda(t), E(t))$ ,  $t \geq 0$ , converging to  $(\lambda_1, E_1)$  as  $t \rightarrow +\infty$ . Let  $\bar{\rho}_1 = \bar{\rho}_{\lambda_1, E_1}$  be the stationary profile associated to  $(\lambda_1, E_1)$  and  $(u(t), j(t))$ ,  $t \geq 0$ , be the solution of (1)–(2) with initial condition  $\rho$ . Since  $u(T)$  converges to  $\bar{\rho}_1$ , the symmetric part of the current,  $J_S(u(T))$ , relaxes as  $T \rightarrow +\infty$  to  $J_S(\bar{\rho}_1) = 0$ . Under suitable assumptions on the transformation, the last integral in the previous formula is convergent as  $T \rightarrow +\infty$ . By letting  $W^{\text{ren}} = \lim_{T \rightarrow \infty} W_{[0,T]}^{\text{ren}}$ , we thus get

$$W^{\text{ren}}(\lambda, E, \rho) = F(\bar{\rho}_1) - F(\rho) + \int_0^{\infty} dt \int_{\Lambda} dx J_S(t, u(t)) \cdot \chi(u(t))^{-1} J_S(t, u(t)) \quad (13)$$

where  $F$  is the equilibrium free energy functional (5). In particular,

$$W^{\text{ren}}(\lambda, E, \rho) \geq F(\bar{\rho}_1) - F(\rho), \quad (14)$$

which is a meaningful version of the Clausius inequality for nonequilibrium states. Furthermore, by considering a sequence of transformations  $(\lambda(t), E(t))$  which vary on a time scale  $1/\delta$ , we realize that the integrand on second term in the right hand side of (13) is of order  $\delta^2$  while the integral essentially extends, due to the finite relaxation time of the system, over an interval of order  $\delta^{-1}$ . Therefore in quasi static limit  $\delta \rightarrow 0$  equality in (14) is achieved. We refer to [15] for more details.

For special transformations, the integral in (13), which represents the *excess work* over a quasi static transformations, can be related to the quasi potential. Consider at time  $t = 0$  a stationary nonequilibrium profile  $\bar{\rho}_0$  corresponding to some driving  $(\lambda_0, E_0)$ . The system is put in contact with new reservoirs at chemical potential  $\lambda_1$  and a new external field  $E_1$ . For  $t > 0$  the system evolves according to the hydrodynamic equation (1)–(2) with initial condition  $\bar{\rho}_0$ , time independent boundary condition  $\lambda_1$  and external field  $E_1$ . In particular, as  $t \rightarrow +\infty$  the system relaxes to  $\bar{\rho}_1$ , the stationary density profile corresponding to  $(\lambda_1, E_1)$ . A simple calculation shows that along such a path

$$\int_0^{\infty} dt \int_{\Lambda} dx J_S(t, u(t)) \cdot \chi(u(t))^{-1} J_S(t, u(t)) = V_{\lambda_1, E_1}(\bar{\rho}_0) - V_{\lambda_1, E_1}(\bar{\rho}_1) = V_{\lambda_1, E_1}(\bar{\rho}_0).$$

The quasi potential  $V_{\lambda_1, E_1}(\bar{\rho}_0)$  thus represents the excess work, with respect to a quasi static transformation, along

the path that solves (1)–(2) with initial condition  $\bar{\rho}_0$  and time-independent driving  $(\lambda_1, E_1)$ .

To connect the above result with classical equilibrium thermodynamics, consider an equilibrium state with vanishing external field and constant chemical potential  $\lambda_0$  and let  $\bar{\rho}_0$  be the corresponding homogeneous density, i.e.  $\lambda_0 = f'(\bar{\rho}_0)$ . The system is put in contact with a new environment with chemical potential  $\lambda_1$ . In this case, recalling that  $f$  is the free energy per unit of volume and that the temperature of the system is the same of the environment, the *availability* per unit of volume is defined, see [18, Ch. 7], by  $a = f(\bar{\rho}_0) - \lambda_1 \bar{\rho}_0$ . The function  $a$ , which depends on state of the system and the environment, can be used to compute the maximal useful work that can be extracted from the system in the given environment. More precisely, by letting  $\bar{\rho}_1$  be such that  $f'(\bar{\rho}_1) = \lambda_1$ , then  $-\Delta a = f(\bar{\rho}_0) - f(\bar{\rho}_1) - \lambda_1(\bar{\rho}_0 - \bar{\rho}_1) \geq 0$  is the maximal useful work per unit of volume that can be extracted from the system in the given environment. By computing the quasi potential for equilibrium states, see [4], we get  $V_{\lambda_1,0}(\bar{\rho}_0) = -|\Lambda|\Delta a$ . Therefore, while a definition of thermodynamic potentials, that is functionals of the state of the system, does not appear possible in nonequilibrium thermodynamics, the quasi potential is the natural extension of the availability.

In terms of the underlying microscopic ensembles, as discussed in [15], the quasi potential  $V_{\lambda_1,E_1}(\bar{\rho}_0)$  can be obtained by computing the relative entropy of the ensemble associated to  $(\lambda_0, E_0)$  with respect to the one associated to  $(\lambda_1, E_1)$ . By considering a Markovian model for such underlying dynamics, it is also possible to give a microscopic definition of the exchanged work which in the hydrodynamic scaling limit converges to (3). The corresponding fluctuations can be deduced from those of the empirical current [3].

The definition of renormalized work we have introduced is natural and ensures, as we have discussed, both its finiteness and the validity of a Clausius inequality. From an operational point of view, the quasi potential, a generically nonlocal quantity, can be obtained from the measurement of the density correlation functions. In fact  $V$  is the Legendre transform of the generating functional of density correlation functions [1]. On the other hand, the identity  $W^{\text{ren}} = \Delta F$ , that is achieved for quasi static transformations, requires the knowledge of the total current in the intermediate stationary states that can be directly measured.

One may ask whether there exist, with respect to (12), alternative renormalizations of the total work. For instance, in the recent work [19], Maes and Netocny considered the topic of a renormalized Clausius inequality in the context of a single Brownian particle in a time dependent environment. To compare the approach in [19] to the present one, consider  $N$  independent diffusions in the thermodynamic limit  $N \rightarrow \infty$ . Each diffusion solves the Langevin equation  $\dot{X} = E(t, X) + \sqrt{2}\dot{w}$ , where  $E$  is

a time dependent vector field and  $\dot{w}$  denotes white noise. The corresponding stationary measure with  $E$  frozen at time  $t$  is denoted by  $\exp\{-v(t, x)\}$ . The scheme discussed here can be now applied, the hydrodynamic equations are (1)–(2) with  $D = 1$  and  $\chi(\rho) = \rho$ . Our renormalized work is given by (12) with  $J_A(t, \rho) = \rho[E(t, x) + \nabla v(t, x)]$ . The renormalization introduced in [19] is instead obtained by introducing a potential field such that the corresponding stationary state has minimal entropy production. Namely, they write  $E = f - \nabla U$  and subtract from the energy exchanged the space-time integral of  $|J_t^\varphi|^2/\rho$  where  $J_t^\varphi = \rho(f - \nabla\varphi) - \nabla\rho$  and  $\varphi = \varphi(t, x; \rho)$  is chosen so that  $\nabla \cdot J_t^\varphi = 0$ . While the two renormalization schemes are different, both satisfy the Clausius inequality (14) with  $F(\rho) = \int dx \rho \log \rho$ . Observe that in this case of independent particles our renormalization is local while the dependence of  $J_t^\varphi$  on  $\rho$  is nonlocal. It is not clear to us how the approach in [19] can be generalized to cover the case of interacting particles in the hydrodynamic scaling limit.

We are in debt to J.L. Lebowitz for continued discussions on the topics here considered. We also acknowledge stimulating exchanges of views with C. Di Castro, T. Komatsu, N. Nakagawa, S. Sasa, and H. Tasaki.

- 
- [1] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, C. Landim; J. Statist. Phys. **107**, 635–675 (2002).
  - [2] C. Appert-Rolland, B. Derrida, V. Lecomte, F. van Wijland; Phys. Rev. E **78**, 021122 (2008).
  - [3] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, C. Landim; J. Statist. Phys. **123**, 237–276 (2006).
  - [4] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, C. Landim; J. Stat. Mech. Theory Exp. 2007, P07014.
  - [5] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, C. Landim; J. Stat. Mech. Theory Exp. 2010, L11001.
  - [6] T. Bodineau, B. Derrida; Phys. Rev. E **72** 066110 (2005).
  - [7] B. Derrida; J. Stat. Mech. Theory Exp. 2007, P07023.
  - [8] C. Jarzynski, Phys. Rev. Lett. **78**, 2690 (1997).
  - [9] G.E. Crooks, Phys. Rev. E **60**, 2731 (1999).
  - [10] T. Hatano, S. Sasa; Phys. Rev. Lett. **86**, 3463 (2001).
  - [11] H.-H. Hasegawa, J. Ishikawa, K. Takara, D.J. Driebe; Phys. Let A **374**, 1001 (2010); K. Takara, H.-H. Hasegawa, D.J. Driebe; Phys. Let A **375**, 88 (2010).
  - [12] M. Esposito, C. Van den Broeck; EPL **95**, 40004 (2011).
  - [13] S. Deffner, E. Lutz; Preprint 2012, [arXiv:1201.3888](#)
  - [14] H. Spohn; *Large scale dynamics of interacting particles*. Springer, Berlin, 1991.
  - [15] L. Bertini, D. Gabrielli, G. Jona-Lasinio, C. Landim; Preprint 2012, [arXiv:1206.2412v1](#)
  - [16] Y. Oono, M. Paniconi; Progr. Theoret. Phys. Suppl. **130**, 29 (1998).
  - [17] T. Komatsu, N. Nakagawa, S. Sasa, H. Tasaki; J. Stat. Phys. **142**, 127 (2011).
  - [18] A.B. Pippard; *Elements of classical thermodynamics for advanced students of physics*. Cambridge University Press, New York 1957.
  - [19] C. Maes, K. Netocny; Preprint 2012, [arXiv:1206.3423](#)