

A Variational Method in Out of Equilibrium Physical Systems

Mario J. Pinheiro^{1, a)}

*Department of Physics, Instituto Superior Técnico, Av. Rovisco Pais,
1049-001 Lisboa, Portugal*

*& Institute for Advanced Studies in the Space, Propulsion and Energy Sciences
265 Ita Ann Ln. Madison, AL 35757 USA^{b)}*

(ΩDated: 13 February 2022)

A variational principle is further developed for out of equilibrium dynamical systems by using the concept of maximum entropy. With this new formulation it is obtained a set of two first-order differential equations, revealing the same formal symplectic structure shared by classical mechanics, fluid mechanics and thermodynamics. In particular, it is obtained an extended equation of motion for a rotating dynamical system, from where it emerges a kind of topological torsion current of the form $\epsilon_{ijk}A_j\omega_k$, with A_j and ω_k denoting components of the vector potential (gravitational or/and electromagnetic) and ω is the angular velocity of the accelerated frame. In addition, it is derived a special form of Umov-Poynting's theorem for rotating gravito-electromagnetic systems, and obtained a general condition of equilibrium for a rotating plasma. The variational method is then applied to clarify the working mechanism of some particular devices, such as the Bennett pinch and vacuum arcs, to calculate the power extraction from an hurricane, and to discuss the effect of transport angular momentum on the radioactive heating of planetary atmospheres. This development is seen to be advantageous and opens options for systematic improvements.

PACS numbers: 45.10.Db, 05.45.-a, 83.10.Ff, 52.30.-q, 52.80.Mg, 47.32.C-, 47.32.Ef, 95.30.Qd, 52.55.Fa

Keywords: Variational methods in classical physics; Nonlinear dynamics; Classical mechanics continuous media; Plasma dynamics; Arc discharges; Vortex dynamics; Rotating flows; Astrophysical plasma, Tokamaks

^{a)}Electronic mail: mpinheiro@ist.utl.pt; <http://web.ist.utl.pt/ist12493>

^{b)}<http://www.ias-spes.org>;

I. INTRODUCTION

In the years 1893-96, the Norwegian explorer Fridtjof Nansen, while he were traveling in the Arctic region, noticed the drift of surface ice across the polar sea making an angle of 20 to 40 degrees to the right related to the wind direction. Nansen has advanced an explanation, that in addition to the wind's force, it was also actuating the Coriolis force. In 1905, Vagn Walfrid Ekman¹ introduced a theory of wind currents in open seas showing that sea current change of direction with depth due to Coriolis force appearing in a rotating based coordinate system. Besides attracting considerable interest in geophysical flow problems², these discoveries stimulated further investigation in such fields as magnetic geodynamics, binary stars, new-born planetary systems, trying to unveil the important problem of angular momentum transport, and as well, this work.

The problem of enhanced angular moment transport in accretion disks³, and the breakdown of Keplerian rotation, as well as the removing of angular momentum from a vortex due to moving spiral waves, playing an important role on the total angular momentum balance of the core and the intensification of a tropical cyclone⁴, are all examples of problems that demand a clear understanding of the dynamics of electromagnetic-gravitational rotating systems. Furthermore, special attention must be dedicated to the role of flux of angular momentum, and its conservation. Notwithstanding all attempts to take an accurate account of the transport angular momentum⁵, it lacks an additional equation of conservation (besides continuity, momentum and energy equations), which is related to local density of angular momentum, and its related flux. In order to develop a logically closed foundation, the equation for the angular momentum balance must be included and, to our best knowledge, these problems were treated by Curtiss⁶ and Livingston and Curtiss⁷, and they are also addressed in the present work but using a variational method.

In this article, we develop a standard technique for treating a physical system on the basis of an information-theoretic framework previously developed⁸⁻¹⁰, which constitutes an entropy maximizing method leading to a set of two first-order differential equations revealing the same formal symplectic structure shared by classical mechanics and thermodynamics^{11,12}. Our method bears some resemblance with isoentropic but energy-nonconserving variational principle proposed by R. Jackiw *et al.*¹³ allowing to study nonequilibrium evolution in the context of quantum field theory, but with various classical analogous such as the Schrödinger

equation giving rise to reflectionless transmission.

The plan of the article is as follows. In Sec. II, we include the extended mathematical formalism to nonequilibrium information theory. In Sec. III, we analysis the equilibrium and stability of a rotating plasma. In Sec. IV, we apply our formalism to angular momentum transport, obtaining in particular the Umov-Poynting's theorem for rotating electromagnetic-gravitational systems (e.g., rotating plasmas, magnetic geodynamics, vortex motion and accretion disks in astrophysics), whose applications might contribute to clarify still poorly understood phenomena.

II. MATHEMATICAL PROCEDURE

Let us consider here a simple dynamical system constituted by a set of N discrete interacting mass-points $m^{(\alpha)}$ ($\alpha = 1, 2, \dots, N$) with x_i^α and v_i^α ($i = 1, 2, 3; \alpha = 1, \dots, N$) denoting the coordinates and velocities of the mass point in a given inertial frame of reference. The inferior Latin index refers to the Cartesian components and the superior Greek index distinguishes the different mass-points.

The gravitational potential $\phi^{(\alpha)}$ associated to the mass-point α is given by

$$\phi^{(\alpha)} = G \sum_{\substack{\beta=1 \\ \beta \neq \alpha}}^N \frac{m^{(\alpha)}}{|\mathbf{x}^{(\alpha)} - \mathbf{x}^{(\beta)}|}, \quad (1)$$

with G denoting the gravitational constant and $\mathbf{x}^{(\alpha)}$ and $\mathbf{x}^{(\beta)}$ representing the instantaneous positions of the mass-points (α) and (β). \sum denotes summation over all the particles of the system. Energy, momentum and angular momentum conservation laws must be verified for a totally isolated system:

$$E = \sum_{\alpha=1}^N E_{(\alpha)}, \quad (2)$$

$$\mathbf{P} = \sum_{\alpha=1}^N \mathbf{p}^{(\alpha)}, \quad (3)$$

the particles total angular momentum (sum of the orbital angular momentum plus the

intrinsic angular momentum or spin)

$$\mathbf{L} = \sum_{\alpha=1}^N \mathbf{L}^{(\alpha)} = \sum_{\alpha=1}^N ([\mathbf{r}^{(\alpha)} \times \mathbf{p}^{(\alpha)}] + \mathbf{J}^{(\alpha)}). \quad (4)$$

In the above equations, $\mathbf{r}^{(\alpha)}$ is the position vector relatively to a fixed frame of reference \mathcal{R} , $\mathbf{p}^{(\alpha)}$ is the total momentum (particle + field) and $\mathbf{L}^{(\alpha)}$ is the total angular momentum of the particle, comprising a vector sum of the particle orbital angular momentum and the intrinsic angular momentum \mathbf{J} (e.g., contributed by the electron spin and/or nuclear spin, since the electromagnetic momentum is already included in the preceding term through $\mathbf{p}^{(\alpha)}$). The maximum entropy principle introduces Lagrange multipliers from which, as we will see, ponderomotive forces are obtained.

It is necessary to find the conditional extremum; they are set up not for the function S itself but rather for the changed function \bar{S} . Following the mathematical procedure proposed in Ref.¹⁰ the total entropy of the system \bar{S} is thus given by

$$\begin{aligned} \bar{S} = \sum_{\alpha=1}^N \left\{ S^{(\alpha)} \left(E^{(\alpha)} - \frac{(\mathbf{p}^{(\alpha)})^2}{2m^{(\alpha)}} - \frac{(\mathbf{J}^{(\alpha)})^2}{2I^{(\alpha)}} - q^{(\alpha)}V^{(\alpha)} + q^{(\alpha)}(\mathbf{A}^{(\alpha)} \cdot \mathbf{v}^{(\alpha)}) - U_{mec}^{(\alpha)} \right) \right. \\ \left. + (\mathbf{a} \cdot \mathbf{p}^{(\alpha)} + \mathbf{b} \cdot ([\mathbf{r}^{(\alpha)} \times \mathbf{p}^{(\alpha)}] + \mathbf{J}^{(\alpha)})) \right\} = \sum_{\alpha=1}^N \mathfrak{S}^{(\alpha)}. \end{aligned} \quad (5)$$

where \mathbf{a} and \mathbf{b} are (vectors) Lagrange multipliers. It can be shown that $\mathbf{v}_{rel} = \mathbf{a}T$ and $\boldsymbol{\omega} = \mathbf{b}T$ (see also Ref.¹⁴). The conditional extremum points form the dynamical equations of motion of a general physical system (the equality holds whenever the physical system is in thermodynamic and mechanical equilibrium), defined by two first order differential equations:

$$\frac{\partial \bar{S}}{\partial \mathbf{p}^{(\alpha)}} \geq 0 \blacktriangleright \text{canonical momentum}; \quad (6a)$$

$$\frac{\partial \bar{S}}{\partial \mathbf{r}^{(\alpha)}} = -\frac{1}{T} \nabla_{r^{(\alpha)}} U^{(\alpha)} - \frac{1}{T} m^{(\alpha)} \frac{\partial \mathbf{v}^{(\alpha)}}{\partial t} \geq 0 \blacktriangleright \text{fundamental equation of dynamics}. \quad (6b)$$

Eq. 6b gives the fundamental equation of dynamics and has the form of a general local

balance equation having as source term the spatial gradient of entropy, $\nabla_a S > 0$, whilst Eq. 6a gives the canonical momentum (see also Eq. 12). At thermodynamic equilibrium the total entropy of the body has a maximum value, constrained through the supplementary conditions 2, 3, and 4, which ensues typically from the minimization techniques associated to the Lagrange multipliers. In the more general case of a non-equilibrium process, the entropic gradient must be positive in Eq. 6b, according to Vanderlinde's proposition¹⁵, a condition required for the gravitational force to exist. However, new physics may be brought by the set of two first order differential equations, related to the interplay between the tendency of energy to attain a minimum, whilst entropy seeks to maximize its value.

In non-equilibrium processes the gradient of the total entropy in momentum space multiplied by factor T is given by

$$T \frac{\partial \bar{S}}{\partial \mathbf{p}^{(\alpha)}} = \left\{ -\frac{\mathbf{p}^{(\alpha)}}{m^{(\alpha)}} + \frac{q^{(\alpha)}}{m^{(\alpha)}} \mathbf{A} + \mathbf{v}_e + [\boldsymbol{\omega} \times \mathbf{r}^{(\alpha)}] \right\}, \quad (7)$$

so that maximizing entropy change in Eq. 6a leads to the well-known total (canonical) momentum:

$$\mathbf{p}^{(\alpha)} = m^{(\alpha)} \mathbf{v}_e + m^{(\alpha)} [\boldsymbol{\omega} \times \mathbf{r}^{(\alpha)}] + q^{(\alpha)} \mathbf{A}. \quad (8)$$

The above formulation bears some resemblance with Hamiltonian formulation of dynamics which expresses first-order constraints of the Hamiltonian H in a $2n$ dimensional phase space, $\dot{\mathbf{p}} = -\partial H / \partial \mathbf{q}$ and $\dot{\mathbf{q}} = \partial H / \partial \mathbf{p}$, and can be solved along trajectories as quasistatic processes, revealing the same formal symplectic structure shared by classical mechanics and thermodynamics^{11,12}.

The *internal mechanical energy* term, $U_{mec}^{(\alpha)}$, appearing in Eq. 5 may be defined by:

$$U_{mec}^{(\alpha)} = m^{(\alpha)} \phi^{(\alpha)}(\mathbf{r}) + m^{(\alpha)} \sum_{\substack{\beta=1 \\ \beta \neq \alpha}}^N \phi^{(\alpha, \beta)}. \quad (9)$$

Considering that by definition of thermodynamic temperature, $\partial S^{(\alpha)} / \partial U^{(\alpha)} \equiv 1/T^{(\alpha)}$, then it follows

$$\nabla_{r^{(\alpha)}} U^{(\alpha)} = -m^{(\alpha)} \nabla \phi^{(\alpha)} - \frac{\mathbf{p}^{(\alpha)}}{m^{(\alpha)}} \cdot \nabla \mathbf{p}^{(\alpha)} - \nabla_{r^{(\alpha)}} \left(\frac{J^{(\alpha)} 2}{2I^{(\alpha)}} \right) - q^{(\alpha)} \nabla V^{(\alpha)} + q^{(\alpha)} \nabla (\mathbf{A}^\alpha \cdot \mathbf{v}^{(\alpha)}). \quad (10)$$

Eq. 10 contains the particle's self-energy and the particle interaction energy for the gravitational and electromagnetic fields, but it may also include other terms representing different occurring phenomena (energy as a bookkeeping concept), such as terms included in Eq. 9. We may here recall that the entropic flux in space is a kind of generalized force X_α ^{16,17}, and it can be shown that the following equation holds:

$$\begin{aligned} T\nabla_{r^{(\alpha)}}\mathfrak{S}^{(\alpha)} &= -q^{(\alpha)}\nabla_{r^{(\alpha)}}V^{(\alpha)} + q^{(\alpha)}\nabla_{r^{(\alpha)}}(\mathbf{A}^{(\alpha)} \cdot \mathbf{v}^{(\alpha)}) \\ &+ m^{(\alpha)}\mathbf{v}^{(\alpha)} \cdot \nabla\mathbf{v}^{(\alpha)} - \nabla_{r^{(\alpha)}}\left(\frac{(\mathbf{J}^{(\alpha)})^2}{2I^{(\alpha)}} - \boldsymbol{\omega} \cdot \mathbf{J}^{(\alpha)}\right), \end{aligned} \quad (11)$$

Now, it make sense to write the fundamental equation of thermodynamics under the form of a spacetime differential equation:

$$T\nabla\bar{S} + \sum_{\alpha} m^{(\alpha)}\frac{\partial\mathbf{v}^{(\alpha)}}{\partial t} = \sum_{\alpha} \nabla\mathbf{U}^{(\alpha)}. \quad (12)$$

Taking into account the convective derivative, $d\mathbf{v}^{(\alpha)}/dt \equiv \partial\mathbf{v}^{(\alpha)}/\partial t + \mathbf{v}^{(\alpha)} \cdot \nabla\mathbf{v}^{(\alpha)}$, after some algebra we obtain:

$$\begin{aligned} m^{(\alpha)}\frac{d\mathbf{v}^{(\alpha)}}{dt} &= -T\nabla_{r^{(\alpha)}}\mathfrak{S}^\alpha - m^{(\alpha)}\nabla\phi^{(\alpha)} - q^{(\alpha)}\nabla V^{(\alpha)} \\ &+ q^{(\alpha)}\nabla(\mathbf{A}^{(\alpha)} \cdot \mathbf{v}^{(\alpha)}) - \nabla_{r^{(\alpha)}}\left(\frac{(\mathbf{J}^{(\alpha)})^2}{2I^{(\alpha)}} - \boldsymbol{\omega} \cdot \mathbf{J}^{(\alpha)}\right) + \mathbf{F}_{ext}^{(\alpha)}. \end{aligned} \quad (13)$$

For conciseness, the term $U^{(\alpha)}$ now includes all forms of energy inserted into the above Eq. 5. On the right-hand side (r.h.s.), the first term must be present whenever the mechanical and thermodynamical equilibrium conditions are not fulfilled; the second term is the gravitational force term; the third and fourth terms constitute the Lorentz force; the fifth term is a new term which represents the transport of angular momentum; while the last term represents other external forces not explicitly included and actuating over the particle (α) .

A. Examples:

1. *Extended fundamental equation of dynamics*

The present formalism was applied in a previous work¹⁰, and therein we obtained the ponderomotive forces acting on a charged particle. For a neutral particle (body) in a gravitational field, Eq. 13 points to a kind of extended fundamental equation of dynamics for a given species (α) at equilibrium and at a given point of space-time (Eulerian description):

$$m^{(\alpha)} \frac{\partial \mathbf{v}^{(\alpha)}}{\partial t} = -m^{(\alpha)} \nabla \phi^{(\alpha)} - \nabla_{r^{(\alpha)}} \left(\frac{\mathbf{J}^{(\alpha)2}}{2I^{(\alpha)}} - \boldsymbol{\omega} \cdot \mathbf{J}^{(\alpha)} \right). \quad (14)$$

Eq. 14 has a new term because the body possess an intrinsic angular momentum. In a non-rotating frame of reference we must put $\omega = 0$, and from the work-energy theorem it is obtained the total mechanical energy of the system, $\mathcal{E}_{mec} = K + U + J_c^2/2I_c$. This is the common approach in classical mechanics. We are interested in the effect of a given force in a given point of space-time, not in its effect along the particle trajectory. It is worth to point out that Eq. 14 was obtained through a variational procedure in contrast to the usual conservation theorem used, for example, in Ref.^{18,19}.

Included in the internal energy term are the interpressure term, see Eq. 9 (we consider here a homogeneous and isotropic fluid). The above described framework (see also Ref.¹⁰ for additional information) leads us to the well-known hydrodynamic equation for a given species (α):

$$\begin{aligned} m^{(\alpha)} \frac{d\mathbf{v}^{(\alpha)}}{dt} = & -\nabla_{r^{(\alpha)}} (m^{(\alpha)} \phi^{(\alpha)}) - m^{(\alpha)} \nabla_{r^{(\alpha)}} \sum_{\substack{\beta=1 \\ \beta \neq \alpha}}^N \phi^{(\alpha,\beta)} \\ & - \nabla_{r^{(\alpha)}} \left(\frac{J^{(\alpha)2}}{2I^{(\alpha)}} - \boldsymbol{\omega} \cdot \mathbf{J}^{(\alpha)} + T \mathfrak{S}^{(\alpha)} \right). \end{aligned} \quad (15)$$

Here, in the r.h.s. of the Eq. 15 we introduce explicitly external forces terms, eventually present in open systems.

Using the following correspondence from particle to fluid description

$$\sum_{\alpha} m^{(\alpha)} \rightarrow \int_V d^3x' \rho_v(\mathbf{x}'), \quad (16)$$

and, as well, a analogue relationship for the electric charge

$$\sum_{\alpha} q^{(\alpha)} \rightarrow \int_V d^3x' \rho(\mathbf{x}'), \quad (17)$$

we can can rewrite Eq. 15 under the form of the Euler (governing) equation:

$$\rho_v \frac{d\mathbf{v}}{dt} = -\rho_v \nabla_r \phi - \nabla_r p - \nabla_r \Phi_J - \nabla_r f. \quad (18)$$

Here, as usual, the total interparticle pressure term (e.g., Ref.²⁰) is given by:

$$p(\mathbf{r}) = \sum_{\alpha} m^{(\alpha)} \sum_{\substack{\beta=1 \\ \beta \neq \alpha}}^N \phi^{(\alpha, \beta)}(\mathbf{r}). \quad (19)$$

To simplify, we introduce a functional integral in the form of an intrinsic angular momentum energy density (comprising the “interaction energy term”, $\boldsymbol{\omega} \cdot \mathbf{J}$), Φ_J :

$$\sum_{\alpha} \left[\frac{J^{(\alpha)2}}{2I^{(\alpha)}} - \boldsymbol{\omega} \cdot \mathbf{J}^{(\alpha)} - (\Delta F)^{(\alpha)} \right] \rightarrow \int [\Phi_J(\mathbf{x}') + f(\mathbf{x}')] d^3x', \quad (20)$$

considering that the intrinsic angular momentum density refers to a given blob of fluid (with inertial momentum I , a measure of the local rotation, or spin, of the fluid element), and its associated free energy $f = f_0 - Ts$ (per unit of volume). Eq. 18 also means that function $S^{(\alpha)}$ (the *field integral of* $\mathbf{r}^{(\alpha)}$) is constant along the integrals curves of the space field $\mathbf{r}^{(\alpha)}$. The gradient of the free energy f of the out of equilibrium state is the source of spontaneous change from an unstable state to a more stable state, while performing work. For example, a common source of free energy in a collisionless plasma is an electric current²¹; in a magnetically confined plasma, several classes of free energy sources are available to drive instabilities, e.g., relaxation of a non-Maxwellian, nonisotropic velocity distribution²². At this stage, it is worthwhile to refer that our procedure includes the effect of angular momentum (through Eq. 5), as it should be in a consistent theory, according to Curtiss⁶.

Using, furthermore, the mathematical identity:

$$\nabla(\mathbf{A}^{(\alpha)} \cdot \mathbf{v}^{(\alpha)}) = (\mathbf{A}^{(\alpha)} \cdot \nabla) \mathbf{v}^{(\alpha)} + (\mathbf{v}^{(\alpha)} \cdot \nabla) \mathbf{A}^{(\alpha)} + [\mathbf{A}^{(\alpha)} \times [\nabla \times \mathbf{v}^{(\alpha)}]] + [\mathbf{v}^{(\alpha)} \times [\nabla \times \mathbf{A}^{(\alpha)}]], \quad (21)$$

we obtain after some algebra the following expression:

$$\nabla (\mathbf{A}^{(\alpha)} \cdot \mathbf{v}^{(\alpha)}) = -\frac{\partial \mathbf{A}^{(\alpha)}}{\partial t} - [\boldsymbol{\omega} \times \mathbf{A}^{(\alpha)}] + [\mathbf{v}^{(\alpha)} \times \mathbf{B}^{(\alpha)}]. \quad (22)$$

Here, $\mathbf{B} = [\nabla \times \mathbf{A}]$. In addition, we may notice that the following equality holds:

$$(\mathbf{A}^{(\alpha)} \cdot \nabla) \mathbf{v}^{(\alpha)} = [\boldsymbol{\omega} \times \mathbf{A}^{(\alpha)}], \quad (23)$$

where

$$\mathbf{A}^{(\alpha)} = \sum_{\substack{\beta=1 \\ \beta \neq \alpha}} q^{(\beta)} \frac{\mathbf{v}^{(\beta)}}{r_{\alpha\beta}}, \quad (24)$$

denotes the vector potential actuating on (α) particle due to all the other particles, and the vorticity is defined by

$$\boldsymbol{\Omega}^{(\alpha)} = [\nabla_{\mathbf{r}^{(\alpha)}} \times \mathbf{v}^{(\alpha)}] = 2\boldsymbol{\omega}^{(\alpha)}. \quad (25)$$

Therefore, it follows the general equation of dynamics for a physical system (Lagrangian description):

$$\rho \frac{d\mathbf{v}}{dt} = \rho \mathbf{E} + [\mathbf{J} \times \mathbf{B}] - \nabla \phi - \nabla p + \rho [\mathbf{A} \times \boldsymbol{\omega}]. \quad (26)$$

The last term in the r.h.s. of the Eq. 26 is a new term that represents a kind of topological spin vector²³, an artifact of nonequilibrium process. As it will be shown subsequently, the topological spin vector plays a role in plasma arcs, and as well in magnetocumulative generators²⁴, and suggests a new method to obtain the helicity transport equation²⁵.

2. *Rigid body rolling down an inclined plane with holonomic constraint*

We may check Eq. 14 in a standard example of classical mechanics: a rigid body of mass M rolling down an inclined plane making an angle θ with the horizontal (see, e.g., p. 97 of Ref.²⁶). Eq. 15 can be applied to solve the problem, with $\omega = 0$ (there is no rotation of the frame of reference) and considering that only the gravitational force acts on the rolling body, with inertial moment relative to its own center of mass given by $I_c = \beta M R^2$. Hence, we obtain:

$$M\ddot{x} = Mg \sin \theta - \partial_x w. \quad (27)$$

Here, $w \equiv \frac{(\mathbf{J}_c)^2}{2I_c}$. Assuming that the x-axis is directed along the inclined plane, and considering that the angular momentum relative to the rigid body center of mass is given by $J_c = I_c \omega'$, with $\omega' = d\theta/dt$, and noticing that $dx = v_x dt$ (holonomic constraint), it is readily obtained

$$M\ddot{x} = Ma_x = Mg \sin \theta - I_c \omega' \frac{d\omega'}{v_x dt} = Mg \sin \theta - \beta M R^2 \frac{\omega'}{\omega' R} \alpha. \quad (28)$$

Since $\alpha = a/R$, then it results the well-known equation

$$a_x = \frac{g \sin \theta}{(1 + \beta)}. \quad (29)$$

III. EQUILIBRIUM AND STABILITY OF A ROTATING PLASMA

Extremum conditions imposed on entropy or internal energy, does not only provide criteria for the evolution of the system, but, in addition, they determine the stability of thermodynamic systems, at equilibrium. Further, it has been shown²⁷ that a state of mechanical equilibrium can be reached, if the entropy increases with distance:

$$T L_r \bar{S} > 0, \quad (30)$$

where L_r is the Lie derivative along the vector field \mathbf{r} acting on scalar \bar{S} . Therefore, we can search another extremal condition through the general expression

$$T \frac{\partial \mathfrak{G}^{(\alpha)}}{\partial \mathbf{r}^{(\alpha)}} = -q^{(\alpha)} \nabla_{\mathbf{r}^{(\alpha)}} V^{(\alpha)} + q^{(\alpha)} \nabla_{\mathbf{r}^{(\alpha)}} (\mathbf{v}^{(\alpha)} \cdot \mathbf{A}^{(\alpha)}) - m^{(\alpha)} \frac{\partial \mathbf{v}^{(\alpha)}}{\partial t} - \nabla_{\mathbf{r}^{(\alpha)}} \left(\frac{(\mathbf{J}^{(\alpha)})^2}{2I^{(\alpha)}} - \boldsymbol{\omega} \cdot \mathbf{J}^{(\alpha)} \right). \quad (31)$$

According to Noether's theorem, the total canonical momentum is conserved in a closed system, and hence we can state the *closure relation*:

$$\sum_{\alpha=1}^N [\mathbf{p}^{(\alpha)} + q^{(\alpha)} \mathbf{A}^{(\alpha)}] = 0. \quad (32)$$

It can be shown (see Ref.^{9,10} for details) that the relationship that prevails in equilibrium on a rotating plasma is given by

$$q^{(\alpha)}\mathbf{E}^{(\alpha)} + q^{(\alpha)}[\mathbf{v} \times \mathbf{B}^{(\alpha)}] = m^{(\alpha)}\nabla_{\mathbf{r}^{(\alpha)}}\phi^{(\alpha)} + m^{(\alpha)}\sum_{\substack{\beta=1 \\ \beta \neq \alpha}}\nabla_{\mathbf{r}^{(\alpha)}}\phi^{(\alpha,\beta)} - q^{(\alpha)}[\mathbf{A}^{(\alpha)} \times \boldsymbol{\omega}^{(\alpha)}]. \quad (33)$$

We may develop the correspondence drawn from Eqs. 16- 17 and to obtain the *general condition of equilibrium of a rotating plasma*, in the presence of gravitational and electromagnetic interactions (e.g., Ref.²⁸):

$$\rho\mathbf{E} + [\mathbf{J} \times \mathbf{B}] = \nabla\Phi + \nabla p - \rho[\mathbf{A} \times \boldsymbol{\omega}]. \quad (34)$$

Hereinabove, to simplify the algebra, we took the averaged angular frequency $\boldsymbol{\omega}$, for the all system. Embedded into Eq. 34, it is present the vector potential \mathbf{A} , on the same footing as the E and B -fields. The importance of the vector potential when compared to the others, depends on the kind of B -field prevalent on the system; for instance, if the B -field is homogeneous, the vector potential field predominates on the region near the axis, since $\rho\omega A/JB \sim 1/r^2$. The topological spin vector term is fundamental, since it produces work, responsible for the system angular momentum modification, producing a rocket-like rotation effect on the plasma. The theoretical framework delineated here, may help to clarify problems related to rotating-plasma systems^{29,30}, and controlled thermonuclear plasma confinement³¹.

A. Bennett pinch

The compression of an electric current by a magnetic field, the z-pinch effect, can be studied on the basis of Eq. 34, which gives the condition for dynamic equilibria. Let us here assume a typical geometry for an infinitely long axisymmetric cylindrical arc (Fig. 1), with axial current density $J_z = J_z(r)$. Since the current density is assumed constant, Maxwell's equation in the steady state yields the azimuthal component $B_\theta = \mu_0 J_z r/2$, for $r \leq R$, with R the outer boundary of the cylindrical arc. The vector potential is purely radial, given by $A_z(r) = -\mu_0 R^2 J_z/4$, for $r < R$, and the Coriolis term plays no role. We can write Eq. 34 under the form:

$$-J_z B_\theta = \frac{dp}{dr}. \quad (35)$$




FIG. 1. Geometry and vectorial fields in the Bennett pinch generated by an axial current J_z creating a toroidal field B_ϕ . If, instead, we consider a vacuum arc discharge with radial current J_r and magnetic field B_z , we may have rotating arc with angular velocity ω .

and it follows that

$$p(r) = \int_r^R \frac{dp}{dr} dr = \frac{1}{4} \mu_0 J_z^2 (R^2 - r^2), \quad (36)$$

which is a well-known result.

The interaction between vacuum arcs and transverse magnetic fields, is used in switching devices (see e.g. Refs.^{32,33}). We can consider here instead, a coaxial configuration with a cathode on axis with a stabilizing magnetic induction field \mathbf{B} , directed along the symmetry axis, and an arc current density \mathbf{J} flowing radially (and assuming a "filamentary" current with radius R' , $A_r = -\mu_0 R'^2 J_r / 4$, with μ_0 the permeability of the vacuum). In this case, we may apply Eq. 34 to obtain the pressure differential, from the axis to the wall (at R):

$$\Delta p(r = R) = 2\pi R \left[-J_r B_z + |\rho_c| \pi R^2 \frac{\mu_0}{4\pi} J_r \omega \right] \Delta \theta. \quad (37)$$

$$|\rho_c| > \frac{\mu_0}{4\pi} \frac{B_z}{S} \omega, \quad \text{retrograde rotation} \quad (38a)$$

$$|\rho_c| < \frac{\mu_0}{4\pi} \frac{B_z}{S} \omega, \quad \text{amperian rotation.} \quad (38b)$$

Here, S denotes the filamentary current cross-section, and $\rho = \rho_c$. For negative charge carriers $\rho_c = -|\rho_c| = -en_e$, and we obtain an amperian (clockwise) rotation for high magnetic fields, and relatively weak arc currents, or instead, a retrograde rotation for higher intensity arcs (higher S) and small transverse magnetic fields, in agreement with experimental evidence (e.g., Ref.³²). We may see in this effect the compromise between two different tendencies, the energy which tends to a minimum and the entropy which tends to a maximum. From Eq. 37 it is obtained an expression for the spot velocity in a transverse magnetic field:

$$v = \omega R = \frac{4\pi}{\mu_0} \frac{1}{|\rho_c|} \left(\frac{\epsilon_0 i}{2\pi\sigma_c} + \frac{B_z R}{S} \right). \quad (39)$$

Here, we have made use of the Bernoulli relation, $\Delta p = \epsilon_0 E^2/2$, and of the constitutive equation $J_r = \sigma_c E$, σ_c denoting the plasma electrical conductivity. Although Eq. 39 is not self-consistent, it shows that the force term $[\mathbf{J} \times \mathbf{B}]$, is not playing alone its key role, and the spot velocity depends linearly of the arc current (see, e.g., Ref.³⁴). A better understanding of this phenomena is crucial, since arc discharges are powerful generators of non-equilibrium atmospheric pressure plasma.

IV. FLUX OF ANGULAR MOMENTUM

We now address the transport of angular momentum, a phenomena of huge importance in several phenomena, such as in the working mechanism of an accretion disk, the formation of a tornado, or the atmospheric circulation.

The equation of conservation of angular momentum can be obtained from the following equation:

$$m^{(\alpha)} \frac{d\mathbf{v}^{(\alpha)}}{dt} = -T \frac{\partial \mathfrak{S}^{(\alpha)}}{\partial \mathbf{r}^{(\alpha)}} - \nabla_{\mathbf{r}^{(\alpha)}} (m^{(\alpha)} \Phi^{(\alpha)}) - q^{(\alpha)} \nabla_{\mathbf{r}^{(\alpha)}} V^{(\alpha)}$$

$$+ q^{(\alpha)} \nabla_{\mathbf{r}^{(\alpha)}} (\mathbf{v}^{(\alpha)} \cdot \mathbf{A}^{(\alpha)}) - \nabla_{\mathbf{r}^{(\alpha)}} \left(\frac{(\mathbf{J}^{(\alpha)})^2}{2I^{(\alpha)}} - \boldsymbol{\omega}^\alpha \cdot \mathbf{J}^{(\alpha)} \right). \quad (40)$$

Multiplying the previous Eq. 40 by the particle velocity $v_i^{(\alpha)}$ ($i = x, y, z$), and after rearranging the terms, we obtain:

$$\begin{aligned} \frac{1}{2} \sum_{\alpha} m^{(\alpha)} \frac{d}{dt} |v_i^{(\alpha)}|^2 &= -\frac{1}{2} G \sum_{\alpha} \sum_{\beta \neq \alpha} m^{(\alpha)} m^{(\beta)} \frac{(x_i^{(\alpha)} - x_i^{(\beta)})(v_i^{(\alpha)} - v_i^{(\beta)})}{|\mathbf{x}^{(\alpha)} - \mathbf{x}^{(\beta)}|^3} \\ &- \frac{1}{2} \sum_{\alpha} \sum_{\beta \neq \alpha} \frac{1}{4\pi\epsilon_0} q^{(\alpha)} q^{(\beta)} \frac{(x_i^{(\alpha)} - x_i^{(\beta)})(v_i^{(\alpha)} - v_i^{(\beta)})}{|\mathbf{x}^{(\alpha)} - \mathbf{x}^{(\beta)}|^3} - \frac{1}{2} \sum_{\alpha} (q^{(\alpha)} v_i^{(\alpha)} \frac{\partial A_i^{(\alpha)}}{\partial t}) \\ &+ \frac{3}{2} \sum_{\alpha} \varepsilon_{ijk} \omega_k (q^{(\alpha)} A_j^{(\alpha)} v_i^{(\alpha)}) + \frac{1}{2} \sum_{\alpha} \varepsilon_{ijk} (q^{(\alpha)} v_i^{(\alpha)} v_j^{(\alpha)} B_k^{(\alpha)}) \\ &+ \frac{1}{2} \sum_{\alpha} v_i^{(\alpha)} \frac{\partial}{\partial x_i^{(\alpha)}} \left[\frac{|\mathbf{J}^{(\alpha)}|^2}{2I^{(\alpha)}} - (\boldsymbol{\omega}^\alpha \cdot \mathbf{J}^{(\alpha)}) \right] + T \sum_{\alpha} v_i^{(\alpha)} \frac{\partial \mathfrak{S}^{(\alpha)}}{\partial x_i^{(\alpha)}}. \end{aligned} \quad (41)$$

Eq. 41 can be written in a more comprehensive form, if we separate the terms from different contributions. For this purpose, let's define the total kinetic energy by means of the expression

$$\mathfrak{T} \equiv \frac{1}{2} \sum_{\alpha} m^{(\alpha)} |\mathbf{v}^{(\alpha)}|^2, \quad (42)$$

and let us denote by

$$\mathfrak{P}_g \equiv \frac{1}{2} \sum_{\alpha} m^{(\alpha)} \varphi^{(\alpha)} = \frac{1}{2} G \sum_{\alpha} \sum_{\beta \neq \alpha} \frac{m^{(\alpha)} m^{(\beta)}}{|\mathbf{x}^{(\alpha)} - \mathbf{x}^{(\beta)}|}, \quad (43)$$

the overall gravitational energy. Similarly, the total electrostatic energy is given by:

$$\mathfrak{P}_v \equiv \frac{1}{2} \frac{1}{4\pi\epsilon_0} \sum_{\alpha} \sum_{\beta \neq \alpha} \frac{q^{(\alpha)} q^{(\beta)}}{|\mathbf{x}^{(\alpha)} - \mathbf{x}^{(\beta)}|}. \quad (44)$$

Following now the Umov-Poynting procedure, it can be shown that the previous Eq. 40 reduces after some algebra to the following one:

$$\frac{d}{dt} (\mathfrak{T}) = -\frac{d}{dt} (\mathfrak{P}_g) - \frac{d}{dt} \mathfrak{P}_v + \frac{3}{2} \sum_{\alpha} q^\alpha [\mathbf{v}^{(\alpha)} \times [\mathbf{A}^{(\alpha)} \times \boldsymbol{\omega}]]$$

$$+ \frac{1}{2} \sum_{\alpha} \mathbf{v}^{(\alpha)} \cdot \nabla_{\mathbf{r}^{(\alpha)}} \left[\frac{|\mathbf{J}^{\alpha}|^2}{2I^{(\alpha)}} - (\boldsymbol{\omega} \cdot \mathbf{J}^{(\alpha)}) \right] + T \sum_{\alpha} v_i^{(\alpha)} \frac{\partial \mathfrak{S}^{(\alpha)}}{\partial x_i^{(\alpha)}}. \quad (45)$$

As we did before, it is convenient to introduce a new physical quantity, representing the rotational energy:

$$\mathfrak{P}_{rot} = -\frac{3}{2} \sum_{\alpha} q^{\alpha} [\mathbf{v}^{(\alpha)} \times [\mathbf{A}^{(\alpha)} \times \boldsymbol{\omega}]]. \quad (46)$$

Then, the above Eq. 45, can be written under the more general form:

$$\frac{d}{dt} (\mathfrak{J} + \mathfrak{P}_g + \mathfrak{P}_v + \mathfrak{P}_{rot}) = \frac{1}{2} \mathbf{v} \cdot \nabla_{\mathbf{r}} \left[\frac{|\mathbf{J}|^2}{2I} - (\boldsymbol{\omega} \cdot \mathbf{J}) - \Delta F \right]. \quad (47)$$

Here, \mathbf{v} represents now the velocity of an “element of fluid”, and the last term in brackets is an average local value. A simple analysis of the Eq.47 shows that the system is stable in rotatory motion, provided that the following condition is satisfied:

$$\nabla_r \left[\frac{|\mathbf{J}|^2}{2I} - (\boldsymbol{\omega} \cdot \mathbf{J}) - \Delta F \right] < 0. \quad (48)$$

The last term is the free energy per unit of volume, $F = F_0 - TS$. Remark that Eq. 48 is consistent with Gibbs distribution in a rotating body (see, e.g., Ref. ¹⁴), which means that the radial flux of energy must be positive, flowing out radially from the system’s boundary. In addition, we notice that the equilibrium of a gravito-electromagnetic system depends upon its mechanical rotational properties, but also on the free energy available, linking intrinsically any mechanical process to thermodynamic variables, and opening options for possible unconventional mechanisms to control instabilities. In the domain of astrophysical plasmas, gravity and rotation usually dominate over the magnetic field of force, being crucial for the development of instabilities.

From Eq. 47 we may see that equilibrium ensues (neglecting thermal and configurational effects) when the velocity of rotation of the fluid satisfies the local condition (e.g., Ref.³⁵):

$$\frac{d\Omega(r)^2}{dr} \geq 0, \quad (49)$$

where we identified ω with the bulk angular velocity. Eq. 49 is related to the conservation of energy. However, when condition 48 is not fulfilled, it gives rise to the magnetorotational instability (MIR)^{35,36}, that appears as the result of the interplay of its three different terms:

i) the angular momentum acquired by the fluid (or particles); ii) an interaction term due to the coupling between the fluid angular momentum with the driven angular velocity; iii) the fluid thermal energy and configurational entropy.

A typical experiment consists in the rotation of a fluid between two concentric cylinders - related to the so called Taylor-Couette instability - driven by velocity gradients. In the presence of an axial magnetic field, the Taylor-Couette instability is onset when Eq. 49 is not satisfied. Also, we may expect that, owing to the fact that for two different species with different inertial moment, $I^\alpha \neq I^\beta$, it can be expected that at some point $r = r_c$ of the radial axis an inversion of the sign of the inequality of Eq. 48 must take place, and instability is onset. In particular, in the presence of two different species with different inertial moments, the fluid may well be intrinsically unstable at enough high angular speed. MIR instabilities concur against the stability of plasma configurations and in the 1960's, when MHD power plants were considered to be an economic process to convert thermal energy into electrical energy, E. Velikhov, in 1962, discovered the electrothermal instability which is at the source of strong magnetohydrodynamic turbulence^{35,37,38}.

The above Eq. 47 can be written in the form of a conservative equation for energy:

$$\begin{aligned} \frac{\partial}{\partial t}(\mathfrak{J} + \mathfrak{P}_g + \mathfrak{P}_v + \mathfrak{P}_{rot}) = & -\frac{1}{2} \sum_{\alpha} \nabla_{\mathbf{r}^{(\alpha)}} \cdot \left\{ \mathbf{v}^{(\alpha)} \left[-\frac{|\mathbf{J}^{(\alpha)}|^2}{2I^{(\alpha)}} + (\boldsymbol{\omega} \cdot \mathbf{J}^{(\alpha)}) - T\mathfrak{S}^{(\alpha)} \right] \right\} \\ & + \frac{1}{2} \sum_{\alpha} \left[-\frac{|\mathbf{J}^{(\alpha)}|^2}{2I^{(\alpha)}} + (\boldsymbol{\omega} \cdot \mathbf{J}^{(\alpha)}) - T\mathfrak{S}^{(\alpha)} \right] (\nabla_{\mathbf{r}^{(\alpha)}} \cdot \mathbf{v}^{(\alpha)}). \end{aligned} \quad (50)$$

Finally, we can transform the above Eq. 50 into a *Poynting's Theorem for rotating fluids*:

$$-\frac{\partial}{\partial t}(\mathcal{U}) = \nabla \cdot \mathbf{S} + \mathcal{P}', \quad (51)$$

after defining a kind of *Poynting vector for rotational fluids*, which gives the rate of rotational energy flow:

$$\mathbf{S} = \frac{1}{2} \sum_{\alpha} \mathbf{v}^{(\alpha)} \left[-\frac{|\mathbf{J}^{(\alpha)}|^2}{2I^{(\alpha)}} + (\boldsymbol{\omega}^{(\alpha)} \cdot \mathbf{J}^{(\alpha)}) - T\mathfrak{S}^{(\alpha)} \right]. \quad (52)$$

The *power input* driving the process (source/sink term) is given by:

$$\mathcal{P}' \equiv \frac{1}{2} \sum_{\alpha} (\nabla \cdot \mathbf{v}^{(\alpha)}) \left[\frac{|\mathbf{J}^{(\alpha)}|^2}{2I^{(\alpha)}} - (\boldsymbol{\omega} \cdot \mathbf{J}^{(\alpha)}) + T\mathfrak{S}^{(\alpha)} \right], \quad (53)$$

and the total energy is defined by summing up the different contributions:

$$\mathcal{U} = \mathfrak{I} + \mathfrak{P}_g + \mathfrak{P}_v + \mathfrak{P}_{rot}. \quad (54)$$

Here, the term $T\mathfrak{S}^{(\alpha)}$ represents the thermal energy associated to specie (α) , equal to $-\Delta F$, the free energy of the physical system. For a system in contact with a reservoir at constant temperature this is the maximum work which can be done, the free energy tends to decrease for a system in thermal contact with a heat reservoir. In particular, notice that when the angular velocity ω is multiplied by Eq. 40, it is obtained the driving power. It is worth to point out the presence of the term $\omega \cdot \mathbf{J}$ which plays an analogue role to the slip in electrical induction motors, that is, the lag between the rotor speed and the magnetic field's speed, provided by the stator's rotating speed. Furthermore, we may notice that the power input depends on the fluid compressibility $\nabla \cdot \mathbf{v}$. This means that compressibility is a factor that determines the amount of transported angular momentum through the stress-tensor τ_{ij} , and may be responsible of new driving mechanism in addition to the well-known MRI. The driving energy of the rotating system can be expressed in the form:

$$\mathcal{E}_{driv} = \frac{J^2}{2I_p} - (\omega \cdot \mathbf{J}) - \Delta F. \quad (55)$$

Next, we will discuss several examples, illustrating the application of the variational method.

A. Power extraction from an hurricane

A hurricane is a natural airborne structure that converts all forms of kinetic and potential energy by means of the transport of angular momentum from the inner core to the outer regions, conveyed either directly by moving matter, or by non-material stresses such as those exerted by electric fields or magnetic fields³⁹. We may apply Eq. 53 to this specific problem and obtain:

$$\mathcal{P}' = \omega \left(\frac{J^2}{2I_p} - \omega \cdot \mathbf{J} - \Delta F \right) = -\frac{\partial U_e}{\partial t} \quad (56)$$

or

$$\mathcal{P}' = \omega \mathcal{E}_{driv}. \quad (57)$$

Let us consider specifically the case of a hurricane in an axisymmetric configuration, with $J = \omega I$. Likely we have $\omega^2 I^2 / I_p \gg \omega^2 I + \Delta F$. We can envisage a simple and rough model of a hurricane, with total mass M and radius R , turning like a solid cylinder, and $I = MR^2/2$. Hence, the total power driving the hurricane is given by

$$\mathcal{P}' = \frac{1}{8} \frac{\omega^3}{2} \frac{M^2 R^4}{I_p} \propto \omega^3 \frac{M^2 R^4}{I_p}, \quad (58)$$

or, as a function of the fluid density ρ , in the form:

$$\mathcal{P}' = \frac{\pi^2}{4} \omega^3 \frac{R^6 L^2}{I_p} \rho^2. \quad (59)$$

Our result shows the same kind of dependency as shown by Chow & Chey⁴⁰, and, in particular, it shows that the intrinsic inertial momentum of the particles constituting the fluid plays a substantial role.

B. Radiative heating of planetary atmosphere and net angular momentum

It is experimentally shown⁴¹ that periodic radiative heating of the Earth's atmosphere transmits angular momentum to it due to the Earth-atmosphere coupling through frictional interactions⁴². The images sent by ESA's Venus Express confirms this fact on Venus (Earth's twin planet), with the presence of a "double-eye" atmospheric vortex at the planet's south pole and high velocity winds whirling westwards around the planet completing a full rotation in four days.

Schubert and Withehead's⁴¹ did an experiment with the purpose to provide an explanation for the high velocities of apparent cloud formation, in the upper atmosphere of Venus. In this experiment, a bunsen flame rotating below a cylindrical annulus, filled with liquid mercury, induced the rotation of the liquid mercury, in a counter-direction to that of the rotating flame. The speed of the flame was 1 mm/s, and the temperature of the mercury increased from room temperature, at the rate of about 3°C per minute. After 5 minutes, a steady-state flow was established, with the mercury rotating in the counter-direction of the flame, with a speed about 4 mm/s. Considering that the entire mercury liquid as a spinning

body, we can estimate from Eq. 49 that

$$\frac{d}{dr} \left(\frac{I_p \omega'^2}{2} - \frac{I_p \omega^2}{2} + T\mathfrak{S} \right) > 0 \Rightarrow I\Delta(\omega' + \omega)(\omega' - \omega) + \Delta T\mathfrak{S} > 0. \quad (60)$$

Hence, the following result is obtained:

$$\omega' \Delta\omega \gtrsim -\frac{1}{2} \frac{\Delta T\mathfrak{S}}{I_p}. \quad (61)$$

Here, ω is the angular speed imposed on the system; ω' the induced angular speed, due to a sustained source of energy; $\Delta\omega \equiv \omega' - \omega$. We use the following tabulated data: $\mathfrak{S} = 16.6 \text{ J.K}^{-1}\text{mol}^{-1}$ for mercury at temperature of the experiment, $\rho \approx 13.6 \times 10^{-3} \text{ kg/m}^3$, and consider that the volume of mercury is contained in the channel of the experimental apparatus, forming a rim with an average radius $R = 30 \text{ cm}$. With Eq. 49 we estimate that, after one minute, the speed must be of the order 3.7 mm/s . It is clear that the sense of rotation, and its speed, depend on the latent heat stored in the planetary atmosphere and its temperature (through ΔT). Although the initial assumptions taken in the present formulation require further research, considering that interactive terms with the medium, conveyed for example through thermal diffusion coefficient, are not covered by Eq. 49, the agreement is reasonable, and offers a possible explanation for this effect. Once again, the problem of radiative atmosphere heating reveals the interesting interplay between energy and entropy, each one attempting a different equilibrium condition.

V. CONCLUSION

The information-theoretic method proposed in this paper constitute an alternate approach applying the concept of maximizing entropy to the problem of out of equilibrium physical systems. It bears some resemblance with Hamiltonian formulation of dynamics, which expresses first-order constraints of the Hamiltonian H in a $2n$ dimensional phase space, revealing the same formal symplectic structure shared by classical mechanics and thermodynamics. Although the simplifying assumption of isothermal system rules out its ability to explain accurately such problems as, for example, the coherent transport of angular momentum in astrophysics, or a certain kind of laboratory devices (e.g., the Ranque-Hilsch effect), the present method open to understanding certain trends, in particular, predicting

theoretically for electromagnetic-gravitational vortex the angular momentum forced transport outwardly from a symmetry axis of rotation. The kind of Umov-Poynting theorem obtained is revealing of the interplay between entropy and energy, the energy tending to a minimum, and the entropy tending to a maximum, explaining the formation of physical structures. In particular, it was shown that compressibility is an important property in the transport of angular momentum, and a possible driving mechanism for instability. This development is seen to be advantageous and opens options for systematic improvements.

ACKNOWLEDGMENTS

We are grateful to Professor Roman W. Jackiw, MIT Center for Theoretical Physics, for keeping us informed of the fundamental paper referred to in [13]. The author gratefully acknowledge partial financial support by the International Space Science Institute (ISSI) as visiting scientist, and express special thanks to Professor Roger Maurice-Bonnet.

REFERENCES

- ¹V. W. Ekman, *On the Influence of Earth Rotation on Ocean Currents*, Arkiv. Math. Astro. Fys. **2** 1-52 (1905)
- ²H. P. Greenspan, *The Theory of Rotating Fluids*, (Breukelen Press, Brookline-MA, 1990)
- ³Steven A. Balbus, *Enhanced Angular Momentum Transport in Accretion Disks*, Annu. Rev. Astron. Astrophys. **41** 555 (2003)
- ⁴K. C. Chow and Kwing L. Chan, *Angular Momentum Transports by Moving Spiral Waves*, J. Atmos. Sci. **60** 2004 (2003)
- ⁵Hantao Ji, Philipp Kronberg, Stewart C. Prager, and Dmitri A. Uzdensky, Princeton Plasma Physics Laboratory Report PPPL-4311, May 2008
- ⁶C. F. Curtiss, *Kinetic Theory of Nonspherical Molecules*, J. Chem. Phys. **24** (2) pp. 225-241 (1956)
- ⁷Peter M. Livingstone and C. F. Curtiss, *Kinetic Theory of Nonspherical Molecules. IV. Angular Momentum Transport Coefficient* J. Chem. Phys. **31** (6) pp. 1643-1645 (1959)
- ⁸E. T. Jaynes, Phys. Rev. **106** (4) 620 (1957);
- ⁹M. J. Pinheiro, Europhys. Lett. **57**, 305 (2002)

- ¹⁰M. J. Pinheiro, *Physica Scripta* **70** (2-3) 86 (2004)
- ¹¹Mark A. Peterson, *Am. J. Phys.* **47**(6) 488 (1979)
- ¹²V. P. Pavlov and V. M. Sergeev, *Theor. Math. Phys.* **157**(1) 1484 (2008)
- ¹³O. Éboli, R. Jackiw and So-Young Pi, *Phys. Rev. D* **37**(12) 3557-3581 (1988)
- ¹⁴L. Landau and E. M. Lifschitz, *Physique Statistique*, p. 134, (Mir, Moscow, 1960)
- ¹⁵Erik Verlinde, *JHEP* **2011** (4) 29 (2011)
- ¹⁶P. Glansdorff and I. Prigogine, *Structure, Stabilité et Fluctuations* (Masson Éditeurs, Paris, 1971)
- ¹⁷A. I. Khinchin, *Mathematical Foundations of Statistical Mechanics* (Dover Publications, New York, 1949)
- ¹⁸Richard L. Liboff, *Kinetic Theory - Classical, Quantum, and Relativistic Descriptions* (Prentice-Hall, New Jersey, 1990)
- ¹⁹Kerson Huang, *Statistical Mechanics* (Wiley, Hoboken, 1987)
- ²⁰S. Chandrasekhar, *Selected Papers, Vol.4 - Plasma Physics, Hydrodynamic and Hydromagnetic stability, and Applications of the Tensor-Virial Theorem*, (University of Chicago Press, Chicago, 1989)
- ²¹S. Peter Gary, *Theory of Space Plasma Microinstabilities* (Cambridge University Press, Cambridge, 1993)
- ²²Weston M. Stacey *Fusion Plasma Physics* (Wiley-VCH, Weinheim, 2005)
- ²³R. M. Kiehn, *Ann. Fond. L. de Broglie* **32** (2-3) 389-408 (2007)
- ²⁴V. Onoichin and T. E. Phipps Jr, *Eur. J. Elect. Phen.* **3**(2) 256 (2003)
- ²⁵L. C. Steinhauer and A. Ishida, *Phys. Rev. Lett.* **79** (18) 3423 (1997)
- ²⁶Horace Lamb, *Higher Mechanics*, (Cambridge University Press, London, 1920)
- ²⁷B. H. Lavende, *Phys. Rev. A* **9** (2) 929 (1974)
- ²⁸Bishwanath Chakraborty, *Principles of Plasma Mechanics*, p. 286 (Wiley, New Delhi, 1978)
- ²⁹John M. Wilcox, *Rev. Mod. Phys.* **31**(4) pp. 1045-1051 (1959); O. Anderson, W. R. Baker, A. Bratenahl, H. P. Furth, and W. B. Kunkel, *J. Appl. Phys.* **30** (2) pp. 188-196 (1959)
- ³⁰Joseph Slepian, *Rev. Mod. Phys.* **32**(4) pp. 1032-1033 (1960)
- ³¹O. A. Hurricane, *Phys. Plasmas* **5**(6) pp. 2197-2202 (1998)
- ³²Antoni Klajn, *IEEE Trans. Plasma Science* **27** (4) 977 (1999)

- ³³*Power circuit breaker: theory and design* Ed. C. H. Flurschein (Peter Peregrinus, London, 1985)
- ³⁴R. L. Boxman, Philip J. Martin and David Sanders, *Handbook of vacuum arc science and technology: fundamentals and applications* (Noyes Publications, New Jersey, 1995)
- ³⁵Steven A. Balbus and John F. Hawley, *A powerful local shear instability in weakly magnetized disks. I. Linear Analysis*, Astrophys. J. **376** 214-222 (1991)
- ³⁶E. P. Velikhov, *Magnetic Geodynamics* JETP Letters **82** (11) pp. 690-695 (2005)
- ³⁷E. P. Velikhov, Soviet Phys. - JETP, **36**, 1398 (1959)
- ³⁸S. Chandrasekhar, Proc. Nat. Acad. Sci. **46**, 253 (1960)
- ³⁹V. Boořirevich, Ya. Freibergs, E. I. Shilova and E. V. Shcherbinin, *Electrically Induced Vortical Flows* (Kluwer Academic Publishers, Dordrecht, 1989)
- ⁴⁰K. C. Chow and Kwing L. Chan, J. Atmos. Sci. **60** 2004-2009 (2003)
- ⁴¹G. Schubert and J. A. Whitehead, Science, New Series, **163** (3862) pp. 71-72 (1969)
- ⁴²A. Chandrasekar, *Basics of Atmospheric Science* (PHI Learning Private Limited, New Deli, 2010)