Positive Least Energy Solutions and Phase Separation for Coupled Schrödinger Equations with Critical Exponent: Higher Dimensional Case*

Zhijie Chen, Wenming Zou

Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China

Abstract

We study the following nonlinear Schrödinger system which is related to Bose-Einstein condensate:

$$\begin{cases} -\Delta u + \lambda_1 u = \mu_1 u^{2^* - 1} + \beta u^{\frac{2^*}{2} - 1} v^{\frac{2^*}{2}}, & x \in \Omega, \\ -\Delta v + \lambda_2 v = \mu_2 v^{2^* - 1} + \beta v^{\frac{2^*}{2} - 1} u^{\frac{2^*}{2}}, & x \in \Omega, \\ u \ge 0, v \ge 0 \text{ in } \Omega, & u = v = 0 \text{ on } \partial\Omega. \end{cases}$$

Here $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain, $2^* := \frac{2N}{N-2}$ is the Sobolev critical exponent, $-\lambda_1(\Omega) < \lambda_1, \lambda_2 < 0$, $\mu_1, \mu_2 > 0$ and $\beta \neq 0$, where $\lambda_1(\Omega)$ is the first eigenvalue of $-\Delta$ with the Dirichlet boundary condition. When $\beta = 0$, this is just the well-known Brezis-Nirenberg problem. The special case N=4 was studied by the authors in (Arch. Ration. Mech. Anal. 205: 515-551, 2012). In this paper we consider the higher dimensional case $N \geq 5$. It is interesting that we can prove the existence of a positive least energy solution (u_β, v_β) for any $\beta \neq 0$ (which can not hold in the special case N=4). We also study the limit behavior of (u_β, v_β) as $\beta \to -\infty$ and phase separation is expected. In particular, $u_\beta - v_\beta$ will converge to sign-changing solutions of the Brezis-Nirenberg problem, provided $N \geq 6$. In case $\lambda_1 = \lambda_2$, the classification of the least energy solutions is also studied. It turns out that some quite different phenomena appear comparing to the special case N=4.

^{*}Supported by NSFC (11025106). E-mail: chenzhijie1987@sina.com(Chen); wzou@math.tsinghua.edu.cn (Zou)

1 Introduction

In this paper we consider the following coupled nonlinear Schrödinger equations:

$$\begin{cases}
-\Delta u + \lambda_1 u = \mu_1 u^{2p-1} + \beta u^{p-1} v^p, & x \in \Omega, \\
-\Delta v + \lambda_2 v = \mu_2 v^{2p-1} + \beta v^{p-1} u^p, & x \in \Omega, \\
u \ge 0, v \ge 0 \text{ in } \Omega, & u = v = 0 \text{ on } \partial\Omega,
\end{cases}$$
(1.1)

where $\Omega = \mathbb{R}^N$ or $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain, p > 1 and $p \le 2^*/2$ if $N \ge 3$, $\mu_1, \mu_2 > 0$ and $\beta \ne 0$ is a coupling constant. In the case p = 2, the cubic system (1.1) appears in many physical problems, especially in nonlinear optics and Bose-Einstein condensation. We refer for this to [2, 19, 20, 23], which also contain information about the physical relevance of non-cubic nonlinearities. In the subcritical case $p < 2^*/2$, the existence of solutions have received great interest recently, see [4, 7, 18, 24, 25, 27, 29, 34] and references therein.

All the papers mentioned above deal with the subcritical case. To the best of our knowledge, there are no existence results for (1.1) in the critical case $2p = 2^*$ in the literature. Critical exponent problems, which have received great attention in the past thirty years since the cerebrated work by Brezis and Nirenberg [9], are very interesting in view of mathematics. Recently, the authors studied the special critical case p = 2 and N = 4 in [16].

In this paper, we study the existence and properties of least energy solutions to (1.1) in the higher dimensional case. In the sequel we assume that

$$N \ge 5$$
 and $2p = 2^*$. (1.2)

It turns out that different phenomena happen comparing to the special case N=4 ([16]), see Remarks 1.1, 1.3, 1.4 and 1.5 below. If $\Omega=\mathbb{R}^N$ and (u,v) is any a solution of (1.1), then by the Pohozaev identity, it is easy to get that $\int_{\mathbb{R}^N} \lambda_1 u^2 + \lambda_2 v^2 dx = 0$, so $(u,v) \equiv (0,0)$ if $\lambda_1 \lambda_2 > 0$. Therefore, in the sequel we assume that $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain. We call a solution (u,v) nontrivial if both $u \not\equiv 0$ and $v \not\equiv 0$, a solution (u,v) semi-trivial if (u,v) is type of (u,0) or (0,v). We are concerned with nontrivial solutions of (1.1).

Let $\lambda_1(\Omega)$ be the first eigenvalue of $-\Delta$ with the Dirichlet boundary condition. Clearly (1.1) has semi-trivial solutions $(u_{\mu_1}, 0)$ and $(0, u_{\mu_2})$, where u_{μ_i} is a positive least energy solution of the well-known Brezis-Nirenberg problem

$$-\Delta u + \lambda_i u = \mu_i |u|^{2^* - 2} u, \ u \in H_0^1(\Omega), \tag{1.3}$$

if $-\lambda_1(\Omega) < \lambda_1, \lambda_2 < 0$ (see [9]). Hence, system (1.1) can be seen as a critically coupled Brezis-Nirenberg problem. As we will see in Theorems 1.4-1.5, system (1.1) is closely related to the Brezis-Nirenberg problem (1.3). The Brezis-Nirenberg problem (1.3) has been studied intensively, and we refer the readers to [12, 13, 15, 17, 32] and references therein.

Denote $H:=H^1_0(\Omega)\times H^1_0(\Omega)$. It is well known that solutions of (1.1) correspond to the critical points of C^1 functional $E:H\to\mathbb{R}$ given by

$$E(u,v) = \frac{1}{2} \int_{\Omega} (|\nabla u|^2 + \lambda_1 u^2) + \frac{1}{2} \int_{\Omega} (|\nabla v|^2 + \lambda_2 v^2)$$

$$-\frac{1}{2p}\int_{\Omega} (\mu_1|u|^{2p} + 2\beta|u|^p|v|^p + \mu_2|v|^{2p}). \tag{1.4}$$

We say a solution (u, v) of (1.1) is a least energy solution, if (u, v) is nontrivial and $E(u, v) \leq E(\varphi, \psi)$ for any other nontrivial solution (φ, ψ) of (1.1). As in [24], we define a Nehari type manifold

$$\mathcal{M} = \{(u, v) \in H : u \not\equiv 0, v \not\equiv 0, E'(u, v)(u, 0) = E'(u, v)(0, v) = 0\}.$$

Then any nontrivial solutions of (1.1) belong to \mathcal{M} . Similarly as [16], it is trivial to see that $\mathcal{M} \neq \emptyset$. Define the least energy

$$B := \inf_{(u,v) \in \mathcal{M}} E(u,v) = \inf_{(u,v) \in \mathcal{M}} \frac{1}{N} \int_{\Omega} (|\nabla u|^2 + \lambda_1 u^2 + |\nabla v|^2 + \lambda_2 v^2) \, dx. \quad (1.5)$$

First we consider the symmetric case $-\lambda_1(\Omega) < \lambda_1 = \lambda_2 = \lambda < 0$. By [9] the Brezis-Nirenberg problem

$$-\Delta u + \lambda u = |u|^{2^* - 2} u, \ u \in H_0^1(\Omega)$$
 (1.6)

has a positive least energy solution ω with energy

$$B_1 := \frac{1}{N} \int_{\Omega} (|\nabla \omega|^2 + \lambda \omega^2) \, dx = \frac{1}{N} \int_{\Omega} \omega^{2^*} \, dx. \tag{1.7}$$

Moreover,

$$\int_{\Omega} (|\nabla u|^2 + \lambda u^2) \, dx \ge (NB_1)^{2/N} \left(\int_{\Omega} |u|^{2^*} \, dx \right)^{2/2^*}, \quad \forall \ u \in H_0^1(\Omega).$$
 (1.8)

Consider the following nonlinear problem $(p = \frac{N}{N-2} < 2 \text{ since } N \ge 5)$

$$\begin{cases} \mu_1 k^{p-1} + \beta k^{p/2-1} l^{p/2} = 1, \\ \beta k^{p/2} l^{p/2-1} + \mu_2 l^{p-1} = 1, \\ k > 0, \ l > 0. \end{cases}$$
 (1.9)

We will prove in Lemma 2.1 that there exists (k_0, l_0) such that

$$(k_0, l_0)$$
 satisfies (1.9) and $k_0 = \min\{k : (k, l) \text{ is a solution of } (1.9)\}.$ (1.10)

Our first result deals with the symmetric case $\lambda_1 = \lambda_2$.

Theorem 1.1. Assume that $-\lambda_1(\Omega) < \lambda_1 = \lambda_2 = \lambda < 0$. Let (k_0, l_0) in (1.10). Then for any $\beta > 0$, $(\sqrt{k_0}\omega, \sqrt{l_0}\omega)$ is a positive solution of (1.1). Moreover, if $\beta \geq \frac{2}{N-2} \max\{\mu_1, \mu_2\}$, then $E(\sqrt{k_0}\omega, \sqrt{l_0}\omega) = B$, that is, $(\sqrt{k_0}\omega, \sqrt{l_0}\omega)$ is a positive least energy solution of (1.1).

Remark 1.1. (1) In the special case N=4 and $2p=2^*$, [16, Theorem 1.1] said that (1.1) has no nontrivial nonnegative solution if $\beta \in [\min\{\mu_1, \mu_2\}, \max\{\mu_1, \mu_2\}]$ and $\mu_1 \neq \mu_2$. Therefore, the general case $N \geq 5$ is quite different from the case N=4. As we will see in Section 2, the proof of Theorem 1.1 is much more delicate than [16] because of the nonlinearity of problem (1.9).

(2) Similarly as in [16, Remark 1.1], we can prove that, if Ω is starshaped, the assumption $-\lambda_1(\Omega) < \lambda < 0$ in Theorem 1.1 is optimal.

Our second result deals with the classification of the least energy solutions.

Theorem 1.2. Let assumptions in Theorem 1.1 hold. There exists $\beta_0 \geq \frac{2}{N-2} \max\{\mu_1, \mu_2\}$ determined by (μ_1, μ_2) , and assume that $\beta > \beta_0$. Let (u, v) be any a positive least energy solution of (1.1), then $(u, v) = (\sqrt{k_0}U, \sqrt{l_0}U)$, where U is a positive least energy solution of (1.6). In particular, the positive least energy solution of (1.1) is unique if $\Omega \subset \mathbb{R}^N$ is a ball.

Remark 1.2. (1) We can give a precise definition of β_0 (see (4.2) in Section 4). In particular, if $\mu_1 = \mu_2$, then $\beta_0 = \frac{2}{N-2} \max\{\mu_1, \mu_2\}$. (2) For the case p = 2 and $N \leq 3$, some uniqueness results about system

(2) For the case p=2 and $N \leq 3$, some uniqueness results about system (1.1) were introduced in [36]. However, their proofs heavily depend on p=2, and so can not work here.

Now, let us consider the general case $-\lambda_1(\Omega) < \lambda_1, \lambda_2 < 0$. Without loss of generality, we may assume that $\lambda_1 \leq \lambda_2$.

Theorem 1.3. Assume that $-\lambda_1(\Omega) < \lambda_1 \le \lambda_2 < 0$. Then system (1.1) has a positive least energy solution (u, v) with E(u, v) = B for any $\beta \ne 0$.

Remark 1.3. For the general case $\lambda_1 \leq \lambda_2$, when N=4 and $2p=2^*$, [16, Theorem 1.3] said that (1.1) has a positive least energy solution for any

$$\beta \subset (-\infty, 0) \cup (0, \beta_1) \cup (\beta_2, +\infty),$$

where β_i , i = 1, 2 are some positive constants satisfying

$$\beta_1 \le \min\{\mu_1, \mu_2\} \le \max\{\mu_1, \mu_2\} \le \beta_2.$$

That is, we do not know whether the least energy solution exists or not if $\beta \in [\beta_1, \beta_2]$ (In the symmetric case $\lambda_1 = \lambda_2$, Remark 1.1-(1) says that nontrivial positive solutions do not exist if $\beta \in [\min\{\mu_1, \mu_2\}, \max\{\mu_1, \mu_2\}]$ and $\mu_1 \neq \mu_2$). Comparing this with Theorem 1.3, it turns out that the general case $N \geq 5$ is completely different from the special case N = 4.

Now, we study the limit behavior of the positive least energy solutions in the repulsive case $\beta \to -\infty$. It is expected that components of the limiting profile tend to separate in different regions of the underlying domain Ω . This phenomena, called *phase separation*, has been well studied for L^{∞} -bounded positive solutions of system (1.1) in subcritical case $2p < 2^*$ by [34, 35, 28]. The critical case N = 4 and p = 2 was studied by [16]. For other kinds of elliptic systems with strong competition, phase separation has also been well studied, we refer to [10, 11, 14] and references therein. Denote $\{u > 0\} := \{x \in \Omega : u(x) > 0\}$ and $u^{\pm} := \max\{\pm u, 0\}$. Then we have the following result.

Theorem 1.4. Assume that $-\lambda_1(\Omega) < \lambda_1 \leq \lambda_2 < 0$. Let $\beta_n < 0$, $n \in \mathbb{N}$, satisfy $\beta_n \to -\infty$ as $n \to \infty$, and (u_n, v_n) be the positive least energy solutions of (1.1) with $\beta = \beta_n$ which exists by Theorem 1.3. Then $\int_{\Omega} \beta_n u_n^p v_n^p dx \to 0$ as $n \to \infty$, and passing to a subsequence, one of the following conclusions holds.

(1) $u_n \to u_\infty$ strongly in $H_0^1(\Omega)$ and $v_n \to 0$ weakly in $H_0^1(\Omega)$ (so $v_n \to 0$ for almost every $x \in \Omega$), where u_∞ is a positive least energy solution of

$$-\Delta u + \lambda_1 u = \mu_1 |u|^{2^* - 2} u, \ u \in H_0^1(\Omega).$$

(2) $v_n \to v_\infty$ strongly in $H_0^1(\Omega)$ and $u_n \to 0$ weakly in $H_0^1(\Omega)$ (so $u_n \to 0$ for almost every $x \in \Omega$), where v_∞ is a positive least energy solution of

$$-\Delta v + \lambda_2 v = \mu_2 |v|^{2^* - 2} v, \ v \in H_0^1(\Omega).$$

(3) $(u_n, v_n) \to (u_\infty, v_\infty)$ strongly in H and $u_\infty \cdot v_\infty \equiv 0$, where $u_\infty \in C(\overline{\Omega})$ is a positive least energy solution of

$$-\Delta u + \lambda_1 u = \mu_1 |u|^{2^* - 2} u, \ u \in H_0^1(\{u_\infty > 0\}),$$

and $v_{\infty} \in C(\overline{\Omega})$ is a positive least energy solution of

$$-\Delta v + \lambda_2 v = \mu_2 |v|^{2^* - 2} v, \ v \in H_0^1(\{v_\infty > 0\}).$$

Furthermore, both $\{v_{\infty} > 0\}$ and $\{u_{\infty} > 0\}$ are connected domains, and $\{v_{\infty} > 0\} = \Omega \setminus \overline{\{u_{\infty} > 0\}}$.

In particular, if $N \ge 6$, then only conclusion (3) holds, and $u_{\infty} - v_{\infty}$ is a least energy sign-changing solution to problem

$$-\Delta u + \lambda_1 u^+ - \lambda_2 u^- = \mu_1 (u^+)^{2^* - 1} - \mu_2 (u^-)^{2^* - 1}, \quad u \in H_0^1(\Omega). \tag{1.11}$$

Here a sign-changing solution u of (1.11) is called a least energy sign-changing solution, if u attains the minimal functional energy among all sign-changing solutions of (1.11). As an application of Theorem 1.4, we turn to consider the Brezis-Nirenberg problem

$$-\Delta u + \lambda_1 u = \mu_1 |u|^{2^* - 2} u, \ u \in H_0^1(\Omega), \tag{1.12}$$

where $-\lambda_1(\Omega) < \lambda_1 < 0$. Its corresponding functional is

$$J(u) := \frac{1}{2} \int_{\Omega} (|\nabla u|^2 + \lambda_1 u^2) \, dx - \frac{1}{2^*} \int_{\Omega} \mu_1 |u|^{2^*} \, dx, \ u \in H_0^1(\Omega).$$

Then we have the following result.

Theorem 1.5. Assume $N \geq 6$. Let (u_{∞}, v_{∞}) be in Theorem 1.4 in the symmetric case where $\lambda_2 = \lambda_1$ and $\mu_2 = \mu_1$. Then $u_{\infty} - v_{\infty}$ is a least energy sign-changing solution of (1.12), and

$$J(u_{\infty} - v_{\infty}) < B_{\mu_1} + \frac{1}{N} \mu_1^{-\frac{N-2}{2}} S^{\frac{N}{2}}, \tag{1.13}$$

where B_{μ_1} is the least energy of problem (1.12) (see (3.1) in Section 3).

- Remark 1.4. (i) Theorem 1.4 has been proved for the special case N=4 and $2p=2^*$ by the authors([16]), where we raised an open question: Can one show that only conclusion (3) holds? The reviewer of [16] pointed out that this question may be related to the existence of sign-changing solutions to the Brezis-Nirenberg problem (1.12). Motivated by the reviewer's comment, it is natural for us to consider (1.1) under assumption (1.2) in this paper. Here in the case $N \geq 6$, we exclude conclusions (1)-(2) and verifies the reviewer's comment successfully, and so system (1.1) is closely related to the Brezis-Nirenberg problem. Unfortunately, we do not know whether only Theorem 1.4-(3) holds for N=4,5, which still remains as an interesting open question.
- (ii) In the proof of Theorem 1.4-(3), a key point is to prove the continuity of u_{∞} and v_{∞} . We remark here that, our proof of the continuity of u_{∞} and v_{∞} is completely different from that in [16] for the special case N=4, and can also be used to the special case N=4.
- (iii) The existence of least energy sign-changing solutions to the Brezis-Nirenberg problem (1.12) in the case N ≥ 6 was proved in [13] in 1986. Here, Theorem 1.5 is a direct corollary of Theorem 1.4, and so the proof of Theorem 1.5 is completely different from [13].

Since the nonlinearity and the coupling term are both critical in (1.1), the existence of nontrivial solutions of (1.1) depends heavily on the existence of the least energy solution of the following limit problem

$$\begin{cases}
-\Delta u = \mu_1 |u|^{2p-2} u + \beta |u|^{p-2} u |v|^p, & x \in \mathbb{R}^N, \\
-\Delta v = \mu_2 |v|^{2p-2} v + \beta |v|^{p-2} v |u|^p, & x \in \mathbb{R}^N, \\
u, v \in D^{1,2}(\mathbb{R}^N),
\end{cases} (1.14)$$

where $D^{1,2}(\mathbb{R}^N) := \{ u \in L^2(\mathbb{R}^N) : |\nabla u| \in L^2(\mathbb{R}^N) \}$ with norm $||u||_{D^{1,2}} := (\int_{\mathbb{R}^N} |\nabla u|^2 dx)^{1/2}$. Let S be the sharp constant of $D^{1,2}(\mathbb{R}^N) \hookrightarrow L^{2^*}(\mathbb{R}^N)$

$$\int_{\mathbb{R}^N} |\nabla u|^2 \, dx \ge S \left(\int_{\mathbb{R}^N} |u|^{2^*} \, dx \right)^{\frac{2}{2^*}}. \tag{1.15}$$

For $\varepsilon > 0$ and $y \in \mathbb{R}^N$, we consider the Aubin-Talenti instanton [1, 33] $U_{\varepsilon,y} \in D^{1,2}(\mathbb{R}^N)$ defined by

$$U_{\varepsilon,y}(x) := \left[N(N-2)\right]^{\frac{N-2}{4}} \left(\frac{\varepsilon}{\varepsilon^2 + |x-y|^2}\right)^{\frac{N-2}{2}}.$$
 (1.16)

Then $U_{\varepsilon,y}$ satisfies $-\Delta u = |u|^{2^*-2}u$ in \mathbb{R}^N and

$$\int_{\mathbb{R}^N} |\nabla U_{\varepsilon,y}|^2 dx = \int_{\mathbb{R}^N} |U_{\varepsilon,y}|^{2^*} dx = S^{N/2}.$$
 (1.17)

Furthermore, $\{U_{\varepsilon,y}: \varepsilon>0, y\in\mathbb{R}^N\}$ contains all positive solutions of the equation $-\Delta u=|u|^{2^*-2}u$ in \mathbb{R}^N .

Clearly (1.14) has semi-trivial solutions $(\mu_1^{-\frac{N-2}{4}}U_{\varepsilon,y},0)$ and $(0,\mu_2^{\frac{N-2}{4}}U_{\varepsilon,y})$. Here, we are only interested in nontrivial solutions of (1.14). Define $D:=D^{1,2}(\mathbb{R}^N)\times D^{1,2}(\mathbb{R}^N)$ and a C^1 functional $I:D\to\mathbb{R}$ given by

$$I(u,v) := \frac{1}{2} \int_{\mathbb{R}^N} (|\nabla u|^2 + |\nabla v|^2) - \frac{1}{2p} \int_{\mathbb{R}^N} (\mu_1 |u|^{2p} + 2\beta |u|^p |v|^p + \mu_2 |v|^{2p}).$$
(1.18)

As in [24], we consider the set

$$\mathcal{N} = \{(u, v) \in D : u \not\equiv 0, v \not\equiv 0, I'(u, v)(u, 0) = I'(u, v)(0, v) = 0\}.$$

Then any nontrivial solutions of (1.14) belong to \mathcal{N} . Similarly $\mathcal{N} \neq \emptyset$. We set

$$A := \inf_{(u,v) \in \mathcal{N}} I(u,v) = \inf_{(u,v) \in \mathcal{N}} \frac{1}{N} \int_{\mathbb{R}^N} (|\nabla u|^2 + |\nabla v|^2) \, dx. \tag{1.19}$$

Then we have the following theorem, which plays an important role in the proof of Theorem 1.3.

Theorem 1.6. (1) If $\beta < 0$, then A is not attained.

- (2) If $\beta > 0$, then (1.14) has a positive least energy solution (U, V) with I(U, V) = A, which is radially symmetric decreasing. Moreover,
 - (2-1) if $\beta \geq \frac{2}{N-2} \max\{\mu_1, \mu_2\}$, then $I(\sqrt{k_0}U_{\varepsilon,y}, \sqrt{l_0}U_{\varepsilon,y}) = A$, where (k_0, l_0) in (1.10). That is, $(\sqrt{k_0}U_{\varepsilon,y}, \sqrt{l_0}U_{\varepsilon,y})$ is a positive least energy solution of (1.14).
 - (2-2) there exists $0 < \beta_1 \le \frac{2}{N-2} \max\{\mu_1, \mu_2\}$, and for any $0 < \beta < \beta_1$, there exists a solution $(k(\beta), l(\beta))$ of (1.9), such that

$$I(\sqrt{k(\beta)}U_{\varepsilon,u}, \sqrt{l(\beta)}U_{\varepsilon,u}) > A = I(U, V).$$

That is, $(\sqrt{k(\beta)}U_{\varepsilon,y}, \sqrt{l(\beta)}U_{\varepsilon,y})$ is a different positive solution of (1.14) with respect to (U,V).

Remark 1.5. In the case N=4 and $2p=2^*$, [16, Theorem 1.5] said that (1.14) has no nontrivial nonnegative solution if $\beta \in [\min\{\mu_1, \mu_2\}, \max\{\mu_1, \mu_2\}]$ and $\mu_1 \neq \mu_2$; $(\sqrt{k(\beta)}U_{\varepsilon,y}, \sqrt{l(\beta)}U_{\varepsilon,y})$ is a positive least energy solution of (1.14) if $0 < \beta < \min\{\mu_1, \mu_2\}$. Hence the general case $N \geq 5$ is completely different from the case N=4. As we will see in Section 2, the idea of proving Theorem 1.6-(2) in case $0 < \beta < \frac{2}{N-2} \max\{\mu_1, \mu_2\}$, which also works for the case $\beta \geq \frac{2}{N-2} \max\{\mu_1, \mu_2\}$, is completely different from that in case N=4 ([16]).

We can also study the uniqueness of the positive least energy solutions of (1.14) just as Theorem 1.2.

Theorem 1.7. Let β_0 be in Theorem 1.2 and assume that $\beta > \beta_0$. Let (u, v) be any a positive least energy solution of (1.14). Then $(u, v) = (\sqrt{k_0}U_{\varepsilon, y}, \sqrt{l_0}U_{\varepsilon, y})$ for some $\varepsilon > 0$ and $y \in \mathbb{R}^N$.

The rest of this paper proves these theorems, and some ideas of the proofs are similar to those in [16]. However, as pointed out above, the general case $N \geq 5$ is quite different from N=4, and some new ideas are needed. We give some notations here. Throughout this paper, we denote the norm of $L^q(\Omega)$ by $|u|_q = (\int_{\Omega} |u|^q \, dx)^{\frac{1}{q}}$, the norm of $H^1_0(\Omega)$ by $||u|| = |\nabla u|_2$ and positive constants (possibly different) by C. The paper is organized as follows. Theorems 1.1 and 1.6 are proved in Section 2, and we will see that these proofs are more delicate than those in case N=4 ([16]). In Section 3, we use Nehari manifold approach and Ekeland variational principle to prove Theorem 1.3 for the case $\beta < 0$, and use mountain pass argument to Theorem 1.3 for the case $\beta > 0$. In Section 4, we use an elementary approach to prove Theorems 1.2 and 1.7. Finally, we use energy estimate methods to prove Theorems 1.4 and 1.5 in Section 5, where some different ideas are needed.

After the completion of this paper (see arXiv:1209.2522v1 for the original version), we learned that (1.1) has also been studied in [22], where the author showed the existence of least energy solutions for $\beta > \beta_0 > 0$, where β_0 is an unknown constant. For $|\beta|$ sufficiently small, the existence of nontrivial solutions was studied in [22] via a perturbation method, but the solutions obtained there seem not necessary to be least energy solutions. Theorem 1.3 in this paper is much more general that those results in [22]. We also remark that any other results in our paper can not be found in [22].

2 Proof of Theorems 1.1 and 1.6

Define functions

$$\alpha_1(k,l) := \mu_1 k^{p-1} + \beta k^{\frac{p}{2} - 1} l^{\frac{p}{2}} - 1, \quad k > 0, \ l \ge 0; \tag{2.1}$$

$$\alpha_2(k,l) := \mu_2 l^{p-1} + \beta l^{\frac{p}{2}-1} k^{\frac{p}{2}} - 1, \quad l > 0, \ k \ge 0;$$
 (2.2)

$$h_1(k) := \beta^{-2/p} \left(k^{1-p/2} - \mu_1 k^{p/2} \right)^{2/p}, \quad 0 < k \le \mu_1^{-\frac{1}{p-1}};$$
 (2.3)

$$h_2(l) := \beta^{-2/p} \left(l^{1-p/2} - \mu_2 l^{p/2} \right)^{2/p}, \quad 0 < l \le \mu_2^{-\frac{1}{p-1}}.$$
 (2.4)

Then $\alpha_1(k, h_1(k)) \equiv 0$ and $\alpha_2(h_2(l), l) \equiv 0$.

Lemma 2.1. Assume that $\beta > 0$, then equation

$$\alpha_1(k,l) = 0, \quad \alpha_2(k,l) = 0, \quad k,l > 0$$
 (2.5)

has a solution (k_0, l_0) , which satisfies

$$\alpha_2(k, h_1(k)) < 0, \quad \forall \ 0 < k < k_0,$$
 (2.6)

that is, (k_0, l_0) satisfies (1.10). Similarly, (2.5) has a solution (k_1, l_1) such that

$$\alpha_1(h_2(l), l) < 0, \quad \forall \ 0 < l < l_1.$$
 (2.7)

Proof. Equation $\alpha_1(k,l) = 0$, k,l > 0 imply that

$$l = h_1(k), \quad 0 < k < \mu_1^{-\frac{1}{p-1}}.$$

While, $\alpha_2(k,l) = 0$ implies that $\mu_2 l^{p/2} + \beta k^{p/2} = l^{1-p/2}$. Therefore, we turn to prove that

$$\mu_2 \frac{1 - \mu_1 k^{p-1}}{\beta k^{p/2 - 1}} + \beta k^{p/2} = \left(\frac{1 - \mu_1 k^{p-1}}{\beta k^{p/2 - 1}}\right)^{\frac{2 - p}{p}}, \quad 0 < k^{p-1} < \frac{1}{\mu_1}$$
 (2.8)

have a solution. Note that (2.8) is equivalent to

$$f(k) := \left(\frac{1}{\beta k^{p-1}} - \frac{\mu_1}{\beta}\right)^{\frac{2-p}{p}} - \frac{\mu_2}{\beta} - \frac{\beta^2 - \mu_1 \mu_2}{\beta} k^{p-1} = 0, \ 0 < k^{p-1} < \frac{1}{\mu_1}. \ (2.9)$$

Recall that $N \geq 5$ and $2p = 2^*$, we have 2 - p > 0 and so

$$\lim_{k \to 0+} f(k) = +\infty, \quad f(\mu_1^{-\frac{1}{p-1}}) = -\frac{\beta}{\mu_1} < 0.$$

Therefore, there exists $k_0 \in (0, \mu_1^{-\frac{1}{p-1}})$ such that $f(k_0) = 0$ and f(k) > 0 for $k \in (0, k_0)$. Let $l_0 = h_1(k_0)$, then (k_0, l_0) is a solution of (2.5). Moreover, (2.6) follows directly from f(k) > 0 for $k \in (0, k_0)$. The existence of (k_1, l_1) that satisfy (2.5) and (2.7) is similar.

Lemma 2.2. Assume that $\beta \geq (p-1) \max\{\mu_1, \mu_2\}$, then $h_1(k) + k$ is strictly increasing for $k \in [0, \mu_1^{-\frac{1}{p-1}}]$ and $h_2(l) + l$ is strictly increasing for $l \in [0, \mu_2^{-\frac{1}{p-1}}]$.

Proof. Since for k > 0

$$h_1'(k) = \frac{2}{p} \beta^{-2/p} \left(k^{1-p/2} - \mu_1 k^{p/2} \right)^{2/p-1} \left((1-p/2) k^{-p/2} - \frac{p}{2} \mu_1 k^{p/2-1} \right),$$

we see that $h_1'(k) \geq 0$ for $0 < \mu_1 k^{p-1} \leq \frac{2-p}{p}$ or $\mu_1 k^{p-1} = 1$, and $h_1'(k) < 0$ for $\frac{2-p}{p} < \mu_1 k^{p-1} < 1$. By direct computations, we deduce from $h_1''(k) = 0$, $\frac{2-p}{p} < \mu_1 k^{p-1} < 1$ that $k = (\mu_1 p)^{-\frac{1}{p-1}}$. Since $\beta \geq (p-1) \max\{\mu_1, \mu_2\}$, we have

$$\min_{0 < k^{p-1} < \mu_1^{-1}} h_1'(k) = h_1' \left((\mu_1 p)^{-\frac{1}{p-1}} \right) = -\beta^{-2/p} \mu_1^{2/p} (p-1)^{2/p} \ge -1,$$

and so $h_1'(k) > -1$ for $0 < k \le \mu_1^{-\frac{1}{p-1}}$ with $k \ne (\mu_1 p)^{-\frac{1}{p-1}}$. This implies that $h_1(k) + k$ is strictly increasing for $k \in [0, \mu_1^{-\frac{1}{p-1}}]$. Similarly, $h_2(l) + l$ is strictly increasing for $l \in [0, \mu_2^{-\frac{1}{p-1}}]$.

Lemma 2.3. Assume that $\beta \geq (p-1) \max\{\mu_1, \mu_2\}$. Let (k_0, l_0) be in Lemma 2.1. Then $\max\{\mu_1(k_0 + l_0)^{p-1}, \mu_2(k_0 + l_0)^{p-1}\} < 1$ and

$$\alpha_2(k, h_1(k)) < 0, \ \forall \ 0 < k < k_0; \quad \alpha_1(h_2(l), l) < 0, \ \forall \ 0 < l < l_0.$$
 (2.10)

Proof. By Lemma 2.2 we have

$$h_1(\mu_1^{-\frac{1}{p-1}}) + \mu_1^{-\frac{1}{p-1}} = \mu_1^{-\frac{1}{p-1}} > h_1(k_0) + k_0 = k_0 + l_0,$$

that is, $\mu_1(k_0 + l_0)^{p-1} < 1$. Similarly, $\mu_2(k_0 + l_0)^{p-1} < 1$. By Lemma 2.1, to prove (2.10), it suffices to prove that $(k_0, l_0) = (k_1, l_1)$. By (2.6)-(2.7) we see that $k_1 \ge k_0, l_0 \ge l_1$. If $k_1 > k_0$, then $k_1 + h_1(k_1) > k_0 + h_1(k_0)$, that is, $h_2(l_1) + l_1 = k_1 + l_1 > k_0 + l_0 = h_2(l_0) + l_0$, and so $l_1 > l_0$, a contradiction. Therefore, $k_1 = k_0$ and $l_0 = l_1$. This completes the proof.

Lemma 2.4. Assume that $\beta \geq (p-1) \max{\{\mu_1, \mu_2\}}$. Then

$$\begin{cases} k+l \le k_0 + l_0, \\ \alpha_1(k,l) \ge 0, \quad \alpha_2(k,l) \ge 0, \\ k,l \ge 0, \quad (k,l) \ne (0,0) \end{cases}$$
 (2.11)

has a unique solution (k_0, l_0) .

Proof. Note that (k_0, l_0) satisfies (2.11). Let (\tilde{k}, \tilde{l}) be any a solution of (2.11). Without loss of generality, we assume that $\tilde{k} > 0$. If $\tilde{l} = 0$, then by $\tilde{k} \le k_0 + l_0$ and $\alpha_1(\tilde{k}, 0) \ge 0$ we get that

$$1 \le \mu_1 \tilde{k}^{p-1} \le \mu_1 (k_0 + l_0)^{p-1},$$

which contradicts with Lemma 2.3. Therefore l > 0.

Assume by contradiction that $\tilde{k} < k_0$. Similarly to the proof of Lemma 2.2, by (2.4) it is easy to see that $h_2(l)$ is strictly increasing for $0 < \mu_2 l^{p-1} \le \frac{2-p}{p}$, and strictly decreasing for $\frac{2-p}{p} \le \mu_2 l^{p-1} \le 1$. Moreover, $h_2(0) = h_2(\mu_2^{-\frac{1}{p-1}}) = 0$. Since $0 < \tilde{k} < k_0 = h_2(l_0)$, there exists $0 < l_2 < l_3 < \mu_2^{-\frac{1}{p-1}}$ such that $h_2(l_2) = h_2(l_3) = \tilde{k}$ and

$$\alpha_2(\tilde{k}, l) < 0 \iff h_2(l) > \tilde{k} \iff l_2 < l < l_3. \tag{2.12}$$

Since $\alpha_2(\tilde{k},\tilde{l}) \geq 0$, we have $\tilde{l} \leq l_2$ or $\tilde{l} \geq l_3$. Since $\alpha_1(\tilde{k},\tilde{l}) \geq 0$, we have $\tilde{l} \geq h_1(\tilde{k})$. By Lemma 2.3 we have $\alpha_2(\tilde{k},h_1(\tilde{k})) < 0$, and so $l_2 < h_1(\tilde{k}) < l_3$. These imply

$$\tilde{l} \ge l_3. \tag{2.13}$$

On the other hand, since $l_1 := k_0 + l_0 - \tilde{k} > l_0$, we have

$$h_2(l_1) + k_0 + l_0 - \tilde{k} = h_2(l_1) + l_1 > h_2(l_0) + l_0 = k_0 + l_0$$

that is, $h_2(l_1) > \tilde{k}$. By (2.12) we have $l_2 < l_1 < l_3$. By $\tilde{k} + \tilde{l} \le k_0 + l_0$ we have $\tilde{l} \le l_1 < l_3$.

which contradicts with (2.13). Therefore, $\tilde{k} \geq k_0$. By a similar argument, we also have $\tilde{l} \geq l_0$. Therefore, $(\tilde{k}, \tilde{l}) = (k_0, l_0)$. This completes the proof.

Proof of Theorem 1.1. Assume that $-\lambda_1(\Omega) < \lambda_1 = \lambda_2 = \lambda < 0$. By the Sobolev inequality (1.15) it is standard to see that B > 0. Since $\beta > 0$, by Lemma 2.1 equation (1.9) has a solution (k_0, l_0) . Recall (1.7), we see that $(\sqrt{k_0}\omega, \sqrt{l_0}\omega)$ is a nontrivial solution of (1.1) and

$$0 < B \le E(\sqrt{k_0}\omega, \sqrt{l_0}\omega) = (k_0 + l_0)B_1. \tag{2.14}$$

Now we assume that $\beta \geq (p-1) \max\{\mu_1, \mu_2\}$, and we shall prove that $B = E(\sqrt{k_0}\omega, \sqrt{l_0}\omega)$. Let $\{(u_n, v_n)\} \subset \mathcal{M}$ be a minimizing sequence for B, that is, $E(u_n, v_n) \to B$. Define

$$c_n = \left(\int_{\Omega} |u_n|^{2p} dx \right)^{1/p}, \quad d_n = \left(\int_{\Omega} |v_n|^{2p} dx \right)^{1/p}.$$

By (1.8) we have

$$(NB_1)^{2/N}c_n \le \int_{\Omega} (|\nabla u_n|^2 + \lambda u_n^2) = \int_{\Omega} (\mu_1 |u_n|^{2p} + \beta |u_n|^p |v_n|^p)$$

$$\le \mu_1 c_n^p + \beta c_n^{p/2} d_n^{p/2}, \qquad (2.15)$$

$$(NB_1)^{2/N}d_n \le \int_{\Omega} (|\nabla v_n|^2 + \lambda v_n^2) = \int_{\Omega} (\mu_2 |v_n|^{2p} + \beta |u_n|^p |v_n|^p)$$

$$\le \mu_2 d_n^p + \beta c_n^{p/2} d_n^{p/2}. \tag{2.16}$$

Since $E(u_n, v_n) = \frac{1}{N} \int_{\Omega} (|\nabla u_n|^2 + \lambda u_n^2 + |\nabla v_n|^2 + \lambda v_n^2)$, by (2.14) we have

$$(NB_1)^{2/N}(c_n+d_n) \le NE(u_n, v_n) \le N(k_0+l_0)B_1 + o(1), \tag{2.17}$$

$$\mu_1 c_n^{p-1} + \beta c_n^{p/2-1} d_n^{p/2} \ge (NB_1)^{2/N},$$
 (2.18)

$$\mu_2 d_n^{p-1} + \beta c_n^{p/2} d_n^{p/2-1} \ge (NB_1)^{2/N}.$$
 (2.19)

First, this means c_n, d_n are uniformly bounded. Passing to a subsequence, we assume that $c_n \to c$ and $d_n \to d$. Then by (2.15)-(2.16) we have $\mu_1 c^p + 2\beta c^{p/2} d^{p/2} + \mu_2 d^p \ge NB > 0$. Hence, without loss of generality, we assume that c > 0. If d = 0, then (2.17) implies $c \le (NB_1)^{1-2/N} (k_0 + l_0)$. By (2.18) and Lemma 2.3 we get

$$(NB_1)^{2/N} \le \mu_1 c^{p-1} \le \mu_1 (k_0 + l_0)^{p-1} (NB_1)^{2/N} < (NB_1)^{2/N},$$

a contradiction. Therefore, c > 0 and d > 0. Let $k = \frac{c}{(NB_1)^{1-2/N}}$ and $l = \frac{d}{(NB_1)^{1-2/N}}$, then by (2.17)-(2.19) we see that (k, l) satisfies (2.11). By Lemma

2.4 we see that $(k,l)=(k_0,l_0)$. It follows that $c_n\to k_0(NB_1)^{1-2/N}$ and $d_n\to l_0(NB_1)^{1-2/N}$ as $n\to +\infty$, and

$$NB = \lim_{n \to +\infty} NE(u_n, v_n) \ge \lim_{n \to +\infty} (NB_1)^{2/N} (c_n + d_n) = N(k_0 + l_0)B_1.$$

Combining this with (2.14), one has that

$$B = (k_0 + l_0)B_1 = E(\sqrt{k_0\omega}, \sqrt{l_0\omega}),$$

and so $(\sqrt{k_0}\omega, \sqrt{l_0}\omega)$ is a positive least energy solution of (1.1).

Now we turn to the proof of Theorem 1.6. By the Sobolev inequality (1.15) it is standard to see that

$$A = \inf_{(u,v) \in \mathcal{N}} \frac{1}{4} \int_{\mathbb{R}^N} (|\nabla u|^2 + |\nabla v|^2) \, dx > 0.$$
 (2.20)

Lemma 2.5. If A (resp. B) is attained by a couple $(u, v) \in \mathcal{N}$ (resp. $(u, v) \in \mathcal{M}$), then this couple is a critical point of I (resp. E), provided $-\infty < \beta < 0$.

Proof. Let $\beta < 0$. Assume that $(u, v) \in \mathcal{N}$ such that A = I(u, v). Define

$$G_1(u,v) := I'(u,v)(u,0) = \int_{\mathbb{R}^N} |\nabla u|^2 - \int_{\mathbb{R}^N} (\mu_1 |u|^{2p} + \beta |u|^p |v|^p),$$

$$G_2(u,v) := I'(u,v)(0,v) = \int_{\mathbb{R}^N} |\nabla v|^2 - \int_{\mathbb{R}^N} (\mu_2 |v|^{2p} + \beta |u|^p |v|^p).$$

Then there exists two Lagrange multipliers $L_1, L_2 \in \mathbb{R}$ such that

$$I'(u,v) + L_1 G'_1(u,v) + L_2 G'_2(u,v) = 0.$$

Acting on (u,0) and (0,v) respectively, we obtain

$$\left((2p-2) \int_{\mathbb{R}^N} \mu_1 |u|^{2p} - (2-p) \int_{\mathbb{R}^N} \beta |u|^p |v|^p \right) L_1 + L_2 p \int_{\mathbb{R}^N} \beta |u|^p |v|^p = 0,
\left((2p-2) \int_{\mathbb{R}^N} \mu_2 |v|^{2p} - (2-p) \int_{\mathbb{R}^N} \beta |u|^p |v|^p \right) L_2 + L_1 p \int_{\mathbb{R}^N} \beta |u|^p |v|^p = 0.$$

Since $\beta < 0$, we deduce from $G_1(u, v) = G_2(u, v) = 0$ that

$$\left((2p-2) \int_{\mathbb{R}^N} \mu_1 |u|^{2p} - (2-p) \int_{\mathbb{R}^N} \beta |u|^p |v|^p \right)
\times \left((2p-2) \int_{\mathbb{R}^N} \mu_2 |v|^{2p} - (2-p) \int_{\mathbb{R}^N} \beta |u|^p |v|^p \right) > \left(p \int_{\mathbb{R}^N} \beta |u|^p |v|^p \right)^2.$$

From this we have $L_1 = L_2 = 0$ and so I'(u, v) = 0. Similarly, if $(u, v) \in \mathcal{M}$ such that E(u, v) = B, then E'(u, v) = 0.

Proof of (1) in Theorem 1.6. This proof is similar to the proof of [16, Theorem 1.5-(1)] in case N=4, but the details are more delicate. By (1.16) we see that $\omega_{\mu_i} := \mu_i^{-\frac{N-2}{4}} U_{1,0}$ satisfies equation $-\Delta u = \mu_i |u|^{2^*-2} u$ in \mathbb{R}^N . Let $e_1 = (1, 0, \dots, 0) \in \mathbb{R}^N$ and

$$(u_R(x), v_R(x)) = (\omega_{\mu_1}(x), \omega_{\mu_2}(x + Re_1)).$$

Then $v_R \rightharpoonup 0$ weakly in $D^{1,2}(\mathbb{R}^N)$ and so $v_R^p \rightharpoonup 0$ weakly in $L^2(\mathbb{R}^N)$ as $R \to +\infty$. That is,

$$\lim_{R \to +\infty} \int_{\mathbb{R}^N} u_R^p v_R^p \, dx = 0.$$

Note that $\beta < 0$. Then for R > 0 sufficiently large, by a similar argument as that of Lemma 2.1 (or see the argument of existing $(t_{\varepsilon}, s_{\varepsilon})$ in the proof of Lemma 3.1 below), we see that

$$\begin{cases} t^2 \int_{\mathbb{R}^N} |\nabla u_R|^2 dx = t^2 \mu_1 \int_{\mathbb{R}^N} u_R^{2p} dx = t^{2p} \mu_1 \int_{\mathbb{R}^N} u_R^{2p} dx + t^p s^p \beta \int_{\mathbb{R}^N} u_R^p v_R^p dx, \\ s^2 \int_{\mathbb{R}^N} |\nabla v_R|^2 dx = s^2 \mu_2 \int_{\mathbb{R}^N} v_R^{2p} dx = s^{2p} \mu_2 \int_{\mathbb{R}^N} v_R^{2p} dx + t^p s^p \beta \int_{\mathbb{R}^N} u_R^p v_R^p dx, \end{cases}$$

have a solution (t_R, s_R) with $t_R > 1$ and $s_R > 1$. Denote

$$D_{1} := \mu_{1} \int_{\mathbb{R}^{N}} u_{R}^{2p} dx = \mu_{1} \int_{\mathbb{R}^{N}} \omega_{\mu_{1}}^{2p} dx > 0,$$

$$D_{2} := \mu_{2} \int_{\mathbb{R}^{N}} v_{R}^{2p} dx = \mu_{2} \int_{\mathbb{R}^{N}} \omega_{\mu_{2}}^{2p} dx > 0,$$

$$F_{R} := |\beta| \int_{\mathbb{R}^{N}} u_{R}^{p} v_{R}^{p} dx \to 0, \quad \text{as } R \to +\infty.$$

Then

$$t_R^2 D_1 = t_R^{2p} D_1 - t_R^p s_R^p F_R, \quad s_R^2 D_2 = s_R^{2p} D_2 - t_R^p s_R^p F_R. \tag{2.21}$$

Assume that, up to a subsequence, $t_R \to +\infty$ as $R \to \infty$, then by

$$t_R^{2p}D_1 - t_R^2D_1 = s_R^{2p}D_2 - s_R^2D_2$$

we also have $s_R \to +\infty$. Note that 2 - p < p, we have

$$t_R^p D_1 - t_R^{2-p} D_1 \ge \frac{1}{2} t_R^p D_1, \ s_R^p D_2 - s_R^{2-p} D_2 \ge \frac{1}{2} s_R^p D_2, \text{ for } R \text{ large enough,}$$

and so

$$F_R = \frac{t_R^p - t_R^{2-p}}{s_R^p} D_1 \ge \frac{t_R^p}{2s_R^p} D_1, \quad F_R = \frac{s_R^p - s_R^{2-p}}{t_R^p} D_2 \ge \frac{s_R^p}{2t_R^p} D_2,$$

which implies that

$$0 < \frac{1}{4}D_1D_2 \le F_R^2 \to 0$$
, as $R \to +\infty$,

a contradiction. Therefore, t_R and s_R are uniformly bounded. Then by (2.21) and $F_R \to 0$ as $R \to \infty$, we get that

$$\lim_{R \to +\infty} (|t_R - 1| + |s_R - 1|) = 0.$$

Note that $(t_R u_R, s_R v_R) \in \mathcal{N}$, we see from (1.17) that

$$A \leq I(t_R u_R, s_R v_R) = \frac{1}{N} \left(t_R^2 \int_{\mathbb{R}^N} |\nabla u_R|^2 dx + s_R^2 \int_{\mathbb{R}^N} |\nabla v_R|^2 dx \right)$$
$$= \frac{1}{N} \left(t_R^2 \mu_1^{-\frac{N-2}{2}} + s_R^2 \mu_2^{-\frac{N-2}{2}} \right) S^{N/2}.$$

Letting $R \to +\infty$, we get that $A \leq \frac{1}{N}(\mu_1^{-\frac{N-2}{2}} + \mu_2^{-\frac{N-2}{2}})S^{N/2}$. On the other hand, for any $(u,v) \in \mathcal{N}$, we see from $\beta < 0$ and (1.15) that

$$\int_{\mathbb{R}^N} |\nabla u|^2 dx \le \mu_1 \int_{\mathbb{R}^N} |u|^{2p} dx \le \mu_1 S^{-p} \left(\int_{\mathbb{R}^N} |\nabla u|^2 dx \right)^p,$$

and so $\int_{\mathbb{R}^N} |\nabla u|^2 dx \ge \mu_1^{-\frac{N-2}{2}} S^{N/2}$. Similarly, $\int_{\mathbb{R}^N} |\nabla v|^2 dx \ge \mu_2^{-\frac{N-2}{2}} S^{N/2}$. Combining these with (1.19), we get that $A \ge \frac{1}{N} (\mu_1^{-\frac{N-2}{2}} + \mu_2^{-\frac{N-2}{2}}) S^{N/2}$. Hence,

$$A = \frac{1}{N} \left(\mu_1^{-\frac{N-2}{2}} + \mu_2^{-\frac{N-2}{2}} \right) S^{N/2}. \tag{2.22}$$

Now, assume that A is attained by some $(u,v) \in \mathcal{N}$, then $(|u|,|v|) \in \mathcal{N}$ and I(|u|,|v|) = A. By Lemma 2.5, we get that (|u|,|v|) is a nontrivial solution of (1.14). By the maximum principle, we may assume that u > 0, v > 0 and so $\int_{\mathbb{R}^N} u^p v^p \, dx > 0$. That is,

$$\int_{\mathbb{R}^{N}} |\nabla u|^{2} dx < \mu_{1} \int_{\mathbb{R}^{N}} |u|^{2p} dx \le \mu_{1} S^{-p} \left(\int_{\mathbb{R}^{N}} |\nabla u|^{2} dx \right)^{p},$$

Therefore, it is easy to see that

$$A = I(u, v) = \frac{1}{N} \int_{\mathbb{R}^N} (|\nabla u|^2 + |\nabla v|^2) \, dx > \frac{1}{N} \left(\mu_1^{-\frac{N-2}{2}} + \mu_2^{-\frac{N-2}{2}} \right) S^{N/2},$$

which is a contradiction. This completes the proof.

Proof of (2-1) in Theorem 1.6. This proof is similar to the proof of Theorem 1.1. Since $\beta > 0$, by Lemma 2.1 equation (1.9) has a solution (k_0, l_0) . Then $(\sqrt{k_0}U_{\varepsilon,y}, \sqrt{l_0}U_{\varepsilon,y})$ is a nontrivial solution of (1.14) and

$$A \le I\left(\sqrt{k_0}U_{\varepsilon,y}, \sqrt{l_0}U_{\varepsilon,y}\right) = \frac{1}{N}(k_0 + l_0)S^{N/2}.$$
 (2.23)

Assume that $\beta \geq (p-1) \max\{\mu_1, \mu_2\}$. Let $\{(u_n, v_n)\} \subset \mathcal{N}$ be a minimizing sequence for A, that is, $I(u_n, v_n) \to A$. Define $c_n = \left(\int_{\mathbb{R}^N} |u_n|^{2p} dx\right)^{1/p}$, $d_n = \left(\int_{\mathbb{R}^N} |v_n|^{2p} dx\right)^{1/p}$, we have

$$Sc_n \le \int_{\mathbb{R}^N} |\nabla u_n|^2 \, dx = \int_{\mathbb{R}^N} \mu_1 |u_n|^{2p} + \beta |u_n|^p |v_n|^p \, dx \le \mu_1 c_n^p + \beta c_n^{p/2} d_n^{p/2},$$

$$Sd_n \le \int_{\mathbb{R}^N} |\nabla v_n|^2 dx = \int_{\mathbb{R}^N} \mu_2 |v_n|^{2p} + \beta |u_n|^p |v_n|^p dx \le \mu_2 d_n^p + \beta c_n^{p/2} d_n^{p/2}.$$

This means

$$S(c_n + d_n) \le NI(u_n, v_n) \le (k_0 + l_0)S^{N/2} + o(1),$$

$$\mu_1 c_n^{p-1} + \beta c_n^{p/2-1} d_n^{p/2} \ge S, \quad \beta c_n^{p/2} d_n^{p/2-1} + \mu_2 d_n^{p-1} \ge S.$$

Similarly as in the proof of Theorem 1.1, we see that $c_n \to k_0 S^{N/2-1}$ and $d_n \to l_0 S^{N/2-1}$ as $n \to +\infty$, and

$$NA = \lim_{n \to +\infty} NI(u_n, v_n) \ge \lim_{n \to +\infty} S(c_n + d_n) = (k_0 + l_0)S^{N/2}.$$

This implies that

$$A = \frac{1}{N}(k_0 + l_0)S^{N/2} = I(\sqrt{k_0}U_{\varepsilon,y}, \sqrt{l_0}U_{\varepsilon,y}), \tag{2.24}$$

and so
$$(\sqrt{k_0}U_{\varepsilon,y}, \sqrt{l_0}U_{\varepsilon,y})$$
 is a positive least energy solution of (1.14).

To finish the proof of Theorem 1.6, we need to show that (1.14) has a positive least energy solution for any $0 < \beta < (p-1) \max\{\mu_1, \mu_2\}$. The following proof works for all $\beta > 0$. Therefore, we assume that $\beta > 0$, and define

$$A' := \inf_{(u,v) \in \mathcal{N}'} I(u,v), \tag{2.25}$$

where

$$\mathcal{N}' := \{(u, v) \in D \setminus \{(0, 0)\}, \ I'(u, v)(u, v) = 0\}.$$
(2.26)

Note that $\mathcal{N} \subset \mathcal{N}'$, one has that $A' \leq A$. By Sobolev inequality, we have A' > 0. Define $B(0,R) := \{x \in \mathbb{R}^N : |x| < R\}$ and $H(0,R) := H_0^1(B(0,R)) \times H_0^1(B(0,R))$. Consider

$$\begin{cases}
-\Delta u = \mu_1 |u|^{2p-2} u + \beta |u|^{p-2} u |v|^p, & x \in B(0, R), \\
-\Delta v = \mu_2 |v|^{2p-2} v + \beta |v|^{p-2} v |u|^p, & x \in B(0, R), \\
u, v \in H_0^1(B(0, R)),
\end{cases} (2.27)$$

and define

$$A'(R) := \inf_{(u,v) \in \mathcal{N}'(R)} I(u,v), \tag{2.28}$$

where

$$\mathcal{N}'(R) := \left\{ (u, v) \in H(0, R) \setminus \{(0, 0)\}, \int_{B(0, R)} (|\nabla u|^2 + |\nabla v|^2) - \int_{B(0, R)} (\mu_1 |u|^{2p} + 2\beta |u|^p |v|^p + \mu_2 |v|^{2p}) = 0 \right\}.$$
 (2.29)

Lemma 2.6. $A'(R) \equiv A'$ for all R > 0.

Proof. Take any $R_1 > R_2$. By $\mathcal{N}'(R_2) \subset \mathcal{N}'(R_1)$, we have $A'(R_1) \leq A'(R_2)$. On the other hand, for any $(u, v) \in \mathcal{N}'(R_1)$, we define

$$(u_1(x), v_1(x)) := \left(\left(\frac{R_1}{R_2} \right)^{\frac{N-2}{2}} u \left(\frac{R_1}{R_2} x \right), \ \left(\frac{R_1}{R_2} \right)^{\frac{N-2}{2}} v \left(\frac{R_1}{R_2} x \right) \right),$$

then it is standard to see that $(u_1, v_1) \in \mathcal{N}'(R_2)$, and so

$$A'(R_2) \le I(u_1, v_1) = I(u, v), \quad \forall (u, v) \in \mathcal{N}'(R_1).$$

That is, $A'(R_2) \leq A'(R_1)$ and so $A'(R_1) = A'(R_2)$.

Clearly $A' \leq A'(R)$. Let $(u_n, v_n) \in \mathcal{N}'$ be a minimizing sequence of A'. Moreover, we may assume that $u_n, v_n \in H^1_0(B(0, R_n))$ for some $R_n > 0$. Then $(u_n, v_n) \in \mathcal{N}'(R_n)$ and

$$A' = \lim_{n \to \infty} I(u_n, v_n) \ge \lim_{n \to \infty} A'(R_n) \equiv A'(R).$$

Therefore, $A'(R) \equiv A'$ for all R > 0.

Let $0 \le \varepsilon . Consider$

$$\begin{cases}
-\Delta u = \mu_1 |u|^{2p-2-2\varepsilon} u + \beta |u|^{p-2-\varepsilon} u |v|^{p-\varepsilon}, & x \in B(0,1), \\
-\Delta v = \mu_2 |v|^{2p-2-2\varepsilon} v + \beta |v|^{p-2-\varepsilon} v |u|^{p-\varepsilon}, & x \in B(0,1), \\
u, v \in H_0^1(B(0,1)),
\end{cases}$$
(2.30)

and define

$$A_{\varepsilon} := \inf_{(u,v) \in \mathcal{N}_{\varepsilon}'} I_{\varepsilon}(u,v), \tag{2.31}$$

where

$$\begin{split} I_{\varepsilon}(u,v) := & \frac{1}{2} \int_{B(0,1)} (|\nabla u|^2 + |\nabla v|^2) \\ & - \frac{1}{2p - 2\varepsilon} \int_{B(0,1)} (\mu_1 |u|^{2p - 2\varepsilon} + 2\beta |u|^{p - \varepsilon} |v|^{p - \varepsilon} + \mu_2 |v|^{2p - 2\varepsilon}), \\ \mathcal{N}'_{\varepsilon} := & \{(u,v) \in H(0,1) \setminus \{(0,0)\}, \ H_{\varepsilon}(u,v) := I'_{\varepsilon}(u,v)(u,v) = 0\} \,. \end{split}$$

Lemma 2.7. For any $0 < \varepsilon < p-1$, there holds

$$A_{\varepsilon} < \min \left\{ \inf_{(u,0) \in \mathcal{N}'_{\varepsilon}} I_{\varepsilon}(u,0), \inf_{(0,v) \in \mathcal{N}'_{\varepsilon}} I_{\varepsilon}(0,v) \right\}.$$

Proof. Fix any $0 < \varepsilon < p-1$. Recall that $2 < 2p-2\varepsilon < 2^*$, we may let u_i be a least energy solution of

$$-\Delta u = \mu_i |u|^{2p-2-2\varepsilon} u, \quad u \in H_0^1(B(0,1)),$$

Then

$$I_{\varepsilon}(u_1,0) = c_1 := \inf_{(u,0) \in \mathcal{N}_{\varepsilon}'} I_{\varepsilon}(u,0), \quad I_{\varepsilon}(0,u_2) = c_2 := \inf_{(0,v) \in \mathcal{N}_{\varepsilon}'} I_{\varepsilon}(0,v).$$

The following proof is inspired by [3]. For any $s \in \mathbb{R}$, there exists a unique t(s) > 0 such that $(t(s)u_1, t(s)su_2) \in \mathcal{N}'_{\varepsilon}$. In fact,

$$\begin{split} t(s)^{2p-2\varepsilon-2} &= \frac{\int_{B(0,1)} (|\nabla u_1|^2 + s^2 |\nabla u_2|^2)}{\int_{B(0,1)} (\mu_1 |u_1|^{2p-2\varepsilon} + 2\beta |u_1|^{p-\varepsilon} |su_2|^{p-\varepsilon} + \mu_2 |su_2|^{2p-2\varepsilon})} \\ &= \frac{p'c_1 + s^2 p'c_2}{p'c_1 + |s|^{2p-2\varepsilon} p'c_2 + |s|^{p-\varepsilon} \int_{B(0,1)} 2\beta |u_1|^{p-\varepsilon} |u_2|^{p-\varepsilon}}, \end{split}$$

where $p' = \frac{2p-2\varepsilon}{p-1-\varepsilon}$. Note that t(0) = 1. Recall that 1 , by direct computations we have

$$\lim_{s \to 0} \frac{t'(s)}{|s|^{p-2-\varepsilon}s} = -\frac{(p-\varepsilon) \int_{B(0,1)} 2\beta |u_1|^{p-\varepsilon} |u_2|^{p-\varepsilon}}{(2p-2\varepsilon-2)p'c_1},$$

that is,

$$t'(s) = -\frac{(p-\varepsilon)\int_{B(0,1)} 2\beta |u_1|^{p-\varepsilon} |u_2|^{p-\varepsilon}}{(2p-2\varepsilon-2)p'c_1} |s|^{p-\varepsilon-2} s(1+o(1)), \text{ as } s \to 0,$$

and so

$$t(s) = 1 - \frac{\int_{B(0,1)} 2\beta |u_1|^{p-\varepsilon} |u_2|^{p-\varepsilon}}{(2p - 2\varepsilon - 2)p'c_1} |s|^{p-\varepsilon} (1 + o(1)), \text{ as } s \to 0.$$

This implies that

$$t(s)^{2p-2\varepsilon} = 1 - \frac{(2p-2\varepsilon) \int_{B(0,1)} 2\beta |u_1|^{p-\varepsilon} |u_2|^{p-\varepsilon}}{(2p-2\varepsilon-2)p'c_1} |s|^{p-\varepsilon} (1+o(1))$$

$$= 1 - \frac{\int_{B(0,1)} 2\beta |u_1|^{p-\varepsilon} |u_2|^{p-\varepsilon}}{2c_1} |s|^{p-\varepsilon} (1+o(1)), \text{ as } s \to 0.$$

Therefore, we see from $1/2 - 1/p' = 1/(2p - 2\varepsilon) > 0$ that

$$\begin{split} A_{\varepsilon} &\leq I_{\varepsilon}\left(t(s)u_{1},t(s)su_{2}\right) \\ &= \frac{t(s)^{2p-2\varepsilon}}{p'}\left(p'c_{1}+|s|^{2p-2\varepsilon}p'c_{2}+|s|^{p-\varepsilon}\int_{B(0,1)}2\beta|u_{1}|^{p-\varepsilon}|u_{2}|^{p-\varepsilon}\right) \\ &= c_{1}-\left(\frac{1}{2}-\frac{1}{p'}\right)|s|^{p-\varepsilon}\int_{B(0,1)}2\beta|u_{1}|^{p-\varepsilon}|u_{2}|^{p-\varepsilon}+o(|s|^{p-\varepsilon}) \\ &< c_{1}=\inf_{(u,0)\in\mathcal{N}_{\varepsilon}'}I_{\varepsilon}(u,0)\quad\text{as }|s|>0\text{ small enough,} \end{split}$$

By a similar argument, we have $A_{\varepsilon} < \inf_{(0,v) \in \mathcal{N}'_{\varepsilon}} I_{\varepsilon}(0,v)$.

Recalling ω_{μ_i} in the proof of Theorem 1.6 -(1), similarly as Lemma 2.7, we have

$$A' < \min \left\{ \inf_{(u,0) \in \mathcal{N}'} I(u,0), \inf_{(0,v) \in \mathcal{N}'} I(0,v) \right\} = \min \left\{ I(\omega_{\mu_1},0), I(0,\omega_{\mu_2}) \right\}$$
$$= \min \left\{ \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{N/2}, \frac{1}{N} \mu_1^{-\frac{N-2}{2}} S^{N/2} \right\}. \tag{2.32}$$

Theorem 2.1. For any $0 < \varepsilon < p-1$, (2.30) has a classical least energy solution $(u_{\varepsilon}, v_{\varepsilon})$, and $u_{\varepsilon}, v_{\varepsilon}$ are both positive radially symmetric decreasing.

Proof. Fix any $0 < \varepsilon < p - 1$, it is easy to see that $A_{\varepsilon} > 0$. For $(u, v) \in \mathcal{N}'_{\varepsilon}$ with $u \geq 0, v \geq 0$, we denote by (u^*, v^*) as its Schwartz symmetrization. Then by the properties of Schwartz symmetrization and $\beta > 0$, we have

$$\int_{B(0,1)} (|\nabla u^*|^2 + |\nabla v^*|^2) \le \int_{B(0,1)} (\mu_1 |u^*|^{2p-2\varepsilon} + 2\beta |u^*|^{p-\varepsilon} |v^*|^{p-\varepsilon} + \mu_2 |v^*|^{2p-2\varepsilon}).$$

Therefore, there exists $0 < t^* \le 1$ such that $(t^*u^*, t^*v^*) \in \mathcal{N}'_{\varepsilon}$, and then

$$I_{\varepsilon}(t^*u^*, t^*v^*) = \left(\frac{1}{2} - \frac{1}{2p - 2\varepsilon}\right)(t^*)^2 \int_{B(0,1)} (|\nabla u^*|^2 + |\nabla v^*|^2)$$

$$\leq \left(\frac{1}{2} - \frac{1}{2p - 2\varepsilon}\right) \int_{B(0,1)} (|\nabla u|^2 + |\nabla v|^2) = I_{\varepsilon}(u, v). \tag{2.33}$$

Therefore, we may take a minimizing sequence $(u_n, v_n) \in \mathcal{N}_{\varepsilon}'$ of A_{ε} such that $(u_n, v_n) = (u_n^*, v_n^*)$ and $I_{\varepsilon}(u_n, v_n) \to A_{\varepsilon}$. We see from (2.33) that u_n, v_n are uniformly bounded in $H_0^1(B(0,1))$. Passing to a subsequence, we may assume that $u_n \to u_{\varepsilon}, v_n \to v_{\varepsilon}$ weakly in $H_0^1(B(0,1))$. By the compactness of the embedding $H_0^1(B(0,1)) \hookrightarrow L^{2p-2\varepsilon}(B(0,1))$, we have

$$\int_{B(0,1)} (\mu_1 |u_{\varepsilon}|^{2p-2\varepsilon} + 2\beta |u_{\varepsilon}|^{p-\varepsilon} |v_{\varepsilon}|^{p-\varepsilon} + \mu_2 |v_{\varepsilon}|^{2p-2\varepsilon})$$

$$= \lim_{n \to \infty} \int_{B(0,1)} (\mu_1 |u_n|^{2p-2\varepsilon} + 2\beta |u_n|^{p-\varepsilon} |v_n|^{p-\varepsilon} + \mu_2 |v_n|^{2p-2\varepsilon})$$

$$= \frac{2p-2\varepsilon}{p-1-\varepsilon} \lim_{n \to \infty} I_{\varepsilon}(u_n, v_n) = \frac{2p-2\varepsilon}{p-1-\varepsilon} A_{\varepsilon} > 0,$$

which implies $(u_{\varepsilon}, v_{\varepsilon}) \neq (0, 0)$. Moreover, $u_{\varepsilon} \geq 0$, $v_{\varepsilon} \geq 0$ are radially symmetric. Meanwhile, $\int_{B(0,1)} (|\nabla u_{\varepsilon}|^2 + |\nabla v_{\varepsilon}|^2) \leq \lim_{n \to \infty} \int_{B(0,1)} (|\nabla u_n|^2 + |\nabla v_n|^2)$, so

$$\int_{B(0,1)} (|\nabla u_{\varepsilon}|^2 + |\nabla v_{\varepsilon}|^2) \le \int_{B(0,1)} (\mu_1 u_{\varepsilon}^{2p-2\varepsilon} + 2\beta u_{\varepsilon}^{p-\varepsilon} v_{\varepsilon}^{p-\varepsilon} + \mu_2 v_{\varepsilon}^{2p-2\varepsilon}).$$

Therefore, there exists $0 < t_{\varepsilon} \le 1$ such that $(t_{\varepsilon}u_{\varepsilon}, t_{\varepsilon}v_{\varepsilon}) \in \mathcal{N}'_{\varepsilon}$, and then

$$A_{\varepsilon} \leq I_{\varepsilon}(t_{\varepsilon}u_{\varepsilon}, t_{\varepsilon}v_{\varepsilon}) = \left(\frac{1}{2} - \frac{1}{2p - 2\varepsilon}\right)(t_{\varepsilon})^{2} \int_{B(0,1)} (|\nabla u_{\varepsilon}|^{2} + |\nabla v_{\varepsilon}|^{2})$$

$$\leq \lim_{n \to \infty} \left(\frac{1}{2} - \frac{1}{2p - 2\varepsilon}\right) \int_{B(0,1)} (|\nabla u_{n}|^{2} + |\nabla v_{n}|^{2})$$

$$= \lim_{n \to \infty} I_{\varepsilon}(u_{n}, v_{n}) = A_{\varepsilon}.$$

Therefore, $t_{\varepsilon} = 1$ and $(u_{\varepsilon}, v_{\varepsilon}) \in \mathcal{N}'_{\varepsilon}$ with $I(u_{\varepsilon}, v_{\varepsilon}) = A_{\varepsilon}$. Moreover,

$$\int_{B(0,1)} (|\nabla u_{\varepsilon}|^2 + |\nabla v_{\varepsilon}|^2) = \lim_{n \to \infty} \int_{B(0,1)} (|\nabla u_n|^2 + |\nabla v_n|^2),$$

that is, $u_n \to u_{\varepsilon}$ and $v_n \to v_{\varepsilon}$ strongly in $H_0^1(B(0,1))$. There exists a Lagrange multiplier $\gamma \in \mathbb{R}$ such that

$$I_{\varepsilon}'(u_{\varepsilon}, v_{\varepsilon}) - \gamma H_{\varepsilon}'(u_{\varepsilon}, v_{\varepsilon}) = 0.$$

Since $I'_{\varepsilon}(u_{\varepsilon}, v_{\varepsilon})(u_{\varepsilon}, v_{\varepsilon}) = H_{\varepsilon}(u_{\varepsilon}, v_{\varepsilon}) = 0$ and

$$H'_{\varepsilon}(u_{\varepsilon}, v_{\varepsilon})(u_{\varepsilon}, v_{\varepsilon}) = (2 + 2\varepsilon - 2p) \int_{B(0,1)} (\mu_1 u_{\varepsilon}^{2p - 2\varepsilon} + 2\beta u_{\varepsilon}^{p - \varepsilon} v_{\varepsilon}^{p - \varepsilon} + \mu_2 v_{\varepsilon}^{2p - 2\varepsilon}) < 0,$$

we get that $\gamma=0$ and so $I'_{\varepsilon}(u_{\varepsilon},v_{\varepsilon})=0$. By Lemma 2.7, we see that $u_{\varepsilon}\not\equiv 0$ and $v_{\varepsilon}\not\equiv 0$. This means that $(u_{\varepsilon},v_{\varepsilon})$ is a least energy solution of (2.28). Recall that $u_{\varepsilon},v_{\varepsilon}\geq 0$ are radially symmetric non-increasing. By regularity theory and the maximum principle, we see that $u_{\varepsilon},v_{\varepsilon}>0$ in $B(0,1),\,u_{\varepsilon},v_{\varepsilon}\in C^2(B(0,1))$ and are radially symmetric decreasing.

Completion of the proof of (2) in Theorem 1.6. Recalling (2.29), for any $(u,v) \in \mathcal{N}'(1)$, there exists $t_{\varepsilon} > 0$ such that $(t_{\varepsilon}u, t_{\varepsilon}v) \in \mathcal{N}'_{\varepsilon}$ with $t_{\varepsilon} \to 1$ as $\varepsilon \to 0$. Then

$$\limsup_{\varepsilon \to 0} A_{\varepsilon} \le \limsup_{\varepsilon \to 0} I_{\varepsilon}(t_{\varepsilon}u, t_{\varepsilon}v) = I(u, v), \ \forall (u, v) \in \mathcal{N}'(1).$$

By Lemma 2.6 we have

$$\limsup_{\varepsilon \to 0} A_{\varepsilon} \le A'(1) = A'. \tag{2.34}$$

By Theorem 2.1, let $(u_{\varepsilon}, v_{\varepsilon})$ be a positive least energy solution of (2.30), which is radially symmetric decreasing. By $I'_{\varepsilon}(u_{\varepsilon}, v_{\varepsilon})(u_{\varepsilon}, v_{\varepsilon}) = 0$ and Sobolev inequality, it is easily seen that

$$\frac{2p - 2\varepsilon}{p - \varepsilon - 1} A_{\varepsilon} = \int_{B(0,1)} (|\nabla u_{\varepsilon}|^2 + |\nabla v_{\varepsilon}|^2) \ge C_0, \quad \forall \ 0 < \varepsilon \le \frac{p - 1}{2}, \tag{2.35}$$

where C_0 is a positive constant independent of ε . Then $u_{\varepsilon}, v_{\varepsilon}$ are uniformly bounded in $H_0^1(B(0,1))$. Passing to a subsequence, we may assume that $u_{\varepsilon} \rightharpoonup u_0$ and $v_{\varepsilon} \rightharpoonup v_0$ weakly in $H_0^1(B(0,1))$. Then (u_0, v_0) is a solution of

$$\begin{cases}
-\Delta u = \mu_1 |u|^{2p-2} u + \beta |u|^{p-2} u |v|^p, & x \in B(0,1), \\
-\Delta v = \mu_2 |v|^{2p-2} v + \beta |v|^{p-2} v |u|^p, & x \in B(0,1), \\
u, v \in H_0^1(B(0,1)).
\end{cases} (2.36)$$

Assume by contradiction that $||u_{\varepsilon}||_{\infty} + ||v_{\varepsilon}||_{\infty}$ is uniformly bounded, then by the Dominated Convergent Theorem, we get that

$$\lim_{\varepsilon \to 0} \int_{B(0,1)} u_{\varepsilon}^{2p-2\varepsilon} = \int_{B(0,1)} u_{0}^{2p}, \quad \lim_{\varepsilon \to 0} \int_{B(0,1)} v_{\varepsilon}^{2p-2\varepsilon} = \int_{B(0,1)} v_{0}^{2p},$$

$$\lim_{\varepsilon \to 0} \int_{B(0,1)} u_{\varepsilon}^{p-\varepsilon} v_{\varepsilon}^{p-\varepsilon} = \int_{B(0,1)} u_{0}^{p} v_{0}^{p}.$$

Combining these with $I'_{\varepsilon}(u_{\varepsilon}, v_{\varepsilon}) = I'(u_0, v_0) = 0$, it is standard to show that $u_{\varepsilon} \to u_0$ and $v_{\varepsilon} \to v_0$ strongly in $H^1_0(B(0,1))$. Then by (2.35), we see that $(u_0, v_0) \neq (0, 0)$. Moreover, $u_0 \geq 0$, $v_0 \geq 0$. We may assume that $u_0 \not\equiv 0$. By the strong maximum principle, $u_0 > 0$ in B(0, 1). Note that $2p = 2^*$. Combining these with Pohozaev identity, we have

$$0 < \int_{\partial B(0,1)} (|\nabla u_0|^2 + |\nabla v_0|^2)(x \cdot \nu) \, d\sigma = 0,$$

a contradiction. Here, ν denotes the outward unit normal vector on $\partial B(0,1)$. Therefore, $\|u_{\varepsilon}\|_{\infty} + \|v_{\varepsilon}\|_{\infty} \to \infty$ as $\varepsilon \to 0$. We will use a blowup analysis. Note that $u_{\varepsilon}(0) = \max_{B(0,1)} u_{\varepsilon}(x)$ and $v_{\varepsilon}(0) = \max_{B(0,1)} v_{\varepsilon}(x)$, we define $K_{\varepsilon} := \max\{u_{\varepsilon}(0), v_{\varepsilon}(0)\}$, then $K_{\varepsilon} \to +\infty$. Define

$$U_{\varepsilon}(x) = K_{\varepsilon}^{-1} u_{\varepsilon}(K_{\varepsilon}^{-\alpha_{\varepsilon}} x), \quad V_{\varepsilon}(x) = K_{\varepsilon}^{-1} v_{\varepsilon}(K_{\varepsilon}^{-\alpha_{\varepsilon}} x), \quad \alpha_{\varepsilon} = p - 1 - \varepsilon.$$

Then

$$1 = \max\{U_{\varepsilon}(0), V_{\varepsilon}(0)\} = \max\left\{\max_{x \in B(0, K_{\varepsilon}^{\alpha_{\varepsilon}})} U_{\varepsilon}(x), \max_{x \in B(0, K_{\varepsilon}^{\alpha_{\varepsilon}})} V_{\varepsilon}(x)\right\}$$
(2.37)

and $U_{\varepsilon}, V_{\varepsilon}$ satisfy

$$\begin{cases} -\Delta U_{\varepsilon} = \mu_1 U_{\varepsilon}^{2p-2\varepsilon-1} + \beta U_{\varepsilon}^{p-1-\varepsilon} V_{\varepsilon}^{p-\varepsilon}, & x \in B(0, K_{\varepsilon}^{\alpha_{\varepsilon}}), \\ -\Delta V_{\varepsilon} = \mu_2 V_{\varepsilon}^{2p-2\varepsilon-1} + \beta V_{\varepsilon}^{p-1-\varepsilon} U_{\varepsilon}^{p-\varepsilon}, & x \in B(0, K_{\varepsilon}^{\alpha_{\varepsilon}}). \end{cases}$$

Since

$$\int_{\mathbb{R}^N} |\nabla U_\varepsilon|^2 \, dx = K_\varepsilon^{-(N-2)\varepsilon} \int_{\mathbb{R}^N} |\nabla u_\varepsilon|^2 \, dx \leq \int_{\mathbb{R}^N} |\nabla u_\varepsilon|^2 \, dx,$$

we see that $\{(U_{\varepsilon}, V_{\varepsilon})\}_{n\geq 1}$ is bounded in $D^{1,2}(\mathbb{R}^N) \times D^{1,2}(\mathbb{R}^N) = D$. By elliptic estimates, for a subsequence we have $(U_{\varepsilon}, V_{\varepsilon}) \to (U, V) \in D$ uniformly in every

compact subset of \mathbb{R}^N as $\varepsilon \to 0$, and (U, V) satisfies (1.14), that is I'(U, V) = 0. Moreover, $U \geq 0, V \geq 0$ are radially symmetric non-increasing. By (2.37) we have $(U, V) \neq (0, 0)$, and so $(U, V) \in \mathcal{N}'$. Then we deduce from (2.34) that

$$A' \leq I(U, V) = \left(\frac{1}{2} - \frac{1}{2p}\right) \int_{\mathbb{R}^N} (|\nabla U|^2 + |\nabla V|^2) \, dx$$

$$\leq \liminf_{\varepsilon \to 0} \left(\frac{1}{2} - \frac{1}{2p - 2\varepsilon}\right) \int_{B(0, K_{\varepsilon}^{\alpha_{\varepsilon}})} (|\nabla U_{\varepsilon}|^2 + |\nabla V_{\varepsilon}|^2) \, dx$$

$$\leq \liminf_{\varepsilon \to 0} \left(\frac{1}{2} - \frac{1}{2p - 2\varepsilon}\right) \int_{B(0, 1)} (|\nabla u_{\varepsilon}|^2 + |\nabla v_{\varepsilon}|^2) \, dx$$

$$= \liminf_{\varepsilon \to 0} A_{\varepsilon} \leq A'.$$

This implies that I(U,V)=A'. By (2.32) we have that $U\not\equiv 0$ and $V\not\equiv 0$. By the strong maximum principle, U>0 and V>0 are radially symmetric decreasing. We also have $(U,V)\in\mathcal{N}$, and so $I(U,V)\geq A\geq A'$, that is,

$$I(U,V) = A = A',$$
 (2.38)

and (U, V) is a positive least energy solution of (1.14), which is radially symmetric decreasing.

Finally, we show the existence of $(k(\beta), l(\beta))$ for $\beta > 0$ small. Recall (2.1)-(2.2), We denote $\alpha_i(k, l)$ by $\alpha_i(k, l, \beta)$ here. Define $k(0) = \mu_1^{-\frac{1}{p-1}}$ and $l(0) = \mu_2^{-\frac{1}{p-1}}$, then $\alpha_i(k(0), l(0), 0) = 0, i = 1, 2$. Note that

$$\partial_k \alpha_1(k(0), l(0), 0) = (p-1)\mu_1 k(0)^{p-2} > 0,$$

$$\partial_l \alpha_2(k(0), l(0), 0) = (p-1)\mu_2 l(0)^{p-2} > 0,$$

$$\partial_l \alpha_1(k(0), l(0), 0) = \partial_k \alpha_2(k(0), l(0), 0) = 0,$$

which implies that

$$\det \left(\begin{array}{cc} \partial_k \alpha_1(k(0), l(0), 0) & \partial_l \alpha_1(k(0), l(0), 0) \\ \partial_k \alpha_2(k(0), l(0), 0) & \partial_l \alpha_2(k(0), l(0), 0) \end{array} \right) > 0.$$

Therefore, by the implicit function theorem, $k(\beta)$, $l(\beta)$ are well defined and class C^1 on $(-\beta_2, \beta_2)$ for some $\beta_2 > 0$, and $\alpha_i(k(\beta), l(\beta), \beta) \equiv 0, i = 1, 2$. This implies that $(\sqrt{k(\beta)}U_{\varepsilon,y}, \sqrt{l(\beta)}U_{\varepsilon,y})$ is a positive solution of (1.14). Note that $2p = 2^*$, This implies that

$$\lim_{\beta \to 0} \left(k(\beta) + l(\beta) \right) = k(0) + l(0) = \mu_1^{-\frac{N-2}{2}} + \mu_2^{-\frac{N-2}{2}},$$

that is, there exists $0 < \beta_1 \le \beta_2$, such that

$$k(\beta)+l(\beta) > \min\left\{\mu_1^{-\frac{N-2}{2}}, \quad \mu_2^{-\frac{N-2}{2}}\right\}, \quad \forall \ \beta \in (0,\beta_1).$$

Combining this with (2.23) and (2.32), we have

$$I(U,V) = A' = A < I(\sqrt{k(\beta)}U_{\varepsilon,y}, \sqrt{l(\beta)}U_{\varepsilon,y}), \quad \forall \beta \in (0,\beta_1),$$

that is, $(\sqrt{k(\beta)}U_{\varepsilon,y}, \sqrt{l(\beta)}U_{\varepsilon,y})$ is different positive solution of (1.14) with respect to (U, V). This completes the proof.

Before ending this section, we need to study the following properties of (U, V) obtained in Theorem 1.6.

Proposition 2.1. Assume that $\beta > 0$. Let (U, V) be a positive radially symmetric least energy solution of (1.14) obtained in Thoerem 1.6. Then there exists C > 0 such that

$$U(x) + V(x) \le C(1+|x|)^{2-N}, \quad |\nabla U(x)| + |\nabla V(x)| \le C(1+|x|)^{1-N}.$$

Proof. Define the Kelvin transformation:

$$U^*(x) := |x|^{2-N} U\left(\frac{x}{|x|^2}\right), \quad V^*(x) := |x|^{2-N} V\left(\frac{x}{|x|^2}\right).$$

Then $U^*, V^* \in D^{1,2}(\mathbb{R}^N)$ and (U^*, V^*) satisfies the same system (1.14). Then by a standard Brezis-Kato type argument ([8]), we see that $U^*, V^* \in L^{\infty}(\mathbb{R}^N)$. Therefore, there exists C > 0 such that

$$U(x) + V(x) \le C|x|^{2-N}. (2.39)$$

On the other hand, note that U, V are radially symmetric decreasing. We also have $U, V \in L^{\infty}(\mathbb{R}^N)$, and so

$$U(x) + V(x) \le C(1 + |x|)^{2-N}$$
.

Moreover, standard elliptic regularity theory implies that $U, V \in C^2(\mathbb{R}^N)$. We write U(|x|) = U(x) for convenience. Then

$$(r^{N-1}U_r)_r = -r^{N-1}(\mu_1 U^{2^*-1} + \beta U^{2^*/2-1} V^{2^*/2}),$$

and so for any $R \geq 1$, we see from (2.39) that

$$R^{N-1}|U_r(R)| \le |U_r(1)| + \int_1^R r^{N-1} (\mu_1 U^{2^*-1} + \beta U^{2^*/2-1} V^{2^*/2}) dr$$

$$\le C + C \int_1^{+\infty} r^{N-1} r^{-N-2} dr \le C.$$

Therefore, it is easy to see that $|\nabla U(x)| \leq C(1+|x|)^{1-N}$ for some C>0. Similarly, $|\nabla V(x)| \leq C(1+|x|)^{1-N}$.

3 Proof of Theorem 1.3

In this section, we assume that $-\lambda_1(\Omega) < \lambda_1 \le \lambda_2 < 0$. Recalling the definition of B in (1.5), since

$$\int_{\Omega} (|\nabla u|^2 + \lambda_i u^2) \ge \left(1 + \frac{\lambda_i}{\lambda_1(\Omega)}\right) \int_{\Omega} |\nabla u|^2, \quad i = 1, 2,$$

it is standard to see that B > 0. As has been pointed out in Section 1, by [9] the Brezis-Nirenberg problem (1.3)

$$-\Delta u + \lambda_i u = \mu_i |u|^{2^* - 2} u, \qquad u \in H_0^1(\Omega)$$

has a positive least energy solution $u_{\mu_i} \in C^2(\Omega) \cap C(\overline{\Omega})$ with energy

$$\frac{1}{N} \left(\frac{\lambda_1(\Omega) + \lambda_i}{\lambda_1(\Omega)} \right)^{\frac{N}{2}} \mu_i^{-\frac{N-2}{2}} S^{N/2} \le B_{\mu_i} := \frac{1}{2} \int_{\Omega} (|\nabla u_{\mu_i}|^2 + \lambda_i u_{\mu_i}^2) - \frac{1}{2^*} \int_{\Omega} \mu_i u_{\mu_i}^{2^*} < \frac{1}{N} \mu_i^{-\frac{N-2}{2}} S^{N/2}, \quad i = 1, 2.$$
(3.1)

The next lemma is very important, where we need the assumption $\lambda_1, \lambda_2 < 0$.

Lemma 3.1. Let $\beta < 0$, then

$$B < \min \left\{ B_{\mu_1} + \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{N/2}, \ B_{\mu_2} + \frac{1}{N} \mu_1^{-\frac{N-2}{2}} S^{N/2}, \ A \right\}.$$

Proof. The idea of this proof comes from [16], but some arguments are more delicate. Let $\beta < 0$. Let $t_0 > 0$ such that

$$\frac{N}{2}B_{\mu_1}t^2 - \frac{N}{4p}B_{\mu_1}t^{2p} + \frac{1}{N}(\mu_1/2)^{-\frac{N-2}{2}}S^{N/2} < 0, \quad \forall \ t > t_0. \eqno(3.2)$$

Since $u_{\mu_1} \in C(\overline{\Omega})$ and $u_{\mu_1} \equiv 0$ on $\partial \Omega$, there exists $B(y_0, 2R) := \{x : |x - y_0| \le 2R\} \subset \Omega$ such that

$$\delta := \max_{B(y_0, 2R)} u_{\mu_1} \le \min \left\{ \left(\frac{\mu_2}{2|\beta|} \right)^{\frac{1}{p-1}}, \left(\frac{\lambda_1 + \lambda_1(\Omega)}{2|\beta|} \right)^{\frac{1}{p-1}} \right\}. \tag{3.3}$$

Let $\psi \in C_0^1(B(y_0, 2R))$ be a function with $0 \le \psi \le 1$ and $\psi \equiv 1$ for $|x - y_0| \le R$. Define $v_{\varepsilon} = \psi U_{\varepsilon, y_0}$, where U_{ε, y_0} is defined in (1.16) and (1.17). Then by [9] or [37, Lemma 1.46], we have the following inequalities

$$\int_{\Omega} |\nabla v_{\varepsilon}|^2 = S^{N/2} + O(\varepsilon^{N-2}), \quad \int_{\Omega} |v_{\varepsilon}|^{2^*} = S^{N/2} + O(\varepsilon^N), \tag{3.4}$$

$$\int_{\Omega} |v_{\varepsilon}|^2 \ge C\varepsilon^2 + O(\varepsilon^{N-2}). \tag{3.5}$$

Moreover, since $N \geq 5$, we have

$$\int_{\Omega} v_{\varepsilon}^{\frac{N}{N-2}} dx \le \int_{B(y_0, 2R)} U_{\varepsilon, y_0}^{\frac{N}{N-2}} dx = C \int_{B(0, 2R)} \left(\frac{\varepsilon}{\varepsilon^2 + |x|^2} \right)^{N/2} dx
\le C \varepsilon^{N/2} \left(\ln \frac{2R}{\varepsilon} + 1 \right) = o(\varepsilon^2).$$
(3.6)

Since $supp(v_{\varepsilon}) \subset B(y_0, 2R)$, by (3.3) we have for t, s > 0 that

$$2|\beta|t^{p}s^{p}\int_{\Omega}u_{\mu_{1}}^{p}v_{\varepsilon}^{p} \leq 2|\beta|\delta^{p-1}t^{p}s^{p}\int_{\Omega}u_{\mu_{1}}v_{\varepsilon}^{p}$$

$$\leq |\beta|\delta^{p-1}t^{2p}\int_{\Omega}u_{\mu_{1}}^{2} + |\beta|\delta^{p-1}s^{2p}\int_{\Omega}v_{\varepsilon}^{2p}$$

$$\leq \frac{|\beta|\delta^{p-1}}{\lambda_{1} + \lambda_{1}(\Omega)}t^{2p}\int_{\Omega}(|\nabla u_{\mu_{1}}|^{2} + \lambda_{1}u_{\mu_{1}}^{2}) + |\beta|\delta^{p-1}s^{2p}\int_{\Omega}v_{\varepsilon}^{2p}$$

$$\leq \frac{1}{2}t^{2p}\int_{\Omega}\mu_{1}u_{\mu_{1}}^{2p} + \frac{1}{2}s^{2p}\int_{\Omega}\mu_{2}v_{\varepsilon}^{2p}, \tag{3.7}$$

and so

$$E(tu_{\mu_{1}}, sv_{\varepsilon}) = \frac{1}{2}t^{2} \int_{\Omega} (|\nabla u_{\mu_{1}}|^{2} + \lambda_{1}u_{\mu_{1}}^{2}) + \frac{1}{2}s^{2} \int_{\Omega} (|\nabla v_{\varepsilon}|^{2} + \lambda_{2}v_{\varepsilon}^{2})$$

$$- \frac{1}{2p} \int_{\Omega} (t^{2p} \mu_{1}u_{\mu_{1}}^{2p} + 2t^{p}s^{p}\beta u_{\mu_{1}}^{p}v_{\varepsilon}^{p} + s^{2p}\mu_{2}v_{\varepsilon}^{2p})$$

$$\leq \frac{1}{2}t^{2} \int_{\Omega} (|\nabla u_{\mu_{1}}|^{2} + \lambda_{1}u_{\mu_{1}}^{2}) - \frac{1}{4p}t^{2p} \int_{\Omega} \mu_{1}u_{\mu_{1}}^{2p}$$

$$+ \frac{1}{2}s^{2} \int_{\Omega} (|\nabla v_{\varepsilon}|^{2} + \lambda_{2}v_{\varepsilon}^{2}) - \frac{1}{2p}s^{2p} \frac{\mu_{2}}{2} \int_{\Omega} v_{\varepsilon}^{2p}$$

$$= f(t) + g(s). \tag{3.8}$$

By (3.4)-(3.5), it is standard to check that (cf. [9, 37])

$$\max_{s>0} g(s) < \frac{1}{N} (\mu_2/2)^{-\frac{N-2}{2}} S^{N/2} \quad \text{for } \varepsilon \text{ small enough.}$$
 (3.9)

By (3.1) we see that

$$f(t) = \frac{N}{2}B_{\mu_1}t^2 - \frac{N}{4p}B_{\mu_1}t^{2p}.$$

Combining these with (3.2), we get that

$$f(t) + g(s) < 0, \quad \forall t > t_0, \ s > 0,$$

and so it follows from (3.8) that

$$\max_{t,s>0} E(tu_{\mu_1},sv_{\varepsilon}) = \max_{0 < t \leq t_0,s>0} E(tu_{\mu_1},sv_{\varepsilon}).$$

Define

$$g_{\varepsilon}(s) := \frac{1}{2}s^2 \int_{\Omega} (|\nabla v_{\varepsilon}|^2 + \lambda_2 v_{\varepsilon}^2) \, dx - \frac{s^{2p}}{2p} \int_{\Omega} \mu_2 v_{\varepsilon}^{2p} \, dx, \quad s > 0.$$

Then there exists a unique $s(\varepsilon) > 0$, such that $g'_{\varepsilon}(s(\varepsilon)) = 0$ with

$$\begin{split} s(\varepsilon)^{2p-2} &= \frac{\int_{\Omega} (|\nabla v_{\varepsilon}|^2 + \lambda_2 v_{\varepsilon}^2) \, dx}{\int_{\Omega} \mu_2 v_{\varepsilon}^{2p} \, dx} \geq \left(1 + \frac{\lambda_2}{\lambda_1(\Omega)}\right) \frac{\int_{\Omega} |\nabla v_{\varepsilon}|^2 \, dx}{\mu_2 \int_{\mathbb{R}^N} U_{\varepsilon,y_0}^{2p} \, dx} \\ &= \left(1 + \frac{\lambda_2}{\lambda_1(\Omega)}\right) \frac{S^{N/2} + O(\varepsilon^{N-2})}{\mu_2 S^{N/2}} \\ &\geq \frac{1}{2\mu_2} \left(1 + \frac{\lambda_2}{\lambda_1(\Omega)}\right) =: s_0^{2p-2}, \quad \text{for } \varepsilon \text{ small enough.} \end{split}$$

Therefore, since g_{ε} is increasing for $0 < s \le s(\varepsilon)$, for any $0 < s < s_0$, we have $g_{\varepsilon}(s) < g_{\varepsilon}(s_0)$ and so $E(tu_{\mu_1}, sv_{\varepsilon}) < E(tu_{\mu_1}, s_0v_{\varepsilon})$. That is,

$$\max_{t,s>0} E(tu_{\mu_1}, sv_{\varepsilon}) = \max_{0 < t \le t_0, s \ge s_0} E(tu_{\mu_1}, sv_{\varepsilon}). \tag{3.10}$$

For $0 < t \le t_0, s \ge s_0$, we see from (3.6) that

$$|\beta| t^p s^p \int_{\Omega} u^p_{\mu_1} v^p_{\varepsilon} \le |\beta| t^p_0 \delta^p s^{p-2}_0 s^2 \int_{\Omega} v^p_{\varepsilon} \le C s^2 \cdot o(\varepsilon^2),$$

and so

$$E(tu_{\mu_{1}}, sv_{\varepsilon}) = \frac{1}{2}t^{2} \int_{\Omega} (|\nabla u_{\mu_{1}}|^{2} + \lambda_{1}u_{\mu_{1}}^{2}) + \frac{1}{2}s^{2} \int_{\Omega} (|\nabla v_{\varepsilon}|^{2} + \lambda_{2}v_{\varepsilon}^{2})$$

$$- \frac{1}{2p} \int_{\Omega} (t^{2p} \mu_{1}u_{\mu_{1}}^{2p} + 2t^{p}s^{p}\beta u_{\mu_{1}}^{p}v_{\varepsilon}^{p} + s^{2p}\mu_{2}v_{\varepsilon}^{2p})$$

$$\leq \frac{1}{2}t^{2} \int_{\Omega} (|\nabla u_{\mu_{1}}|^{2} + \lambda_{1}u_{\mu_{1}}^{2}) - \frac{1}{2p}t^{2p} \int_{\Omega} \mu_{1}u_{\mu_{1}}^{2p}$$

$$+ \frac{1}{2}s^{2} \left(\int_{\Omega} (|\nabla v_{\varepsilon}|^{2} + \lambda_{2}v_{\varepsilon}^{2}) + o(\varepsilon^{2}) \right) - \frac{1}{2p}s^{2p}\mu_{2} \int_{\Omega} v_{\varepsilon}^{2p}$$

$$=: f_{1}(t) + g_{1}(s). \tag{3.11}$$

Note that $\max_{t>0} f_1(t) = f_1(1) = B_{\mu_1}$. By (3.4)-(3.5) and $\lambda_2 < 0$, it is easy to show that

$$\max_{s>0} g_1(s) < \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{N/2} \quad \text{for } \varepsilon \text{ small enough.}$$

Combining these with (3.10) and (3.11), we obtain that

$$\max_{t,s>0} E(tu_{\mu_1}, sv_{\varepsilon}) = \max_{0 < t \le t_0, s \ge s_0} E(tu_{\mu_1}, sv_{\varepsilon})$$

$$\leq \max_{t>0} f_1(t) + \max_{s>0} g_1(s)$$

$$< B_{\mu_1} + \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{N/2} \quad \text{for } \varepsilon \text{ small enough.}$$
(3.12)

Now, we claim that there exists $t_{\varepsilon}, s_{\varepsilon} > 0$ such that $(t_{\varepsilon}u_{\mu_1}, s_{\varepsilon}v_{\varepsilon}) \in \mathcal{M}$. Similarly as (3.7), we have

$$\left(\int_{\Omega} \beta u_{\mu_{1}}^{p} v_{\varepsilon}^{p} dx\right)^{2} \leq |\beta|^{2} \delta^{2p-2} \left(\int_{\Omega} u_{\mu_{1}} v_{\varepsilon}^{p} dx\right)^{2}
\leq |\beta|^{2} \delta^{2p-2} \int_{\Omega} u_{\mu_{1}}^{2} dx \int_{\Omega} v_{\varepsilon}^{2p} dx
\leq \frac{|\beta|^{2} \delta^{2p-2}}{(\lambda_{1}(\Omega) + \lambda_{1})\mu_{2}} \int_{\Omega} \mu_{1} u_{\mu_{1}}^{2p} dx \int_{\Omega} \mu_{2} v_{\varepsilon}^{2p} dx
< \int_{\Omega} \mu_{1} u_{\mu_{1}}^{2p} dx \int_{\Omega} \mu_{2} v_{\varepsilon}^{2p} dx.$$

For convenience we denote

$$D_1 = \int_{\Omega} \mu_1 u_{\mu_1}^{2p} dx, \quad D_2 = \int_{\Omega} \beta u_{\mu_1}^p v_{\varepsilon}^p dx,$$
$$D_3 = \int_{\Omega} \mu_2 v_{\varepsilon}^{2p} dx, \quad D_4 = \int_{\Omega} (|\nabla v_{\varepsilon}|^2 + \lambda_2 v_{\varepsilon}^2) dx.$$

Then $D_2 < 0$ and $D_1D_3 - D_2^2 > 0$. Furthermore, $(tu_{\mu_1}, sv_{\varepsilon}) \in \mathcal{M}$ for some t, s > 0 is equivalent to

$$t^{2-p}D_1 = t^p D_1 + s^p D_2, \quad s^{2-p}D_4 = s^p D_3 + t^p D_2, \quad s, t > 0.$$
 (3.13)

Note that $1 , by <math>s^p = (t^{2-p} - t^p)D_1/D_2 > 0$ we have t > 1. Therefore, (3.13) is equivalent to

$$f_3(t) := D_4 \left(\frac{D_1}{|D_2|} (1 - t^{2-2p}) \right)^{\frac{2-p}{p}} - \frac{D_1 D_3 - D_2^2}{|D_2|} t^{2p-2} + \frac{D_1 D_3}{|D_2|}, \ t > 1. \ (3.14)$$

Since $f_3(1) > 0$ and $\lim_{t \to +\infty} f_3(t) < 0$, (3.14) has a solution t > 1. Hence (3.13) has a solution $t_{\varepsilon} > 0$, $s_{\varepsilon} > 0$. That is, $(t_{\varepsilon}u_{\mu_1}, s_{\varepsilon}v_{\varepsilon}) \in \mathcal{M}$ and from (3.12) we get

$$B \le E(t_{\varepsilon}u_{\mu_1}, s_{\varepsilon}v_{\varepsilon}) \le \max_{t, s > 0} E(tu_{\mu_1}, sv_{\varepsilon}) < B_{\mu_1} + \frac{1}{N}\mu_2^{-\frac{N-2}{2}}S^{N/2}$$

By a similar argument, we can also prove that $B < B_{\mu_2} + \frac{1}{N}\mu_1^{-\frac{N-2}{2}}S^{N/2}$. By (2.22) and (3.1), we have

$$A > \max \left\{ B_{\mu_1} + \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{N/2}, \ B_{\mu_2} + \frac{1}{N} \mu_1^{-\frac{N-2}{2}} S^{N/2} \right\}$$

This completes the proof.

Lemma 3.2. Assume that $\beta < 0$, then there exists $C_2 > C_1 > 0$, such that for any $(u, v) \in \mathcal{M}$ with $E(u, v) \leq A$, there holds

$$C_1 \le \int_{\Omega} |u|^{2p} dx, \int_{\Omega} |v|^{2p} dx \le C_2.$$

Proof. This follows directly from

$$\frac{\lambda_1(\Omega) + \lambda_1}{\lambda_1(\Omega)} S\left(\int_{\Omega} |u|^{2p}\right)^{\frac{1}{p}} \le \int_{\Omega} (|\nabla u|^2 + \lambda_1 u^2) \le \mu_1 \int_{\Omega} |u|^{2p},$$
$$\frac{\lambda_1(\Omega) + \lambda_2}{\lambda_1(\Omega)} S\left(\int_{\Omega} |v|^{2p}\right)^{\frac{1}{p}} \le \int_{\Omega} (|\nabla v|^2 + \lambda_2 v^2) \le \mu_1 \int_{\Omega} |v|^{2p},$$

$$E(u,v) \leq A \text{ and } (1.5).$$

Lemma 3.3. Let $u_n \rightharpoonup u, v_n \rightharpoonup v$ in $H_0^1(\Omega)$ as $n \to \infty$, then passing to a subsequence, there holds

$$\lim_{n \to \infty} \int_{\Omega} (|u_n|^p |v_n|^p - |u_n - u|^p |v_n - v|^p - |u|^p |v|^p) \ dx = 0.$$

Proof. Note that $2p = 2^*$, we have

$$u_n \to u$$
, $v_n \to v$ strongly in $L^q(\Omega)$, $\forall 0 < q < 2p$, $u_n \rightharpoonup u$, $v_n \rightharpoonup v$ weakly in $L^{2p}(\Omega)$.

Fix any $t \in [0,1]$. First, we claim that

$$|u_n - tu|^{p-2}(u_n - tu)|v_n|^p \rightharpoonup (1-t)^{p-1}|u|^{p-2}u|v|^p$$
 weakly in $L^{\frac{2p}{2p-1}}(\Omega)$. (3.15)

Since the map $h: L^{q_1}(\Omega) \to L^{q_1/q_2}(\Omega)$ with $h(s) = |s|^{q_2-1}s$ is continuous, so

$$|u_n - tu|^{p-2}(u_n - tu) \to (1-t)^{p-1}|u|^{p-2}u$$
 strongly in $L^q(\Omega)$, $\forall \ 0 < q < \frac{2p}{p-1}$, $|v_n|^p \to |v|^p$ strongly in $L^q(\Omega)$, $\forall \ 0 < q < 2$.

Then for any $1 \le q < \frac{2p}{2p-1}$, one has

$$|u_n - tu|^{p-2}(u_n - tu)|v_n|^p \to (1-t)^{p-1}|u|^{p-2}u|v|^p$$
 strongly in $L^q(\Omega)$.

Since $|u_n - tu|^{p-2}(u_n - tu)|v_n|^p$ is uniformly bounded in $L^{\frac{2p}{2p-1}}(\Omega)$, passing to a subsequence, we may assume that $|u_n - tu|^{p-2}(u_n - tu)|v_n|^p \rightharpoonup w$ weakly in $L^{\frac{2p}{2p-1}}(\Omega)$. Then for any $\varphi \in C_0^{\infty}(\mathbb{R}^N)$, we have

$$\int_{\Omega} w\varphi = \lim_{n \to \infty} \int_{\Omega} |u_n - tu|^{p-2} (u_n - tu) |v_n|^p \varphi = \int_{\Omega} (1 - t)^{p-1} |u|^{p-2} u |v|^p \varphi,$$

which implies $w=(1-t)^{p-1}|u|^{p-2}u|v|^p$, that is, (3.15) holds. Similarly, we can show that $|u_n-u|^p|v_n-tv|^{p-2}(v_n-tv) \to 0$ weakly in $L^{\frac{2p}{2p-1}}(\Omega)$. Therefore, by (3.15), the Fubini Theorem and the Dominated Convergent Theorem,

$$\int_{\Omega} (|u_n|^p |v_n|^p - |u_n - u|^p |v_n - v|^p) \ dx$$

$$= p \int_{\Omega} \int_{0}^{1} |u_{n} - tu|^{p-2} (u_{n} - tu) |v_{n}|^{p} u \, dt \, dx$$

$$+ p \int_{\Omega} \int_{0}^{1} |u_{n} - u|^{p} |v_{n} - tv|^{p-2} (v_{n} - tv) v \, dt \, dx$$

$$= p \int_{0}^{1} \int_{\Omega} |u_{n} - tu|^{p-2} (u_{n} - tu) |v_{n}|^{p} u \, dx \, dt$$

$$+ p \int_{0}^{1} \int_{\Omega} |u_{n} - u|^{p} |v_{n} - tv|^{p-2} (v_{n} - tv) v \, dx \, dt$$

$$\to p \int_{0}^{1} \int_{\Omega} (1 - t)^{p-1} |u|^{p} |v|^{p} \, dx \, dt = \int_{\Omega} |u|^{p} |v|^{p} \, dx, \text{ as } n \to \infty.$$

This completes the proof.

Proof of Theorem 1.3 for the case $\beta < 0$. The main idea of the proof is similar to the proof of [16, Theorem 1.3-(1)] in case N = 4, but as we will see, some new ideas are needed. Assume that $\beta < 0$. Note that E is coercive and bounded from below on \mathcal{M} . Then by the Ekeland variational priciple (cf. [30]), there exists a minimizing sequence $\{(u_n, v_n)\} \subset \mathcal{M}$ satisfying

$$E(u_n, v_n) \le \min\left\{B + \frac{1}{n}, A\right\},\tag{3.16}$$

$$E(u,v) \ge E(u_n,v_n) - \frac{1}{n} \|(u_n,v_n) - (u,v)\|, \quad \forall (u,v) \in \mathcal{M}.$$
 (3.17)

Here, $\|(u,v)\| := (\int_{\Omega} (|\nabla u|^2 + |\nabla v|^2) dx)^{1/2}$ is the norm of H. Then $\{(u_n, v_n)\}$ is bounded in H. For any $(\varphi, \phi) \in H$ with $\|\varphi\|, \|\phi\| \le 1$ and each $n \in \mathbb{N}$, we define the functions h_n and $g_n : \mathbb{R}^3 \to \mathbb{R}$ by

$$h_{n}(t,s,l) = \int_{\Omega} |\nabla(u_{n} + t\varphi + su_{n})|^{2} + \lambda_{1} \int_{\Omega} |u_{n} + t\varphi + su_{n}|^{2}$$

$$- \mu_{1} \int_{\Omega} |u_{n} + t\varphi + su_{n}|^{2p} - \beta \int_{\Omega} |u_{n} + t\varphi + su_{n}|^{p} |v_{n} + t\varphi + lv_{n}|^{p}, \quad (3.18)$$

$$g_{n}(t,s,l) = \int_{\Omega} |\nabla(v_{n} + t\varphi + lv_{n})|^{2} + \lambda_{2} \int_{\Omega} |v_{n} + t\varphi + lv_{n}|^{2}$$

$$- \mu_{2} \int_{\Omega} |v_{n} + t\varphi + lv_{n}|^{2p} - \beta \int_{\Omega} |u_{n} + t\varphi + su_{n}|^{p} |v_{n} + t\varphi + lv_{n}|^{p}. \quad (3.19)$$

Let $\mathbf{0} = (0, 0, 0)$. Then $h_n, g_n \in C^1(\mathbb{R}^3, \mathbb{R}), h_n(\mathbf{0}) = g_n(\mathbf{0}) = 0$ and

$$\frac{\partial h_n}{\partial s}(\mathbf{0}) = -(2p-2) \int_{\Omega} (|\nabla u_n|^2 + \lambda_1 u_n^2) + p\beta \int_{\Omega} |u_n|^p |v_n|^p,$$

$$\frac{\partial h_n}{\partial l}(\mathbf{0}) = \frac{\partial g_n}{\partial s}(\mathbf{0}) = -p\beta \int_{\Omega} |u_n|^p |v_n|^p dx,$$

$$\frac{\partial g_n}{\partial l}(\mathbf{0}) = -(2p-2) \int_{\Omega} (|\nabla v_n|^2 + \lambda_2 v_n^2) + p\beta \int_{\Omega} |u_n|^p |v_n|^p.$$

Define the matrix

$$F_n := \begin{pmatrix} \frac{\partial h_n}{\partial s}(\mathbf{0}) & \frac{\partial h_n}{\partial l}(\mathbf{0}) \\ \frac{\partial g_n}{\partial s}(\mathbf{0}) & \frac{\partial g_n}{\partial l}(\mathbf{0}) \end{pmatrix}.$$

Since $\beta < 0$, it follows from Lemma 3.2 that

$$\det(F_n) \ge (2p - 2)^2 \int_{\Omega} (|\nabla u_n|^2 + \lambda_1 |u_n|^2) \int_{\Omega} (|\nabla v_n|^2 + \lambda_2 |v_n|^2)$$

$$\ge CS^2 \left(\int_{\Omega} |u_n|^{2p} \right)^{\frac{1}{p}} \left(\int_{\Omega} |v_n|^{2p} \right)^{\frac{1}{p}} \ge C > 0, \tag{3.20}$$

where C is independent of n. By the implicit function theorem, functions $s_n(t)$ and $l_n(t)$ are well defined and class C^1 on some interval $(-\delta_n, +\delta_n)$ for $\delta_n > 0$. Moreover, $s_n(0) = l_n(0) = 0$ and

$$h_n(t, s_n(t), l_n(t)) \equiv 0, \quad q_n(t, s_n(t), l_n(t)) \equiv 0, \quad t \in (-\delta, +\delta).$$

With these, it is standard to prove that (see [16, Theorem 1.3-(1)] for instance)

$$\lim_{n \to +\infty} E'(u_n, v_n) = 0. \tag{3.21}$$

Since $\{(u_n, v_n)\}$ is bounded in H, we may assume that $(u_n, v_n) \rightharpoonup (u, v)$ weakly in H. Passing to a subsequence, we may assume that

$$u_n \rightharpoonup u, \quad v_n \rightharpoonup v, \quad \text{weakly in } L^{2p}(\Omega),$$
$$|u_n|^{q-1}u_n \rightharpoonup |u|^{q-1}u, \quad |v_n|^{q-1}v_n \rightharpoonup |v|^{q-1}v, \quad \text{weakly in } L^{2p/q}(\Omega), \ 1 < q < 2p,$$
$$u_n \rightarrow u, \quad v_n \rightarrow v, \quad \text{strongly in } L^2(\Omega).$$

Thus, by (3.21) we have E'(u, v) = 0. Set $\omega_n = u_n - u$ and $\sigma_n = v_n - v$. Then by Brezis-Lieb Lemma (cf. [37]), there holds

$$|u_n|_{2p}^{2p} = |u|_{2p}^{2p} + |\omega_n|_{2p}^{2p} + o(1), \quad |v_n|_{2p}^{2p} = |v|_{2p}^{2p} + |\sigma_n|_{2p}^{2p} + o(1).$$
 (3.22)

Note that $(u_n, v_n) \in \mathcal{M}$ and E'(u, v) = 0. Combining these with (3.22), Lemma 3.3, we get that

$$\int_{\Omega} |\nabla \omega_n|^2 - \int_{\Omega} (\mu_1 |\omega_n|^{2p} + \beta |\omega_n|^p |\sigma_n|^p) = o(1), \tag{3.23}$$

$$\int_{\Omega} |\nabla \sigma_n|^2 - \int_{\Omega} (\mu_2 |\sigma_n|^{2p} + \beta |\omega_n|^p |\sigma_n|^p) = o(1), \tag{3.24}$$

$$E(u_n, v_n) = E(u, v) + I(\omega_n, \sigma_n) + o(1).$$
(3.25)

Passing to a subsequence, we may assume that

$$\lim_{n \to +\infty} \int_{\Omega} |\nabla \omega_n|^2 = b_1, \quad \lim_{n \to +\infty} \int_{\Omega} |\nabla \sigma_n|^2 = b_2.$$

Then by (3.23) and (3.24) we have $I(\omega_n, \sigma_n) = \frac{1}{N}(b_1 + b_2) + o(1)$. Letting $n \to +\infty$ in (3.25), we get that

$$0 \le E(u, v) \le E(u, v) + \frac{1}{N}(b_1 + b_2) = \lim_{n \to +\infty} E(u_n, v_n) = B.$$
 (3.26)

Case 1. $u \equiv 0, v \equiv 0$.

By Lemma 3.2, (3.22) and (3.26), we have $0 < b_1 < +\infty$ and $0 < b_2 < +\infty$, and we may assume that both $\omega_n \not\equiv 0$ and $\sigma_n \not\equiv 0$ for n large. Then by (3.23) and (3.24) we have

$$\int_{\Omega} \mu_1 |\omega_n|^{2p} \int_{\Omega} \mu_2 |\sigma_n|^{2p} - \left(\beta \int_{\Omega} |\omega_n|^p |\sigma_n|^p\right)^2 > 0, \quad \text{for } n \text{ large.}$$

Then by a similar argument as Lemma 3.1, for n large, there exists $t_n, s_n > 0$ such that $(t_n \omega_n, s_n \sigma_n) \in \mathcal{N}$. Up to a subsequence, we claim that

$$\lim_{n \to +\infty} (|t_n - 1| + |s_n - 1|) = 0. \tag{3.27}$$

This conclusion is obvious in case N=4 and p=2 (see [16]), but it is not trivial in our general case $N\geq 5$ here. Denote

$$B_{n,1} = \int_{\Omega} |\nabla \omega_n|^2 \to b_1, \quad B_{n,2} = \int_{\Omega} |\nabla \sigma_n|^2 \to b_2,$$

$$C_{n,1} = \int_{\Omega} \mu_1 |\omega_n|^{2p}, \quad C_{n,2} = \int_{\Omega} \mu_2 |\sigma_n|^{2p},$$

$$D_n = |\beta| \int_{\Omega} |\omega_n|^p |\sigma_n|^p.$$

Passing to a subsequence, we may assume that $C_{n,1} \to c_1 < +\infty$, $C_{n,2} \to c_2 < +\infty$ and $D_n \to d < +\infty$. By (3.23)-(3.24) we have

$$c_1 = b_1 + d > b_1 > 0, \quad c_2 = b_2 + d > b_2 > 0,$$
 (3.28)

$$t_n^2 B_{n,1} = t_n^{2p} C_{n,1} - t_n^p s_n^p D_n, \quad s_n^2 B_{n,2} = s_n^{2p} C_{n,2} - t_n^p s_n^p D_n.$$
 (3.29)

This implies that

$$t_n^{2p-2} \ge \frac{B_{n,1}}{C_{n,1}} \to \frac{b_1}{c_1} > 0, \quad s_n^{2p-2} \ge \frac{B_{n,2}}{C_{n,2}} \to \frac{b_2}{c_2} > 0.$$
 (3.30)

Assume that, up to a subsequence, $t_n \to +\infty$ as $n \to \infty$, then by

$$t_n^{2p}C_{n,1} - t_n^2B_{n,1} = s_n^{2p}C_{n,2} - s_n^2B_{n,2},$$

we also have $s_n \to +\infty$. Then

$$d^{2} = \lim_{n \to \infty} D_{n}^{2} = \lim_{n \to \infty} \frac{t_{n}^{p} C_{n,1} - t_{n}^{2-p} B_{n,1}}{s_{n}^{p}} \cdot \frac{s_{n}^{p} C_{n,2} - s_{n}^{2-p} B_{n,2}}{t_{n}^{p}}$$

$$= \lim_{n \to \infty} (C_{n,1} - t_n^{2-2p} B_{n,1}) (C_{n,2} - s_n^{2-2p} B_{n,2})$$

= $c_1 c_2 = (b_1 + d)(b_2 + d) > d^2$,

a contradiction. Therefore, t_n, s_n are uniformly bounded. Passing to a subsequence, by (3.30) we may assume that $t_n \to t_\infty \ge (b_1/c_1)^{\frac{1}{2p-2}} > 0$ and $s_n \to s_\infty \ge (b_2/c_2)^{\frac{1}{2p-2}} > 0$. Then we see from (3.29) that

$$s_{\infty}^{p}d = t_{\infty}^{p}c_{1} - t_{\infty}^{2-p}b_{1}, \quad t_{\infty}^{p}d = s_{\infty}^{p}c_{2} - s_{\infty}^{2-p}b_{2}.$$

If d=0, then $c_i=b_i$, and so $t_{\infty}=s_{\infty}=1$. That is, (3.27) holds. Now we consider the case d>0. Define $f(t)=t^pc_1-t^{2-p}b_1$, then for $t\geq (b_1/c_1)^{\frac{1}{2p-2}}$, we have

$$f'(t) = pc_1t^{p-1} - (2-p)b_1t^{1-p} > (2-p)t^{1-p}(c_1t^{2p-2} - b_1) \ge 0,$$

that is, f is increasing with respect to $t \geq (b_1/c_1)^{\frac{1}{2p-2}}$. If $t_{\infty} < 1$, then

$$s_{\infty}^{p} d = f(t_{\infty}) < f(1) = c_1 - b_1 = d,$$

that is, $s_{\infty} < 1$, and we see from (3.28) that

$$d^{2} = \frac{t_{\infty}^{p} c_{1} - t_{\infty}^{2-p} b_{1}}{s_{\infty}^{p}} \cdot \frac{s_{\infty}^{p} c_{2} - s_{\infty}^{2-p} b_{2}}{t_{\infty}^{p}} = (c_{1} - t_{\infty}^{2-2p} b_{1})(c_{2} - s_{\infty}^{2-2p} b_{2})$$
$$= (d + b_{1} - t_{\infty}^{2-2p} b_{1})(d + b_{2} - s_{\infty}^{2-2p} b_{2}) < d^{2},$$

a contradiction. If $t_{\infty} > 1$, since $1 \ge (b_1/c_1)^{\frac{1}{2p-2}}$, we have

$$s_{\infty}^{p}d = f(t_{\infty}) > f(1) = c_1 - b_1 = d,$$

that is, $s_{\infty} > 1$, and so

$$d^2 = (d + b_1 - t_{\infty}^{2-2p}b_1)(d + b_2 - s_{\infty}^{2-2p}b_2) > d^2,$$

a contradiction. Therefore, $t_{\infty}=s_{\infty}=1$ and (3.27) holds. This implies that

$$\frac{1}{N}(b_1+b_2) = \lim_{n \to +\infty} I(\omega_n, \sigma_n) = \lim_{n \to +\infty} I(t_n \omega_n, s_n \sigma_n) \ge A.$$

Combining this with (3.26) we get that $B \geq A$, a contradiction with Lemma 3.1. Therefore, Case 1 is impossible.

Case 2. $u \not\equiv 0, v \equiv 0 \text{ or } u \equiv 0, v \not\equiv 0.$

Without loss of generality, we assume that $u \not\equiv 0, v \equiv 0$. Then $b_2 > 0$. By Case 1 we may assume that $b_1 = 0$. Then $\lim_{n \to +\infty} \int_{\Omega} |\omega_n|^p |\sigma_n|^p = 0$, and so

$$\int_{\Omega} |\nabla \sigma_n|^2 = \int_{\Omega} \mu_2 |\sigma_n|^{2p} + o(1) \le \mu_2 S^{-p} \left(\int_{\Omega} |\nabla \sigma_n|^2 \right)^p + o(1).$$

This implies that $b_2 \ge \mu_2^{-\frac{N-2}{2}} S^{N/2}$. Note that u is a nontrivial solution of $-\Delta u + \lambda_1 u = \mu_1 |u|^{2^*-2} u$, we have from (3.1) that $E(u,0) \ge B_{\mu_1}$. By (3.26) we get that

 $B \ge B_{\mu_1} + \frac{1}{N} b_2 \ge B_{\mu_1} + \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{N/2},$

a contradiction with Lemma 3.1. Therefore, Case 2 is impossible.

Since Cases 1 and 2 are both impossible, we have that $u \not\equiv 0, v \not\equiv 0$, that is, $(u,v) \in \mathcal{M}$. By (3.26) we have E(u,v) = B. Then $(|u|,|v|) \in \mathcal{M}$ and E(|u|,|v|) = B. By Lemma 2.5, (|u|,|v|) is a solution of (1.1). Then, using the maximum principle, we see that |u|,|v| > 0 in Ω . Therefore, (|u|,|v|) is a positive least energy solution of (1.1). This completes the proof.

It remains to prove Theorem 1.3 for the case $\beta > 0$. Let $\beta > 0$ and define

$$\mathcal{B} := \inf_{h \in \Gamma} \max_{t \in [0,1]} E(h(t)), \tag{3.31}$$

where $\Gamma = \{h \in C([0,1], H) : h(0) = (0,0), E(h(1)) < 0\}$. By (1.4), we see that for any $(u,v) \in H$, $(u,v) \neq (0,0)$,

$$\max_{t>0} E(tu, tv) = E(t_{u,v}u, t_{u,v}v)
= \frac{1}{N} t_{u,v}^2 \int_{\Omega} (|\nabla u|^2 + \lambda_1 u^2 + |\nabla v|^2 + \lambda_2 v^2)
= \frac{1}{N} t_{u,v}^{2^*} \int_{\Omega} (\mu_1 |u|^{2p} + 2\beta |u|^p |v|^p + \mu_2 |v|^{2p}),$$
(3.32)

where $t_{u,v} > 0$ satisfies

$$t_{u,v}^{2p-2} = \frac{\int_{\Omega} (|\nabla u|^2 + \lambda_1 u^2 + |\nabla v|^2 + \lambda_2 v^2)}{\int_{\Omega} (\mu_1 |u|^{2p} + 2\beta |u|^p |v|^p + \mu_2 |v|^{2p})}.$$
 (3.33)

Note that $(t_{u,v}u, t_{u,v}v) \in \mathcal{M}'$, where

$$\mathcal{M}' := \left\{ (u, v) \in H \setminus \{(0, 0)\}, \ G(u, v) := \int_{\Omega} (|\nabla u|^2 + \lambda_1 u^2 + |\nabla v|^2 + \lambda_2 v^2) - \int_{\Omega} (\mu_1 |u|^{2p} + 2\beta |u|^p |v|^p + \mu_2 |v|^{2p}) = 0 \right\}, \quad (3.34)$$

it is easy to check that

$$\mathcal{B} = \inf_{H \ni (u,v) \neq (0,0)} \max_{t>0} E(tu, tv) = \inf_{(u,v) \in \mathcal{M}'} E(u,v).$$
(3.35)

Note that $\mathcal{M} \subset \mathcal{M}'$, one has that $\mathcal{B} \leq B$. Similarly as (2.20), we have $\mathcal{B} > 0$.

Lemma 3.4. Let $\beta > 0$, then

$$\mathcal{B} < \min\{B_{\mu_1}, B_{\mu_2}, A\}.$$

Proof. Step 1. We prove that $\mathcal{B} < A$. Without loss of generality, we may assume that $0 \in \Omega$. Then there exists $\rho > 0$ such that $B(0, 2\rho) := \{x : |x| \le 2\rho\} \subset \Omega$. Let $\psi \in C_0^1(B(0, 2\rho))$ be a nonnegative function with $0 \le \psi \le 1$ and $\psi \equiv 1$ for $|x| \le \rho$. Recall that (U, V) in Theorem 1.6. we define

$$(U_{\varepsilon}(x), V_{\varepsilon}(x)) := \left(\varepsilon^{-\frac{N-2}{2}} U\left(\frac{x}{\varepsilon}\right), \ \varepsilon^{-\frac{N-2}{2}} V\left(\frac{x}{\varepsilon}\right)\right).$$

Then it is easy to see that

$$\int_{\mathbb{R}^N} |\nabla U_{\varepsilon}|^2 = \int_{\mathbb{R}^N} |\nabla U|^2, \quad \int_{\mathbb{R}^N} |U_{\varepsilon}|^{2^*} = \int_{\mathbb{R}^N} |U|^{2^*},$$
$$\int_{\mathbb{R}^N} |\nabla V_{\varepsilon}|^2 = \int_{\mathbb{R}^N} |\nabla V|^2, \quad \int_{\mathbb{R}^N} |V_{\varepsilon}|^{2^*} = \int_{\mathbb{R}^N} |V|^{2^*}.$$

Define

$$(u_{\varepsilon}, v_{\varepsilon}) := (\psi U_{\varepsilon}, \psi V_{\varepsilon}). \tag{3.36}$$

First we claim the following inequalities

$$\int_{\Omega} |\nabla u_{\varepsilon}|^2 \le \int_{\mathbb{R}^N} |\nabla U|^2 + O(\varepsilon^{N-2}), \tag{3.37}$$

$$\int_{\Omega} |u_{\varepsilon}|^{2^*} \ge \int_{\mathbb{R}^N} |U|^{2^*} + O(\varepsilon^N), \tag{3.38}$$

$$\int_{\Omega} |u_{\varepsilon}|^{\frac{2^{*}}{2}} |v_{\varepsilon}|^{\frac{2^{*}}{2}} \ge \int_{\mathbb{R}^{N}} |U|^{\frac{2^{*}}{2}} |V|^{\frac{2^{*}}{2}} + O(\varepsilon^{N}), \tag{3.39}$$

$$\int_{\Omega} |u_{\varepsilon}|^2 \ge C\varepsilon^2 + O(\varepsilon^{N-2}),\tag{3.40}$$

where C is a positive constant.

Let $0 < \varepsilon \ll \rho$. By Proposition 2.1 we have

$$\begin{split} \int_{\Omega} |\nabla \psi|^2 |U_{\varepsilon}|^2 \, dx &\leq C \int_{\rho \leq |x| \leq 2\rho} \varepsilon^{2-N} U^2(x/\varepsilon) \, dx \\ &\leq C \varepsilon^2 \int_{\rho/\varepsilon \leq |x| \leq 2\rho/\varepsilon} U^2(x) \, dx \\ &\leq C \varepsilon^2 \int_{\rho/\varepsilon \leq |x| \leq 2\rho/\varepsilon} |x|^{4-2N} \, dx = O(\varepsilon^{N-2}); \\ \int_{\Omega} |\nabla U_{\varepsilon}|^2 |\psi|^2 \, dx &\leq \int_{\mathbb{R}^N} |\nabla U_{\varepsilon}|^2 = \int_{\mathbb{R}^N} |\nabla U|^2; \\ \left| \int_{\Omega} \psi U_{\varepsilon} \nabla \psi \nabla U_{\varepsilon} \right| &\leq C \int_{\rho \leq |x| \leq 2\rho} |\nabla U_{\varepsilon}| |U_{\varepsilon}| \, dx \\ &\leq C \int_{\rho \leq |x| \leq 2\rho} \varepsilon^{1-N} |\nabla_x U(x/\varepsilon)| |U(x/\varepsilon)| \, dx \\ &= C \varepsilon \int_{\rho/\varepsilon \leq |x| \leq 2\rho/\varepsilon} |\nabla U(x)| |U(x)| \, dx \end{split}$$

$$\leq C\varepsilon \int_{
ho/\varepsilon \leq |x| \leq 2\rho/\varepsilon} |x|^{3-2N} dx = O(\varepsilon^{N-2}).$$

Therefore,

$$\int_{\Omega} |\nabla u_{\varepsilon}|^{2} dx = \int_{\mathbb{R}^{N}} |\nabla U_{\varepsilon}|^{2} |\psi|^{2} + \int_{\Omega} |\nabla \psi|^{2} |U_{\varepsilon}|^{2} + 2 \int_{\Omega} \psi U_{\varepsilon} \nabla \psi \nabla U_{\varepsilon}
\leq \int_{\mathbb{R}^{N}} |\nabla U|^{2} dx + O(\varepsilon^{N-2}),$$

that is, (3.37) holds. Note that

$$\int_{\mathbb{R}^N} (1 - \psi^{2^*}) |U_{\varepsilon}|^{2^*} dx \le \int_{|x| \ge \rho} \varepsilon^{-N} |U(x/\varepsilon)|^{2^*} dx = \int_{|x| \ge \rho/\varepsilon} |U(x)|^{2^*} dx$$
$$\le C \int_{|x| \ge \rho/\varepsilon} |x|^{-2N} dx = O(\varepsilon^N),$$

then

$$\int_{\Omega} |u_{\varepsilon}|^{2^{*}} dx = \int_{\mathbb{R}^{N}} |U_{\varepsilon}|^{2^{*}} dx - \int_{\mathbb{R}^{N}} (1 - \psi^{2^{*}}) |U_{\varepsilon}|^{2^{*}} dx$$
$$\geq \int_{\mathbb{R}^{N}} |U|^{2^{*}} dx + O(\varepsilon^{N}),$$

that is, (3.38) holds. Similarly, (3.39) holds. Note that

$$\int_{\Omega} |u_{\varepsilon}|^{2} dx \ge \int_{|x| \le \rho} \varepsilon^{2-N} |U(x/\varepsilon)|^{2} dx
= \varepsilon^{2} \int_{\mathbb{R}^{N}} U^{2} dx - \varepsilon^{2} \int_{|x| \ge \rho/\varepsilon} U^{2}(x) dx
\ge C\varepsilon^{2} - C\varepsilon^{2} \int_{|x| \ge \rho/\varepsilon} |x|^{4-2N} dx = C\varepsilon^{2} + O(\varepsilon^{N-2}),$$

that is, (3.40) holds. Similarly, we have

$$\int_{\Omega} |\nabla v_{\varepsilon}|^2 \le \int_{\mathbb{R}^N} |\nabla V|^2 + O(\varepsilon^{N-2}), \tag{3.41}$$

$$\int_{\Omega} |v_{\varepsilon}|^{2^*} \ge \int_{\mathbb{R}^N} |V|^{2^*} + O(\varepsilon^N), \tag{3.42}$$

$$\int_{\Omega} |v_{\varepsilon}|^2 \ge C\varepsilon^2 + O(\varepsilon^{N-2}). \tag{3.43}$$

Recall that I(U, V) = A, we have

$$NA = \int_{\mathbb{R}^N} |\nabla U|^2 + |\nabla V|^2 = \int_{\mathbb{R}^N} \mu_1 U^{2^*} + 2\beta U^{\frac{2^*}{2}} V^{\frac{2^*}{2}} + \mu_2 V^{2^*}.$$

Combining this with (3.37)-(3.43) and recalling that $\lambda_1, \lambda_2 < 0, 2p = 2^*, N \ge 5$, we have for any t > 0 that

$$E(tu_{\varepsilon}, tv_{\varepsilon}) = \frac{1}{2}t^{2} \int_{\Omega} (|\nabla u_{\varepsilon}|^{2} + \lambda_{1}u_{\varepsilon}^{2} + |\nabla v_{\varepsilon}|^{2} + \lambda_{2}v_{\varepsilon}^{2})$$

$$- \frac{1}{2p}t^{2p} \int_{\Omega} (\mu_{1}u_{\varepsilon}^{2p} + 2\beta u_{\varepsilon}^{p}v_{\varepsilon}^{p} + \mu_{2}v_{\varepsilon}^{2p})$$

$$\leq \frac{1}{2} \left(\int_{\mathbb{R}^{N}} (|\nabla U|^{2} + |\nabla V|^{2}) - C\varepsilon^{2} + O(\varepsilon^{N-2}) \right) t^{2}$$

$$- \frac{1}{2^{*}} \left(\int_{\mathbb{R}^{N}} \left(\mu_{1}U^{2^{*}} + 2\beta U^{\frac{2^{*}}{2}}V^{\frac{2^{*}}{2}} + \mu_{2}V^{2^{*}} \right) + O(\varepsilon^{N}) \right) t^{2^{*}}$$

$$= \frac{1}{2} \left(NA - C\varepsilon^{2} + O(\varepsilon^{N-2}) \right) t^{2} - \frac{1}{2^{*}} \left(NA + O(\varepsilon^{N}) \right) t^{2^{*}}$$

$$\leq \frac{1}{N} \left(NA - C\varepsilon^{2} + O(\varepsilon^{N-2}) \right) \left(\frac{NA - C\varepsilon^{2} + O(\varepsilon^{N-2})}{NA + O(\varepsilon^{N})} \right)^{\frac{N-2}{2}}$$

$$< A \quad \text{for } \varepsilon > 0 \text{ small enough.}$$

$$(3.44)$$

Hence, for $\varepsilon > 0$ small enough, there holds

$$\mathcal{B} \le \max_{t>0} E(tu_{\varepsilon}, tv_{\varepsilon}) < A. \tag{3.45}$$

Step 2. we shall prove that $\mathcal{B} < B_{\mu_1}$. This proof is similar to Lemma 2.7. Recall (3.1) and (3.33), we define $t(s) := t_{u_{\mu_1}, su_{\mu_2}}$, that is,

$$t(s)^{2p-2} = \frac{NB_{\mu_1} + s^2 NB_{\mu_2}}{NB_{\mu_1} + |s|^{2p} NB_{\mu_2} + |s|^p \int_{\Omega} 2\beta |u_{\mu_1}|^p |u_{\mu_2}|^p}.$$

Note that t(0) = 1. Recall that 1 , by direct computations we get that

$$\lim_{s \to 0} \frac{t'(s)}{|s|^{p-2}s} = -\frac{p \int_{\Omega} 2\beta |u_{\mu_1}|^p |u_{\mu_2}|^p}{(2p-2)NB_{\mu_1}},$$

that is,

$$t'(s) = -\frac{p \int_{\Omega} 2\beta |u_{\mu_1}|^p |u_{\mu_2}|^p}{(2p-2)NB_{\mu_1}} |s|^{p-2} s (1+o(1)), \text{ as } s \to 0,$$

and so

$$t(s) = 1 - \frac{\int_{\Omega} 2\beta |u_{\mu_1}|^p |u_{\mu_2}|^p}{(2p-2)NB_{\mu_1}} |s|^p (1+o(1)), \text{ as } s \to 0.$$

This implies that

$$t(s)^{2p} = 1 - \frac{2p \int_{\Omega} 2\beta |u_{\mu_1}|^p |u_{\mu_2}|^p}{(2p-2)NB_{\mu_1}} |s|^p (1+o(1)), \text{ as } s \to 0.$$

Therefore, we deduce from (3.32) and $\frac{2p}{2p-2} = N/2$ that

$$\mathcal{B} \leq E(t(s)u_{\mu_{1}}, t(s)su_{\mu_{2}})$$

$$= \frac{t(s)^{2p}}{N} \left(NB_{\mu_{1}} + |s|^{2p}NB_{\mu_{2}} + |s|^{p} \int_{\Omega} 2\beta |u_{\mu_{1}}|^{p} |u_{\mu_{2}}|^{p} \right)$$

$$= B_{\mu_{1}} - \left(\frac{1}{2} - \frac{1}{N} \right) |s|^{p} \int_{\Omega} 2\beta |u_{\mu_{1}}|^{p} |u_{\mu_{2}}|^{p} + o(|s|^{p})$$

$$< B_{\mu_{1}} \quad \text{as } |s| > 0 \text{ small enough,}$$

that is, $\mathcal{B} < B_{\mu_1}$. By a similar argument, we can prove that $\mathcal{B} < B_{\mu_2}$. This completes the proof.

Proof of Theorem 1.3 for the case $\beta > 0$. Assume that $\beta > 0$. Since the functional E has a mountain pass structure, by the mountain pass theorem (cf. [5, 37]) there exists $\{(u_n, v_n)\} \subset H$ such that

$$\lim_{n \to +\infty} E(u_n, v_n) = \mathcal{B}, \quad \lim_{n \to +\infty} E'(u_n, v_n) = 0.$$

It is standard to see that $\{(u_n, v_n)\}$ is bounded in H, and so we may assume that $(u_n, v_n) \rightharpoonup (u, v)$ weakly in H. Set $\omega_n = u_n - u$ and $\sigma_n = v_n - v$ and use the same symbols as in the proof of Theorem 1.3 for the case $\beta < 0$, we see that E'(u, v) = 0 and (3.23)-(3.25) also hold. Moreover,

$$0 \le E(u, v) \le E(u, v) + \frac{1}{N}(b_1 + b_2) = \lim_{n \to +\infty} E(u_n, v_n) = \mathcal{B}.$$
 (3.46)

Case 1. $u \equiv 0, v \equiv 0$.

By (3.46), we have $b_1 + b_2 > 0$. Then we may assume that $(\omega_n, \sigma_n) \neq (0, 0)$ for n large. Recall \mathcal{N}' in (2.26), by (3.23)-(3.24), it is easy to check that there exists $t_n > 0$ such that $(t_n \omega_n, t_n \sigma_n) \in \mathcal{N}'$ and $t_n \to 1$ as $n \to \infty$. Then by (2.38) and (3.46) we have

$$\mathcal{B} = \frac{1}{N}(b_1 + b_2) = \lim_{n \to +\infty} I(\omega_n, \sigma_n) = \lim_{n \to +\infty} I(t_n \omega_n, t_n \sigma_n) \ge A' = A,$$

a contradiction with Lemma 3.4. Therefore, Case 1 is impossible.

Case 2. $u \not\equiv 0, v \equiv 0 \text{ or } u \equiv 0, v \not\equiv 0.$

Without loss of generality, we may assume that $u \neq 0, v \equiv 0$. Then u is a nontrivial solution of $-\Delta u + \lambda_1 u = \mu_1 |u|^{2^*-2} u$, and so $\mathcal{B} \geq E(u,0) \geq B_{\mu_1}$, a contradiction with Lemma 3.4. Therefore, Case 2 is also impossible.

Since Cases 1 and 2 are both impossible, we have that $u \not\equiv 0, v \not\equiv 0$. Since E'(u,v)=0, we have $(u,v)\in \mathcal{M}$. By $\mathcal{B}\leq B$ and (3.46) we have $E(u,v)=\mathcal{B}=B$. This means $(|u|,|v|)\in \mathcal{M}\subset \mathcal{M}'$ and $E(|u|,|v|)=\mathcal{B}=B$. By (3.34) and (3.35), there exists a Lagrange multiplier $\gamma\in\mathbb{R}$ such that

$$E'(|u|,|v|) - \gamma G'(|u|,|v|) = 0.$$

Since E'(|u|, |v|)(|u|, |v|) = G(|u|, |v|) = 0 and

$$G'(|u|,|v|)(|u|,|v|) = -(2p-2) \int_{\Omega} (\mu_1|u|^{2p} + 2\beta|u|^p|v|^p + \mu_2|v|^{2p}) \neq 0,$$

we get that $\gamma = 0$ and so E'(|u|, |v|) = 0. This means that (|u|, |v|) is a least energy solution of (1.1). By the maximum principle, we see that |u|, |v| > 0 in Ω . Therefore, (|u|, |v|) is a positive least energy solution of (1.1).

4 Proof of Theorems 1.2 and 1.7

In this section, we assume that $-\lambda_1(\Omega) < \lambda_1 = \lambda_2 = \lambda < 0$ and $\beta \ge (p-1) \max\{\mu_1, \mu_2\}$. Define $g: [(p-1) \max\{\mu_1, \mu_2\}, +\infty)$ by

$$g(\beta) := (p-1)\mu_1 \mu_2 \beta^{2/p-2} + \beta^{2/p}. \tag{4.1}$$

Then

$$g'(\beta) = \frac{2}{p} \beta^{2/p-3} \left(\beta^2 - (p-1)^2 \mu_1 \mu_2 \right) > 0, \quad \forall \ \beta > (p-1) \max\{\mu_1, \mu_2\}.$$

By direct computations, we have

$$g((p-1)\max\{\mu_1,\mu_2\}) \le p(p-1)^{\frac{2}{p}-1}\max\{\mu_1^{2/p}, \mu_2^{2/p}\}.$$

Therefore, there exists a unique $\beta_0 \ge (p-1) \max\{\mu_1, \mu_2\}$ such that

$$g(\beta_0) = p(p-1)^{\frac{2}{p}-1} \max\left\{\mu_1^{2/p}, \ \mu_2^{2/p}\right\}, \text{ and } (4.2)$$

$$g(\beta) > p(p-1)^{\frac{2}{p}-1} \max \left\{ \mu_1^{2/p}, \ \mu_2^{2/p} \right\}, \quad \forall \ \beta > \beta_0.$$
 (4.3)

Moreover,

$$\beta_0 = (p-1) \max\{\mu_1, \mu_2\}, \quad \text{if } \mu_1 = \mu_2.$$
 (4.4)

Lemma 4.1. Assume that $\beta > \beta_0$, where β_0 is defined in (4.2). Let (k_0, l_0) be in Lemma 2.1. Then $p\mu_1k_0^{p-1} < 1$ and $p\mu_2l_0^{p-1} < 1$.

Proof. Let $k_1 = (p\mu_1)^{\frac{1}{p-1}}$, then by (2.3) we have

$$l_1 := h_1(k_1) = \left[\frac{p-1}{p\beta(p\mu_1)^{\frac{2-p}{2(p-1)}}}\right]^{2/p}.$$

By (4.3) and direct computations, we get that

$$\alpha_2(k_1, l_1) = \mu_2 l_1^{p-1} + \beta k_1^{p/2} l_1^{p/2-1} - 1$$

$$= \frac{1}{l_1} \left[\mu_2 l_1^p + k_1 (1 - \mu_1 k_1^{p-1}) \right] - 1 = \frac{1}{l_1} \left[\mu_2 l_1^p + \frac{p-1}{p} k_1 \right] - 1$$

$$= \left[\frac{p\beta(p\mu_1)^{\frac{2-p}{2(p-1)}}}{p-1} \right]^{\frac{2}{p}} \left\{ \mu_2 \left[\frac{p-1}{p\beta(p\mu_1)^{\frac{2-p}{2(p-1)}}} \right]^2 + \frac{p-1}{p} (p\mu_1)^{-\frac{1}{p-1}} \right\} - 1$$
$$= (p-1)^{1-2/p} p^{-1} \mu_1^{-2/p} g(\beta) - 1 > 0.$$

Combining this with Lemma 2.3 we have $k_1 > k_0$, that is, $p\mu_1 k_0^{p-1} < 1$. Similarly, let $l_2 = (p\mu_2)^{\frac{1}{p-1}}$, then

$$\alpha_1(h_2(l_2), l_2) = (p-1)^{1-2/p} p^{-1} \mu_2^{-2/p} g(\beta) - 1 > 0.$$

By Lemma 2.3 again, we have $l_2 > l_0$, and so $p\mu_2 l_0^{p-1} < 1$.

Lemma 4.2. Assume that $\beta > \beta_0$, where β_0 is defined in (4.2). Recall α_1, α_2 defined in (2.1)-(2.2), and (k_0, l_0) obtained in Lemma 2.1. Then

$$F(k_0, l_0) := \det \begin{pmatrix} \partial_k \alpha_1(k_0, l_0) & \partial_l \alpha_1(k_0, l_0) \\ \partial_k \alpha_2(k_0, l_0) & \partial_l \alpha_2(k_0, l_0) \end{pmatrix} < 0.$$

Proof. By $\alpha_1(k_0, l_0) = \alpha_2(k_0, l_0) = 0$ we have

$$\beta k_0^{p/2-2} l_0^{p/2} = k_0^{-1} - \mu_1 k_0^{p-2}, \quad \beta l_0^{p/2-2} k_0^{p/2} = l_0^{-1} - \mu_2 l_0^{p-2}.$$

Then

$$\begin{split} \partial_k \alpha_1(k_0, l_0) &= (p-1)\mu_1 k_0^{p-2} + (p/2-1)\beta k_0^{p/2-2} l_0^{p/2} \\ &= \frac{p}{2}\mu_1 k_0^{p-2} - (1-p/2)k_0^{-1}; \\ \partial_l \alpha_2(k_0, l_0) &= (p-1)\mu_2 l_0^{p-2} + (p/2-1)\beta l_0^{p/2-2} k_0^{p/2} \\ &= \frac{p}{2}\mu_2 l_0^{p-2} - (1-p/2)l_0^{-1}; \\ \partial_l \alpha_1(k_0, l_0) &= \partial_k \alpha_2(k_0, l_0) = \frac{p}{2}\beta k_0^{p/2-1} l_0^{p/2-1} \\ &= \frac{p}{2}\sqrt{(k_0^{-1} - \mu_1 k_0^{p-2})(l_0^{-1} - \mu_2 l_0^{p-2})}. \end{split}$$

Therefore,

$$F(k_0, l_0) = \left[\frac{p}{2}\mu_1 k_0^{p-2} - (1 - p/2)k_0^{-1}\right] \left[\frac{p}{2}\mu_2 l_0^{p-2} - (1 - p/2)l_0^{-1}\right]$$
$$-\frac{p^2}{4} \left(k_0^{-1} - \mu_1 k_0^{p-2}\right) \left(l_0^{-1} - \mu_2 l_0^{p-2}\right)$$
$$= \frac{p}{2} (p - 1)k_0^{-1} l_0^{-1} \left(\mu_1 k_0^{p-1} + \mu_2 l_0^{p-1} - \frac{2}{p}\right) < 0$$

from Lemma 4.1.

Lemma 4.3. Fix any $\mu_1, \mu_2 > 0$ and $\beta > \beta_0$. Let (u_0, v_0) be a least energy solution of (1.1) with (μ_1, μ_2, β) which exists by Theorem 1.3. Recall $(\sqrt{k_0}\omega, \sqrt{l_0}\omega)$ in Theorem 1.1. Then

$$\int_{\Omega} |u_0|^{2p} dx = k_0^p \int_{\Omega} \omega^{2p} dx. \tag{4.5}$$

Proof. Fix any $\mu_1, \mu_2 > 0$ and $\beta > \beta_0$. We remark from (4.1)-(4.2) that $\beta_0(\mu_1, \mu_2) := \beta_0$ is completely determined by μ_1, μ_2 . Hence there exists $0 < \varepsilon < \mu_1$ such that for any $\mu \in (\mu_1 - \varepsilon, \mu_1 + \varepsilon)$, one has $\beta > \beta_0(\mu, \mu_2)$. Then by Lemmas 2.1, 4.2 and the implicit function theorem, when μ_1 is replaced by μ , functions $k_0(\mu)$ and $l_0(\mu)$ are well defined and class C^1 for $\mu \in (\mu_1 - \varepsilon_1, \mu_1 + \varepsilon_1)$ for some $0 < \varepsilon_1 \le \varepsilon$. Recall the definition of E, \mathcal{M} and E, E, \mathcal{M} and

$$B(\mu) = \inf_{H \ni (u,v) \neq (0,0)} \max_{t>0} E_{\mu}(tu, tv).$$

Denote

$$C = \int_{\Omega} (|\nabla u_0|^2 + \lambda_1 u_0^2 + |\nabla v_0|^2 + \lambda_2 v_0^2),$$

$$D = \int_{\Omega} (2\beta |u_0|^p |v_0|^p + \mu_2 |v_0|^{2p}), \quad G = \int_{\Omega} |u_0|^{2p} dx.$$

There exists $t(\mu) > 0$ such that

$$\max_{t>0} E_{\mu}(tu_0, tv_0) = E_{\mu}\Big(t(\mu)u_0, t(\mu)v_0\Big),$$

where $t(\mu) > 0$ satisfies $f(\mu, t(\mu)) = 0$, and

$$f(\mu, t) := t^{2p-2}(\mu G + D) - C.$$

Note that $f(\mu_1, 1) = 0$, $\frac{\partial}{\partial t} f(\mu_1, 1) = (2p - 2)(\mu_1 G + D) > 0$, and $f(\mu, t(\mu)) \equiv 0$. By the implicit function theorem, there exists $0 < \varepsilon_2 \le \varepsilon_1$, such that $t(\mu) \in C^{\infty}((\mu_1 - \varepsilon_2, \mu_1 + \varepsilon_2), \mathbb{R})$. By $f(\mu, t(\mu)) \equiv 0$ we see that

$$t'(\mu_1) = -\frac{G}{(2p-2)(\mu_1 G + D)}.$$

By Taylor expansion one has $t(\mu) = 1 + t'(\mu_1)(\mu - \mu_1) + O((\mu - \mu_1)^2)$, and so

$$t^{2}(\mu) = 1 + 2t'(\mu_{1})(\mu - \mu_{1}) + O((\mu - \mu_{1})^{2}).$$

Note that $C = \mu_1 G + D = NB(\mu_1)$. Then by (3.32) that

$$B(\mu) \le E_{\mu}(t(\mu)u_0, t(\mu)v_0) = \frac{1}{N}t^2(\mu)C = t^2(\mu)B(\mu_1)$$

$$= B(\mu_1) - \frac{2GB(\mu_1)}{(2p-2)(\mu_1G+D)}(\mu-\mu_1) + O((\mu-\mu_1)^2)$$

= $B(\mu_1) - \frac{G}{2p}(\mu-\mu_1) + O((\mu-\mu_1)^2),$

It follows that $\frac{B(\mu) - B(\mu_1)}{\mu - \mu_1} \ge -\frac{G}{2p} + O((\mu - \mu_1))$, as $\mu \nearrow \mu_1$, and so $B'(\mu_1) \ge -\frac{G}{2p}$. Similarly, we have

$$\frac{B(\mu) - B(\mu_1)}{\mu - \mu_1} \le -\frac{G}{2p} + O((\mu - \mu_1)), \text{ as } \mu \searrow \mu_1,$$

that is, $B'(\mu_1) \leq -\frac{G}{2p}$. Hence, $B'(\mu_1) = -\frac{G}{2p} = -\frac{1}{2p} \int_{\Omega} |u_0|^{2p} dx$. By Theorem 1.1, $(\sqrt{k_0}\omega, \sqrt{l_0}\omega)$ is also a positive least energy solution of (1.1). Therefore, $B'(\mu_1) = -\frac{k_0^p}{2p} \int_{\Omega} \omega^{2p} dx$, that is, (4.5) holds.

Proof of Theorem 1.2. Let (u, v) be any a positive least energy solution of (1.1). By Lemma 4.3, we have

$$\int_{\Omega} |u|^{2p} dx = k_0^p \int_{\Omega} \omega^{2p} dx.$$

By a similar proof of Lemma 4.3, that is, by computing $B'(\mu_2)$ and $B'(\beta)$ respectively, we can show that

$$\int_{\Omega} |v|^{2p} dx = l_0^p \int_{\Omega} \omega^{2p} dx, \quad \text{and} \quad \int_{\Omega} |u|^p |v|^p dx = k_0^{p/2} l_0^{p/2} \int_{\Omega} \omega^{2p} dx.$$

Therefore.

$$\int_{\Omega} |u|^p |v|^p \, dx = l_0^{p/2} k_0^{-p/2} \int_{\Omega} |u|^{2p} \, dx, \quad \int_{\Omega} |u|^p |v|^p \, dx = l_0^{-p/2} k_0^{p/2} \int_{\Omega} |v|^{2p} \, dx.$$

$$(4.6)$$

Define $(\tilde{u}, \tilde{v}) := (\frac{1}{\sqrt{k_0}} u, \frac{1}{\sqrt{l_0}} v)$. By $\alpha_1(k_0, l_0) = \alpha_2(k_0, l_0) = 0$ and (4.6) we get

$$\int_{\Omega} |\nabla \tilde{u}|^2 + \lambda \tilde{u}^2 dx = \int_{\Omega} |\tilde{u}|^{2p} dx, \quad \int_{\Omega} |\nabla \tilde{v}|^2 + \lambda \tilde{v}^2 dx = \int_{\Omega} |\tilde{v}|^{2p} dx. \tag{4.7}$$

Then by (1.8) we have

$$\frac{1}{N} \int_{\Omega} |\nabla \tilde{u}|^2 + \lambda \tilde{u}^2 \, dx \ge B_1, \quad \frac{1}{N} \int_{\Omega} |\nabla \tilde{v}|^2 + \lambda \tilde{v}^2 \, dx \ge B_1,$$

and so

$$B = (k_0 + l_0)B_1 = \frac{1}{N} \int_{\Omega} (|\nabla u|^2 + \lambda_1 u^2 + |\nabla v|^2 + \lambda_2 v^2)$$

$$= \frac{1}{N} k_0 \int_{\Omega} (|\nabla \tilde{u}|^2 + \lambda_1 \tilde{u}^2) + \frac{1}{N} l_0 \int_{\Omega} (|\nabla \tilde{v}|^2 + \lambda_1 \tilde{v}^2)$$

$$\geq (k_0 + l_0) B_1.$$

This implies that

$$\frac{1}{N} \int_{\Omega} |\nabla \tilde{u}|^2 + \lambda \tilde{u}^2 dx = B_1, \quad \frac{1}{N} \int_{\Omega} |\nabla \tilde{v}|^2 + \lambda \tilde{v}^2 dx = B_1.$$

Combining this with (4.7), we see from [9] that \tilde{u} and \tilde{v} are both postive least energy solutions of (1.6). Then we see from (u, v) satisfies (1.1) that

$$-\Delta \tilde{u} + \lambda \tilde{u} = \mu_1 k_0^{p-1} \tilde{u}^{2p-1} + \beta k_0^{p/2-1} l_0^{p/2} \tilde{u}^{p-1} \tilde{v}^p = \tilde{u}^{2p-1}$$

that is, $\tilde{u}^{p-1}\tilde{v}^p = \tilde{u}^{2p-1}$ and so $\tilde{u} = \tilde{v}$. Denote $U = \tilde{u}$, then $(u, v) = (\sqrt{k_0}U, \sqrt{l_0}U)$, where U is a positive least energy solution of (1.6).

Now we assume that Ω is a ball in \mathbb{R}^N , then the positive least energy solution of the Brezis-Nirenberg problem (1.6) is unique (cf. [6]). Therefore, the positive least energy solution of (1.1) is unique.

Proof of Theorem 1.7. The proof is the same as that of Theorem 1.2. \Box

5 Proof of Theorems 1.4 and 1.5

This section is devoted to the proof of Theorems 1.4 and 1.5. Recall the definition of E, \mathcal{M} and B, they both depend on β , and we use notations $E_{\beta}, \mathcal{M}_{\beta}, B_{\beta}$ in this section. Define $B(x_0, R) := \{x \in \mathbb{R}^N : |x - x_0| < R\}$. Consider the problem

$$\begin{cases}
-\Delta u + \lambda_2 u = \mu_2 u^{2^* - 1} & \text{in } B(0, R), \\
u > 0 & \text{in } B(0, R), \quad u = 0 & \text{on } \partial B(0, R),
\end{cases}$$
(5.1)

and the corresponding functional is $J_R: H^1_0(B(0,R)) \to \mathbb{R}$ given by

$$J_R(u) = \frac{1}{2} \int_{B(0,R)} (|\nabla u|^2 + \lambda_2 u^2) \, dx - \frac{1}{2^*} \mu_2 \int_{B(0,R)} |u|^{2^*} \, dx. \tag{5.2}$$

We need the following energy estimates from the authors' paper [15].

Theorem 5.1. (see [15]) Let $N \ge 5$. Then there exists $R_0 > 0$ and $C_1, C_2 > 0$, such that for any $0 < R < R_0$, (5.1) has a least energy solution U_R and

$$\frac{1}{N}\mu_2^{-\frac{N-2}{2}}S^{N/2} - C_1R^{\frac{2N-4}{N-4}} \le J_R(U_R) \le \frac{1}{N}\mu_2^{-\frac{N-2}{2}}S^{N/2} - C_2R^{\frac{2N-4}{N-4}}.$$
 (5.3)

With the help of Theorem 5.1, we have the following lemma, which improves Lemma 3.1 in case $N \ge 6$.

Lemma 5.1. Let $N \geq 6$. Then

$$\sup_{\beta < 0} B_{\beta} < \min \left\{ B_{\mu_1} + \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{N/2}, \ B_{\mu_2} + \frac{1}{N} \mu_1^{-\frac{N-2}{2}} S^{N/2} \right\}.$$

Proof. Let $N \geq 6$. For any R > 0 small, we take $x_R \in \Omega$ with $\operatorname{dist}(x_R, \partial \Omega) = 3R$. Then

$$|u_{\mu_1}(x)| \le CR, \quad x \in B(x_R, 3R).$$
 (5.4)

Let $\psi \in C_0^{\infty}(B(0,2))$ with $0 \le \psi \le 1$ and $\psi \equiv 1$ in B(0,1). Define $\varphi_R(x) := 1 - \psi(\frac{x - x_R}{B})$, then

$$\varphi_R(x) := \begin{cases} 0 & \text{if} \quad x \in B(x_R, R), \\ 1 & \text{if} \quad x \in \mathbb{R}^N \backslash B(x_R, 2R), \end{cases} \quad |\nabla \varphi_R(x)| \le C/R. \tag{5.5}$$

Define $u_R := \varphi_R u_{\mu_1}$, then by (5.4) and (5.5), it is easy to prove that

$$\int_{\Omega} |\nabla u_R|^2 dx \le \int_{\Omega} |\nabla u_{\mu_1}|^2 dx + CR^N;$$

$$\int_{\Omega} |u_R|^2 dx \ge \int_{\Omega} |u_{\mu_1}|^2 dx - CR^{N+2};$$

$$\int_{\Omega} |u_R|^{2^*} dx \ge \int_{\Omega} |u_{\mu_1}|^{2^*} dx - CR^{N+2^*}.$$

Therefore, there exists $t_R > 0$ independent of $\beta < 0$ such that

$$\max_{t>0} E_{\beta}(tu_{R}, 0) = E_{\beta}(t_{R}u_{R}, 0) = \frac{1}{N} \left(\frac{\int_{\Omega} (|\nabla u_{R}|^{2} + \lambda_{1}u_{R}^{2})}{(\mu_{1} \int_{\Omega} |u_{R}|^{2^{*}})^{2/2^{*}}} \right)^{N/2} \\
\leq \frac{1}{N} \left(\frac{\int_{\Omega} (|\nabla u_{\mu_{1}}|^{2} + \lambda_{1}u_{\mu_{1}}^{2}) + CR^{N} + CR^{N+2}}{(\int_{\Omega} \mu_{1} |u_{\mu_{1}}|^{2^{*}} - CR^{N+2^{*}})^{2/2^{*}}} \right)^{N/2} \\
= \frac{1}{N} \left(\frac{NB_{\mu_{1}} + CR^{N} + CR^{N+2}}{(NB_{\mu_{1}} - CR^{N+2^{*}})^{2/2^{*}}} \right)^{N/2} \\
\leq B_{\mu_{1}} + CR^{N} \quad \text{for } R > 0 \text{ small enough.}$$

Recall U_R in Theorem 5.1, we have $U_R(\cdot - x_R) \cdot u_R \equiv 0$, and so $(t_R u_R, U_R(\cdot - x_R)) \in \mathcal{M}_\beta$ for all $\beta < 0$. Since $N \geq 6$, one has $N > \frac{2N-4}{N-4}$. Then we see from Theorem 5.1 that

$$\begin{split} \sup_{\beta < 0} B_{\beta} & \leq E_{\beta}(t_R u_R, U_R(\cdot - x_R)) = E_{\beta}(t_R u_R, 0) + E_{\beta}(0, U_R(\cdot - x_R)) \\ & = E_{\beta}(t_R u_R, 0) + J_R(U_R) \\ & \leq B_{\mu_1} + CR^N + \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{N/2} - C_2 R^{\frac{2N-4}{N-4}} \\ & < B_{\mu_1} + \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{N/2} \quad \text{for } R > 0 \text{ small enough.} \end{split}$$

By a similar argument, we also have $\sup_{\beta < 0} B_{\beta} < B_{\mu_2} + \frac{1}{N} \mu_1^{-\frac{N-2}{2}} S^{N/2}$.

Proof of Theorem 1.4. This proof is similar to the proof of [16, Theorem 1.4] in case N=4. The novelty here is that, with the help of Lemma 5.1, we can exclude conclusions (1)-(2) in case $N\geq 6$. Let $\beta_n<0$, $n\in\mathbb{N}$ satisfy $\beta_n\to-\infty$ as $n\to\infty$, and (u_n,v_n) be the positive least energy solutions of (1.1) with $\beta=\beta_n$. By Lemma 3.1, $E_{\beta_n}(u_n,v_n)\leq A$ and so (u_n,v_n) is uniformly bounded in H by (1.5). Passing to a subsequence, we may assume that

$$u_n \rightharpoonup u_\infty$$
, $v_n \rightharpoonup v_\infty$ weakly in $H_0^1(\Omega)$,

Then, by following the proof of [16, Theorem 1.4] in case N=4, we can prove that $\int_{\Omega} \beta_n u_n^p v_n^p dx \to 0$ as $n \to \infty$, and passing to a subsequence, one of the following conclusions holds.

(1) $u_n \to u_\infty$ strongly in $H_0^1(\Omega)$ and $v_n \to 0$ weakly in $H_0^1(\Omega)$ (so $v_n \to 0$ for almost every $x \in \Omega$), where u_∞ is a positive least energy solution of

$$-\Delta u + \lambda_1 u = \mu_1 |u|^{2^* - 2} u, \ u \in H_0^1(\Omega).$$

Moreover,

$$\lim_{n \to \infty} B_{\beta_n} = B_{\mu_1} + \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{N/2}.$$
 (5.6)

(2) $v_n \to v_\infty$ strongly in $H_0^1(\Omega)$ and $u_n \to 0$ weakly in $H_0^1(\Omega)$ (so $u_n \to 0$ for almost every $x \in \Omega$), where v_∞ is a positive least energy solution of

$$-\Delta v + \lambda_2 v = \mu_2 |v|^{2^* - 2} v, \ v \in H_0^1(\Omega).$$

Moreover,

$$\lim_{n \to \infty} B_{\beta_n} = B_{\mu_2} + \frac{1}{N} \mu_1^{-\frac{N-2}{2}} S^{N/2}.$$
 (5.7)

(3) $(u_n, v_n) \to (u_\infty, v_\infty)$ strongly in $H_0^1(\Omega) \times H_0^1(\Omega)$ and $u_\infty \cdot v_\infty = 0$ for almost $x \in \Omega$, where $u_\infty \not\equiv 0$, $v_\infty \not\equiv 0$ satisfy

$$\int_{\Omega} (|\nabla u_{\infty}|^2 + \lambda_1 u_{\infty}^2) = \int_{\Omega} \mu_1 u_{\infty}^{2p}, \tag{5.8}$$

$$\int_{\Omega} (|\nabla v_{\infty}|^2 + \lambda_2 v_{\infty}^2) = \int_{\Omega} \mu_2 v_{\infty}^{2p}, \tag{5.9}$$

$$\lim_{n \to \infty} B_{\beta_n} = E(u_{\infty}, v_{\infty}). \tag{5.10}$$

Moreover, if u_{∞} and v_{∞} are both continuous (we will prove this later), then $u_{\infty} \cdot v_{\infty} \equiv 0$, $u_{\infty} \in C(\overline{\Omega})$ is a positive least energy solution of

$$-\Delta u + \lambda_1 u = \mu_1 |u|^{2^* - 2} u, \ u \in H_0^1(\{u_\infty > 0\}),$$

and $v_{\infty} \in C(\overline{\Omega})$ is a positive least energy solution of

$$-\Delta v + \lambda_2 v = \mu_2 |v|^{2^* - 2} v, \ v \in H_0^1(\{v_\infty > 0\}).$$

Furthermore, both $\{v_{\infty} > 0\}$ and $\{u_{\infty} > 0\}$ are connected domains, and $\{v_{\infty} > 0\} = \Omega \setminus \overline{\{u_{\infty} > 0\}}$.

Note that (5.6)-(5.7) imply that one of (1) and (2) in Theorem 1.4 does not hold in some cases. For example, if we assume that $-\lambda_1(\Omega) < \lambda_1 < \lambda_2 < 0$ and $\mu_1 = \mu_2$ in Theorem 1.4, then $B_{\mu_1} + \frac{1}{N}\mu_2^{-\frac{N-2}{2}}S^{N/2} < B_{\mu_2} + \frac{1}{N}\mu_1^{-\frac{N-2}{2}}S^{N/2}$, and so (2) in Theorem 1.4 does not hold, since (5.7) contradicts with Lemma 3.1.

In particular, Lemma 5.1 implies that neither (1) nor (2) hold in case $N \ge 6$. That is, only (3) holds if $N \ge 6$. Therefore, the proof is complete by combining Lemma 5.2 below.

From the previous proof, it suffices to prove that u_{∞} and v_{∞} are continuous and $u_{\infty} - v_{\infty}$ is a least energy sign-changing solution of (1.11). As pointed out in Remark 1.4, the following proof is completely different from that in [16] for the case N=4.

Lemma 5.2. Let (u_{∞}, v_{∞}) be in conclusion (3). Then $u_{\infty} - v_{\infty}$ is a least energy sign-changing solution of (1.11), and u_{∞}, v_{∞} are both continuous.

Proof. Consider the problem (1.11). Its related functional is

$$P(u) = \frac{1}{2} \int_{\Omega} (|\nabla u|^2 + \lambda_1 (u^+)^2 + \lambda_2 (u^-)^2) - \frac{1}{2^*} \int_{\Omega} (\mu_1 (u^+)^{2^*} + \mu_2 (u^-)^{2^*}).$$

It is standard to prove that $P \in C^1$ and its critical points are solutions of (1.11). Define

$$J_{i}(u) := \int_{\Omega} (|\nabla u|^{2} + \lambda_{i}u^{2} - \mu_{i}|u|^{2^{*}}), \quad i = 1, 2,$$

$$S := \left\{ u \in H_{0}^{1}(\Omega) : u^{\pm} \not\equiv 0, J_{1}(u^{+}) = 0, J_{2}(u^{-}) = 0 \right\},$$

$$m := \inf_{u \in S} P(u).$$

Then any sign-changing solutions of (1.11) belong to S. By (5.8)-(5.9), we have $u_{\infty} - v_{\infty} \in S$ and so $m \leq P(u_{\infty} - v_{\infty}) = E(u_{\infty}, v_{\infty})$. For any $u \in S$, we have $(u^+, u^-) \in \mathcal{M}_{\beta}$ for all β . Then by (5.10) we see that

$$P(u_{\infty} - v_{\infty}) = E(u_{\infty}, v_{\infty}) = \lim_{n \to \infty} B_{\beta_n} \le E(u^+, u^-) = P(u), \quad \forall u \in \mathcal{S},$$

and so $P(u_{\infty} - v_{\infty}) \leq m$. Combining these with Lemma 5.1, we obtain

$$P(u_{\infty} - v_{\infty}) = m < \min \left\{ B_{\mu_1} + \frac{1}{N} \mu_2^{-\frac{N-2}{2}} S^{\frac{N}{2}}, \ B_{\mu_2} + \frac{1}{N} \mu_1^{-\frac{N-2}{2}} S^{\frac{N}{2}} \right\}.$$
 (5.11)

Step 1. We show that $P'(u_{\infty} - v_{\infty}) = 0$, and so $u_{\infty} - v_{\infty}$ is a least energy sign-changing solution of (1.11).

Thanks to (5.11), the following argument is standard (see [26, 31] for example), and we give the details here for completeness.

Assume that $u_{\infty}-v_{\infty}$ is not a critical point of P, then there exists $\phi \in C_0^{\infty}(\Omega)$ such that $P'(u_{\infty}-v_{\infty})\phi \leq -1$. Then there exists $0 < \varepsilon_0 < 1/10$, such that for $|t-1| \leq \varepsilon_0$, $|s-1| \leq \varepsilon_0$, $|\sigma| \leq \varepsilon_0$, there holds

$$P'(tu_{\infty} - sv_{\infty} + \sigma\phi)\phi \le -\frac{1}{2}.$$

Consider a function $0 \le \eta \le 1$ defined for $(t,s) \in T = [\frac{1}{2}, \frac{3}{2}] \times [\frac{1}{2}, \frac{3}{2}]$, such that

$$\begin{split} & \eta(t,s) = 1, \quad \text{for } |t-1| \leq \frac{\varepsilon_0}{2}, \ |s-1| \leq \frac{\varepsilon_0}{2}, \\ & \eta(t,s) = 0, \quad \text{for } |t-1| \geq \varepsilon_0 \text{ or } |s-1| \geq \varepsilon_0. \end{split}$$

Then for $|t-1| \le \varepsilon_0, |s-1| \le \varepsilon_0$, we have

$$\begin{split} &P(tu_{\infty} - sv_{\infty} + \varepsilon_{0}\eta(t,s)\phi) \\ = &P(tu_{\infty} - sv_{\infty}) + \int_{0}^{1} P'(tu_{\infty} - sv_{\infty} + \theta\varepsilon_{0}\eta(t,s)\phi)[\varepsilon_{0}\eta(t,s)\phi]d\theta \\ \leq &P(tu_{\infty} - sv_{\infty}) - \frac{1}{2}\varepsilon_{0}\eta(t,s). \end{split}$$

Note that

$$\sup_{t,s>0} P(tu_{\infty} - sv_{\infty}) = P(u_{\infty} - v_{\infty}) = m,$$

and for $|t-1| \ge \frac{\varepsilon_0}{2}$ or $|s-1| \ge \frac{\varepsilon_0}{2}$, there exists $0 < \delta < \frac{\varepsilon_0}{2}$ such that

$$P(tu_{\infty} - sv_{\infty}) \le m - \delta.$$

We have, for $|t-1| \leq \frac{\varepsilon_0}{2}, |s-1| \leq \frac{\varepsilon_0}{2}$, that

$$P(tu_{\infty} - sv_{\infty} + \varepsilon_0 \eta(t, s)\phi) \le m - \frac{\varepsilon_0}{2};$$

for $\frac{\varepsilon_0}{2} \le |t-1| \le \varepsilon_0, |s-1| \le \varepsilon_0$ or $\frac{\varepsilon_0}{2} \le |s-1| \le \varepsilon_0, |t-1| \le \varepsilon_0$,

$$P(tu_{\infty} - sv_{\infty} + \varepsilon_0 \eta(t, s)\phi) \le P(tu_{\infty} - sv_{\infty}) \le m - \delta;$$

for $|t-1| \ge \varepsilon_0$ or $|s-1| \ge \varepsilon_0$,

$$P(tu_{\infty} - sv_{\infty} + \varepsilon_0 \eta(t, s)\phi) = P(tu_{\infty} - sv_{\infty}) < m - \delta.$$

So

$$\sup_{(t,s)\in T} P(tu_{\infty} - sv_{\infty} + \varepsilon_0 \eta(t,s)\phi) \le m - \delta.$$
 (5.12)

On the other hand, for $\varepsilon \in [0, \varepsilon_0]$, let $h_{\varepsilon} : T \to H_0^1(\Omega)$ by $h_{\varepsilon}(t, s) = tu_{\infty} - sv_{\infty} + \varepsilon \eta(t, s) \phi$, and $H_{\varepsilon} : T \to \mathbb{R}^2$ by

$$H_{\varepsilon}(t,s) = (J_1(h_{\varepsilon}(t,s)^+), J_2(h_{\varepsilon}(t,s)^-)).$$

Note that for any $(t,s) \in \partial T$, we have $\eta(t,s) = 0$ and so $h_{\varepsilon}(t,s) \equiv h_0(t,s) = tu_{\infty} - sv_{\infty}$. Moreover, $H_0(t,s) = (J_1(tu_{\infty}), J_2(sv_{\infty}))$. Then it is easy to see that

$$\deg(H_{\varepsilon_0}(t,s), T, (0,0)) = \deg(H_0(t,s), T, (0,0)) = 1,$$

that is, there exists $(t_0, s_0) \in T$ such that $h_{\varepsilon_0}(t_0, s_0) \in \mathcal{S}$, which is a contradiction with (5.12).

Step 2. We show that u_{∞} and v_{∞} are continuous.

By Step 1, $u_{\infty} - v_{\infty}$ is a nontrivial solution of (1.11). Then by a Brezis-Kato argument (see [8]), we see that $u_{\infty} - v_{\infty} \in L^{q}(\Omega), \forall q \geq 2$. In particular,

$$\mu_1 u_{\infty}^{2^*-1} - \mu_2 v_{\infty}^{2^*-1} - \lambda_1 u_{\infty} + \lambda_2 v_{\infty} \in L^q(\Omega), \ \forall q > N.$$

Then by elliptic regularity theory, $u_{\infty} - v_{\infty} \in W^{2,q}(\Omega)$ with q > N. By Sobolev embedding, we have $u_{\infty} - v_{\infty} \in C(\overline{\Omega})$. Since $u_{\infty} = (u_{\infty} - v_{\infty})^+$ and $v_{\infty} = (u_{\infty} - v_{\infty})^-$, we see that u_{∞} and v_{∞} are both continuous. This completes the proof and so completes the proof of Theorem 1.4.

Proof of Theorem 1.5. Let $N \geq 6$. Actually, by Theorem 1.4 and Lemma 5.2, we have proved that the problem (1.11) has a least energy sign-changing solution $u_{\infty}-v_{\infty}$. Obviously, Theorem 1.5 is a direct corollary by letting $\lambda_1 = \lambda_2$ and $\mu_1 = \mu_2$, and (1.13) follows directly from (5.11).

References

- [1] T. Aubin, Problemes isoperimetriques et espaces de Sobolev, *J. Differ. Geom.*, **11** (1976), 573-598.
- [2] N. Akhmediev, A. Ankiewicz, Partially coherent solitons on a finite background, *Phys. Rev. Lett.*, **82** (1999), 2661-2664.
- [3] B. Abdellaoui, V. Felli, I. Peral, Some remarks on systems of elliptic equations doubly critical in the whole \mathbb{R}^N , Calc. Var. PDE., **34** (2009), 97-137.
- [4] A. Ambrosetti, E. Colorado, Standing waves of some coupled nonlinear Schrödinger equations, *J. London. Math. Soc.*, **75** (2007), 67–82.
- [5] A. Ambrosetti, P. H. Rabinowitz, Dual variational methods in critical point theory and applications, *J. Funct. Anal.*, **14** (1973), 349–381.
- [6] Adimurthi, S. L. Yadava, An elementary proof of the uniqueness of positive radial solutions of a quasilinear dirichlet problem, *Arch. Ration. Mech. Anal.*, **127** (1994), 219-229.
- [7] T. Bartsch, N. Dancer, Z.-Q. Wang, A Liouville theorem, a priori bounds, and bifurcating branches of positive solutions for a nonlinear elliptic system, *Calc. Var. PDE.*, **37** (2010), 345-361.
- [8] H. Brezis, T. Kato, Remarks on the Schrodinger operator with singular complex potentials, *J. Math. Pures Appl.*, **58** (1979), 137-151.
- [9] H. Brezis, L. Nirenberg, Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, Comm. Pure Appl. Math., 36 (1983), 437-477.
- [10] L.A. Caffarelli, F.-H. Lin, Singularly perturbed elliptic systems and multivalued harmonic functions with free boundaries, J. Amer. Math. Soc., 21 (2008), 847-862.

- [11] L. A. Caffarelli, J. M. Roquejoffre, Uniform Höder estimates in a class of elliptic systems and applications to singular limits in models for diffusion flames, *Arch. Ration. Mech. Anal.*. **183** (2007), 457-487.
- [12] G. Cerami, D. Fortunato, M. Struwe, Bifurcation and multiplicity results for nonlinear elliptic problems involving critical Sobolev exponents, Ann. Inst. H. Poincare-Analyse, Nonlin., 1 (1984), 341-350.
- [13] G. Cerami, S. Solimini, M. Struwe, Some existence results for superlinear elliptic boundary value problems involving critical exponents, *J. Funct. Anal.*, **69** (1986), 289-306.
- [14] M. Conti, S. Terracini, G. Verzini, Asymptotic estimates for the spatial segregation of competitive systems, *Adv. Math.*, **195** (2005), 524-560.
- [15] Z. Chen, W. Zou, On the Brezis-Nirenberg problem in a ball, Differential and Integral Equations, 25 (2012), 527-542.
- [16] Z. Chen, W. Zou, Positive least energy solutions and phase separation for coupled Schrödinger equations with critical exponent, Arch. Ration. Mech. Anal., 205 (2012), 515-551.
- [17] G. Devillanova, S. Solimini, Concentration estimates and multiple solutions to elliptic problems at critical growth, *Adv. Differ. Equ.*, 7 (2002), 1257-1280.
- [18] N. Dancer, J. Wei, T. Weth, A priori bounds versus multiple existence of positive solutions for a nonlinear Schrödinger systems, Ann. Inst. H. Poincaré AN. 27 (2010), 953-969.
- [19] B. Esry, C. Greene, J. Burke, J. Bohn, Hartree-Fock theory for double condesates, Phys. Rev. Lett., 78 (1997), 3594-3597.
- [20] D. J. Frantzeskakis, Dark solitons in atomic Bose-Einstein condesates: from theory to experiments, J. Phys. A, 43 (2010) 213001.
- [21] D. Gilbarg, N. S. Trudinger, Elliptic partial differential equations of second order. Second edition, Grundlehren 224, Springer, Berlin, (1983).
- [22] S. Kim, On vector solutions for coupled nonlinear Schrödinger equations with critical exponents, *Comm. Pure Appl. Anal.*, **12** (2013), 1259-1277.
- [23] Yu. S. Kivshar, B. Luther-Davies, Dark optical solitons: physics and applications, *Physics Reports* **298** (1998), 81-197.
- [24] T. Lin, J. Wei, Ground state of N coupled nonlinear Schrödinger equations in \mathbb{R}^n , $n \leq 3$, Commun. Math. Phys., **255** (2005), 629-653.
- [25] Z. Liu, Z.-Q. Wang, Multiple bound states of nonlinear Schrödinger systems, Commun. Math. Phys., 282 (2008), 721-731.
- [26] Z. Liu, Z.-Q. Wang, On the Ambrosetti-Rabinowitz superlinear condition, Adv. Nonl. Stud., 4 (2004), 563-574.
- [27] L. Maia, E. Montefusco, B. Pellacci, Positive solutions for a weakly coupled nonlinear Schrödinger systems, *J. Differ. Equ.*, **229** (2006), 743-767.

- [28] B. Noris, H. Tavares, S. Terracini, G. Verzini, Uniform Hölder bounds for nonlinear Schrödinger systems with strong competition, *Comm. Pure Appl. Math.*, **63** (2010), 267-302.
- [29] B. Sirakov, Least energy solitary waves for a system of nonlinear Schrödinger equations in \mathbb{R}^n , Commun. Math. Phys., **271** (2007), 199-221.
- [30] M. Struwe, Variational methods- applications to nonlinear partial differential equations and Hamiltonian systems, springer, 1996.
- [31] A. Szukin, T. Weth, the method of Nehari manifolds, Handbook of Nonconvex Analysis and Applications, D.Y. Gao and D. Motreanu eds., International Press, Boston, 2010, 597-632.
- [32] M. Schechter, W. Zou, On the Brezis-Nirenberg Problem, Arch. Ration. Mech. Anal., 197 (2010), 337-356.
- [33] G. Talenti, Best constant in Sobolev inequality, Ann. Mat. Pure Appl., 110 (1976), 352-372.
- [34] J. Wei, T. Weth, Radial solutions and phase separation in a system of two coupled Schrödinger equations, Arch. Ration. Mech. Anal., 190 (2008), 83-106.
- [35] J. Wei, T. Weth, Asymptotic behaviour of solutions of planar elliptic systems with strong competition, *Nonlinearity*, **21** (2008), 305-317.
- [36] J. Wei, W. Yao, uniqueness of positive solutions to some coupled nonlinear Schrödinger equations, *Comm. Pure. Appl. Anal.*, **11** (2012), 1003-1011.
- [37] M. Willem, Minimax theorems, Birkhäuser, 1996.