

“Super”–Dilatation Symmetry of the Top-Higgs System

Christopher T. Hill*

Fermi National Accelerator Laboratory

P.O. Box 500, Batavia, Illinois 60510, USA

(Dated: February 22, 2022)

Abstract

The top-Higgs system, consisting of top quark (LH doublet, RH singlet) and Higgs boson kinetic terms, with gauge fields set to zero, has an *exact* (modulo total divergences) symmetry where both fermion and Higgs fields are shifted and mixed in a supersymmetric fashion. The full Higgs-Yukawa interaction and Higgs-potential, including additional $\sim 1/\Lambda^2$ NJL-like interactions, also has this symmetry to $\mathcal{O}(1/\Lambda^4)$, up to null-operators. Thus the interaction lagrangian can be viewed as a power series in $1/\Lambda^2$. The symmetry involves interplay of the Higgs quartic interaction with the Higgs-Yukawa interaction and implies the relationship, $\lambda = \frac{1}{2}g^2$ between the top-Yukawa coupling, g , and Higgs quartic coupling, λ , at a high energy scale $\Lambda \gtrsim \text{few TeV}$. We interpret this to be a new physics scale. The top quark is massless in the symmetric phase, satisfying the Nambu-Goldstone theorem. The fermionic shift part of the current is $\propto (1 - H^\dagger H/v^2)$, owing to the interplay of λ and g , and vanishes in the broken phase. Hence the Nambu-Goldstone theorem is trivially evaded in the broken phase and the top quark becomes heavy (it is not a Goldstino). We have $m_t = m_h$, subject to radiative corrections that can in principle pull the Higgs into concordance with experiment.

PACS numbers: 12.60.Cn, 12.60.Fr, 14.65.Ha, 03.70.+k, 11.10.-z, 11.30.-j

*Electronic address: hill@fnal.gov

I. INTRODUCTION

One of the remarkable features of the newly discovered Higgs boson is the proximity of three of the defining quantities of the standard model:

$$m_{Higgs}^2 \approx \frac{1}{2}m_{top}^2 \quad m_{top} \approx v_{weak} \quad (1)$$

where $v_{weak} \approx 175$ GeV is the Higgs VEV (Note: in the following m_h will denote the physical particle mass of the Higgs boson, while $M_H^2 H^\dagger H$ will be the lagrangian mass term) The part of the standard model lagrangian that subsumes these relationships is the top-Higgs system:

$$\mathcal{L} = \mathcal{L}_{kinetic} + g\bar{\psi}_L t_R H + h.c. - \frac{\lambda}{2} (H^\dagger H - v_{weak}^2)^2 \quad (2)$$

where $\psi = (t, b)$ [9]. Eq.(1) corresponds to the following choice of top Higgs-Yukawa, and quartic coupling constants at the scale of the top quark and Higgs boson masses:

$$g \approx 1, \quad \lambda \approx \frac{1}{4} \quad (3)$$

This is astonishingly consistent with experiment to within a few percent. These relationships are suggestive of some hidden dynamical or symmetrical interplay between the top quark and Higgs boson.

Our main interest is in mechanisms that can relate the quantities m_h , m_t and v_{weak} . One idea is that of a composite Higgs boson, *e.g.*, composed of $\bar{t}t$. This is an old approach [1, 2], typically treated in the context of the Nambu-Jona-Lasinio model, which implies some new underlying strong dynamics, such as topcolor [3]. The pure fermion loop approximation NJL model yields $m_h = 2m_t$, while fine-tuning a large hierarchy between the Higgs composite scale and the weak scale leads to a lighter Higgs boson, but $m_h \sim 260$ GeV, and $m_t \sim 220$ GeV, given the input $v_{weak} = 175$ GeV. Still other approaches avoid the fine-tuning, yielding $m_h \sim 1$ TeV. Hence $\bar{t}t$ composite Higgs models do succeed in relating m_h , m_t and v_{weak} , but generally predict a Higgs boson that is significantly heavier than the top quark [2]. We would have expected a light pNGB, such as the top-pion [3], in lieu of a light Higgs boson with the tree-level rates for $h \rightarrow ZZ^*$ (a pNGB would have $\pi \rightarrow ZZ^*$ at the significantly lower, axial anomaly level, rather than at the observed tree-level).

In the present paper we examine an alternative idea: we propose a generalization of dilatation symmetry for the Higgs boson that involves a “super”-symmetric relationship

between the top and Higgs fields. Indeed, this symmetry is exact in the top, with bottom-left, and Higgs kinetic terms, in the gaugeless limit (up to total divergences; it does not rely upon equations of motion).

This symmetry is similar to an ordinary c-number dilatation in the sense that it shifts the Higgs boson field, but the shift of the Higgs is now promoted to an operator. Our transformation also shifts the fermions. Finally, it includes a supersymmetric rotation, or “twist,” between the fermions and the Higgs field. The twist and shifts are locked together by the invariance of the Higgs mass term, the Higgs quartic interaction and the Higgs-Yukawa coupling constant in a nontrivial way. While the symmetry is exact in the (gaugeless) kinetic terms, we do make use of the fermion equations of motion (*i.e.*, we set null-operators to zero) in the transformation of the Higgs-Yukawa interaction. This has the nontrivial consequence of modifying the current in such a way that the fermionic shifts are present in the unbroken phase of the theory, but “turn off” in the broken phase.

Indeed, the symmetry yields $\lambda = g^2/2$ in both phases of the Standard Model. However, the fermionic shift part of the current is nontrivially modified by the quartic-Yukawa interplay and is $\propto (1 - H^\dagger H/v^2)$. It thus vanishes in the broken phase with $\langle H \rangle = v$. Therefore, the top quark can become massive in the broken phase in the usual way, with the relationship $m_h = m_t$. This is consistent with the Nambu-Goldstone theorem that would otherwise naively force the top quark to be a massless Goldstino. This relationship $\lambda = g^2/2$ is the analogue of a Goldberger-Treiman” relationship. It holds at a high scale, Λ , and is subject to renormalization group and higher dimension operator effects that can bring the physical masses into concordance with $m_h^2 \approx m_t^2/2$.

While the usual superalgebra of SUSY does not permit the Higgs to be the superpartner of the (t, b) quarks, the symmetry we present here does accomplish this. We emphasize at the outset that the “super”-dilatation symmetry is not a conventional superalgebra, *i.e.*, it is not a grading of the Lorentz Group, and is not associated with a nontrivial nonabelian closed superalgebra (at least not in our present exploratory formulation). The symmetry is a bosonic $U(1)$ invariance, and therefore closes trivially. As such, there is some *a priori* freedom in the normalization of its generators. The freedom is partially fixed by the exact invariance of the top and Higgs kinetic terms in the limit that gauge bosons are switched off (*i.e.*, global gauge symmetries remain). It is completely fixed by super-mixing between Higgs mass, quartic and Yukawa interactions.

Formally, this is reminiscent of a nonlinearly realized supersymmetry but it isn't: it's actually linearly realized. Operationally it is also similar to a "reparameterization invariance," *e.g.*, as occurs in heavy quark effective field theory (HQET) [4–6]. In the latter case one considers an $M \rightarrow \infty$ limit for a heavy quark and constructs a field theoretic lagrangian for a given four-velocity "supersector," v_μ . The lagrangian takes the form of a series expansion in higher dimension operators weighted by powers of $1/M$. The leading terms in the theory display heavy-spin symmetry (*e.g.*, degenerate 0^- and 1^- mesons). The reparameterization invariance is a residual symmetry that constrains the full operator structure and relates the coefficients of the terms in the lagrangian to higher orders of $1/M$. The reparameterization invariance is essentially the vestige of the underlying hidden Lorentz invariance [6].

We will presently move directly to discuss our symmetry. However, in Appendix A we review the analogous dilatonic Higgs symmetry to remind the reader of the interplay of a "shift" symmetry with a dilatation. The dilaton is inherited from a shift invariance, *e.g.*, the "modular symmetry" of a flat potential, which is then incorporated into the scale transformation. We show explicitly how this works, because we feel it is a confusing point in these schemes. The dilaton current comes from the shift-current and *not* the scale current, which is $S_\mu \sim x^\nu T_{\mu\nu}$. Here $T_{\mu\nu}$ is the "improved stress tensor" and involves only second derivatives of bosons whereas the dilatonic current is first order in derivatives. In Higgs-as-dilaton schemes we can always start in the unbroken phase, and the orientation of the Higgs boson in group space must be specified when a dilatonic shift is performed. This is in analogy to the fermionic orientations we encounter here. The reader may wish to consult Appendix A to see the analogy with our present construction.

In Appendix B we give, for motivation sake, the analogy of our present approach to a "bottoms-up" derivation of a nonlinear chiral lagrangian and the Goldberger-Treiman relation. This is completely parallel to the present derivation. We will initially consider the top and Higgs kinetic terms of the standard model with gauge fields set to zero:

$$\mathcal{L}_K = \bar{\psi}_L i \not{\partial} \psi_L + \bar{t}_R i \not{\partial} t_R + \partial H^\dagger \partial H \quad (4)$$

These kinetic term \mathcal{L}_K will be exactly invariant under our shift symmetry, modulo total divergences. We will then consider the potential terms:

$$\mathcal{L}_V = g(\bar{\psi}_L t_R H + h.c.) - M_H^2 H^\dagger H - \frac{\lambda}{2} (H^\dagger H)^2 \quad (5)$$

We will demand \mathcal{L}_V be invariant under the symmetry. We will be able to impose the symmetry, up to null operators (operators that vanish by equations of motion). This will yield the analogue of the “Goldberger-Treiman” relation, $\lambda = g^2/2$. It will also require the introduction of higher dimension operators $\sum_n \mathcal{O}^n \propto 1/\Lambda^{2n}$. At present we don’t know how to sum this series, and we will truncate at the scale $1/\Lambda^4$. Hence, to order $1/\Lambda^4$ we’ll have a lagrangian that is fully invariant under our symmetry. We will refer to our main relationship, $\lambda = g^2/2$, or $m_h = \sqrt{2}|M_H| = m_t$ as the analogue of a “Goldberger-Treiman” relation in this context.

II. “SUPER”–SHIFT SYMMETRY OF THE TOP-HIGGS KINETIC TERMS

We begin by introducing fixed orientations, $\theta_{L,R}$. These will be treated, with scale factors, as shifts of the quark fields themselves, $\delta\psi_L = \theta_L\epsilon + \dots$ and $\delta t_R = \theta_R\epsilon + \dots$, where $\psi_L = (t, b)_L$. The $\theta_{L,R}$ can be viewed as fixed zero 4-momentum components of the top (and bottom) fermion fields that have an arbitrary alignment in spin-space, and gauge group spaces of color and electroweak interactions. The $\theta_{L,R}$ are assumed to carry the same global color, isospin, weak hypercharge and baryon number quantum numbers as the corresponding ψ_L and t_R fermion fields, ie, $\theta_L \sim (N_c = 3, I = \frac{1}{2}, Y = \frac{1}{3}, B = \frac{1}{3})$ and $\theta_R \sim (N_c = 3, I = 0, Y = \frac{4}{3}, B = \frac{1}{3})$. In a sense, one could view the $\theta_{L,R}$ as constant VEV’s of the corresponding fermion fields, but they have no physical effect since the theory will be assumed symmetric under these small shifts. Our transformation is done for arbitrary fixed values of the $\theta_{L,R}$. The parameter is an infinitesimal bosonic c-number, ϵ . The $\theta_{L,R}$ are considered to have scale dimension- $\frac{3}{2}$, (in analogy to the dilatonic Higgs case of eq.(96) where we have scale-dimension 1). We thus introduce a scale Λ into our transformation, that is to be viewed as “large” compared to the weak scale.

Consider the top and Higgs kinetic terms of the standard model with gauge fields set to zero:

$$\mathcal{L}_K = \bar{\psi}_L i \not{\partial} \psi_L + \bar{t}_R i \not{\partial} t_R + \partial H^\dagger \partial H \quad (6)$$

We define the infinitesimal transformation:

$$\begin{aligned}
\delta\psi_L^{ia} &= \theta_L^{ia}\eta\epsilon - i\frac{\not{\partial} H^i\theta_R^a}{\Lambda^2}\epsilon; & \delta\bar{\psi}_{L\,ia} &= \bar{\theta}_{L\,ia}\eta\epsilon + i\frac{\bar{\theta}_{Ra}\not{\partial} H_i^\dagger}{\Lambda^2}\epsilon; \\
\delta t_R^a &= \theta_R^a\eta\epsilon - i\frac{\not{\partial} H_i^\dagger\theta_L^{ia}}{\Lambda^2}\epsilon; & \delta\bar{t}_{Ra} &= \bar{\theta}_{Ra}\eta\epsilon + i\frac{\bar{\theta}_{Lia}\not{\partial} H^i}{\Lambda^2}\epsilon; \\
\delta H^i &= \frac{\bar{\theta}_{Ra}\psi_L^{ia} + \bar{t}_{Ra}\theta_L^{ia}}{\Lambda^2}\epsilon; & \delta H_i^\dagger &= \frac{\bar{\psi}_{Lai}\theta_R^a + \bar{t}_{Lai}t_R^a}{\Lambda^2}\epsilon.
\end{aligned} \tag{7}$$

where i (a) is an isospin (color) index. η is a relative normalization factor that we determine subsequently. Presently we are not including a c-number shift in H , (as in eq.(96) of Appendix A), into the full transformation. The above transformation is our most general case, but it is convenient to consider more minimal versions of the symmetry where either θ_L or θ_R are set to zero or equal (see section III).

It is readily seen that eq.(7) is an invariance of eq.(6) up to total derivatives:

$$\begin{aligned}
\delta(\bar{\psi}_L i \not{\partial} \psi_L) &= \frac{(\bar{\psi}_L \theta_R) \cdot \partial^2 H}{\Lambda^2} \epsilon + h.c. + t.d. \\
\delta(\bar{t}_R i \not{\partial} t_R) &= \frac{(\bar{\theta}_L t_R) \cdot \partial^2 H}{\Lambda^2} \epsilon + h.c. + t.d. \\
\delta(\partial H^\dagger \partial H) &= -\frac{(\bar{\psi}_L \theta_R + \bar{\theta}_L t_R) \cdot \partial^2 H}{\Lambda^2} \epsilon + h.c. + t.d. \\
\text{hence, } \delta\mathcal{L}_K &= 0 + t.d.
\end{aligned} \tag{8}$$

(we've suppressed indices; "t.d." refers to "total divergence"). The symmetry of the gauge free kinetic terms makes no use of equations of motion or on-shell conditions. At this stage, the shifts in ψ_L and t_R by $\theta_{L,R}$ proportional to η play no role, but will be essential with the Higgs-Yukawa interaction and Higgs mass term. Indeed, shifting $\bar{\psi}_L i \not{\partial} \psi_L \rightarrow \bar{\psi}_L i \not{\partial} \theta_L$ yields a total divergence provided we have switched off the local gauge fields.

This transformation exploits the interplay of the quantum numbers of ψ_L , t_R and H . It resembles a scalar supermultiplet transformation of component fields [8], where the Higgs field is treated as a superpartner of ψ_L . We emphasize that this is not a representation of the supersymmetry algebra (there is no "F" auxillary field, [8]; this is essentially a scalar supermultiplet transformation with fixed $F = 0$ and the superparameters replaced by $\theta\epsilon$). With the assignment of scales of the $\theta_{L,R}$ and the presence of Λ the commutators of subsequent transformations for different $\theta_{L,R}$ cannot close. Also, the $\theta_{L,R}$ carry flavor and color quantum numbers, and the failure of the algebra to close into a superalgebra is presumably a supersymmetric extension of the Coleman-Mandula no-go theorem. In fact, this is a $U(1)$

symmetry with the transformation parameter, ϵ , for fixed background values of $\theta_{L,R}$. As such, the commutator trivially vanishes on the fields:

$$[\delta_{\epsilon'}, \delta_{\epsilon}](\psi, H, t_R) = 0 \quad (9)$$

We presently turn to the full lagrangian of the top-Higgs system in the standard model with gauge fields turned off:

$$\begin{aligned} \mathcal{L}_H = & i\bar{\psi}_L \not{\partial} \psi_L + i\bar{t}_R \not{\partial} t_R + \partial H^\dagger \partial H \\ & + g(\bar{\psi}_L t_R H + h.c.) - M_H^2 H^\dagger H - \frac{\lambda}{2} (H^\dagger H)^2 \end{aligned} \quad (10)$$

From eq.(7) we compute the transformations:

$$\delta(-M_H^2 H^\dagger H) = -\frac{\epsilon}{\Lambda^2} M_H^2 (\bar{\psi}_L \theta_R + \bar{\theta}_L t_R) \cdot H + h.c. \quad (11)$$

$$\delta(-\frac{\lambda}{2} (H^\dagger H)^2) = -\frac{\epsilon}{\Lambda^2} \lambda (\bar{\psi}_L \theta_R + \bar{\theta}_L t_R) \cdot H H^\dagger H + h.c. \quad (12)$$

$$\begin{aligned} \delta(g\bar{\psi}_L t_R H + h.c.) = & g\eta\epsilon(\bar{\psi}_L \theta_R + \bar{\theta}_L t_R) H + g^2 \frac{\epsilon}{2\Lambda^2} (\bar{\theta}_R \psi_L + \bar{t}_R \theta_L) \cdot (H^\dagger H^\dagger H) \\ & + g \frac{\epsilon}{\Lambda^2} \bar{\psi}_L t_R (\bar{\theta}_R \psi_L + \bar{t}_R \theta_L) \\ & + ig \frac{2\epsilon}{\Lambda^2} \bar{\psi}_L \gamma_\mu \frac{\tau^A}{2} \theta_L \left(H^\dagger \overset{\leftrightarrow}{\partial}^\mu \frac{\tau^A}{2} H \right) + ig \frac{\epsilon}{2\Lambda^2} \bar{\psi}_L \gamma_\mu \theta_L \left(H^\dagger \overset{\leftrightarrow}{\partial}^\mu H \right) \\ & - ig \frac{\epsilon}{\Lambda^2} \bar{\theta}_R \gamma_\mu t_R \left(H^\dagger \overset{\leftrightarrow}{\partial}^\mu H \right) + h.c. + t.d. \end{aligned} \quad (13)$$

Eq.(13) is crucial to our construction. It is obtained as follows:

$$\begin{aligned} \delta(g\bar{\psi}_L t_R H + h.c.) = & g \left(\bar{\psi}_L (\theta_R \eta \epsilon - i\epsilon \frac{\not{\partial} H^\dagger \theta_L}{\Lambda^2}) H + (\bar{\theta}_L \eta \epsilon + i\epsilon \frac{\bar{\theta}_R \not{\partial} H^\dagger}{\Lambda^2}) t_R H \right) \\ & + g\epsilon \left(\bar{\psi}_L t_R (\bar{\theta}_R \psi_L + \bar{t}_R \theta_L) \frac{1}{\Lambda^2} \right) + h.c. \\ = & g\eta\epsilon(\bar{\psi}_L \theta_R + \bar{\theta}_L t_R) H + \frac{g\epsilon}{\Lambda^2} \bar{\psi}_L t_R (\bar{\theta}_R \psi_L + \bar{t}_R \theta_L) \\ & + i \frac{g\epsilon}{2\Lambda^2} (\partial^\mu \bar{\psi}_L \gamma_\mu) \cdot H (H^\dagger \cdot \theta_L) - i \frac{g\epsilon}{2\Lambda^2} \bar{\theta}_R (\gamma_\mu \partial^\mu t_R) (H^\dagger \cdot H) \\ & + i \frac{2g\epsilon}{\Lambda^2} \bar{\psi}_L \gamma_\mu \frac{\tau^A}{2} \theta_L \left(H^\dagger \overset{\leftrightarrow}{\partial}^\mu \frac{\tau^A}{2} H \right) + i \frac{g\epsilon}{2\Lambda^2} \bar{\psi}_L \gamma_\mu \theta_L \left(H^\dagger \overset{\leftrightarrow}{\partial}^\mu H \right) \\ & - i \frac{g\epsilon}{\Lambda^2} \bar{\theta}_R \gamma_\mu t_R \left(H^\dagger \overset{\leftrightarrow}{\partial}^\mu H \right) + h.c. + t.d. \end{aligned} \quad (14)$$

where we use the isospin Fierz identity, $[\tau^A]_{ij}[\tau^A]_{kl} = 2\delta_{il}\delta_{kj} - \delta_{ij}\delta_{kl}$, and: $\overset{\leftrightarrow}{\partial}^\mu = \frac{1}{2}(\overset{\rightarrow}{\partial}^\mu - \overset{\leftarrow}{\partial}^\mu)$.

We then apply the fermionic equations of motion to eq.(14):

$$i\not{\partial} t_R + g\psi_L \cdot H^\dagger = 0 \quad i\not{\partial} \psi_L + gt_R H = 0 \quad (15)$$

and eq.(13) follows.

Notice in eq.(13) we have generated higher dimension operator terms of the form:

$$\begin{aligned} & \frac{g\epsilon}{\Lambda^2} \bar{\psi}_L t_R (\bar{\theta}_R \psi_L + \bar{t}_R \theta_L) + i \frac{2g\epsilon}{\Lambda^2} \bar{\psi}_L \gamma_\mu \frac{\tau^A}{2} \theta_L (H^\dagger \overset{\leftrightarrow}{\partial}^\mu \frac{\tau^A}{2} H) \\ & + i \frac{g\epsilon}{2\Lambda^2} \bar{\psi}_L \gamma_\mu \theta_L (H^\dagger \overset{\leftrightarrow}{\partial}^\mu H) - i \frac{g\epsilon}{\Lambda^2} \bar{\theta}_R \gamma_\mu t_R (H^\dagger \overset{\leftrightarrow}{\partial}^\mu H) + h.c. + t.d. \end{aligned} \quad (16)$$

In analogy to the “bottoms up” derivation of a nonlinear chiral lagrangian (see Appendix B), these terms can be cancelled by adding higher dimension operators to the original lagrangian of the form:

$$\begin{aligned} \mathcal{L}_{d=6} = & \frac{\kappa}{\Lambda^2} (\bar{\psi}_L t_R \bar{t}_R \psi_L) + \frac{2\kappa}{\Lambda^2} (\bar{\psi}_L \gamma_\mu \frac{\tau^A}{2} \psi_L) (H^\dagger i \overset{\leftrightarrow}{\partial}^\mu \frac{\tau^A}{2} H) \\ & + \frac{\kappa}{2\Lambda^2} (\bar{\psi}_L \gamma_\mu \psi_L) (H^\dagger i \overset{\leftrightarrow}{\partial}^\mu H) - \frac{\kappa}{\Lambda^2} (\bar{t}_R \gamma_\mu t_R) (H^\dagger i \overset{\leftrightarrow}{\partial}^\mu H) \end{aligned} \quad (17)$$

where we will relate the coupling constant, κ , to M_H , Λ and g below.

We thus obtain the effective lagrangian,

$$\begin{aligned} \mathcal{L}_H = & \bar{\psi}_L i \not{\partial} \psi_L + i \bar{t}_R \not{\partial} t_R + \partial H^\dagger \partial H \\ & + g (\bar{\psi}_L t_R H + h.c.) - M_H^2 H^\dagger H - \frac{\lambda}{2} (H^\dagger H)^2 \\ & + \frac{\kappa}{\Lambda^2} (\bar{\psi}_L t_R \bar{t}_R \psi_L) + \frac{2\kappa}{\Lambda^2} (\bar{\psi}_L \gamma_\mu \frac{\tau^A}{2} \psi_L) (H^\dagger i \overset{\leftrightarrow}{\partial}^\mu \frac{\tau^A}{2} H) \\ & + \frac{\kappa}{2\Lambda^2} (\bar{\psi}_L \gamma_\mu \psi_L) (H^\dagger i \overset{\leftrightarrow}{\partial}^\mu H) - \frac{\kappa}{\Lambda^2} (\bar{t}_R \gamma_\mu t_R) (H^\dagger i \overset{\leftrightarrow}{\partial}^\mu H) \end{aligned} \quad (18)$$

Performing the super-dilatation transformation of eq.(7) we now demand that:

$$\delta \mathcal{L}_H = \mathcal{O} \left(\frac{1}{\Lambda^4} \right) \quad (19)$$

The generated $\mathcal{O}(1/\Lambda^4)$ terms can be compensated by adding additional $1/\Lambda^4$ terms to the lagrangian. By continued application of eq.(7) we would generate a power series of contact interactions that are scaled by $\sim 1/\Lambda^{2n}$.

First we see that the transformation of the Higgs mass term of eq.(18), from eqs.(11–13), cancels against the first term of the transformed Higgs-Yukawa interaction, provided:

$$g\eta = \frac{M_H^2}{\Lambda^2} \quad (20)$$

(beware: M_H^2 is the negative Lagrangian (mass)², while $m_h^2 = -2M_H^2$ is the physical positive Higgs boson (mass)² in the broken phase, and we take the normalization $v^2 = \langle H^\dagger H \rangle =$

(175 GeV)²) This establishes the normalization factor, η . It also establishes the relative sign (we assume Λ^2 positive). If we're in the symmetric (broken) phase, M_H^2 positive (negative), then we have $g\eta > 0$ ($g\eta < 0$). We have the freedom of choosing arbitrary η since the defining kinetic term invariance involves only ϵ .

One might think we can now take Λ^2 to be arbitrarily large compared to M_H^2 by adjusting $|\eta| \ll 1$. However, the second term of eq.(13) must also cancel against the transformation of the first κ term appearing in eq.(18). This requires that:

$$\eta\kappa = -g, \quad \text{or, using eq.(20):} \quad \frac{\kappa}{\Lambda^2} = -\frac{g^2}{M_H^2} \quad (21)$$

This is to us a striking result. In the $d = 6$ operators we have a Nambu-Jona-Lasinio component. The Nambu-Jona-Lasinio attractive interaction corresponds to $\kappa > 0$, and we see in eq.(21) that the super-dilatation is then consistent only if $M_H^2 < 0$. Moreover, to make η small requires taking κ large (or explicitly breaking the super-dilatation symmetry, which we consider in the weak coupling case IV.C).

Finally, the most interesting relationship, which is the analogue of the Goldberger-Treiman relationship in a chiral lagrangian (Appendix A), arises from the cancellation of the $\sim \epsilon(\bar{\psi}_L\theta_R + \bar{\theta}_Lt_R) \cdot HH^\dagger H$ terms of eqs.(12) and (13) under the super-dilatation symmetry. This transformation does not involve η and requires:

$$0 = (\lambda - \frac{1}{2}g^2)\frac{\epsilon}{\Lambda^2}(\bar{\psi}_L\theta_R + \bar{\theta}_Lt_R) \cdot HH^\dagger H + h.c \quad (22)$$

or,

$$\lambda = \frac{1}{2}g^2 \quad (23)$$

We emphasize that the critical aspect of our construction is that the operator shift of δH in the quartic Higgs interaction is cancelling against the super-rotation (*i.e.*, the “twist”) of $\delta\psi$ in the Higgs-Yukawa interaction. Moreover, the pure fermionic shift in $\delta\psi \sim \eta\epsilon\theta$, in the Higgs-Yukawa interaction, *i.e.*, proportional to η , cancels against the δH shift in the Higgs mass term. This ties the transformations together into a single structure. We'll see, in the next section that this leads to a remarkable effect in the current structure. The Nambu-Goldstone theorem for the fermionic shift $\delta\psi \sim \eta\epsilon\theta$, which would naively imply a massless top quark (a “Goldstino”), is evaded in the broken phase.

Note that the $\lambda = g^2/2$ relationship refers to the coefficient of the $D = 6$ operator, $(\bar{\theta}_Lt_R) \cdot HH^\dagger H + h.c$. We therefore assume that it applies at the scale Λ . The low energy

relationship between the couplings g^2 and λ then depends upon the renormalization group running from Λ to $v_{weak} \approx 175$ GeV. If we ignore the RG running then eq.(23) would hold at the weak scale, and implies the supersymmetric relationship $m_h^2 = 2\lambda v_{weak}^2 = m_t^2$ in the broken phase. This is an improvement over the usual NJL result, $m_h^2 = 4m_t^2$.

Some comments on the structure of these higher dimension operators are in order. Note that we can Fierz rearrange the first term of eq.(17):

$$(\bar{\psi}_L^a t_{Ra})_i (\bar{t}_{Rb} \psi^b)^i \rightarrow -(\bar{\psi}_{iL} \gamma_\mu \frac{\lambda^A}{2} \psi_L^i) (\bar{t}_R \gamma^\mu \frac{\lambda^A}{2} t_R) + \mathcal{O}(\infty/\mathcal{N}) \quad (24)$$

where $N = 3$ is the number of colors. This term is a pure Nambu-Jona-Lasinio interaction as arises in topcolor [2, 3] in the form of a (color current) \times (color current). Indeed, massive Yang-Mills boson exchange for a boson of mass M^2 and momentum exchange $q^2 < M^2$ produces the negative sign for (current) \times (current) interactions. A positive sign for the first term of eq.(17) is the attractive sign for the Nambu-Jona-Lasinio model, and we thus see that the attractive sign corresponds to the correct (negative) sign for topgluon exchange. However, we see that the isospin (current) \times (current) interaction (second term of eq.(17)) then has the wrong sign (positive) for a gauge boson exchange. We will discuss this “frustration of signs” further in Section IV.A.

Since all of the higher dimension $d = 6$ operators are of the form (current) \times (current), they preserve the chirality of the fermions. That is, the terms of eq.(17) contain no cross terms of the form $\bar{\psi}_L H t_R (H^\dagger H)^p$. They thus admit the discrete symmetry, $\psi_L \rightarrow (-1)^N \psi_L$ and $t_R \rightarrow (-1)^{N+1} t_R$. Operators of mixed chirality can therefore be excluded, or suppressed, on symmetry grounds, though we consider their inclusion in Section (IV.C) to improve the prediction for m_h .

III. CURRENT STRUCTURE AND NAMBU-GOLDSTONE THEOREM

The critical aspect of our construction is that the operator shift of δH in the quartic Higgs interaction is cancelling against the super-rotation (*i.e.*, the “twist”) of $\delta\psi$ in the Higgs-Yukawa interaction. Moreover, the pure fermionic shift in $\delta\psi \sim \eta\epsilon\theta$, in the Higgs-Yukawa interaction, *i.e.*, proportional to η , cancels against the δH shift in the Higgs mass term. This ties the transformations together into a single structure.

The Nambu-Goldstone theorem for a fermionic shift $\delta\psi \sim \eta\epsilon\theta$, which would naively imply

a massless top quark (a “Goldstino”), is evaded in the broken phase. How does our theory evade the existence of a zero-mode associated with the fermionic shift? Naively, this would seem to prohibit a massless top quark. In fact, this happens in a subtle way. One must carefully construct the currents given our use of equations of motion in $\delta(g\bar{\psi}_L t_R H + h.c.)$. We therefore wish to clarify the the relationship to the Nambu-Goldstone theorem in the present set up.

We consider, for technical simplicity, a simpler “minimal” transformation defined by $\theta_L = 0$:

$$\delta\psi_L^{ia} = -i\frac{\not{\partial} H^i \theta_R^a}{\Lambda^2}\epsilon; \quad \delta\bar{\psi}_L{}_{ia} = i\frac{\bar{\theta}_{Ra}\not{\partial} H_i^\dagger}{\Lambda^2}\epsilon; \quad (25)$$

$$\delta t_R^a = \theta_R^a \eta \epsilon; \quad \delta \bar{t}_{Ra} = \bar{\theta}_{Ra} \eta \epsilon; \quad (26)$$

$$\delta H^i = \frac{\bar{\theta}_{Ra} \psi_L^{ia}}{\Lambda^2} \epsilon; \quad \delta H_i^\dagger = \frac{\bar{\psi}_{Lai} \theta_R^a}{\Lambda^2} \epsilon. \quad (27)$$

The parameter η is still fixed by the symmetry interplay of the Higgs mass term and Yukawa interaction as in eq.(20),

$$g\eta = \frac{M_H^2}{\Lambda^2} = -\frac{\lambda v^2}{\Lambda^2} \quad (28)$$

Consider the top and Higgs system of the standard model with gauge fields set to zero:

$$\mathcal{L}_H = \mathcal{L}_K + g(\bar{\psi}_L t_R H + h.c.) - M_H^2 H^\dagger H - \frac{\lambda}{2}(H^\dagger H) \quad (29)$$

$$\mathcal{L}_K = \bar{\psi}_L i \not{\partial} \psi_L + \bar{t}_R i \not{\partial} t_R + \partial H^\dagger \partial H \quad (30)$$

It is readily seen that eqs.(25–27) is a global invariance of eqs.(30) up to total derivatives. We presently allow ϵ to be a local function of spacetime $\epsilon(x)$ (note that the derivatives in eq.(25) act only upon H and not upon $\epsilon(x)$). We have:

$$\begin{aligned} \delta(\bar{\psi}_L i \not{\partial} \psi_L) &= \frac{(\bar{\psi}_L \theta_R) \cdot \partial^2 H}{\Lambda^2} \epsilon + \frac{(\bar{\psi}_L \gamma_\mu \not{\partial} H \theta_R)}{\Lambda^2} \partial^\mu \epsilon + h.c. + t.d. \\ \delta(\bar{t}_R i \not{\partial} t_R) &= i(\bar{t}_R \not{\partial} \theta_R) \eta \epsilon + i(\bar{t}_R \gamma_\mu \theta_R) \eta \partial^\mu \epsilon + h.c. + t.d. \\ \delta(\partial H^\dagger \partial H) &= -\frac{(\bar{\psi}_L \theta_R) \cdot \partial^2 H}{\Lambda^2} \epsilon + \frac{(\bar{\psi}_L \theta_R) \cdot \partial_\mu H}{\Lambda^2} \partial^\mu \epsilon + h.c. + t.d. \end{aligned} \quad (31)$$

The kinetic terms thus lead to a Noether current:

$$J_\mu^K = \frac{\delta \mathcal{L}_K}{\delta \partial_\mu \epsilon} = i(\bar{t}_R \gamma_\mu \theta_R) \eta + \frac{(\bar{\psi}_L \gamma_\mu \not{\partial} H \theta_R)}{\Lambda^2} + \frac{(\bar{\psi}_L \theta_R)}{\Lambda^2} \partial_\mu H + h.c. \quad (32)$$

The symmetry of the full action, as we have emphasized, involves a cancellation of the shift of eqs.(27) in the Higgs quartic interaction against the “twist” of eq.(25) in the Higgs-Yukawa

interaction. In calculating the transformation of the Higgs-Yukawa interaction, however, we make use of an “integration by parts” and discard total divergences (and subsequently use the fermion equations of motion). This integration by parts in the “twist” of eq.(25) causes the derivative to act upon the parameter $\epsilon(x)$:

$$\delta(-M_H^2 H^\dagger H) = -\frac{\epsilon}{\Lambda^2} M_H^2 (\bar{\psi}_L \theta_R) \cdot H + h.c \quad (33)$$

$$\delta(-\frac{\lambda}{2} (H^\dagger H)^2) = -\frac{\epsilon}{\Lambda^2} \lambda (\bar{\psi}_L \theta_R) \cdot H H^\dagger H + h.c. \quad (34)$$

$$\begin{aligned} \delta(g \bar{\psi}_L t_R H + h.c.) &= g \eta \epsilon (\bar{\psi}_L \theta_R) H + \frac{g^2 \epsilon}{2 \Lambda^2} (\bar{\theta}_R \psi_L) \cdot (H^\dagger H^\dagger H) \\ &\quad + \frac{g \epsilon}{\Lambda^2} (\bar{\psi}_L t_R) (\bar{\theta}_R \psi_L) - i \frac{g \epsilon}{\Lambda^2} \bar{\theta}_R \gamma_\mu t_R \left(H^\dagger \overset{\leftrightarrow}{\partial}^\mu H \right) \\ &\quad - i \frac{g}{2 \Lambda^2} \bar{\theta}_R \gamma_\mu t_R (H^\dagger H) \partial^\mu \epsilon + h.c. + t.d. \end{aligned} \quad (35)$$

The last term in eq.(35) shows explicitly that the result of the integration by parts leads to an additional term $\propto \partial^\mu \epsilon$. This, in turn, modifies the current, which now becomes:

$$J_\mu = \frac{\delta \mathcal{L}_H}{\delta \partial_\mu \epsilon} = i(\bar{t}_R \gamma_\mu \theta_R) \left(\eta + \frac{g H^\dagger H}{2 \Lambda^2} \right) + \frac{(\bar{\psi}_L \gamma_\mu \not{\partial} H \theta_R)}{\Lambda^2} + \frac{(\bar{\psi}_L \theta_R)}{\Lambda^2} \partial_\mu H + h.c. \quad (36)$$

Using the relationship of eq.(28), $g \eta = -\lambda v^2 / \Lambda^2$, and the “Goldberger-Treiman” analogue, $\lambda = g^2 / 2$, the current can be written:

$$J_\mu = \frac{\delta \mathcal{L}_H}{\delta \partial_\mu \epsilon} = i \eta (\bar{t}_R \gamma_\mu \theta_R) \left(1 - \frac{H^\dagger H}{v^2} \right) + \frac{(\bar{\psi}_L \gamma_\mu \not{\partial} H \theta_R)}{\Lambda^2} + \frac{(\bar{\psi}_L \theta_R)}{\Lambda^2} \partial_\mu H + h.c. \quad (37)$$

The modification of the current occurs in the first term which is associated with the fermionic “shift”-symmetry of t_R . Again, this arises from the crucial linkage of the δH shift in the quartic Higgs interaction to the $\delta \psi_L$ shift in the Higgs-Yukawa interaction.

Note that, in the broken phase where $\langle H \rangle = v$ the modification of the current has the effect of “turning off” the fermionic shift. Indeed, we will now see that this has a remarkable effect in evading the Nambu-Goldstone theorem in the broken phase, and permitting the top quark to be massive.

Consider the two-point function of our current of eq.(37) with t_R :

$$S(y) = \int d^4 x e^{i q \cdot x} \partial^\mu \langle 0 | T^* J_\mu(x) t_R(y) | 0 \rangle \quad (38)$$

(T^* implies anti-commutation in the ordering of fermion fields). Formally, with $\partial^\mu J_\mu = 0$,

we have, from the ∂_0 acting upon the T^* ordering a $\delta(x^0 - y^0)$, and:

$$\begin{aligned}
S(y) &= \int d^3x e^{iq \cdot x} \langle 0 | \{J_0(x), t_R(y)\} | 0 \rangle \\
&= \int d^3x e^{-i\vec{q} \cdot \vec{x}} \langle 0 | \{J_0(\vec{x}), t_R(\vec{y})\}_{e.t.} | 0 \rangle \\
&= \langle 0 | \{Q, t_R(\vec{y})\} | 0 \rangle
\end{aligned} \tag{39}$$

where the charge operator Q is:

$$Q = \int d^3x J_0(\vec{x}). \tag{40}$$

In the symmetric phase of the standard model we have the Higgs VEV, $\langle H \rangle = 0$, and we can neglect all terms in the current that involve H . The charge operator then involves only the first term in $J_\mu^K = i\eta \bar{t}_R \gamma_\mu \theta_R + h.c.$, whence it generates a shift in the fermion field:

$$\langle 0 | \{Q, t_R(\vec{y})\} | 0 \rangle = \eta \theta_R \tag{41}$$

On the other hand we have:

$$\begin{aligned}
S(y) &= \int d^4x e^{iq \cdot x} \partial^\mu \langle 0 | T^* i \bar{t}_R(x) \gamma_\mu \theta_R \eta \quad t_R(y) | 0 \rangle \\
&= - \int d^4x e^{iq \cdot x} i \partial^\mu \gamma_\mu S_F(x - y) \theta_R \eta \\
&= \left. \frac{q^2 + \not{q} m}{q^2 - m^2} \theta_R \eta \right|_{q \rightarrow 0}
\end{aligned} \tag{42}$$

In the $q^2 \rightarrow 0$ limit the consistency of eq.(42) with eqs.(39, 41) requires that the fermion mass satisfy $m = 0$. This is the fermionic Nambu-Goldstone theorem and it informs us that any fermionic action which has a pure fermionic shift symmetry, must correspond to a massless fermion. This is, indeed, the case in the symmetric phase in which the top quark is massless and $\langle H \rangle = 0$.

Naively we might conclude that the top quark is forced by our symmetry to be a Goldstino and remain massless, even in the broken phase. However, we have seen that the current is modified in a significant way in the present case:

$$J_\mu^{shift} = i\eta (\bar{t}_R \gamma_\mu \theta_R) \left(1 - \frac{H^\dagger H}{v^2} \right) + h.c. \tag{43}$$

In the broken phase, when $\langle H \rangle = v \neq 0$, this implies that the pure fermionic shift operator in the current “turns off:”

$$J_\mu^{shift} = 0 \tag{44}$$

This is a consequence of the interplay between the quartic interaction and the Higgs-Yukawa interaction in our construction. It implies that there can exist dynamical situations in which a Goldstino is massless in a symmetric phase, but acquires mass in a broken phase of a theory. The underlying fermionic shift, $\delta\psi = \eta\theta$ is intact, but the current is modified dynamically to evade the naive Nambu-Goldstone theorem. One might wonder what happens for $\langle H^\dagger H \rangle \neq v^2$ and $\neq 0$? This is, of course, an unstable vacuum and the S-matrix derivation fails. Of course, the above current algebra analysis serves only as a consistency check on our original Lagrangian analysis, which showed that a vacuum with massive top and Higgs, with $m_t = m_h$, exists.

Indeed, the symmetry yields $\lambda = g^2/2$ in both phases of the Standard Model. However, the fermionic shift part of the current is nontrivially modified by the quartic-Yukawa interplay and is $\propto (1 - H^\dagger H/v^2)$. It thus vanishes in the broken phase with $\langle H \rangle = v$. Therefore, the top quark becomes massive in the broken phase in the usual way, with the relationship $m_h = m_t$. This is consistent with the Nambu-Goldstone theorem that would otherwise naively force the top quark to be a massless Goldstino. This relationship $\lambda = g^2/2$ is the analogue of a Goldberger-Treiman" relationship. It holds at a high scale, Λ , and is subject to renormalization group and higher dimension operator effects that can bring the physical masses into concordance with $m_h^2 \approx m_t^2/2$.

Super-Dilatation in the Broken Phase

The result of eq.(44) helps us to understand the symmetry when written in the broken phase of the theory. We will presently treat the broken phase in a vectorlike case, defining $\theta_L = \theta_R$, and we'll ignore the Nambu-Goldstone bosons of the Higgs field. With the Higgs potential defined as $V(H) = (\lambda/2)(H^\dagger H - v^2)^2$ we write:

$$H = \begin{pmatrix} v + h/\sqrt{2} \\ 0 \end{pmatrix} \quad (45)$$

The lagrangian then takes the form:

$$\mathcal{L}_H = \mathcal{L}_K + m_t \bar{t}t + \frac{g}{\sqrt{2}} \bar{t}t h - \lambda v^2 h^2 - \frac{\lambda}{\sqrt{2}} h^3 - \frac{\lambda}{8} h^4 + \mathcal{O}\left(\frac{1}{\Lambda^2}\right) \quad (46)$$

and the physical Higgs boson mass is $m_h^2 = 2v^2\lambda$. The kinetic terms become:

$$\mathcal{L}_K = i\bar{t}\not{\partial}t + \frac{1}{2}\partial_\mu h \partial^\mu h \quad (47)$$

As noted above, the fermionic shift “shuts off” in the broken phase. We assume the symmetry of the broken phase effective lagrangian now involves only the (“twist”) super-transformation on t and an operator shift of h :

$$\delta t = -i \frac{\not{\partial} h \theta}{\sqrt{2} \Lambda^2} \epsilon; \quad \delta \bar{t} = i \frac{\bar{\theta} \not{\partial} h}{\sqrt{2} \Lambda^2} \epsilon; \quad (48)$$

$$\delta h = \frac{\bar{\theta} t}{\sqrt{2} \Lambda^2} \epsilon + h.c. \quad (49)$$

where θ is a fixed orientation carrying color and spin. This is readily seen to be a symmetry, modulo total divergences, of the kinetic terms:

$$\delta \bar{t} i \not{\partial} t = \bar{t} \theta \frac{\partial^2 h}{\sqrt{2} \Lambda^2} \epsilon + h.c. + t.d. \quad (50)$$

$$\frac{1}{2} \delta \partial_\mu h \partial^\mu h = -\bar{t} \theta \frac{\partial^2 h}{\sqrt{2} \Lambda^2} \epsilon + h.c. + t.d. \quad (51)$$

No use of equations of motion is involved in the invariance of the kinetic terms. The transformations of eqs.(48,49) also define a symmetry of the full theory:

$$\delta(-\lambda v^2 h^2) = -\sqrt{2} \lambda v^2 h \left(\frac{\bar{\theta} t}{\Lambda^2} \epsilon + h.c. \right) \quad (52)$$

$$\delta\left(-\frac{\lambda v}{\sqrt{2}} h^3\right) = -\frac{3\lambda}{2} h^2 v \left(\frac{\bar{\theta} t}{\Lambda^2} \epsilon + h.c. \right) \quad (53)$$

$$\delta\left(-\frac{\lambda}{8} h^4\right) = -\frac{\lambda}{2\sqrt{2}} h^3 \left(\frac{\bar{\theta} t}{\Lambda^2} \epsilon + h.c. \right) \quad (54)$$

and:

$$\delta(m_t \bar{t} t) = \left(\frac{g^2 v^2 h}{\sqrt{2}} + \frac{g^2 v h^2}{2} \right) \left(\frac{\bar{\theta} t}{\Lambda^2} \epsilon + h.c. \right) \quad (55)$$

$$\begin{aligned} \delta\left(\frac{g}{\sqrt{2}} \bar{t} t h\right) &= \left(\frac{g^2 v h^2}{4} + \frac{g^2 h^3}{4\sqrt{2}} \right) \left(\frac{\bar{\theta} t}{\Lambda^2} \epsilon + h.c. \right) \\ &\quad + \frac{g \bar{t} t}{2} \left(\frac{\bar{\theta} t}{\Lambda^2} \epsilon + h.c. \right) \end{aligned} \quad (56)$$

where in eqs.(55 – 56) we have used $m_t = gv$ and the fermionic equation of motion:

$$i \not{\partial} t + m_t t + \frac{g}{\sqrt{2}} h t = 0 \quad - i \partial^\mu \bar{t} \gamma_\mu + m_t \bar{t} + \frac{g}{\sqrt{2}} h \bar{t} = 0 \quad (57)$$

One can readily check, using $\lambda = g^2/2$, that eq.(46) is invariant under eqs.(48–56). This implies $m_t = m_h$. The four-fermion term in eq.(55) requires the addition of the $(\bar{t} t)^2/\Lambda^2$ operator into the action, which in turn modifies the fermion equation of motion, and will also

be invariant under eqs.(48 – 49). Note that we are always demanding an exact symmetry of kinetic terms, without use of equations of motion. We use only the fermion equation of motion in the interaction terms which is equivalent to setting null operators in the action to zero. The resulting symmetry has the nontrivial consequence that $\lambda = g^2/2$, equivalently $m_t = m_h$.

IV. PHYSICAL INTERPRETATION

A. Minimal Symmetry and Correct-Sign (current) \times (current) Interactions

As we noted in section II, the full set of κ terms of eq.(18) have an inconsistency if we want to interpret them all in terms of gauge boson mediated (current) \times (current) interactions. In particular, the first two terms conflict if we Fierz the first κ term into a (color current) \times (color current) interaction, and compare to the second (isospin current) \times (isospin current) interaction. We see that the interactions have opposite signs; hence we can choose a given sign for κ and one of the two (current) \times (current) will have the wrong-sign (positive) for a Yang-Mills gauge boson mediated interaction term (negative; note that the $(U(1) \text{ current})\times(U(1) \text{ current})$ terms can have either sign, depending upon the $U(1)$ charge assignments of the fields).

However, there is a simple remedy: use the minimal symmetry. The (isospin current) \times (isospin current) interaction arises from the presence of θ_L in our transformation eq.(13), while the NJL term arises from both θ_L and θ_R . We can thus use “minimal symmetry” that contains only the iso-singlet θ_R :

$$\begin{aligned}\delta\psi_L^{ia} &= -i\frac{\not{\partial} H^i\theta_R^a}{\Lambda^2}\epsilon; & \delta\bar{\psi}_L{}_{ia} &= i\frac{\bar{\theta}_{Ra}\not{\partial} H_i^\dagger}{\Lambda^2}\epsilon; \\ \delta t_R^a &= \theta_R^a\eta\epsilon; & \delta\bar{t}_{Ra} &= \bar{\theta}_{Ra}\eta\epsilon; \\ \delta H^i &= \frac{\bar{\theta}_{Ra}\psi_L^{ia}}{\Lambda^2}\epsilon; & \delta H_i^\dagger &= \frac{\bar{\psi}_{La}i\theta_R^a}{\Lambda^2}\epsilon.\end{aligned}\tag{58}$$

The effective lagrangian we now obtain is simpler:

$$\begin{aligned}\mathcal{L}_H &= \bar{\psi}_L i\not{\partial} \psi_L + i\bar{t}_R \not{\partial} t_R + \partial H^\dagger \partial H \\ &\quad + g(\bar{\psi}_L t_R H + h.c.) - M_H^2 H^\dagger H - \frac{\lambda}{2}(H^\dagger H)^2 \\ &\quad + \frac{\kappa}{\Lambda^2}(\bar{\psi}_L t_R \bar{t}_R \psi_L) - \frac{\kappa}{\Lambda^2}(\bar{t}_R \gamma_\mu t_R)(H^\dagger i \overleftrightarrow{\partial}^\mu H)\end{aligned}\tag{59}$$

The frustrated (current) \times (current) signs are now absent given that we have banished the isospin interaction. We retain the relationships of eqs.(20,21,23). The minimal symmetry may be a more natural framework, and is also suggested by the weakly interacting case discussed in Section IV.D below.

B. Strong Coupling

Let us consider a super-dilatation symmetry in the context of a UV completion model involving a presumed new strong dynamics at the (multi-) TeV scale. We assume the broken phase and hence $M_H^2 < 0$, and $\kappa > 0$ and large.

The largest physically acceptable value we might consider for the coupling constant κ is of order a critical value $\kappa < \kappa_c \sim (4\pi)^2$. Thus, we obtain (recall the physical Higgs boson mass, $125 \text{ GeV} = \sqrt{2}M_H$):

$$\Lambda^2 < \kappa_c |M_H^2|/g^2 \quad \Lambda \lesssim 4\pi|M_H| \sim 1 \text{ TeV for } g \sim 1. \quad (60)$$

(RG effects allow $g \lesssim 1$ and Λ could thus be upwards of \sim a few TeV within our approximations).

We assume a strong interaction at the high scale Λ , containing new physics described by the κ terms in eq.(18). All of these terms are current-current interactions. We interpret this as a compositeness scale for the Higgs boson, and the low energy theory is then the standard model. This differs from the usual NJL model, where g and λ diverge at the compositeness scale, which results in $m_h = 2m_t$ [2], and typically very large g and λ near Λ . λ is now locked to g^2 at the scale Λ by our symmetry.

If we also included the normal scalar dilation transformation of the Higgs, eq.(96), then λ is an explicit symmetry breaking scale, and can be arguably small, *i.e.*, the dilation symmetry becomes a custodial symmetry for small λ . Our basic hypothesis is that the super-dilatation is then a surviving spectrum symmetry of the solution to the theory. We've reverse engineered the strong UV theory as a power series in higher dimension operators from the symmetry. We have not explicitly shown that such a UV theory has, in fact, such a solution.

We can estimate the radiative effects on the Higgs mass using the renormalization group (RG) equations for λ and g . We include the QCD effects, and integrate in the approximation

of constant rh sides of the RG equations. This yields the leading log effect:

$$\begin{aligned} 16\pi^2 \frac{\partial \lambda}{\partial \ln(\mu)} &= 12\lambda^2 + 4N_c \lambda g^2 - 4N_c g^4 \approx (3 - 2N_c)g^4 \approx -3, \\ 16\pi^2 \frac{\partial g^2}{\partial \ln(\mu)} &= (2N_c + 3)g^4 - 2(N_c^2 - 1)g^2 g_{QCD}^2 \approx 9g^4 - 16 \times (4\pi\alpha_{QCD})g^2 \approx -13 \end{aligned} \quad (61)$$

In the last expressions we've substituted $\lambda = g^2/2$, $\alpha_{QCD} = 0.11$ and $g = 1$. This yields, upon imposing the boundary condition $\lambda(\Lambda) - g^2(\Lambda)/2 = 0$:

$$\lambda(v_{weak}) - \frac{1}{2}g^2(v_{weak}) \approx -\frac{3.5}{16\pi^2} \ln\left(\frac{\Lambda}{v_{weak}}\right) \approx -0.04 \quad (62)$$

This implies $m_h = \sqrt{2\lambda}v_{weak} \approx 167$ GeV. This is an improvement over Nambu-Jona-Lasinio inspired models, ~ 260 GeV [2], but still too high. The radiative corrections slightly nudge the mass relationship in the right direction, $m_h < m_t$. Naively, with $\Lambda \sim 10^7$ GeV we bring $m_h \sim 125$ GeV.

C. Chiral Higher Dimension Operators

We consider presently modifications to eq.(18) consisting of towers of higher dimension chiral operators:

$$\mathcal{L}_H = \mathcal{L}_{KT} + g(\bar{\psi}_L t_R H + h.c.)P(H^\dagger H) + M_H^2 H^\dagger H - \frac{\lambda}{2}Q(H^\dagger H) \quad (63)$$

where we work in the broken phase and:

$$P(H^\dagger H) = \sum_{n=0} c_n \left(\frac{H^\dagger H}{\Lambda^2}\right)^n \quad c_0 = 1 \quad (64)$$

$$Q(H^\dagger H) = \sum_{n=0} d_n (H^\dagger H)^2 \left(\frac{H^\dagger H}{\Lambda^2}\right)^n \quad d_0 = 1 \quad (65)$$

The P operators break the discrete chiral symmetry, $\psi_L \rightarrow (-1)^N \psi_L$ and $t_R \rightarrow (-1)^{N+1} t_R$. These operators are connected by the super-dilatation to the Q 's as we'll see below. Therefore, we would expect the coefficients of these to be naturally small.

The Higgs potential minimum is now modified by Q , given by:

$$0 = -M_H^2 + \frac{\lambda}{2} \frac{\partial Q(v)}{\partial (v)^2} \Big|_{v=v_{weak}} \quad (66)$$

where $H^\dagger H \equiv v^2$. At the potential minimum, the top quark mass is:

$$m_t = gv_{weak}P(v_{weak}^2) \quad (67)$$

M_H^2 is, of course, the curvature of the potential at the origin, $v \approx 0$, but the physical Higgs boson mass is determined by the curvature at the potential minimum, $v = v_{weak}$:

$$m_h^2 = -M_H^2 + \frac{\lambda}{4} \frac{\partial^2 Q(v)}{(\partial v)^2} \Big|_{v=v_{weak}} \quad (68)$$

For the quartic potential we have $Q(v) = v^4$, and the usual results obtain:

$$\text{Eq.(66): } 0 = -M_H^2 + \lambda v^2, \quad \text{Eq.(68): } m_h^2 = -M_H^2 + 3\lambda v^2. \quad (69)$$

This gives the usual relationship, $m_h^2 = 2M_H^2 = 2\lambda v^2$, however, this is modified by the presence of the additional terms in $Q(v)$.

If we perform the super-dilatation on the Yukawa and potential terms of eq.(63) we see that the previous relationship of eq.(22) is now modified:

$$0 = \frac{\epsilon}{\Lambda^2} (\bar{\psi}_L \theta_R + \bar{\theta}_L t_R) \cdot H \left(-\frac{1}{2} g^2 H^\dagger H (P(H^\dagger H))^2 + \frac{\lambda}{2} \frac{\partial Q(H^\dagger H)}{\partial (H^\dagger H)} \right) + h.c. \quad (70)$$

Thus, to maintain the super-dilatation symmetry in the full lagrangian for all values of Higgs fields we must demand the symmetry condition on the operator towers:

$$\frac{1}{2} g^2 v^2 (P(v^2))^2 = \frac{\lambda}{2} \frac{\partial Q(v^2)}{\partial v^2} \quad (71)$$

At the potential minimum, $v = v_{weak}$ we also have from eq.(66):

$$m_t^2 = g^2 v_{weak}^2 (P(v_{weak}^2))^2 = \lambda \frac{\partial Q(v^2)}{\partial v^2} \Big|_{v=v_{weak}} = 2M_H^2. \quad (72)$$

hence, even in the presence of the tower of operators we get the result $m_t^2 = 2M_H^2$. But we emphasize that this is the *curvature at the origin of the potential*, and does not give the physical Higgs mass, which is *the curvature at the minimum*. For the quartic potential this condition implies $m_h^2 = 2M_H^2 = m_t^2$, but in general, $2M_H^2 \neq m_h^2$, by virtue of the higher dimension terms.

To estimate the size of the effect, consider:

$$\frac{\lambda}{2} Q(v^2) = \frac{\lambda}{2} \left(v^4 + \sigma \kappa \frac{v^6}{\Lambda^2} \right) \quad (73)$$

where the higher dimension operator is assumed to have the strong coupling κ . The minimum $v = v_{weak}$ and physical Higgs mass now satisfy from eqs.(66,68):

$$M_H^2 = \lambda v_{weak}^2 + \sigma \kappa \lambda \frac{3v_{weak}^4}{2\Lambda^2}, \quad m_h^2 = -M_H^2 + 3\lambda v_{weak}^2 + \sigma \kappa \lambda \frac{15v_{weak}^4}{2\Lambda^2}. \quad (74)$$

Hence:

$$m_t^2 = 2M_H^2 = 2\lambda v_{weak}^2 + 3\sigma \kappa \lambda \frac{v_{weak}^4}{\Lambda^2}, \quad m_h^2 = 2\lambda v_{weak}^2 + 6\sigma \kappa \lambda \frac{v_{weak}^4}{\Lambda^2}, \quad (75)$$

and we thus predict:

$$m_h^2 \approx m_t^2 + \frac{3}{2} \sigma \kappa m_t^2 \frac{v_{weak}^2}{\Lambda^2} \quad (76)$$

Thus, we determine σ by demanding $m_h^2 \approx 0.5m_t^2$,

$$\sigma \approx -\frac{1}{3} \frac{\Lambda^2}{\kappa v_{weak}^2} \approx -6.5/\kappa \approx -0.04\kappa_c/\kappa \quad (77)$$

This is a relatively small coefficient compared to $\mathcal{O}(1)\kappa_c$. It's effect is helped by the strong coupling constant κ .

Of course, the Higgs potential must be stabilized, which can be done by inclusion of still higher dimension operators. This result is a sketch to demonstrate that an phenomenologically acceptable Higgs mass can emerge from the present scheme. The main strategy is to reduce the curvature of the potential at the minimum relative to that at the origin.

D. Weakly Coupled Theory

We can also raise Λ and allow $\kappa \sim g^2$ to be perturbatively small. Eq.(21) then requires $\Lambda \sim M_H$, and thus the Higgs mass floats up to a large scale.

To have a small Higgs boson mass term, we can then break the super-dilatation invariance by adding an explicit symmetry breaking Higgs mass term. This leads to a situation similar to the fine-tuning of the mass of the lightest Higgs boson in MSSM. We imagine $\kappa \sim g^2 \sim 1$ and choose $\Lambda \sim M_H$ large. We then add a symmetry breaking term of order $(-M_H^2 + \delta M_H^2)H^\dagger H$ to the lagrangian. Thus we are fine-tuning a small Higgs boson mass to a level of $\sim \delta M_H^2/M_H^2$ by demanding a cancellation between a large Higgs boson mass that respects the symmetry and another that breaks it. If we take $\Lambda \sim 10^4$ TeV, then our above RG estimate of eq.(62) yields the phenomenologically acceptable result $m_h^2 \approx m_t^2/2$.

This is more naturally framed in the minimal version of the symmetry, eq.(58). We are essentially violating the δt_R transformation (corresponding to $\eta\epsilon$ while retaining the ϵ transformation. That is, the “most minimal” transformation is:

$$\begin{aligned}\delta\psi_L^{ia} &= -i\frac{\not{\partial} H^i\theta_R^a}{\Lambda^2}\epsilon; & \delta\bar{\psi}_{L\,ia} &= i\frac{\bar{\theta}_{Ra}\not{\partial} H_i^\dagger}{\Lambda^2}\epsilon; \\ \delta H^i &= \frac{\bar{\theta}_{Ra}\psi_L^{ia}}{\Lambda^2}\epsilon; & \delta H_i^\dagger &= \frac{\bar{\psi}_{Lai}\theta_R^a}{\Lambda^2}\epsilon.\end{aligned}\tag{78}$$

The effective lagrangian we now obtain is:

$$\begin{aligned}\mathcal{L}_H &= \bar{\psi}_L i\not{\partial} \psi_L + i\bar{t}_R \not{\partial} t_R + \partial H^\dagger \partial H \\ &+ g(\bar{\psi}_L t_R H + h.c.) - M_H^2 H^\dagger H - \frac{\lambda}{2}(H^\dagger H)^2 \\ &+ \frac{\kappa}{\Lambda^2}(\bar{\psi}_L t_R \bar{t}_R \psi_L) - \frac{\kappa}{\Lambda^2}(\bar{t}_R \gamma_\mu t_R)(H^\dagger i \overset{\leftrightarrow}{\partial}^\mu H)\end{aligned}\tag{79}$$

The symmetry is broken, but we retain only the most interesting relationship eq.(23).

V. CONCLUSIONS

In summary, we've shown that the following lagrangian is invariant under the transformation of eq.(7) up to total divergences to order $1/\Lambda^4$:

$$\begin{aligned}
\mathcal{L}_H = & \bar{\psi}_L i \not{\partial} \psi_L + i \bar{t}_R \not{\partial} t_R + \partial H^\dagger \partial H \\
& + g(\bar{\psi}_L t_R H + h.c.) - M_H^2 H^\dagger H - \frac{\lambda}{2} (H^\dagger H)^2 \\
& + \frac{\kappa}{\Lambda^2} (\bar{\psi}_L t_R \bar{t}_R \psi_L) + \frac{2\kappa}{\Lambda^2} (\bar{\psi}_L \gamma_\mu \frac{\tau^A}{2} \psi_L) (H^\dagger i \overleftrightarrow{\partial}^\mu \frac{\tau^A}{2} H) \\
& + \frac{\kappa}{2\Lambda^2} (\bar{\psi}_L \gamma_\mu \psi_L) (H^\dagger i \overleftrightarrow{\partial}^\mu H) - \frac{\kappa}{\Lambda^2} (\bar{t}_R \gamma_\mu t_R) (H^\dagger i \overleftrightarrow{\partial}^\mu H)
\end{aligned} \tag{80}$$

where:

$$\lambda = \frac{1}{2} g^2 \quad \text{and} \quad g^2 \Lambda^2 = -\kappa M_H^2. \tag{81}$$

The super-dilatation symmetry is a bashing together of supersymmetry and dilatation or reparameterization invariance. It involves a super-rotation, or “twist,” amongst the Higgs and top quark fields, essentially forcing the Higgs to become a superpartner $\psi_L = (t, b)_L$. The leading $D = 4$ terms include the invariant gaugeless kinetic terms of Higgs, and top doublet and singlet fields, and the Higgs potential and Higgs-Yukawa interaction.

Our key result is that the top Yukawa coupling constant can be directly related to the Higgs quartic coupling constant by the super-dilatation symmetry, yielding the unrenormalized relationship in the broken phase $m_h = m_t$ (note $m_h^2 = 2|M_H^2|$). We've taken pains to show the consistency of this and the symmetry with the Nambu-Goldstone theorem for fermions. We have also explicitly demonstrated the symmetry in the broken phase where the fermionic shift “turns off.” The broken phase is a simpler case to analyze. Though equations of motion are deployed in interaction terms, this is a bona fide symmetry as it implies the nontrivial relationship $\lambda = g^2/2$ or $m_t = m_h$.

With radiative and/or small higher dimension operator effects, we can in principle obtain the observed $m_h^2 \approx 0.5 m_t^2$. However, we feel that exploring generalizations of the standard model with this kind of symmetry may lead to a tree level relationship, $m_h^2 = m_t^2/2$. A remarkable aspect of this is that the super-dilatation symmetry is already seen to be present in the existing (gaugeless) kinetic terms of the top-Higgs system. This may be a hint of the onset of a larger extended supersymmetry in a non-standard form.

The super-dilatation symmetry seems also to reflect a Nambu-Jona-Lasinio-like structure

at Λ which may be a natural harbinger of a composite Higgs boson, composed of $t\bar{t}$, but with a new dynamics, different than the conventional NJL setting [2].

VI. ACKNOWLEDGEMENTS

I thank Bill Bardeen, Roni Harnik, Adam Martin, Graham Ross and Geban Stavenga for discussions.

Appendix A: Dilatonic Higgs

An idea that is currently receiving revived attention is that the Higgs boson is an approximate dilaton, or “pseudo-dilaton” (see, *e.g.*, [7] and references therein). There is, of course, a fundamental distinction between a “scale invariant Higgs field” and a “dilatonic Higgs field.” A scale invariant Higgs field has a vanishing mass term, but can have a nonvanishing gauge, quartic and Yukawa couplings. To qualify as a (pseudo) dilatonic Higgs boson, the Higgs potential must be (approximately) flat.

Consider the pure Higgs lagrangian (no gauge fields or couplings to fermions):

$$\mathcal{L}_0 = \partial_\mu H^\dagger \partial^\mu H - \frac{\lambda}{2} (H^\dagger H - v^2)^2 \quad (82)$$

As usual, the groundstate has a minimum for:

$$\langle H^i \rangle = \theta^i \quad \text{where} \quad \theta^i = (v, 0) \quad (83)$$

(of course, θ^i can be rotated under $SU(2)_L \times U(1)$ and has an arbitrary orientation, but $\theta^\dagger \theta = v^2$). We can now take the limit $\lambda \rightarrow 0$, corresponding to a flat Higgs potential. The Higgs potential has served the role of an “applied external magnetic field to a spin system,” and aligns the spins. In the present case it pulls H^i to the minimum VEV. But once we take the limit $\lambda \rightarrow 0$ the lagrangian acquires a “shift symmetry,”

$$\delta H^i = \theta^i \epsilon \quad \longrightarrow \quad \delta \partial_\mu H^\dagger \partial^\mu H = 0 \quad (84)$$

Note that the alignment θ^i is held fixed (and remains arbitrary) and the shift is parameterized by the variable ϵ . The Noether current is:

$$J_\mu = \frac{\delta \mathcal{L}_0}{\delta \partial^\mu \epsilon} = \theta^\dagger \partial_\mu H + H^\dagger \partial_\mu \theta \quad (85)$$

We see that θ is a defining part of the current. If we view θ as co-rotating with H under the global $SU(2) \times U(1)$ transformations, the charge $\int d^3x J_0$ commutes with the gauge group.

In the broken phase of the theory we consider the small fluctuations around the Higgs VEV, and we can extract the broken phase Higgs field as:

$$\frac{1}{2v} (\theta^\dagger H + H^\dagger \theta) = v + \frac{h}{\sqrt{2}} \quad (86)$$

In the broken phase of the theory we thus see that:

$$J_\mu \rightarrow \sqrt{2} v \partial_\mu h \quad (87)$$

This becomes the “dilaton current” of the standard model Higgs boson in the broken phase. This is the only possible source of the dilaton current, which has a single derivative, as the usual scale current, $S_\mu = x^\nu T_{\mu\nu}$, with improved stress tensor $T_{\mu\nu}$, is bilinear in derivatives. The Higgs h in the limit $\lambda \rightarrow 0$ is now a Nambu-Goldstone boson of spontaneous scale symmetry breaking, inherited from the shift symmetry of the flat potential, *i.e.*, h has become a “dilaton.”

The dilatonic nature of the Higgs implies that fields that acquire masses $\propto v$ are scale invariant in the sense of spontaneous scale breaking. That is, we can perform a scale transformation which would shift mass terms, but we can then undo this rescaling by a compensating shift in h . To see this, consider the top quark mass term:

$$g\overline{\psi}_L t_R H + h.c. \longrightarrow m_t \bar{t} t \left(1 + \frac{h}{\sqrt{2}v}\right) \quad (88)$$

Under an infinitesimal scale transformation we have $t(x) \rightarrow (1 - \epsilon)^{3/2} t(x')$ and $h(x) \rightarrow (1 - \epsilon) h(x')$ where $x_\mu = (1 + \epsilon) x'_\mu$, $d^4 x = (1 + \epsilon)^4 d^4 x'$. Hence the action transforms as:

$$\begin{aligned} S_0 &= \int d^4 x \, m_t \bar{t} t(x) \left(1 + \frac{h(x)}{\sqrt{2}v}\right) \\ &\rightarrow \int d^4 x' \left((1 + \epsilon) m_t \bar{t} t(x') + m_t \bar{t} t(x') \frac{h(x')}{\sqrt{2}v} \right) \end{aligned} \quad (89)$$

The latter expression exhibits the fact that the $d = 4$ Higgs-Yukawa interaction is scale invariant while the $d = 3$ mass term would scales as $m \bar{t} t \rightarrow (1 - \epsilon) m \bar{t} t$.

However, with the dilatonic shift symmetry we can compensate the rescaled mass term by a shift in the Higgs-dilaton field. That is, under:

$$h(x') \rightarrow h(x') - \sqrt{2}v\epsilon \quad (90)$$

we see that:

$$\int d^4 x' \left((1 + \epsilon) m_t \bar{t} t(x') + m_t \bar{t} t(x') \frac{h(x')}{\sqrt{2}v} \right) \rightarrow \int d^4 x' \left(m_t \bar{t} t(x') + m_t \bar{t} t(x') \frac{h(x')}{\sqrt{2}v} \right) = S_0 \quad (91)$$

Hence, the simultaneous application of the conformal transformation and Higgs shift symmetry allows us to maintain the symmetry of the top quark mass term; the scale symmetry is broken spontaneously with the Higgs boson playing the role of the Nambu-Goldstone mode. The same invariance applies to the masses of all fermions, and of the gauge fields, W and Z . The Higgs self-interactions that involve nonzero λ would not be invariant under

scale transformations with dilatonic shifts in h . The shift symmetry is what generates the Nambu-Goldstone pole.

A consequence of the approximate dilatonic nature of the Higgs are low energy theorems for Higgs boson couplings to gauge and fermion fields. The effective lagrangian for the masses of the W -boson, Z -boson arises from the Higgs kinetic term,

$$\mathcal{L} = (D_\mu H)^\dagger (D^\mu H) \longrightarrow M_W^2 W_\mu^+ W_\mu^- + \frac{1}{2} M_Z^2 Z_\mu Z^\mu \quad (92)$$

where the Standard Model coupling constants, g_1 and g_2 , imply:

$$M_W^2 = \frac{1}{2} g_2^2 v^2 \quad M_Z^2 = \frac{1}{2} (g_1^2 + g_2^2) v^2 \quad (93)$$

The Higgs tree-level couplings are obtained simply by shifting v by the dynamical Higgs field, h :

$$v \rightarrow v + \frac{h}{\sqrt{2}} \quad (94)$$

to obtain:

$$\delta \mathcal{L} = \frac{h}{\sqrt{2}v} (2M_W^2 W_\mu^+ W_\mu^- + M_Z^2 Z_\mu Z^\mu) \quad (95)$$

These are the leading tree-level couplings of the Higgs field to the gauge bosons and are measured at the LHC in concordance with the prediction of a standard model Higgs boson.

The scale anomalies (running coupling constants) break the scale symmetry. However, this leads to a collection of low energy theorems for the Higgs coupled to gauge bosons, such as $h \rightarrow gg$ and $h \rightarrow \gamma\gamma$; γZ , etc.

Note that the $\lambda \rightarrow 0$ limit is the opposite of a nonlinear σ -model limit in which $\lambda \rightarrow \infty$ as v is held fixed. This latter case leads to the Nambu-Goldstone bosons of electroweak symmetry breaking, but an infinitely massive Higgs boson. In the case of a dilatonic Higgs, we package both the standard model $SU(2) \times U(1)$ symmetry together with the scale invariance and both are spontaneously broken.

The Higgs boson is then a (pseudo) dilaton if the shift transformation is a (approximate) symmetry of the action. Fundamentally it stems from the exact shift or modular symmetry of the gaugeless Higgs kinetic term:

$$\delta H^i = \theta^i \epsilon \quad \longrightarrow \quad \delta \partial_\mu H^\dagger \partial^\mu H = 0 \quad (96)$$

A key point we wish to emphasize here is that ϵ is the infinitesimal parameter of the transformation, while θ^i is held fixed. The spurion θ^i defines an arbitrary direction in isospin

space, a “ray,” and the shift moves the field along this direction in field space. We take θ^i to have the same mass dimension as the Higgs, *i.e.*, dimensions of mass. Here i refers to isospin and θ^i is a normalized isospinor spurion, $\theta^\dagger\theta = v^2$, where we, after-the fact, choose the alignment $\theta^i = (v, 0)$. Eq.(96) is a symmetry of the gaugeless Higgs boson kinetic terms, $\partial H^\dagger\partial H$. In such a theory the shift symmetry is exact.

Our present scheme is essentially a “super” generalization of this shift symmetry of eq.(96), involving shifts and rotations together with the top quark doublet and singlet fields. For us, the shift in the Higgs field becomes an operator, $\delta H \propto \bar{\psi}_L\theta_R + \bar{\theta}_Lt_R$, and fermions shift as well, $\delta\psi_L \propto \theta_L$ and $\delta t_R \propto \theta_R$. In addition we have the super-rotation, hence we dub this a “super-dilataion.”

Appendix B: Analogy to a Chiral Lagrangian

To orient the reader to our present strategy we quickly review a familiar derivation of a toy π chiral lagrangian from the “bottoms-up.” Consider the kinetic terms:

$$\mathcal{L}_K = \bar{\psi}_L i \not{\partial} \psi_L + \bar{\psi}_R i \not{\partial} \psi_R + \frac{1}{2} \partial_\mu \pi \partial^\mu \pi \quad (97)$$

We’ll consider the RH-chiral symmetry:

$$\begin{aligned} \delta \psi_L &= 0 \\ \delta \psi_R &= i\theta \psi_R & \delta \bar{\psi}_R &= -i\theta \bar{\psi}_R \\ \delta \pi &= f_\pi \theta \end{aligned} \quad (98)$$

and we demand the lagrangian is invariant under this global transformation:

$$\delta \mathcal{L}_K = 0 \quad (99)$$

The RH-chiral current is:

$$-\frac{\delta \mathcal{L}_K}{\delta \partial^\mu \theta} = \bar{\psi}_R \gamma_\mu \psi_R - f_\pi \partial_\mu \pi \quad (100)$$

and we assume f_π is “determined from experiment,” *e.g.*, $\pi \rightarrow \mu\nu$ (of course, in the real world this is the left-handed current).

Consider the interactions consisting of a massive “nucleon” coupled to pion:

$$\mathcal{L}_V = M \bar{\psi} \psi - ig \pi \bar{\psi} \gamma^5 \psi = M \bar{\psi}_L \psi_R - ig \pi \bar{\psi}_L \psi_R + h.c. \quad (101)$$

We perform the RH-chiral transformation transformation:

$$\delta \mathcal{L}_V = (i\theta M - ig f_\pi \theta + g \pi f_\pi \theta) \bar{\psi}_L \psi_R + h.c. \quad (102)$$

so:

$$\delta \mathcal{L}_V = 0 \quad \longrightarrow \quad g = \frac{M}{f_\pi} \quad (103)$$

which is the Goldberger-Treiman relation.

However, we still must cancel the “higher order term” $\propto \pi \theta \bar{\psi}_L \psi_R$. We thus include an $\mathcal{O}(\pi^2)$ term:

$$\mathcal{L}_V \rightarrow M \left(1 - \frac{i\pi}{f_\pi} + c \frac{\pi^2}{f_\pi^2} \right) \bar{\psi}_L \psi_R + h.c. \quad (104)$$

Now:

$$\delta\mathcal{L}_V \rightarrow M \left(i\theta - i\frac{f_\pi\theta}{f_\pi} + \frac{\pi\theta}{f_\pi} + 2c\frac{\pi}{f_\pi^2}f_\pi\theta + ic\frac{\pi^2}{f_\pi^2}f_\pi\theta \right) \bar{\psi}_L\psi_R h.c. \quad (105)$$

so:

$$\delta\mathcal{L}_V = 0 \quad \longrightarrow \quad g = \frac{M}{f_\pi}, \quad c = -\frac{1}{2} \quad (106)$$

But, now we must cancel higher order term $\propto \pi^2\theta\bar{\psi}_L\psi_R$ which implies an $\mathcal{O}(\pi^3)$ interaction, and so-forth.

We can sum the resulting power series and we find, iteratively, the solution:

$$\mathcal{L}_V = M\bar{\psi}_L U\psi_R + h.c. \quad U = \exp(i\pi/f_\pi) \quad (107)$$

whence,

$$\mathcal{L} = \bar{\psi}_L i\not{\partial}\psi_L + \bar{\psi}_R i\not{\partial}\psi_R + \frac{f_\pi^2}{2}\partial_\mu U^\dagger\partial^\mu U + M\bar{\psi}_L U\psi_R + h.c. \quad (108)$$

and we have obtained the “nonlinear σ -model lagrangian.”

Our present strategy is similar. We begin with the super-dilatational invariance of the top-Higgs kinetic terms. We then analyze the transformation of the Higgs-Yukawa, Higgs mass and quartic interactions. We demand overall invariance of the lagrangian. We thus find the “Goldberger-Treiman” relationship, $\lambda = g^2/2$, which implies $m_t = m_h$ in the broken phase. This induces higher dimension operators. Ultimately, we expect to sum the tower of operators, though in the present case we expect that these arise via new dynamics, such as heavy recurrences of composite Higgs bosons and vector-like top quarks.

-
- [1] V. A. Miransky, M. Tanabashi, and K. Yamawaki, Phys. Lett. **B221**, 177 (1989); Mod. Phys. Lett. **A4**, 1043 (1989);
W. J. Marciano, Phys. Rev. Lett. **62**, 2793 (1989);
Y. Nambu, Enrico Fermi Institute Report No. 89-08, 1989 (unpublished); in *Proceedings of the 1989 Workshop on Dynamical Symmetry Breaking*, edited by T. Muta and K. Yamawaki (Nagoya University, Nagoya, Japan, 1990).
- [2] W. A. Bardeen, C. T. Hill, and M. Lindner, Phys. Rev. **D41**, 1647 (1990).
- [3] C. T. Hill, Phys. Lett. B **266**, 419 (1991); C. T. Hill, Phys. Lett. B **345**, 483 (1995).
- [4] H. Georgi, Nucl. Phys. **B348**, 293 (1991); Phys. Lett. B **240**, 447 (1990);
M. Neubert, Phys. Rept. **245**, 259 (1994)
A. V. Manohar and M. B. Wise, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **10**, 1 (2000).
- [5] M. E. Luke and A. V. Manohar, Phys. Lett. B **286**, 348 (1992).
- [6] J. Heinonen, R. J. Hill and M. P. Solon, arXiv:1208.0601 [hep-ph].
- [7] See: S. Iso and Y. Orikasa, “TeV Scale B-L model with a flat Higgs potential at the Planck scale - in view of the hierarchy problem -,” arXiv:1210.2848 [hep-ph].
T. Abe, R. Kitano, Y. Konishi, K. -y. Oda, J. Sato and S. Sugiyama, “Minimal Dilaton Model,” arXiv:1209.4544 [hep-ph].
K. A. Meissner and H. Nicolai, Acta Phys. Polon. B **40**, 2737 (2009) [arXiv:0809.1338 [hep-th]].
and references therein.
- [8] J. Wess and J. Bagger, Princeton, USA: Univ. Pr. (1992) 259 p; see eqs(3.3-3.9).
- [9] For notational reference: The physical real Higgs boson field h , defined by v at the minimum of the Higgs potential, is given by $H = v + h/\sqrt{2}$, with mass $m_h^2 = 2\lambda v^2$. When the potential is written in the form $-M_H^2 H^\dagger H + (\lambda/2)(H^\dagger H)^2$ we have the correspondence $M_H^2 = \lambda v^2 = m_h^2/2$