

Higgs couplings to t, b , and τ with Flavor Symmetry

Paul H. Frampton*

Department of Physics and Astronomy, UNC-Chapel Hill, NC 27599.

Abstract

Experimental study at LHC of the possible Higgs boson should soon provide accurate estimates of the Yukawa couplings $Y_{H\bar{f}f}$ for $f = t, b, \tau$. In the presence of a non-abelian discrete flavor symmetry G_F (*e.g.* T') the usual prediction that $Y_{H\bar{f}f} \propto m_f$ should be inexact, and departures therefrom will provide through G_F valuable input to an improved derivation of the quark and lepton mixing matrices.

*frampton@physics.unc.edu

The 2012 discovery of a resonance at ~ 125 GeV at the LHC [1, 2], strongly suggestive of the Higgs boson, has naturally caused intense interest. Its preliminary properties are consistent within significant errors with the Higgs particle predicted by the minimal standard model.

The two-body decays which can be measured accurately in the near future include $H \rightarrow \gamma\gamma$, $H \rightarrow \bar{b}b$, and $H \rightarrow \bar{\tau}\tau$. These are respectively sensitive to the Yukawa couplings $Y_{H\bar{t}t}$ (through the top triangle contribution which competes with the W -loop), $Y_{H\bar{b}b}$, and $Y_{H\bar{\tau}\tau}$.

In the minimal standard model, the Yukawa couplings $Y_{H\bar{f}f}$ appear in the simple form

$$Y_{H\bar{f}f}\bar{f}fH \quad (1)$$

so that they are proportional to the masses

$$Y_{H\bar{f}f} \propto m_f \quad (2)$$

with proportionality constant $\langle H \rangle^{-1}$ where $\langle H \rangle$ is the vacuum expectation value of the Higgs field.

In a renormalizable model with a non-trivial flavor symmetry G_F , which we will take here to be non-abelian and discrete, there must be several Higgs and the Yukawa couplings of the lightest one H will generally deviate from the simple proportionality of Eq.(2). Such deviations may likely be small but crucial to understanding how the group G_F operates. One may even say that if the conventional prediction of Eq. (2) would hold exactly at high precision then renormalizable G_F models would be disfavored.

These statements are true for general renormalizable G_F models. To illustrate them, we focus on the choice $G_F = T'$ [3] and the minimal model discussed in [4]. The flavor group is $(T' \times Z_2)$, and we shall concentrate only on the third-generation couplings $Y_{H\bar{f}f}$ for $f = t, b, \tau$.

The leptons are assigned under $(T' \times Z_2)$ as

$$\left. \begin{array}{l} \left(\begin{array}{c} \nu_\tau \\ \tau^- \end{array} \right)_L \\ \left(\begin{array}{c} \nu_\mu \\ \mu^- \end{array} \right)_L \\ \left(\begin{array}{c} \nu_e \\ e^- \end{array} \right)_L \end{array} \right\} L_L(3, +1) \quad \begin{array}{ll} \tau_R^- (1_1, -1) & N_R^{(1)} (1_1, +1) \\ \mu_R^- (1_2, -1) & N_R^{(2)} (1_2, +1) \\ e_R^- (1_3, -1) & N_R^{(3)} (1_3, +1). \end{array} \quad (3)$$

Imposing strict renormalizability on the lepton lagrangian allows as nontrivial terms for the τ mass only

$$Y_\tau (L_L \tau_R H'_3) + \text{h.c.} \quad (4)$$

where H'_3 transforms as $H'_3(3, -1)$.

The left-handed quark doublets $(t, b)_L, (c, d)_L, (u, s)_L$ are assigned under $(T' \times Z_2)$ to

$$\left. \begin{array}{l} \begin{pmatrix} t \\ b \end{pmatrix}_L \\ \begin{pmatrix} c \\ s \end{pmatrix}_L \\ \begin{pmatrix} u \\ d \end{pmatrix}_L \end{array} \right\} \mathcal{Q}_L \quad \begin{array}{l} (\mathbf{1}_1, +1) \\ (\mathbf{2}_1, +1) \end{array} \quad (5)$$

and the six right-handed quarks as

$$\left. \begin{array}{l} t_R \\ b_R \\ c_R \\ u_R \end{array} \right\} \mathcal{C}_R \quad \begin{array}{l} (\mathbf{1}_1, +1) \\ (\mathbf{1}_2, +1) \\ (\mathbf{2}_3, -1) \end{array} \quad (6)$$

$$\left. \begin{array}{l} s_R \\ d_R \end{array} \right\} \mathcal{S}_R \quad (\mathbf{2}_2, +1)$$

We must two new scalars $H_{1_1}(1_1, +1)$ and $H_{1_3}(1_3, +1)$ whose VEVs

$$\langle H_{1_1} \rangle = m_t/Y_t \quad \langle H_{1_3} \rangle = m_b/Y_b \quad (7)$$

provide the (t, b) masses. In particular, no T' doublet $(2_1, 2_2, 2_3)$ scalars have been added. This allows a non-zero value only for Θ_{12} . The other angles vanish making the third family stable

The Yukawa couplings to the third family of quarks are contained in

$$\begin{aligned} \mathcal{L}_Y^{(quarks)} = & Y_t(\{\mathcal{Q}_L\}_{1_1}\{t_R\}_{1_1}H_{1_1}) \\ & + Y_b(\{\mathcal{Q}_L\}_{1_1}\{b_R\}_{1_2}H_{1_3}) \\ & + \text{h.c.} \end{aligned} \quad (8)$$

The use of T' singlets and doublets permits the third family to differ from the first two and thus make plausible the mass hierarchies $m_t \gg m_b$, $m_b > m_{c,u}$ and $m_b > m_{s,d}$ as outlined in [3].

Such a model leads to the formula [4] for the Cabibbo angle

$$\tan 2\Theta_{12} = \left(\frac{\sqrt{2}}{3} \right) \quad (9)$$

or equivalently $\sin \Theta_{12} = 0.218..$ close to the experimental value $\sin \Theta_{12} \simeq 0.227$.

It can also lead to the successful relationship between neutrino mixing angles θ_{ij}

$$\theta_{13} = (\sqrt{2})^{-1} \left| \frac{\pi}{4} - \theta_{23} \right| \quad (10)$$

which is also in excellent agreement with the latest experiments [5].

In such a model, the lightest Higgs H is a linear combination

$$H = aH_{1_1} + bH_{1_3} + cH'_{3'} + \dots \quad (11)$$

and the consequent Yukawa couplings are

$$Y_{H\bar{t}t} = a^{-1}Y_t, \quad Y_{H\bar{b}b} = b^{-1}Y_b, \quad Y_{H\bar{\tau}\tau} = c^{-1}Y_\tau \quad (12)$$

The VEV $\langle H \rangle$ is shared between the $\langle H_\alpha \rangle$ ($\alpha = 1_1, 1_3, 3', \dots$) irreps of T' and there is no reason to expect $a = b = c = \dots = 1$ so that the proportionality of Eq.(2) will be lost. In fact, if Eq.(2) remained exact, the only solution would be a trivial one where all states transform as 1_1 of T' and the G_F is inapplicable. The successes in [4] and [5] would, in such a case, be accidental.

On the other hand, if Eq.(2) is inexact, the evaluations of the coefficients a, b, c, \dots can then be used to understand more perspicuously the derivations of mixing angles for quarks and leptons given respectively in [4] and [5], in a first clear departure from the minimal standard model.

Acknowledgements

This work should have been supported by U.S. Department of Energy grant number DE-FG02-06ER41418.

References

- [1] J. Incalada, talk given at CERN on July 4, 2012. CMS Collaboration. CMS-PAS-HIG-12-020.
- [2] F. Gionetti, talk given at CERN on July 4, 2012. ATLAS Collaboration. ATLAS-CONF-2012-093.
- [3] P.H. Frampton and T.W. Kephart, Int. J. Math. Phys. **A10**, 4689 (1995). hep-ph/9409330.
- [4] P.H. Frampton, T.W. Kephart, and S. Matsuzaki, Phys. Rev. **D78**, 073004 (2008). arXiv:0807.4713 [hep-ph]
- [5] D.A.Eby and P.H. Frampton, Phys. Rev. **D** (in press). arXiv:1112.2675 [hep-ph]