

Discrete symmetries, roots of unity, and lepton mixing

W. Grimus*

University of Vienna, Faculty of Physics
Boltzmanngasse 5, A-1090 Vienna, Austria

August 13, 2014

Abstract

We investigate the possibility that the first column of the lepton mixing matrix U is given by $u_1 = (2, -1, -1)^T/\sqrt{6}$. In a purely group-theoretical approach, based on residual symmetries in the charged-lepton and neutrino sectors and on a theorem on vanishing sums of roots of unity, we discuss the *finite* groups which can enforce this. Assuming that there is only one residual symmetry in the Majorana neutrino mass matrix, we find the almost *unique* solution $\mathbb{Z}_q \times S_4$ where the cyclic factor \mathbb{Z}_q with $q = 1, 2, 3, \dots$ is irrelevant for obtaining u_1 in U . Our discussion also provides a natural mechanism for achieving this goal. Finally, barring vacuum alignment, we realize this mechanism in a class of renormalizable models.

*E-mail: walter.grimus@univie.ac.at

1 Introduction

The recent measurements of a rather large reactor mixing angle θ_{13} [1, 2] disfavour tri-bimaximal mixing [3] and, therefore, also such models—see [4, 5] for reviews on models for neutrino masses and mixing. While the third column of the tri-bimaximal mixing matrix U_{TBM} is now definitively in disagreement with the data, the first or the second column of U_{TBM} could still occur in the mixing matrix U . These cases are denoted by TM_1 and TM_2 , respectively, in [6]. However, in the case of TM_2 the solar mixing angle θ_{12} is related to the reactor mixing angle via $\sin^2 \theta_{12}(1 - \sin^2 \theta_{13}) = 1/3$, which creates a tension with the data but is still compatible at the 3σ level. Therefore, it is more interesting to consider TM_1 [7, 8, 9, 10] where the first column in $U = (u_1, u_2, u_3)$ is given by

$$u_1 = \frac{1}{\sqrt{6}} \begin{pmatrix} 2 \\ -1 \\ -1 \end{pmatrix}. \quad (1)$$

Recently, a purely group-theoretical approach has been developed for the investigation of the effect of *finite* family symmetry groups on the mixing matrix [11, 12, 13, 14, 15, 16, 17]. Apart from assuming that the left-handed neutrino fields and the left-handed charged lepton fields are in the same gauge doublet of the Standard Model gauge group, no other assumption concerning the interactions in the lepton sector is made. On the one hand, taking into account extant data on lepton mixing, such a general approach allows a systematic investigation of the possible symmetry groups, see for instance the scan of groups performed in [18, 19]. On the other hand, this approach has its limitations since it is not entirely clear how its results relate to concrete models [20]; we will address this point later in this paper.

The goal of the present investigation is to find all possible finite family symmetry groups underlying TM_1 , without restricting the other two columns u_2 and u_3 beyond orthonormality. In order to accomplish this task, we will use a theorem on vanishing sums of roots of unity. In the course of this investigation we will also come across a mechanism for the implementation of TM_1 . In the following we will use this mechanism to write down a class of models based on S_4 and the type II seesaw mechanism where TM_1 is realized.

The paper is organized as follows. In section 2 the group-theoretical method of [15] is reviewed using our notation. In order to be as clear as possible, some arguments are emphasized by formulating them as propositions. This also applies to section 3 where the symmetry group for TM_1 is determined; some technical points needed in the course of our argumentation are deferred to appendix A. In the same section we also find a mechanism for the implementation of TM_1 , which is then used in section 4 for the construction of a class of models. The summary of our findings is presented in section 5. As a supporting material, we provide a set of generators of S_4 and the three-dimensional irreducible representations of this group in appendix B.

2 Residual symmetries in the mass matrices

The class of models we have in mind as an application of the following discussion are extensions of the Standard Model in the scalar and fermion sectors. Typical examples would be several Higgs doublets and right-handed neutrino singlets which facilitate the seesaw mechanism [21], or Higgs triplet extensions with the type II seesaw mechanism [22]. We further assume that before spontaneous symmetry breaking (SSB) the theory is invariant under a *finite* family symmetry group G and that there are three lepton families. We want to investigate the case that there are residual symmetries in the charged-lepton and Majorana neutrino mass matrices, left over from SSB of G , and study their effect on the lepton mixing matrix U [11, 15].

Such residual symmetries will occur whenever the vacuum expectation values (VEVs) of the neutral components of scalar gauge multiplets are invariant under some transformations of G . However, we are not interested in the full symmetry of the vacuum, which might very well be trivial, but in the symmetries of the vacua in the respective sectors whose VEVs lead to charged-lepton and neutrino masses. It is well known that the mismatch of these symmetries is responsible for predictions in the mixing matrix—see for instance [4, 23]. Therefore, we must distinguish between the symmetries in $M_\ell M_\ell^\dagger$ where M_ℓ the charged-lepton mass matrix and those in the Majorana neutrino mass matrix \mathcal{M}_ν . These symmetries are supposed to generate the subgroups G_ℓ and G_ν of G , pertaining to $M_\ell M_\ell^\dagger$ and \mathcal{M}_ν , respectively. Obviously, we have the relation

$$G_\ell \subseteq U(1) \times U(1) \times U(1), \quad G_\nu \subseteq \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \quad (2)$$

due to the Dirac and Majorana natures of charged and neutral leptons, respectively. The groups G_ℓ and G_ν will be quite small and very often be generated by just one symmetry. In the following we will assume that this is the case for G_ν , but G_ℓ will in principle be allowed to contain several non-trivial elements though the analysis will be phrased in terms of a single matrix $T \in G_\ell$.

The mass Lagrangian—obtained through SSB—has the form

$$\mathcal{L}_{\text{mass}} = -\bar{\ell}_L M_\ell \ell_R + \frac{1}{2} \nu_L^T C^{-1} \mathcal{M}_\nu \nu_L + \text{H.c.} \quad (3)$$

with the residual symmetries

$$T^\dagger M_\ell M_\ell^\dagger T = M_\ell M_\ell^\dagger, \quad S^T \mathcal{M}_\nu S = \mathcal{M}_\nu, \quad (4)$$

where T and S are unitary matrices and \mathcal{M}_ν is symmetric but complex in general. Furthermore, the diagonalizing matrices U_ℓ and U_ν , and the lepton mixing matrix are given

$$U_\ell^\dagger M_\ell M_\ell^\dagger U_\ell = \text{diag} (m_e^2, m_\mu^2, m_\tau^2), \quad U_\nu^T \mathcal{M}_\nu U_\nu = \text{diag} (m_1, m_2, m_3), \quad U = U_\ell^\dagger U_\nu, \quad (5)$$

respectively. We denote the weak basis of M_ℓ and \mathcal{M}_ν by basis 1. Since we are interested in the effect of T and S on the lepton mixing matrix, the action of the residual symmetry on right-handed lepton fields is irrelevant for us, which is the reason to consider $M_\ell M_\ell^\dagger$ instead of M_ℓ .

It is useful to distinguish basis 1 from the weak basis, where the charged-lepton mass matrix is diagonal, in which case we use the phrase basis 2. All matrices in this basis are indicated by a tilde. Usually, basis 1 is the weak basis where the matrices of the representation $D(G)$ on the left-handed lepton gauge doublets have a “nice” form. Depending on the representations of G used in the Lagrangian it can happen that basis 1 coincides with basis 2, but in general the two bases will be different. Because all charged-lepton masses are different, we conclude that \tilde{T} is a diagonal matrix.

In the following analysis we will assume that there are no accidental symmetries in $M_\ell M_\ell^\dagger$ and \mathcal{M}_ν . Then, since S and T are given in a weak basis, they must both belong to $D(G)$, the group of representation matrices of G acting on the left-handed lepton doublets. The matrices S and T generate a group denoted by \tilde{G} . In this case it is clear that \tilde{G} is a subgroup of the group $D(G)$ and both are subgroups of $U(3)$, i.e. $\tilde{G} \subseteq D(G) \subset U(3)$. In the simplest case \tilde{G} is identical with $D(G)$.

Due to equation (2) and assuming $\det S = 1$ without loss of generality, the form of S is given by

$$S = 2uu^\dagger - \mathbb{1} \quad (6)$$

with a unit vector u and $S^2 = \mathbb{1}$.

Proposition 1 *If $S^T \mathcal{M}_\nu S = \mathcal{M}_\nu$ with $S = 2uu^\dagger - \mathbb{1}$, then $\mathcal{M}_\nu u \propto u^*$.*

Proof: By construction, the matrix S fulfills $Su = u$, with a unique eigenvalue 1. Therefore, $S^T \mathcal{M}_\nu u = \mathcal{M}_\nu u$, and due to the hermiticity of S and $S^2 = \mathbb{1}$ we find $S^*(\mathcal{M}_\nu u) = \mathcal{M}_\nu u$. Since S^* has a unique eigenvalue 1 with corresponding eigenvector u^* , we conclude that $\mathcal{M}_\nu u$ is proportional to u^* . \square

In basis 2 the mixing matrix U diagonalizes the neutrino mass matrix:

$$U^T \tilde{\mathcal{M}}_\nu U = \text{diag}(m_1, m_2, m_3). \quad (7)$$

With column vectors u_j and $U = (u_1, u_2, u_3)$, equation (7) is reformulated as

$$\tilde{\mathcal{M}}_\nu u_j = m_j u_j^*. \quad (8)$$

Comparing with proposition 1 and denoting the unit vector associated with \tilde{S} by \tilde{u} , we immediately come to the following conclusion.

Proposition 2 *If $\tilde{S}^T \tilde{\mathcal{M}}_\nu \tilde{S} = \tilde{\mathcal{M}}_\nu$, then, apart from irrelevant phases, \tilde{u} is one of column vectors of the lepton mixing matrix U .*

This proposition is the basis of all discussions concerning residual symmetries in the mass matrices.

Since we assume that G is a finite group, it is a necessary condition that the matrices T and ST have finite orders:

$$T^m = (ST)^n = \mathbb{1} \quad (9)$$

for some natural numbers m and n . Note that S has order two by construction.¹

¹Groups generated by S and T such that the orders of S , T and ST are finite are called von Dyck groups [15]; such groups are not necessarily finite because finite orders of group generators does in general not imply that the group is finite.

The interesting observation is that equation (9) connects group properties with information on the mixing matrix [11, 15]. In effect, one uses the following proposition.

Proposition 3 *Suppose we have a 3×3 matrix M which is a function of a set of parameters $x = (x_1, \dots, x_r)$ such that for all values of x M is unitary. Then the values of x where M has the eigenvalues λ_k ($k = 1, 2, 3$ and $|\lambda_k| = 1$) are determined by the two equations*

$$\text{Tr } M(x) = \lambda_1 + \lambda_2 + \lambda_3 \quad \text{and} \quad \det M(x) = \lambda_1 \lambda_2 \lambda_3. \quad (10)$$

Proof: The characteristic polynomial of M is

$$P_M(\lambda) = \lambda^3 - M_2 \lambda^2 + M_1 \lambda - M_0.$$

In terms of the eigenvalues of M the coefficients M_i are given by

$$M_1 = \lambda_1 \lambda_2 + \lambda_2 \lambda_3 + \lambda_3 \lambda_1, \quad M_2 = \lambda_1 + \lambda_2 + \lambda_3, \quad M_0 = \lambda_1 \lambda_2 \lambda_3.$$

Unitarity of M means that the eigenvalues are located on the unit circle. Therefore, we find $M_1 = M_2^* M_0$. This means that, as long as equation (10) is satisfied, then automatically M gives the correct coefficient M_2 in $P_M(\lambda)$ leading to the eigenvalues λ_k . \square

We make the identification $M = \tilde{S} \tilde{T}$. With

$$\tilde{T} = \text{diag} (e^{i\phi_e}, e^{i\phi_\mu}, e^{i\phi_\tau}) \quad \text{and} \quad \tilde{u} = \begin{pmatrix} U_{ei} \\ U_{\mu i} \\ U_{\tau i} \end{pmatrix} \quad (11)$$

equation (10) reads [15]

$$\sum_{\alpha=e,\mu,\tau} (2|U_{\alpha i}|^2 - 1) e^{i\phi_\alpha} = \lambda_1 + \lambda_2 + \lambda_3 \quad \text{and} \quad \prod_{\alpha} e^{i\phi_\alpha} = \lambda_1 \lambda_2 \lambda_3, \quad (12)$$

where the $e^{i\phi_\alpha}$ are m -th roots and the λ_k are n -th roots of unity. The parameters in M are the mixing angles, the CP phase and the ϕ_α . Equation (12) provides the necessary and sufficient conditions that ST is of order n .

Equation (12) can be used both ways: Given the group \bar{G} generated by S and T , relations among the mixing parameters are obtained [15]; vice versa, assuming a specific column vector in U , we can infer the group \bar{G} .

3 A symmetry for TM_1

Now we discuss the possible symmetry groups \bar{G} leading to TM_1 . From the discussion in the previous section we have learned that in this case we have to take $\tilde{u} \equiv u_1$ with u_1 given by equation (1), which leads to

$$\tilde{S} = \frac{1}{3} \begin{pmatrix} 1 & -2 & -2 \\ -2 & -2 & 1 \\ -2 & 1 & -2 \end{pmatrix}. \quad (13)$$

Specifying equation (12) to this case, we obtain

$$-e^{i\phi_e} + 2e^{i\phi_\mu} + 2e^{i\phi_\tau} + 3\lambda_1 + 3\lambda_2 + 3\lambda_3 = 0 \quad \text{and} \quad e^{i\phi_e} e^{i\phi_\mu} e^{i\phi_\tau} = \lambda_1 \lambda_2 \lambda_3. \quad (14)$$

Our aim is to find all cases of roots of unity $e^{i\phi_\alpha}$ ($\alpha = e, \mu, \tau$) and λ_k ($k = 1, 2, 3$) which satisfy these two relations. For this purpose we use a theorem by Conway and Jones, theorem 6 in [24].

Some remarks are in order before we reproduce this theorem. Formal sums of roots of unity with rational coefficients form a ring [24]. A sum of roots of unity \mathcal{S}' is *similar* to a sum of roots of unity \mathcal{S} if there is a rational number q and a root of unity δ such that $\mathcal{S}' = q\delta\mathcal{S}$. The length of a sum of roots of unity \mathcal{S} is the number of roots involved. Note, however, that, for the definition of this length, roots of unity which differ only by a sign do not count separately. The roots occurring in the following theorem are

$$\omega = e^{2\pi i/3}, \quad \beta = e^{2\pi i/5}, \quad \gamma = e^{2\pi i/7}. \quad (15)$$

Theorem 1 (Conway and Jones) *Let \mathcal{S} be a non-empty vanishing sum of length at most 9. Then either \mathcal{S} involves $\theta, \theta\omega, \theta\omega^2$ for some root θ , or \mathcal{S} is similar to one of*

- a) $1 + \beta + \beta^2 + \beta^3 + \beta^4$,
- b) $-\omega - \omega^2 + \beta + \beta^2 + \beta^3 + \beta^4$,
- c) $1 + \gamma + \gamma^2 + \gamma^3 + \gamma^4 + \gamma^5 + \gamma^6$,
- d) $1 + \beta + \beta^4 - (\omega + \omega^2)(\beta^2 + \beta^3)$,
- e) $-\omega - \omega^2 + \gamma + \gamma^2 + \gamma^3 + \gamma^4 + \gamma^5 + \gamma^6$,
- f) $\beta + \beta^4 - (\omega + \omega^2)(1 + \beta^2 + \beta^3)$,
- g) $1 + \gamma^2 + \gamma^3 + \gamma^4 + \gamma^5 - (\omega + \omega^2)(\gamma + \gamma^6)$,
- h) $1 - (\omega + \omega^2)(\beta + \beta^2 + \beta^3 + \beta^4)$.

The first relation of equation (14) constitutes a vanishing sum \mathcal{S} whose length is at most 6. It is immediately obvious that the vanishing sums in the theorem cannot be part of \mathcal{S} , because c)–h) are too long, sum b) is not similar to \mathcal{S} and sum a) is a vanishing sum of length 5 and adding one root would make a non-vanishing sum. Therefore, in the case of equation (14), the theorem says that \mathcal{S} must involve ω . Exploiting this fact in appendix A, we find that equation (14) has the solution

$$e^{i\phi_e} = \eta, \quad e^{i\phi_\mu} = \eta\omega, \quad e^{i\phi_\tau} = \eta\omega^2, \quad \lambda_1 = \epsilon, \quad \lambda_2 = -\epsilon, \quad \lambda_3 = \eta, \quad (16)$$

with arbitrary roots of unity ϵ and η . The second condition of equation (14) yields

$$\eta^3 = -\epsilon^2\eta \quad \text{or} \quad \epsilon = \pm i\eta. \quad (17)$$

However, this equation will be of no use in the following.

In summary, our considerations has lead us to

$$\tilde{T} = \eta \text{diag} (1, \omega, \omega^2). \quad (18)$$

Thus \tilde{T} is completely determined up to an unknown root of unity η . We remind the reader that the matrices \tilde{S} of equation (13) and \tilde{T} of equation (18) are given in basis 2.

A suitable basis 1—see appendix B—is given by a basis transformation with

$$U_\omega = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}. \quad (19)$$

The resulting matrices are

$$S = U_\omega \tilde{S} U_\omega^\dagger = S_1 B = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad \text{and} \quad T = U_\omega \tilde{T} U_\omega^\dagger = \eta E = \eta \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad (20)$$

where S_1 is defined in equation (B1) and E and B in equation (B2).

Let us assume that η is a primitive root of order q . The following proposition determines the group \bar{G} .

Proposition 4 *The matrix \tilde{S} of equation (13) and \tilde{T} of equation (18) generate the group $\bar{G} = \mathbb{Z}_q \times S_4$.*

Proof: For simplicity of notation we use basis 1. From $(ST)^2 = \eta^2 S_3$ and $T^\dagger \eta^2 S_3 T = \eta^2 S_1$ we find that $\eta^2 S_1 S = \eta^2 B \in \bar{G}$ and, therefore,

$$(\eta^2 B)^\dagger T (\eta^2 B) = \eta \text{diag} (1, \omega^2, \omega) \in \bar{G}.$$

Eventually, with

$$T(\eta^2 B)^\dagger T(\eta^2 B) = \eta^2 \mathbb{1} \in \bar{G} \quad \text{and} \quad T^3 = \eta^3 \mathbb{1} \in \bar{G}$$

we conclude $\eta \mathbb{1} \in \bar{G}$. Therefore, \bar{G} contains E , S_1 and B , which is a set of generators of S_4 —see appendix B. It is then almost trivial to show that every element of $g \in \bar{G}$ can uniquely be decomposed into $g = \eta^k h$ with $k \in \{0, 1, \dots, q-1\}$ and $h \in S_4$. \square

In summary, we have found *all* finite groups which enforce TM_1 in the mixing matrix because we have seen that any \tilde{T} of finite order satisfying equation (14) leads to S_4 times a cyclic factor. In [18] it was stated that S_4 is the smallest such group. Here we have shown that actually S_4 is unique up to a trivial factor with a cyclic group. We formulate our result as a theorem.

Theorem 2 *Under the premises that G_ν is a \mathbb{Z}_2 generated by \tilde{S} of equation (13) and that G_ℓ contains at least one matrix \tilde{T} which is not proportional to the unit matrix, the only symmetry groups \bar{G} generated by the residual symmetries of the mass matrices which are able to enforce TM_1 in the lepton mixing matrix U are $\mathbb{Z}_q \times S_4$.*

With our derivation of theorem 2 we have also demonstrated that

$$E^\dagger \left(M_\ell M_\ell^\dagger \right) E = M_\ell M_\ell^\dagger. \quad (21)$$

This has the following consequence.

Proposition 5 *For every charged-lepton mass matrix which fulfills equation (21) it follows that $U_\omega^\dagger \left(M_\ell M_\ell^\dagger \right) U_\omega$ is diagonal.*

Proof: The eigenvectors of E are identical with the column vectors of U_ω , i.e. $U_\omega = (x_1, x_2, x_3)$ and $E x_k = \omega^{k-1} x_k$. Therefore, we have

$$\left(M_\ell M_\ell^\dagger \right) E x_k = \omega^{k-1} \left(M_\ell M_\ell^\dagger \right) x_k = E \left(M_\ell M_\ell^\dagger \right) x_k,$$

whence we conclude that $\left(M_\ell M_\ell^\dagger \right) x_k$ is an eigenvector of E to the eigenvalue ω^{k-1} . Since eigenvectors to non-degenerate eigenvalues are unique up to a multiplicative factor, we arrive at $\left(M_\ell M_\ell^\dagger \right) x_k = \nu_k x_k$ where the quantities $|\nu_k|$ are identical with the charged-lepton masses. \square

Turning to the neutrino mass matrix \mathcal{M}_ν , we remember that we have started with the requirement that it is invariant under S , i.e. $S^T \mathcal{M}_\nu S = \mathcal{M}_\nu$. In the present discussion S is given by equation (20). By construction, this matrix is an involution—see also equation (6)—with a unique eigenvalue 1:

$$S u = u = U_\omega u_1 \quad \text{with} \quad u = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}. \quad (22)$$

Then we know from proposition 1 that u is also an eigenvector of \mathcal{M}_ν . Therefore, in our basis 1 the mechanism for achieving TM_1 boils down to

$$U_\omega^\dagger u = \frac{1}{\sqrt{6}} \begin{pmatrix} 2 \\ -1 \\ -1 \end{pmatrix}. \quad (23)$$

This mechanism has recently been used in [10] for the construction of a model which exhibits TM_1 .

4 Realizing TM_1 in a concrete S_4 scheme

Though the mechanism for obtaining TM_1 from the mass matrices is unique, it is not unique how to embed it into an S_4 model. Below we introduce a scheme with the type II seesaw mechanism [22].

Our starting point is the tensor product (see for instance [25])

$$\mathbf{3} \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{2} \oplus \mathbf{3} \oplus \mathbf{3}', \quad (24)$$

where the $\mathbf{1}$ is the trivial one-dimensional representation and the $\mathbf{3}$ and $\mathbf{3}'$ are the two inequivalent irreducible three-dimensional representations of S_4 —see appendix B for the generators and the three-dimensional representations. The $\mathbf{3}'$ and $\mathbf{3}$ correspond to the off-diagonal symmetric and antisymmetric parts, respectively, in the tensor product. The $\mathbf{1}$ and the $\mathbf{2}$ comprise the diagonal part. If we assign to both the left-handed lepton gauge doublets and the right-handed lepton gauge singlets a $\mathbf{3}$ of S_4 , then the right-hand side of equation (24) shows the possible irreducible S_4 representations of Higgs doublets.

Since we require the validity of equation (21), we need the vacuum to be invariant under $s = (123)$, which is mapped in the $\mathbf{3}$ and $\mathbf{3}'$ into E —see appendix B. However, for the two-dimensional irreducible representation we have

$$s = (123) \mapsto \begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix}, \quad (25)$$

which means that no non-trivial VEV is invariant under this representation of s and the $\mathbf{2}$ cannot contribute to M_ℓ . For $\mathbf{3}$ and $\mathbf{3}'$, invariance of the VEVs under s means that the VEVs of the three Higgs doublets have to be equal. Eventually, we arrive at the most general M_ℓ , invariant under E and compatible with equation (24):

$$M_\ell = \begin{pmatrix} a & b+c & b-c \\ b-c & a & b+c \\ b+c & b-c & a \end{pmatrix}. \quad (26)$$

Indeed, here U_ω diagonalizes not only $M_\ell M_\ell^\dagger$ but also M_ℓ :

$$U_\omega^\dagger M_\ell U_\omega = \text{diag} \left(a+2b, a-b+\sqrt{3}ic, a-b-\sqrt{3}ic \right). \quad (27)$$

This result shows that M_ℓ of equation (26) is rich enough to accommodate three different charged-lepton masses, albeit with finetuning.

Using equation (24) in the neutrino sector, we note that the antisymmetric part on the right-hand side, the $\mathbf{3}$, is not allowed according to the assumed Majorana nature of the neutrinos. Dropping also the $\mathbf{2}$, we have four scalar gauge triplets Δ_k ($k = 0, 1, 2, 3$) in $\mathbf{1} \oplus \mathbf{3}'$. From equation (20) we know that \mathcal{M}_ν has to be invariant under $S_1 B$. Therefore, the triplet VEVs w_k ($k = 0, 1, 2, 3$) have to be invariant under the S_4 transformation corresponding to $S_1 B$ in the representation of the scalar triplets. Leaving out the trivial case of w_0 and using that $S_1 B$ acts as $-S_1 B$ on the $\mathbf{3}'$, the VEVs w_k ($k = 1, 2, 3$) are determined by

$$-S_1 B \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} \Rightarrow w_3 = -w_2, \quad (28)$$

and, therefore, the neutrino mass matrix has the form

$$\mathcal{M}_\nu = \begin{pmatrix} A & B & -B \\ B & A & C \\ -B & C & A \end{pmatrix}. \quad (29)$$

The mass matrices M_ℓ and \mathcal{M}_ν of this section have recently been obtained in [10] in a different model.

5 Summary

Before we summarize our findings, we want to point out the caveats and limitations attached to the group-theoretical method reviewed in section 2. It is useful to distinguish between three groups: G is the family symmetry group of the Lagrangian, the group $D(G)$ is the $U(3)$ subgroup given by the representation matrices of G on the three left-handed leptonic gauge doublets, and \bar{G} is the $U(3)$ subgroup generated by the residual symmetries of the mass matrices $M_\ell M_\ell^\dagger$ and \mathcal{M}_ν . The method of section 2 is a prescription for the determination of \bar{G} . How \bar{G} is related to G in a specific model and what \bar{G} tells us about model building, is another matter. This always has to be kept in mind when assessing results obtained by the group-theoretical method of section 2. Below, whenever we use the phrase “mass matrices,” we mean $M_\ell M_\ell^\dagger$ and \mathcal{M}_ν . Our list of caveats is the following:

- The method of section 2 explicitly assumes that the family symmetry group G of the Lagrangian is finite and that neutrinos have Majorana nature.
- Since this method is purely group-theoretical and uses only information contained in the mass matrices, it can yield at most $D(G)$.
- It is well known that accidental symmetries can occur in the mass matrices, which contribute, therefore, to \bar{G} . An accidental symmetry cannot be elevated to a symmetry of the Lagrangian. In this case, \bar{G} is not even a subgroup of $D(G)$ —see for instance [20].
- The method does not apply to models where VEVs break G totally.

Note that it is possible that a model is predictive because of an accidental symmetry.² Even if G is totally broken and there are no accidental symmetries, the model can be predictive because of the restrictions imposed by G on the Yukawa couplings or because the VEVs have an alignment but this does not correspond to a subgroup of G . For instance, the typical neutrino mass matrix resulting from $\Delta(27)$ is a case where the group is completely broken, but has a predictive neutrino mass matrix in specific models [26].

The methods and results of the paper can be summarized as follows:

- i) We have used the group-theoretical method of [15], together with theorem 1 on vanishing sums of roots of unity, to determine all possible groups \bar{G} which result from the requirement that in the lepton mixing matrix U the first column is given by equation (1), i.e. identical with the first column of the tri-bimaximal mixing matrix. This is called TM_1 in [6].
- ii) The result is amazingly simple. Only groups \bar{G} of the form $\mathbb{Z}_q \times S_4$ with $q = 1, 2, 3, \dots$ are capable to enforce TM_1 without fixing the columns u_2 and u_3 in U . Note that we have not only shown that all such groups contain S_4 [18], from our investigation

²The typical A_4 models [4] have a $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry in \mathcal{M}_ν , but only one \mathbb{Z}_2 is the residual symmetry of A_4 .

it follows that any such group larger than S_4 is obtained from S_4 by multiplication with a cyclic factor.³

- iii) Furthermore, in the basis where the Klein four-group, which is a subgroup of S_4 —see appendix B, is represented by diagonal matrices, we have found a unique mechanism for achieving TM_1 : the first column in U_ν must be the vector u of equation (22), while $U_\ell = U_\omega$ —see equation (19). This mechanism was recently used in [10].
- iv) Finally, we have pointed out how to straightforwardly implement the mechanism of the previous item in a class of renormalizable S_4 models with type II seesaw mechanism. It is fair to mention that we have not solved the VEV alignment problem in this context.

Acknowledgments: The author thanks P.O. Ludl for many helpful discussions and L. Lavoura for comments on an early version of the manuscript. Moreover, the author is very much indebted to R.M. Fonseca for pointing out the erroneous theorem 1 in the previous version of the paper, which has been replaced here by the theorem of Conway and Jones.

A Roots of unity and the eigenvalues of T and ST

Here we discuss the general solution of

$$\mathcal{S} \equiv -e^{i\phi_e} + 2e^{i\phi_\mu} + 2e^{i\phi_\tau} + 3\lambda_1 + 3\lambda_2 + 3\lambda_3 = 0, \quad (\text{A1})$$

where $e^{i\phi_\alpha}$ ($\alpha = e, \mu, \tau$) and λ_k ($k = 1, 2, 3$) are roots of unity, *c.f.* equation (14). As argued in section 3, according to theorem 1 (theorem 6 of [24]), any solution of this equation can only involve partial sums $\theta_j (1 + \omega + \omega^2)$ and “empty sums” $\epsilon_\kappa - \epsilon_\kappa$ for some roots of unity θ_j and ϵ_κ . \mathcal{S} has 14 roots, though at most six of them are different. Therefore, \mathcal{S} must have one of the following three forms:

$$\mathcal{S} = \sum_{j=1}^4 \theta_j (1 + \omega + \omega^2) + \epsilon_1 - \epsilon_1 \quad (\text{A2})$$

or

$$\mathcal{S} = \sum_{j=1}^2 \theta_j (1 + \omega + \omega^2) + \sum_{\kappa=1}^4 (\epsilon_\kappa - \epsilon_\kappa) \quad (\text{A3})$$

or

$$\mathcal{S} = \sum_{\kappa=1}^7 (\epsilon_\kappa - \epsilon_\kappa). \quad (\text{A4})$$

³In [11] S_4 was identified as the minimal group \bar{G} for tri-bimaximal mixing, but it is also the group for TM_1 alone [18, 20]; this follows for instance from proposition 4. The reason is that S_4 contains not only \tilde{S}_1 of equation (13) but also $\tilde{S}_i = 2u_i u_i^\dagger - \mathbb{1}$ ($i = 2, 3$) where u_2 and u_3 are the second and third column of U_{TBM} , respectively. When S_4 is the group of TM_1 , then \tilde{S}_i with $i = 2, 3$ is broken and not part of G_ν .

Clearly, in order to obtain solutions of equation (A1), we have to reduce the number of different roots occurring in equations (A2)–(A4) to at most six. To facilitate this task, we define the sets $M_j = \{\theta_j, \theta_j\omega, \theta_j\omega^2\}$ and observe that for $j \neq j'$ it follows that either $M_j = M_{j'}$ or $M_j \cap M_{j'} = \emptyset$. Furthermore, in a sum of roots of unity, we call the *positive* coefficients in front of the roots weights. Thus in equation (A1) the weights are 3, 3, 3, 2, 2 and 1, with the sum over the weights being 14.

First we discuss equation (A2). In order to reduce the number of different roots in \mathcal{S} to at most six, without loss of generality we have to make one of the following identifications:

$$M_1 = M_2 = M_3 = M_4, \quad \text{or} \quad M_1 = M_2 = M_3 \neq M_4, \quad \text{or} \quad M_1 = M_2 \neq M_3 = M_4.$$

In the first case we have $\mathcal{S} = 4\theta_1(1 + \omega + \omega^2) + \epsilon_1 - \epsilon_1$. In order to avoid two weights 1 in \mathcal{S} , we have to identify either ϵ_1 or $-\epsilon_1$ with a root in M_1 . Without loss of generality we put $\epsilon_1 = \theta_1$ and obtain $\mathcal{S} = 5\theta_1 + 4\theta_1\omega + 4\theta_1\omega^2 - \theta_1$. It is then obvious that we cannot achieve three weights 3 and two weights 2. Regarding the second and third case, in order to avoid more than six different roots, we must require $\epsilon_1 \in M_2$ and $-\epsilon_1 \in M_4$; without loss of generality, this requirement is satisfied by $\theta_1 = -\theta_4$. But then the second case leads to $\mathcal{S} = 3\theta_1(1 + \omega + \omega^2) - \theta_1(1 + \omega + \omega^2) + \theta_1 - \theta_1$. Indeed there are three weights 3, but not two weights 2. In the third case we have $\mathcal{S} = 2\theta_1(1 + \omega + \omega^2) - 2\theta_1(1 + \omega + \omega^2) + \theta_1 - \theta_1$. Now there are two weights 3 instead of three. In summary, we find that equation (A2) cannot lead to a solution of equation (A1).

Next we consider equation (A4). Without loss of generality, in order to produce three weights 3, we make the identifications $\epsilon_1 = \epsilon_2 = \epsilon_3$ and $\epsilon_4 = \epsilon_5 = \epsilon_6$. Thus we consider $\mathcal{S} = 3(\epsilon_1 - \epsilon_1 + \epsilon_4 - \epsilon_4) + \epsilon_7 - \epsilon_7$. In order to avoid two weights 1, we furthermore equate, without loss of generality, ϵ_7 with ϵ_4 . In this way, we arrive at $\mathcal{S} = 3(\epsilon_1 - \epsilon_1) + 4(\epsilon_4 - \epsilon_4)$. Now it suffices to consider only the non-trivial case $\epsilon_4 \neq \pm\epsilon_1$. It is then clear that the resulting \mathcal{S} leads to a solution of equation (A1) given by $\lambda_1 = -\lambda_2 = \epsilon_1$, $\lambda_3 = \mp\epsilon_4$, $e^{i\phi_e} = e^{i\phi_\mu} = e^{i\phi_r} = \pm\epsilon_4$ and $T = \pm\epsilon_4\mathbb{1}$. But this T is trivial and we discard equation (A4) as well.

Considering equation (A3), we have either $M_1 = M_2$ or $M_1 \neq M_2$. We begin with the second case. In order to have at most six different roots, without loss of generality, we assume $\epsilon_\kappa \in M_1 \forall \kappa = 1, \dots, 4$, which necessitates $\theta_2 = -\theta_1$. In this case, for every root in M_1 there is a corresponding root with the opposite sign in M_2 . Consequently, all weights occur in even numbers, which is a contradiction to equation (A1). Thus we are left with $M_1 = M_2$. We first envisage the possibility that the ϵ_κ do *not* produce weight 3. Then, in order to obtain three weights 3, we identify ϵ_κ with $\theta_1\omega^{\kappa-1}$ for $\kappa = 1, 2, 3$, which yields the sum $\mathcal{S} = 3\theta_1(1 + \omega + \omega^2) - \theta_1(1 + \omega + \omega^2) + \epsilon_4 - \epsilon_4$. Now there is no choice of ϵ_4 such that the weights of equation (A1) are reproduced. It remains to consider the case that the ϵ_κ do produce weight 3, *i.e.* without loss of generality we assume $\epsilon_1 = \epsilon_2 = \epsilon_3$. With this we obtain the sum $\mathcal{S} = 2\theta_1(1 + \omega + \omega^2) + 3(\epsilon_1 - \epsilon_1) + \epsilon_4 - \epsilon_4$. Since we need a third weight 3, without loss of generality we make the identification $\epsilon_4 = \theta_1$. Now we have indeed a viable solution of equation (A1), given by $\lambda_1 = -\lambda_2 = \epsilon_1$, $\lambda_3 = \theta_1$, $e^{i\phi_e} = \theta_1$, $e^{i\phi_\mu} = \theta_1\omega$, $e^{i\phi_r} = \theta_1\omega^2$. With $\theta_1 \equiv \eta$, $\epsilon_1 \equiv \epsilon$ this is precisely the solution presented in equation (16).

B Generators of S_4

The generators of a finite group are not unique. For our purpose it is useful to consider the Klein four-group, which is an Abelian subgroup of S_4 , given by $k_1 = (12)(34)$, $k_2 = (14)(23)$, $k_3 = (13)(24)$ and the unit element. We further need one three-cycle, say $s = (123)$, and one transposition, say $t = (12)$. All elements of S_4 can be obtained as products of these permutations—see for instance [25]. However, what we are really interested in is a faithful three-dimensional irreducible representation of S_4 . Here we display the $\mathbf{3}$ as derived in [25]:

$$\begin{aligned} k_1 &\mapsto S_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \\ k_2 &\mapsto S_2 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \\ k_3 &\mapsto S_3 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \end{aligned} \tag{B1}$$

and

$$s \mapsto E = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad t \mapsto B = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}. \tag{B2}$$

In the spirit of section 2 we call this basis of the $\mathbf{3}$ of S_4 basis 1. The second three-dimensional irreducible representation, $\mathbf{3}'$, is obtained from $\mathbf{3}$ by a sign change in equation (B2), namely $t \mapsto -B$.

It is also useful to have the above generators in the basis where E is diagonal. With the similarity transformation $U_\omega^\dagger A U_\omega = \tilde{A}$ where U_ω is given by equation (19) we obtain (see for instance [5, 27])

$$\tilde{S}_1 = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}, \quad \tilde{S}_2 = \frac{1}{3} \begin{pmatrix} -1 & 2\omega & 2\omega^2 \\ 2\omega^2 & -1 & 2\omega \\ 2\omega & 2\omega^2 & -1 \end{pmatrix}, \quad \tilde{S}_3 = \frac{1}{3} \begin{pmatrix} -1 & 2\omega^2 & 2\omega \\ 2\omega & -1 & 2\omega^2 \\ 2\omega^2 & 2\omega & -1 \end{pmatrix} \tag{B3}$$

and

$$\tilde{E} = \text{diag}(1, \omega, \omega^2), \quad \tilde{B} = B. \tag{B4}$$

In the spirit of section 2 this is the $\mathbf{3}$ given in basis 2.

References

- [1] Y. Abe *et al.* (Double Chooz Coll.), *Indication for the disappearance of reactor electron antineutrinos in the Double Chooz experiment*, Phys. Rev. Lett. **108** (2012) 131801 [arXiv:1112.6353 [hep-ex]];

- F.P. An *et al.* (Daya Bay Coll.), *Observation of electron-antineutrino disappearance at Daya Bay*, Phys. Rev. Lett. **108** (2012) 171803 [arXiv:1203.1669 [hep-ex]];
 J.K. Ahn *et al.* (RENO Coll.), *Observation of reactor electron antineutrino disappearance in the RENO experiment*, Phys. Rev. Lett. **108** (2012) 191802 [arXiv:1204.0626 [hep-ex]].
- [2] D.V. Forero, M. Tórtola, and J.W.F. Valle, *Global status of neutrino oscillation parameters after recent reactor measurements*, Phys. Rev. D **86** (2012) 073012 [arXiv:1205.4018 [hep-ph]];
 G.L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo, and A.M. Rotunno, *Global analysis of neutrino masses, mixings and phases: entering the era of leptonic CP violation searches*, Phys. Rev. D **86** (2012) 013012 [arXiv:1205.5254 [hep-ph]];
 M.C. Gonzalez-Garcia, M. Maltoni, J. Salvado and T. Schwetz, *Global fit to three neutrino mixing: critical look at present precision*, JHEP **1212** (2012) 123 [arXiv:1209.3023 [hep-ph]].
- [3] P.F. Harrison, D.H. Perkins and W.G. Scott, *Tri-bimaximal mixing and the neutrino oscillation data*, Phys. Lett. B **530** (2002) 167 [hep-ph/0202074].
- [4] G. Altarelli and F. Feruglio, *Discrete flavor symmetries and models of neutrino mixing*, Rev. Mod. Phys. **82** (2010) 2701 [arXiv:1002.0211 [hep-ph]];
 A.Yu. Smirnov, *Discrete symmetries and models of flavor mixing*, J. Phys. Conf. Ser. **335** (2011) 012006 [arXiv:1103.3461 [hep-ph]];
 G. Altarelli, F. Feruglio, L. Merlo and E. Stamou, *Discrete flavour groups, θ_{13} and lepton flavour violation*, JHEP **1208** (2012) 021 [arXiv:1205.4670 [hep-ph]].
- [5] G. Altarelli, F. Feruglio and L. Merlo, *Tri-bimaximal neutrino mixing and discrete flavour symmetries*, Fortschr. Phys. (2012) 1-28 [arXiv:1205.5133 [hep-ph]], review written for the special issue on “Flavor Symmetries and Neutrino Oscillations.”
- [6] C.H. Albright and W. Rodejohann, *Comparing trimaximal mixing and its variants with deviations from tri-bimaximal mixing*, Eur. Phys. J. C **62** (2009) 599 [arXiv:0812.0436 [hep-ph]];
 C.H. Albright, A. Dueck and W. Rodejohann, *Possible alternatives to tri-bimaximal mixing*, Eur. Phys. J. C **70** (2010) 1099 [arXiv:1004.2798 [hep-ph]].
- [7] S. Antusch, S.F. King, C. Luhn and M. Spinrath, *Trimaximal mixing with predicted θ_{13} from a new type of constrained sequential dominance*, Nucl. Phys. B **856** (2012) 328 [arXiv:1108.4278 [hep-ph]].
- [8] W. Rodejohann and H. Zhang, *Simple two parameter description of lepton mixing*, Phys. Rev. D **86** (2012) 093008 [arXiv:1207.1225 [hep-ph]].
- [9] Ernest Ma, *Self-organizing neutrino mixing matrix*, Phys. Rev. D **86** (2012) 117301 [arXiv:1209.3374 [hep-ph]].
- [10] I. de Medeiros Varzielas and L. Lavoura, *Flavour models for TM_1 lepton mixing*, J. Phys. G **40** (2013) 085002 [arXiv:1212.3247 [hep-ph]].

- [11] C.S. Lam, *Determining horizontal symmetry from neutrino mixing*, Phys. Rev. Lett. **101** (2008) 121602 [arXiv:0804.2622 [hep-ph]];
 C.S. Lam, *The unique horizontal symmetry of leptons*, Phys. Rev. D **78** (2008) 073015 [arXiv:0809.1185 [hep-ph]];
 C.S. Lam, *A bottom-up analysis of horizontal symmetry*, arXiv:0907.2206 [hep-ph].
- [12] Shao-Feng Ge, D.A. Dicus and W.W. Repko, *Z_2 symmetry prediction for the leptonic Dirac CP phase*, Phys. Lett. B **702** (2011) 220 [arXiv:1104.0602 [hep-ph]];
 Shao-Feng Ge, D.A. Dicus and W.W. Repko, *Residual symmetries for neutrino mixing with a large θ_{13} and nearly maximal δ_D* , Phys. Rev. Lett. **108** (2012) 041801 [arXiv:1108.0964 [hep-ph]].
- [13] Hong-Jian He and Fu-Rong Yin, *Common origin of μ - τ and CP breaking in neutrino seesaw, baryon asymmetry, and hidden flavor symmetry*, Phys. Rev. D **84** (2011) 033009 [arXiv:1104.2654 [hep-ph]];
 Hong-Jian He and Xun-Jie Xu, *Octahedral symmetry with geometrical breaking: New prediction for neutrino mixing angle θ_{13} and CP violation*, Phys. Rev. D **86** (2012) 111301 (R) [arXiv:1203.2908].
- [14] R. de A. Toorop, F. Feruglio and C. Hagedorn, *Discrete flavour symmetries in light of T2K*, Phys. Lett. B **703** (2011) 447 [arXiv:1107.3486 [hep-ph]];
 R. de Adelhart Toorop, F. Feruglio and C. Hagedorn, *Finite Modular Groups and Lepton Mixing*, Nucl. Phys. B **858** (2012) 437 [arXiv:1112.1340 [hep-ph]].
- [15] D. Hernandez and A.Yu. Smirnov, *Lepton mixing and discrete symmetries*, Phys. Rev. D **86** (2012) 053014 [arXiv:1204.0445 [hep-ph]].
- [16] D. Hernandez and A.Yu. Smirnov, *Discrete symmetries and model-independent patterns of lepton mixing*, Phys. Rev. D **87** (2013) 053005 [arXiv:1212.2149 [hep-ph]].
- [17] Bo Hu, *Neutrino mixing and discrete symmetries*, Phys. Rev. D **87** (2013) 033002 [arXiv:1212.2819 [hep-ph]].
- [18] C.S. Lam, *Finite symmetry of leptonic mass matrices*, Phys. Rev. D **87** (2013) 013001 [arXiv:1208.5527 [hep-ph]].
- [19] M. Holthausen, Kher Sham Lim and M. Lindner, *Lepton mixing patterns from a scan of finite discrete groups*, Phys. Lett. B **721** (2013) 61 [arXiv:1212.2411 [hep-ph]].
- [20] W. Grimus, L. Lavoura and P.O. Ludl, *Is S_4 the horizontal symmetry of tri-bimaximal lepton mixing?*, J. Phys. G **36** (2009) 115007 [arXiv:0906.2689 [hep-ph]].
- [21] P. Minkowski, *$\mu \rightarrow e\gamma$ at a rate of one out of 10^9 muon decays?*, Phys. Lett. **67B** (1977) 421;
 T. Yanagida, *Horizontal gauge symmetry and masses of neutrinos*, in *Proceedings of the workshop on unified theory and baryon number in the universe (Tsukuba, Japan, 1979)*, O. Sawata and A. Sugamoto eds., KEK report **79-18** (Tsukuba, Japan, 1979);

- S.L. Glashow, *The future of elementary particle physics*, in *Quarks and leptons, proceedings of the advanced study institute (Cargèse, Corsica, 1979)*, M. Lévy *et al.* eds. (Plenum Press, New York, U.S.A., 1980);
- M. Gell-Mann, P. Ramond and R. Slansky, *Complex spinors and unified theories*, in *Supergravity*, D.Z. Freedman and F. van Nieuwenhuizen eds. (North Holland, Amsterdam, The Netherlands, 1979);
- R.N. Mohapatra and G. Senjanović, *Neutrino mass and spontaneous parity violation*, Phys. Rev. Lett. **44** (1980) 912.
- [22] M. Magg and C. Wetterich, *Neutrino mass problem and gauge hierarchy*, Phys. Lett. **94B** (1980) 61;
 G. Lazarides, Q. Shafi and C. Wetterich, *Proton lifetime and fermion masses in an $SO(10)$ model*, Nucl. Phys. B **181** (1981) 287;
 R.N. Mohapatra and G. Senjanović, *Neutrino masses and mixings in gauge models with spontaneous parity violation*, Phys. Rev. D **23** (1981) 165;
 R.N. Mohapatra and P. Pal, *Massive neutrinos in physics and astrophysics* (World Scientific, Singapore, 1991), p. 127;
 E. Ma and U. Sarkar, *Neutrino masses and leptogenesis with heavy Higgs triplets*, Phys. Rev. Lett. **80** (1998) 5716 [hep-ph/9802445].
- [23] X.G. He, Y.Y. Keum and R.R. Volkas, *A_4 flavour symmetry breaking scheme for understanding quark and neutrino mixing angles*, JHEP **0604** (2006) 039 [hep-ph/0601001];
 F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, *Tri-bimaximal neutrino mixing and quark masses from a discrete flavour symmetry*, Nucl. Phys. B **775** (2007) 120 [hep-ph/0702194];
 A. Blum, C. Hagedorn and M. Lindner, *Fermion masses and mixings from dihedral flavor symmetries with preserved subgroups*, Phys. Rev. D **77** (2008) 076004 [arXiv:0709.3450 [hep-ph]].
- [24] J.H. Conway and A.J. Jones, *Trigonometric diophantine equations (On vanishing sums of roots of unity)*, Acta Arithmetica **30** (1976) 229.
- [25] W. Grimus and P.O. Ludl, *Finite flavour groups of fermions*, J. Phys. A **45** (2012) 233001 [arXiv:1110.6376 [hep-ph]].
- [26] Ernest Ma, *Neutrino mass matrix from $\Delta(27)$ symmetry*, Mod. Phys. Lett. A **21** (2006) 1917 [hep-ph/0607056];
 Ernest Ma, *Near tribimaximal neutrino mixing with $\Delta(27)$ symmetry*, Phys. Lett. B **660** (2008) 505 [arXiv:0709.0507 [hep-ph]];
 P.M. Ferreira, W. Grimus, L. Lavoura and P.O. Ludl, *Maximal CP violation in lepton mixing from a model with $\Delta(27)$ flavour symmetry*, JHEP **1209** (2012) 128 [arXiv:1206.7072 [hep-ph]].
- [27] Y. H. Ahn, S. Baek and P. Gondolo, *Simple renormalizable flavor symmetry for neutrino oscillations*, Phys. Rev. D **86** (2012) 053004 [arXiv:1207.1229 [hep-ph]].