

Flavor-Nonuniversal Dark Gauge Bosons and the Muon $g - 2$

Christopher D. Carone*

*High Energy Theory Group, Department of Physics,
College of William and Mary, Williamsburg, VA 23187-8795*

(Dated: January 2013)

Abstract

The possibility of explaining the current value of the muon anomalous magnetic moment in models with an additional U(1) gauge symmetry that has kinetic mixing with hypercharge are increasingly constrained by dark photon searches at electron accelerators. Here we present a scenario in which the couplings of new, light gauge bosons to standard model leptons are naturally weak and flavor non-universal. A vector-like state that mixes with standard model leptons serves as a portal between the dark and standard model sectors. The flavor symmetry of the model assures that the induced couplings of the new gauge sector to leptons of the first generation are very small and lepton-flavor-violating processes are adequately suppressed. The model provides a framework for constructing ultraviolet complete theories in which new, light gauge fields couple weakly and dominantly to leptons of the second or third generations.

*cdcaro@wm.edu

I. INTRODUCTION

In recent years, the energy, intensity and cosmic frontiers have come into focus as the three domains in which future discoveries in elementary particle physics are likely to arise. Among the possibilities that might be revealed at the intensity frontier are “dark sectors” [1], with particles much lighter than the electroweak scale that have remained undetected due to their very weak interactions with standard model particles. The gravitational effects of dark matter is indirect evidence that at least one dark particle exists. The possibility that dark matter may also couple to spontaneously broken dark gauge forces has been motivated by the observation that sub-GeV dark gauge boson decays could explain the electron and positron excesses observed in the cosmic ray spectra [2]. More generally, sectors that are only weakly coupled to the standard model arise quite generically in string theory [3], so that investigations of other possible dark sectors [4] continue to be well motivated.

Given that the particle content and gauge structure of a generic dark sector is not well constrained, the largest number of analyses have assumed the simplest construction, a dark U(1) gauge field that has a small kinetic mixing with hypercharge [5]. Interestingly, it has been pointed out that a discrepancy between the observed value of the muon anomalous magnetic moment and the expectation of the standard model can be explained in this scenario [6]. However, the region of the mass-kinetic mixing plane in which this result can be achieved is very rapidly being excluded by bounds from dark photon searches [8], in particular, the most recent results from the Mainz Microtron experiment [9]. Dominant kinetic mixing will always assure that a dark photon couples identically to electrons and muons, so there is no way to escape this bound. The possibility that dark forces might couple differently to muons than electrons has been discussed in a somewhat different context, *i.e.*, the proton charge radius problem, in Ref. [10], but from an effective field theory perspective. Here we will present a complete theory in which such a possibility is naturally realized.

Although an explanation of the muon anomalous magnetic moment discrepancy is certainly worthwhile, it is arguably more important that the present work shows, in a complete theory, how a dark sector can naturally couple both weakly and preferentially to charged leptons of the second or third generation. This may provide motivation for a more intensive phenomenological consideration of processes involving muons and taus as a means of discerning couplings to new dark sector particles. The basic structure of the model is this: the

dark gauge and higgs particles couple directly to a vector-like charged lepton that has gauge quantum numbers that allow mixing with the right-handed electron, muon or tau. However, the lepton sector is also constrained by a discrete, A_4 flavor symmetry that is the origin of neutrino tribimaximal mixing. This symmetry leads to preferential mixing between the vector-like state and only one standard model lepton flavor, while simultaneously keeping lepton-flavor-violating effects suppressed. The couplings between the standard model and the dark sector particles are naturally small since they are proportional to a small mixing angle that is set by the ratio of the dark to vector-like mass scales. Lepton-flavor-universal couplings due to kinetic mixing with hypercharge are loop suppressed, since the dark sector gauge group in the proposed model is non-Abelian.

Our letter is organized as follows. In the next section, we describe the basic structure of the theory and show that the muon anomalous magnetic moment discrepancy can be resolved. In Sec. III, we explain how the pattern of couplings assumed in Sec. II can be achieved via the discrete flavor symmetry A_4 . We briefly discuss other phenomenological bounds in Sec. IV, and in Sec. V we conclude.

II. THE PORTAL

The gauge group of the model is $G_{\text{SM}} \times \text{SU}(2)_D$, where G_{SM} is the standard model (SM) gauge group and the additional factor represents the gauge group of the dark sector. Since the dark gauge group is non-Abelian, kinetic mixing with hypercharge does not appear immediately through a renormalizable term in the Lagrangian. We assume the existence of a vector-like doublet

$$E_L^a, \quad E_R^a, \quad (2.1)$$

where a is an $\text{SU}(2)_D$ index. These fields have the same G_{SM} quantum numbers as a right-handed electron, muon or tau. Note that the theory is free of anomalies, since the new fermions come in a vector-like pair. Since the E fields carry both SM and dark quantum numbers, they serve as a portal between the SM and dark sectors.

We assume that the dark gauge symmetry is completely broken by the vacuum expectation value (vev) of a complex scalar doublet ϕ^a ,

$$\langle \phi \rangle = \begin{pmatrix} 0 \\ v_D/\sqrt{2} \end{pmatrix}. \quad (2.2)$$

A renormalizable coupling between ϕ and the SM Higgs field H , namely $\phi^\dagger \phi H^\dagger H$, can serve as another portal between the SM and dark sectors. However, to simplify the model parameter space in the present analysis, we will only consider the case in which this coupling is negligibly small. This limit is consistent with LHC data on the recently discovered 125 GeV boson which (perhaps aside from the decay rate to two photons) shows no substantial deviations from the expectations for a standard model Higgs boson.

Mass mixing between the E and SM lepton fields can occur when the $SU(2)_D$ gauge symmetry is broken. This mixing will induce a coupling between the dark gauge boson multiplet and ordinary SM leptons that is suppressed by a small mixing angle, one proportional to v_D/v , where $v = 246$ GeV is the electroweak scale. Like most dark photon models, we will typically assume that the dark gauge bosons have masses less than 1 GeV. Mass mixing originates through the Yukawa couplings

$$\mathcal{L} \supset h \bar{\mu}_R \phi_a^* E_L^a + h' \epsilon_{ab} \bar{\mu}_R \phi^a E_L^b + \text{h.c.} \quad , \quad (2.3)$$

Here we have taken the SM lepton field to be of the second generation. Generically one expects similar couplings involving the e and τ . Our model, however, is not generic. We will show in Sec. III that the flavor structure of Eq. (2.3) follows accurately from the flavor symmetries of the model. For the remainder of this section, we will simply assume the mixing given in Eq. (2.3) and study the new contributions to the muon $g - 2$.

In order to specify the model, one must know the vector-like lepton mass M , the mass and gauge coupling of the dark gauge bosons, m_V and g_D respectively, and the mass of the dark higgs, m_ϕ . (The dark higgs is the physical fluctuation about v_D in the doublet field ϕ .) In addition, one must know the mixing angles that are functions of the couplings h and h' in Eq. (2.3). While a complete exploration of this six-dimensional parameter space is impractical, the essential physics can be illustrated by adopting a few simplifying assumptions. We take $m_V = m_\phi$, so that a single mass scale characterizes the dark sector. In addition, we set the coupling $h' = 0$, which renders the analysis of mixing between the SM leptons and the vector-like states more transparent. (This latter assumption can be elevated to a definition of the model if one imposes a discrete symmetry that forbids the h' coupling. A Z_3 symmetry in which $E_{L,R} \rightarrow \omega E_{L,R}$ and $\phi^a \rightarrow \omega \phi^a$, with $\omega^3 = 1$, is a suitable example of such a symmetry.)

Given these assumptions, mass mixing between the SM and vector-like states are given




FIG. 1: New contributions to the muon anomalous magnetic moment.

by a two-by-two mass matrix,

$$\mathcal{L} \supset \bar{\Psi}_L \begin{pmatrix} m_\mu & 0 \\ \delta m & M \end{pmatrix} \Psi_R + \text{h.c.}, \quad (2.4)$$

where $\Psi = (\mu, E^{(2)})^T$ and $\delta m = h v_d / \sqrt{2}$. The relationship between the gauge and mass eigenstate bases is given by

$$\Psi_j = \begin{pmatrix} c_j & s_j \\ -s_j & c_j \end{pmatrix} \Psi_j^{\text{mass}} \quad (2.5)$$

where $s_j = \sin \theta_j$, $c_j = \cos \theta_j$, and j is either L or R . With $\delta m \ll M$, the mixing angles are well approximated by $\theta_R \approx \delta m / M$ and $\theta_L \approx (m_\mu / M)(\delta m / M)$. Due to this mixing, the $SU(2)_D$ gauge bosons, which otherwise would couple only to the vector-like states, now also couple to standard model leptons. In terms of mass eigenstate fields, these couplings include

$$\begin{aligned} \mathcal{L} \supset & \left[-\frac{1}{2} s_R^2 g_D (\bar{\mu}_R \mathcal{A}_0 \mu_R) - \frac{1}{\sqrt{2}} s_R g_D (\bar{\mu}_R \mathcal{A}_- E_R^{(1)} + \text{h.c.}) \right. \\ & \left. + \frac{1}{2} s_R c_R g_D (\bar{\mu}_R \mathcal{A}_0 E_R^{(2)} + \text{h.c.}) \right] + [\text{R} \rightarrow \text{L}] \end{aligned} \quad (2.6)$$

Here we have parameterized the $SU(2)$ gauge multiplet

$$A_D^a T^a = \begin{pmatrix} A_0/2 & A_+/\sqrt{2} \\ A_-/\sqrt{2} & -A_0/2 \end{pmatrix} \quad (2.7)$$

F	X_μ	C_V	C_A
μ	A_0	$-\frac{1}{4}g_D(s_R^2 + s_L^2)$	$-\frac{1}{4}g_D(s_R^2 - s_L^2)$
$E^{(1)}$	A_\pm	$-\frac{1}{2\sqrt{2}}g_D(s_R + s_L)$	$-\frac{1}{2\sqrt{2}}g_D(s_R - s_L)$
$E^{(2)}$	A_0	$\frac{1}{4}g_D(c_R s_R + c_L s_L)$	$\frac{1}{4}g_D(c_R s_R - c_L s_L)$

F	S	C_S	C_P
μ	φ	$-\frac{1}{\sqrt{2}}h(c_R s_L)$	0
$E^{(2)}$	φ	$\frac{1}{2\sqrt{2}}h(c_L c_R - s_L s_R)$	$-\frac{1}{2\sqrt{2}}h(c_L c_R + s_L s_R)$

TABLE I: Couplings relevant to the anomalous magnetic moment calculation.

where the T^a are the Pauli matrices over two. Note that the A_0 , A_+ and A_- are electrically neutral, but the $E^{(1)}$, $E^{(2)}$ and μ fields have identical couplings to the photon. Hence there are number of diagrams that contribute to the muon anomalous magnetic moment, as shown in Fig. 1.

In addition to the gauge-boson exchange diagrams, there are also diagrams involving exchange of the dark Higgs boson φ , also shown in Fig. 1. These follow from Eq. (2.3), which includes the following couplings of the mass eigenstate fields

$$\mathcal{L} \supset \frac{h}{\sqrt{2}}\varphi \left(-s_{LCR}\bar{\mu}\mu + c_{LSR}\overline{E^{(2)}}E^{(2)} + [c_{LCR}\bar{\mu}_R E_L^{(2)} - s_{LSR}\bar{\mu}_L E_R^{(2)} + \text{h.c.}] \right). \quad (2.8)$$

The contribution of new fermions and gauge bosons to the muon anomalous magnetic moment have been computed in a general framework in Ref. [11], assuming couplings of the form

$$\mathcal{L} = \bar{\mu}\gamma^\mu(C_V + C_A\gamma^5)FX_\mu + \text{h.c.}, \quad (2.9)$$

where F is a generic fermion and X_μ a generic gauge boson, and

$$\mathcal{L} = \bar{\mu}(C_S + C_P\gamma^5)FS + \text{h.c.}, \quad (2.10)$$

where S is a generic scalar. These results can be applied to the present model given the identifications shown in Table I. An exact numerical evaluation of the $g - 2$ formulae given in Ref. [11] have been incorporated in Fig. 2. Here we show the $m_V - \theta_R$ plane, for a fixed choice of the mixing parameter h and the dark gauge coupling g_D , with $m_\varphi = m_V$. With m_V and g_D fixed, the dark scale v_D is determined. The mass scale of the vector-like sector,




FIG. 2: Mass-mixing angle plane, for the values of g_D and h shown. The region between the dashed lines is consistent with the current experimental value of the muon anomalous magnetic moment, at plus or minus two standard deviations. The solid lines show the vector-like fermion masses, which increase for smaller values of θ_R .

M , is then fixed by its relation to the right-handed mixing angle

$$\theta_R = \left(\frac{\sqrt{2}h}{g_D M} \right) m_V, \quad (2.11)$$

as indicated by the solid lines in the figure. The parameters of the model are now fixed and the (much smaller) left-handed mixing angle can also be computed.

The region between the dashed lines is where the model accounts for the discrepancy between the experimental and SM values of the muon $(g - 2)/2$ [12]

$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = 287 \pm 80 \times 10^{-11}, \quad (2.12)$$

allowing two-standard deviation variation about the central value of Δa_μ . The solid lines represent vector-like fermion mass contours, for $M = 100, 300$ and 500 GeV. Collider searches for heavy charged leptons require only that $M > 100.8$ GeV at the 95% C.L. [12]. The solid and dashed bands still overlap for h closer to 1, but for values of M closer to its experimental lower bound.

III. THE FLAVOR SYMMETRY

There is a vast literature on flavor symmetries that can account for the very non-generic pattern of standard model fermion masses. In this section, we argue that the same flavor symmetry that might explain standard model lepton masses can also yield the non-generic pattern of couplings assumed in the previous section, with lepton-flavor-violating effects adequately suppressed.

A variety of models based on the discrete flavor group A_4 have been proposed after it was discovered that A_4 symmetry can provide neutrino mass matrix textures that are consistent with tribimaximal mixing, at lowest order. Given these promising results, it is well motivated to consider the consequences of this symmetry in the present model. The group A_4 has a triplet and three singlet representations, $\mathbf{3}$, $\mathbf{1}^0$, $\mathbf{1}^+$, and $\mathbf{1}^-$. Typical A_4 models assign the left-handed SM leptons to the $\mathbf{3}$ and the right-handed leptons to distinct singlet representations:

$$L_L \sim \mathbf{3}, e_R \sim \mathbf{1}^0, \mu_R \sim \mathbf{1}^+, \tau_R \sim \mathbf{1}^-. \quad (3.1)$$

The standard model Higgs doublet H is a trivial A_4 singlet, $\mathbf{1}^0$. For a review of group theory and model building with A_4 symmetry, we refer the reader to the review in Ref. [13]. The symmetry is broken by two flavon fields, φ_S and φ_T , that both transform as $\mathbf{3}$'s:

$$\langle \varphi_S \rangle = \begin{pmatrix} v_S \\ v_S \\ v_S \end{pmatrix}, \quad \langle \varphi_T \rangle = \begin{pmatrix} v_T \\ 0 \\ 0 \end{pmatrix}. \quad (3.2)$$

By imposing additional, model-dependent symmetries [13], it is arranged that the symmetry breaking in the charged lepton sector is a consequence of the φ_T flavon only, at lowest order, while in the neutrino sector the same statement holds for the φ_S flavon. Symmetry breaking due to φ_T alone leads to a charged lepton mass matrix that is *exactly* diagonal,

$$Y_L = \frac{v_T}{\Lambda_F} \begin{pmatrix} y_e & 0 & 0 \\ 0 & y_\mu & 0 \\ 0 & 0 & y_\tau \end{pmatrix}, \quad (3.3)$$

where Λ_F represents the scale at which flavor physics is generated. The reason for this result is that $\langle \varphi_T \rangle$ breaks A_4 down to a Z_3 under which all off-diagonal elements of Eq. (3.3) rotate by a nontrivial phase; hence, these elements remain vanishing until this symmetry is broken.

We now assign A_4 quantum numbers to the vector-like states:

$$E_L^a \sim E_R^a \sim \mathbf{1}^+ \quad (3.4)$$

Since this charge assignment is the same as that for μ_R , the mixing terms in Eq. (2.3) are A_4 invariant. However, the other possible terms are not:

$$\bar{e}_R \phi_a^* E_L^a \sim \mathbf{1}^+, \quad \bar{\tau}_R \phi_a^* E_L^a \sim \mathbf{1}^-. \quad (3.5)$$

The representations in Eq. (3.5) rotate by a nontrivial phase under the Z_3 symmetry that is left unbroken by $\langle \varphi_T \rangle$, and are forbidden until the Z_3 symmetry is broken. Hence, the same symmetry structure that leads to a diagonal charged lepton mass matrix when $A_4 \rightarrow Z_3$ assures that the vector-like states mix only with right-handed muons in the present model. For a different choice in Eq. (3.4), one could just as easily arrive at a different model in which mixing with τ_R is preferred.

We must now consider to what extent our desirable results are spoiled when the Z_3 symmetry is broken by the vacuum expectation value of the other flavon field φ_S . Unwanted couplings between E and e_R or τ_R can occur directly, or via Eq (2.3), which is no longer in the mass eigenstate basis if Y_L is non-diagonal. The value of the Z_3 -symmetry-breaking parameter v_S/Λ , however, is not set by the size of a typical charged lepton Yukawa coupling; it is determined by the mass scale of the right-handed neutrinos that participate in the see-saw mechanism. As is standard in A_4 -based models, we take the right-handed neutrinos (charge conjugated) to transform as a triplet

$$\nu_R^c \sim \mathbf{3}. \quad (3.6)$$

We will now show that it is possible to achieve the proper light neutrino mass scale with $v_S \ll v_T < \Lambda_F$, so that deviations from the Z_3 symmetric results in the charged SM lepton and vector-like sectors of our model can be made adequately small. In this case, contributions to lepton-flavor-violating processes due to the exchange of the non-standard-model states in the theory will also be highly suppressed. In what follows, we will take the flavor cut off Λ_F to be the reduced Planck mass, M_* , which provides the smallest possible values for v_S/Λ_F .

The Lagrangian terms that lead to a see-saw mechanism in neutrino sector are

$$\mathcal{L}_\nu = \frac{1}{2} m_R \bar{\nu}_R^c \nu_R + \frac{1}{2} x \bar{\nu}_R^c \varphi_S \nu_R + [y \bar{L}_L H \nu_R + \text{h.c.}], \quad (3.7)$$

where x and y are dimensionless coupling constants. This leads to the conventional tribimaximal form for the Dirac and Majorana neutrino mass matrices [13] that are input into the see-saw

$$m_{LR} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} y v, \quad m_{RR} = \begin{pmatrix} A + 2B/3 & -B/3 & -B/3 \\ -B/3 & 2B/3 & A - B/3 \\ -B/3 & A - B/3 & 2B/3 \end{pmatrix} m_R, \quad (3.8)$$

where $A = 1$, $B = xv_S/m_R$, and $v \approx 246$ GeV is the electroweak scale. The light neutrino mass matrix then follows from the see-saw formula $m_\nu = m_{LR}^T m_{RR}^{-1} m_{LR}$, and has the eigenvalues

$$m_1 = \frac{y^2 v^2}{m_R} \frac{1}{(A+B)}, \quad m_2 = \frac{y^2 v^2}{m_R} \frac{1}{A}, \quad m_3 = \frac{y^2 v^2}{m_R} \frac{1}{B-A}. \quad (3.9)$$

Neutrino oscillation data indicate the neutrino mass squared splittings $\Delta m_{21}^2 = (7.50 \pm 0.20) \times 10^{-5}$ eV² and $\Delta m_{32}^2 = 2.32_{-0.08}^{+0.12} \times 10^{-3}$ eV² [12]. These can be accommodated by Eq. (3.9) for $y^2 v^2/m_R = 0.97 \times 10^{-11}$ GeV and $B = xv_S/m_R = 1.198$. This solution implies

$$\frac{v_S}{M_*} \approx 3.7 \times 10^{-3} \frac{y^2}{x}. \quad (3.10)$$

The key point is this: Eq. (3.10) determines the deviation from the Z_3 -symmetric limit our theory in which no lepton-flavor-violating couplings appear. For $\mathcal{O}(1)$ values of x and small values of the neutrino Dirac Yukawa coupling y , Z_3 symmetry breaking can be made extremely small. It is easy to arrange this from a symmetry perspective by applying, for example, a Z_N flavor symmetry on the right-handed neutrinos that allows Majorana but not Dirac mass terms at lowest order. Since the size of v_S is near the right-handed neutrino mass scale, the smallness of the parameter in Eq. (3.10) is related to how low one can take this scale. For example, with $m_R \sim v_S \sim 20$ TeV, one would have $v_S/M_* \approx 10^{-14}$, with y comparable in size to the electron Yukawa coupling. This is more than adequate to render the theory safe from lepton-flavor-violating effects from the coupling of the dark and standard model sectors. As an example, loop effects will generate an effective operator that contributes to $\ell_i \rightarrow \ell_j \gamma$ which is of the form [13]

$$\frac{1}{\Lambda_{\text{NP}}^2} \bar{L}_{Li} H \sigma^{\mu\nu} F_{\mu\nu} Z_{ij} e_{Rj} \quad (3.11)$$

where the new physics scale here, Λ_{NP} , is related to the vector-like fermion mass scale by $\Lambda_{\text{NP}} \approx 4\pi M/g_D$, and Z_{ij} is a coupling matrix. Since Z has the same flavor-structure as

Y_L and we assume an additional symmetry prevents φ_S from contributing at lowest order, we expect that $Z \sim \mathcal{O}(v_T v_S / \Lambda_F^2)$. From Eq. (3.3), we see that v_T / Λ_F can be no smaller than approximately 0.01 if the tau Yukawa coupling is to remain perturbative. So, assuming $\Lambda_F = M_*$ and $M_R \approx 20$ TeV, we can achieve $Z_{ij} \sim \mathcal{O}(10^{-16})$. In contrast, the bound on $Z_{\mu e}$ from Ref. [13], updated to take into account the latest PDG value [12] for the $\mu \rightarrow e\gamma$ branching fraction, gives $Z_{\mu e} < 4.5 \times 10^{-9} [\Lambda_{\text{NP}} / 1\text{TeV}]^2$, which implies $Z_{\mu e} < 7.1 \times 10^{-7}$ for $g_D = 0.1$ and $M = 100$ GeV. This bound is easily satisfied. Similar conclusions can be drawn from other possible lepton-flavor-violating processes, which we will not consider in explicit detail here.

IV. OTHER PHENOMENOLOGICAL ISSUES

Before concluding, we comment on a few of the other potential bounds on the model. It has been noted [14] in the context of other models [10] in which a right-handed muon couples to a new gauge field V_μ that the effective coupling $g_R \bar{\mu}_R \mathcal{V} \mu_R$ is bounded by the absence of missing mass events in the leptonic kaon decay $K \rightarrow \mu X$. In the present model, the magnitude of g_R is $s_R^2 g_D / 2$, which ranges in Fig. 2 from 10^{-8} to 10^{-4} between approximately 5 to 300 MeV. We comment on this mass range since it roughly corresponds to the one over which limits on g_R as a function of vector boson mass are given in Fig. 3 of Ref. [14]; the limit is always larger than 10^{-3} . Hence, the model parameter space is not severely constrained based on this consideration.

The bounds on the dark sector states from accelerator experiments [7] depend on how these states decay, and there are many possible scenarios. Here we discuss one interesting case that corresponds to the largest parameter region shown in Fig. 2: if $m_V < 2m_\mu$, the dark gauge fields will decay only to electrons via loop-suppressed kinetic mixing and we can compare to the bounds from dark photon searches. The one-loop diagrams that lead to mixing between the dark gauge bosons and hypercharge lead to effective operators of the form

$$\mathcal{L}_{\text{eff}} = \frac{c_1}{M^2} \phi^\dagger F_D^{\mu\nu} \phi F_{Y\mu\nu} + \frac{c_2}{M^2} \tilde{\phi}^\dagger F_D^{\mu\nu} \tilde{\phi} F_{Y\mu\nu} + \left[\frac{c_3}{M^2} \tilde{\phi}^\dagger F_D^{\mu\nu} \phi F_{Y\mu\nu} + \text{h.c.} \right], \quad (4.1)$$

where $\tilde{\phi} = \epsilon^{ab} \phi_b^*$. The diagrams that generate these operators involve the vector-like fermions (and hence the suppression by M^2) as well as two vertices from the in-

teraction terms in Eq. (2.3). The couplings c_1 , c_2 , and c_3 are of order $h^2 g_D g_Y / (16\pi^2)$, $hh' g_D g_Y / (16\pi^2)$, and $h'^2 g_D g_Y / (16\pi^2)$, respectively. Kinetic mixing is conventionally parameterized as $\chi F_i^{\mu\nu} F_{Y\mu\nu} / 2$, where F_i is the field strength tensor corresponding to the dark gauge field A_i . From the first term in Eq. (4.1), for example, we estimate

$$\chi \sim \frac{1}{4\pi^2} g_Y \frac{h^2}{g_D} \left(\frac{m_V}{M} \right)^2. \quad (4.2)$$

The most constraining beam dump bound [15] for small kinetic mixing is from the SLAC E137 experiment. However, the E137 bound does not exclude χ below $\sim 4 \times 10^{-8}$. For the parameter values $h = 2$ and $g = 0.1$, used in Fig. 2, and $M = 200$ GeV, we find that χ is smaller than this value for $m_V \lesssim 67$ MeV. For $m_V \gtrsim 210$ MeV, E137 gives no further bounds. Of course, the dark sector could include dark matter candidates into which the the dark gauge and Higgs fields decay. In this case, dark photon searches at accelerators would present no further bounds on the model.

V. CONCLUSIONS

In this letter, we have shown how one can construct a dark sector that couples preferentially to a single flavor of standard model charged leptons, without generating large lepton-flavor-violating effects. The dark gauge and higgs bosons couple to $\mathcal{O}(100)$ GeV vector-like fermions that have the correct gauge quantum numbers to mix with either the right-handed electron, muon or tau. However, the lepton sector is also constrained by an A_4 flavor symmetry that is the origin of neutrino tribimaximal mixing. This symmetry assures that the vector-like state mixes with only one flavor of charged lepton while the charged lepton Yukawa matrix remains very accurately diagonal. Deviations from this flavor structure are related to the ratio of the right-handed neutrino mass scale to the cut off of the effective theory, and can be made adequately small. Moreover, lepton-flavor-universal couplings via kinetic mixing are loop suppressed since the dark sector gauge group is non-Abelian.

If the vector-like states mix with leptons of the second generation (which has been the assumption in this letter) contributions from the non-standard-model particles can account for the current deviation of the muon anomalous magnetic moment from the standard model expectation. More importantly, however, the construction described in this letter can be used to create ultraviolet complete theories (at least up to the high scale at which flavor is

generated) in which dark sector gauge bosons couple preferentially to either muons or taus. This provides motivation for exploring a vast realm of phenomenological processes involving second and third generation leptons as another possible window on a dark sector.

Acknowledgments

This work was supported by the NSF under Grant PHY-1068008. In addition, the author thanks Joseph J. Plumeri II for his generous support.

-
- [1] For a review, see J. Jaeckel and A. Ringwald, *Ann. Rev. Nucl. Part. Sci.* **60**, 405 (2010).
 - [2] N. Arkani-Hamed, D. P. Finkbeiner, T. R. Slatyer and N. Weiner, *Phys. Rev. D* **79**, 015014 (2009) [arXiv:0810.0713 [hep-ph]].
 - [3] See, for example, M. Goodsell and A. Ringwald, *Fortsch. Phys.* **58**, 716 (2010).
 - [4] For example, non-Abelian dark sectors are discussed in F. Chen, J. M. Cline and A. R. Frey, *Phys. Rev. D* **80**, 083516 (2009) [arXiv:0907.4746 [hep-ph]]. M. Baumgart, C. Cheung, J. T. Ruderman, L. -T. Wang and I. Yavin, *JHEP* **0904**, 014 (2009) [arXiv:0901.0283 [hep-ph]]. H. Zhang, C. S. Li, Q. -H. Cao and Z. Li, *Phys. Rev. D* **82**, 075003 (2010).
 - [5] B. Holdom, *Phys. Lett. B* **166**, 196 (1986).
 - [6] M. Pospelov, *Phys. Rev. D* **80**, 095002 (2009) [arXiv:0811.1030 [hep-ph]].
 - [7] J. D. Bjorken, R. Essig, P. Schuster and N. Toro, *Phys. Rev. D* **80**, 075018 (2009).
 - [8] See, for example, Fig. 3 in R. D. McKeown, *AIP Conf. Proc.* **1423**, 289 (2012).
 - [9] H. Merkel, “Search for Dark Photons at the Mainz Microtron”, Talk presented at DARK2012, Rome, October 2012.
 - [10] B. Batell, D. McKeen and M. Pospelov, *Phys. Rev. Lett.* **107**, 011803 (2011).
 - [11] J. P. Leveille, *Nucl. Phys. B* **137**, 63 (1978).
 - [12] J. Beringer *et al.* [Particle Data Group Collaboration], *Phys. Rev. D* **86**, 010001 (2012).
 - [13] G. Altarelli and F. Feruglio, *Rev. Mod. Phys.* **82**, 2701 (2010) [arXiv:1002.0211 [hep-ph]].
 - [14] V. Barger, C.-W. Chiang, W.-Y. Keung and D. Marfatia, *Phys. Rev. Lett.* **108**, 081802 (2012).
 - [15] S. Andreas, C. Niebuhr and A. Ringwald, *Phys. Rev. D* **86**, 095019 (2012).