

On the double-ridge effect at the LHC

S.M. Troshin, N.E. Tyurin

*Institute for High Energy Physics,
Protvino, Moscow Region, 142281, Russia*

Abstract

We discuss a possible explanation of the double-ridge observation in pPb-collisions at the LHC emphasizing that this double structure in the two-particle correlation function can result from the rotation of the transient state of matter.

The ridge-like structure in correlation function of two particles has been observed for the first time at RHIC in the peripheral collisions of nuclei in the near-side jet production (cf. recent paper [1] and references therein for the earlier ones). It was observed that the secondary particles have a narrow $\Delta\phi$ two-particle correlation distribution (where ϕ is an azimuthal angle) and wide distribution over $\Delta\eta$ (η is a pseudorapidity). This phenomenon was called ridge effect and associated usually with the collective properties of a medium produced under interaction.

For the first time, a similar effect was revealed by the CMS Collaboration [2] in pp-collisions and it becomes clear that a deconfined phase formation has the qualitative similarity in pp- and AA-collisions. This feature has been confirmed by the observation of the ridge effect in PbPb-collisions by ALICE, ATLAS and CMS [3, 4, 5]. We emphasize further that significant role belongs to peripheral collision dynamics controlled in AA-collisions by the experimental conditions and in pp-collisions by the transition into the reflective scattering mode [6] at the LHC energies. Of course, there should be a quantitative difference between pp- and AA-collisions and the signal of ridge should be more prominent in AA-interactions.

The peripheral interactions and quark-pion liquid formation in the transient state of interaction [7] would lead to the coherent rotation which can provide explanation for the double-ridge effect observed recently by the ALICE and ATLAS collaboration in pPb-interactions at $\sqrt{s_{NN}} = 5.02$ TeV [8, 9]. Those most recent experimental data probe net effects related to the matter formed under interaction since they were obtained by subtraction of the low multiplicity events (ALICE) or contributions from recoiling dijets (ATLAS). Thus, they provide quantitative estimates of the collective effects contribution.

It should be noted that peripheral mechanism of pPb-interactions is controlled dynamically by the reflective scattering mode which is prominent at the LHC energies [6]. The details will be described further. Now we briefly repeat the main points of the production mechanism proposed in [10]. It is based on the geometry of the overlap region and dynamical properties of the transient state in hadron interaction¹. The above picture assumes that the deconfinement takes place at the initial stage of interaction. The transient state of matter appears then as a rotating medium of massive quarks and pions which hadronize and form multiparticle state at the final stage of interaction. The essential point which is needed for the presence of this rotation is a non-zero impact parameter in the collision. We believe that this qualitative picture seems applicable to pp, pA and AA-interactions. However, the peripheral mechanism of pp-interaction (and of pA-interactions also), as it was already mentioned, is controlled dynamically by

¹Of course, this is not unique mechanism leading to the appearance of the ridge, description of the other ones can be found in the recent review paper [11]

the transition to the reflective scattering mode at the LHC energies. Due to the reflective scattering [6] the inelastic overlap function $h_{inel}(s, b)$,

$$h_{inel}(s, b) \equiv \frac{1}{4\pi} \frac{d\sigma_{inel}}{db^2},$$

has a peripheral impact parameter dependence in the region of $\sqrt{s} > 2$ TeV [6]. Unitarity equation for the elastic amplitude $f(s, b)$ in the impact parameter representation rewritten at high energies has the simple form

$$\text{Im}f(s, b) = h_{el}(s, b) + h_{inel}(s, b) \quad (1)$$

and $h_{inel}(s, b)$ is the sum of all inelastic channel contributions to the unitarity equation. Consider for illustration the case of pure imaginary elastic scattering amplitude, i.e. $f(s, b) \rightarrow if(s, b)$. Then the inelastic overlap function is related to the elastic scattering amplitude by the simple relation

$$h_{inel}(s, b) = f(s, b)(1 - f(s, b)).$$

Reflective scattering mode implies the saturation of the unitarity limit at small impact parameters, e.g. $f(s, b = 0) \rightarrow 1$ at $s \rightarrow \infty$. Thus, the peripheral form of the inelastic overlap function is just an effect of the reflective scattering². According to peripheral impact parameter dependence of $h_{inel}(s, b)$, the mean multiplicity

$$\langle n \rangle(s) = \frac{\int_0^\infty b db \langle n \rangle(s, b) h_{inel}(s, b)}{\int_0^\infty b db h_{inel}(s, b)}$$

obtains the maximal input from the collisions with non-zero impact parameter values. Thus, the events with significant values of multiplicity at the LHC energies would correspond to the peripheral hadron collisions [6]. At such high energies there is a dynamical selection mechanism of peripheral region in impact parameter space. In the nuclear collisions such selection is provided by the relevant experimental conditions.

The geometrical picture of hadron collision at non-zero impact parameters implies [7] that the generated massive virtual quarks in the overlap region could obtain very large initial orbital angular momentum at high energies (Fig. 1).

Due to strong interaction between the quarks this orbital angular momentum leads to the coherent rotation of the quark system located in the overlap region in the xz -plane (Fig. 2). This rotation is similar to the rotation of the liquid where strong correlations between particles momenta exist. Thus, the non-zero

²The elastic scattering amplitude $f(s, b)$ is supposed to be a monotonically decreasing function of the impact parameter




Figure 1: *Overlap region in the initial stage of the hadron interaction, valence constituent quarks are located in the central part of hadrons.*

orbital angular momentum should be realized as a coherent rotation of the quark-pion liquid as a whole. The assumed particle production mechanism at moderate transverse momenta is the simultaneous excitations of the parts of the rotating transient state (of massive constituent quarks interacting by pion exchanges) by the valence constituent quarks. We refer here to proton-proton interactions, in pA-interactions the effects should be enhanced due to many collisions but would remain qualitatively the same.

Since the transient matter is strongly interacting, the excited parts should be located closely to the periphery of the rotating transient state otherwise absorption would not allow quarks and pions leave the interaction region (quenching).

The mechanism is sensitive to the particular direction of rotation and to the rotation plane orientation leads to the narrow distribution of the two-particle correlations in $\Delta\phi$. However, these correlation could have a broad distribution versus polar angle ($\Delta\eta$) (Fig. 2). Quarks in the excited part of the cloud could have different values of the two components of the momentum (with its third component lying in the rotation xz -plane) since the excited region is spatially extended. The rotation with orbital angular momentum $L(s, b)$ contributes to the x -component of the transverse momentum and does not contribute to the y -component of the transverse momentum, i.e.

$$\Delta p_x = \kappa L(s, b) \quad (2)$$

while

$$\Delta p_y = 0. \quad (3)$$

It should be noted that the ALICE and ATLAS are observing the double-ridge structure with two maximums in the two-particle correlation function at $\Delta\phi = 0$ and $\Delta\phi = \pi$ after subtraction of the low-multiplicity events or expected dijet




Figure 2: *Interaction of the valence constituent quarks with rotating quark-pion liquid, dashed lines indicate the clockwise rotation of the transient matter.*

events, respectively. In terms of Fig.2 this observation means the presence of the correlations between particles in the upper cluster ($\Delta\phi = 0$) and between particles from the upper and lower clusters ($\Delta\phi = \pi$). Lower cluster of particles is the result of the excitation of the transient matter by the constituent valence quark from the second colliding hadron. Since both particle clusters originate from the coherent rotating quark-pion liquid, it is evident that the particles from lower and upper clusters should be correlated and this will produce the second observed ridge.

Thus, the double-ridge structure observed in the high multiplicity events by the ALICE and ATLAS can serve as an experimental manifestation of the coherent rotation of the transient state of matter generated during the hadron collisions. The narrowness of the two-particle correlation distribution in the azimuthal angle is the distinctive feature of this mechanism which reflects the presence of the rotation plane in every event of particle production at the LHC.

Other experimentally observed effects of this collective effect are described in [7].

It should be noted that we believe that the nature of the state of matter revealed at the LHC in proton collisions is the same as the nature of the state revealed at RHIC under collisions of nuclei. But, since the LHC energies are significantly higher than the RHIC energies, the differences should be expected due to the energy-dependent dynamics related to the peripheral form of inelastic collisions generated by the reflective scattering mode. There should be no ridge effect in pp- and pA-interactions at RHIC, while the double-ridge effect should be observed in the events of particle production in the peripheral AA-collisions at RHIC and in pp-collisions at the LHC (after subtraction procedure of the low-multiplicity or

dijet events being performed). Evidently, the double-ridge effect should be observed by the CMS experiment also in pPb-collisions with the use of the similar subtractions.

References

- [1] STAR Collaboration, B. I. Abelev et al., Phys. Rev. Lett. **105**, 022301 (2010).
- [2] CMS Collaboration, V. Khachatryan et al., JHEP **1009**, 091 (2010).
- [3] ALICE Collaboration, K. Aamodt et al., Phys. Lett. **B708**, 249 (2012).
- [4] ATLAS Collaboration, G. Aad et al., Phys. Rev. **C86**, 014907 (2012).
- [5] CMS Collaboration, S. Chatrchyan et al., Eur. Phys. J. **C72**, 2012 (2012).
- [6] S.M. Troshin, N.E. Tyurin, Int. J. Mod. Phys. A **22**, 4437 (2007).
- [7] S.M. Troshin, N.E. Tyurin, Int. J. Mod. Phys. A **26**, 4703 (2011).
- [8] ALICE Collaboration, B. Abelev et al., arXiv: 1212.2001v1 [nucl-ex].
- [9] ATLAS Collaboration, G. Aad et al., arXiv: 1212.5198v2 [hep-ex].
- [10] S.M. Troshin, N.E. Tyurin, Int. J. Mod. Phys. E **17**, 1619 (2008).
- [11] Wei Li, Mod. Phys. Lett. A **27**, 1230018 (2012).