

Multi-component dark matter with magnetic moments for Fermi-LAT gamma-ray line

Pei-Hong Gu*

Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

We propose a model of multi-component dark matter with magnetic moments to explain the 130 GeV gamma-ray line hinted by the Fermi-LAT data. Specifically, we consider a $U(1)_X$ dark sector which contains two vector-like fermions besides the related gauge and Higgs fields. A very heavy messenger scalar is further introduced to construct the Yukawa couplings of the dark fermions to the heavy $[SU(2)]$ -singlet leptons in the $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ left-right symmetric models for universal seesaw. A heavier dark fermion with a very long lifetime can mostly decay into a lighter dark fermion and a photon at one-loop level. The dark fermions can serve as the dark matter particles benefited from their annihilations into the dark gauge and Higgs fields. In the presence of a $U(1)$ kinetic mixing, the dark matter fermions can be verified by the ongoing and forthcoming dark matter direct detection experiments.

PACS numbers: 95.35.+d, 12.60.Cn, 12.60.Fr

I. INTRODUCTION

Astronomical and cosmological observations indicate the existence of dark matter in the present universe. Many dark matter candidates have been suggested in various scenarios beyond the $SU(3)_c \times SU(2)_L \times U(1)_Y$ standard model (SM). The dark matter particles can be directly detected through their scatterings off the familiar nucleons and/or indirectly detected through their annihilations/decays into the SM species. Recently, the Fermi-LAT data on the cosmic gamma-ray spectrum from the Galactic center (GC) have revealed a tentative evidence for a line-like feature at an energy around 130 GeV [1–16]. Such monochromatic photon can be induced by a dark matter annihilation or decay. There have been a lot of models [17–53] realizing the required dark matter annihilations or decays.

Usually, we need some particles heavier than the dark matter to mediate the significant annihilations or decays of the dark matter into the monochromatic photons at loop level since the dark matter is only allowed to have an extremely tiny electric charge [54]. On the other hand, in the $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ left-right symmetric models [55] for universal [56] seesaw [57, 58], some heavy $[SU(2)]$ -singlet leptons and quarks are well motivated to alleviate the hierarchy among the SM fermion masses and solve the strong CP problem [59]. Such heavy fermions could take part in mediating the radiative dark matter annihilations or decays for the 130 GeV gamma-ray line. In this paper, we shall demonstrate this possibility in a multi-component dark matter [60–69] scenario. Specifically, we shall introduce a $U(1)_X$ dark sector which contains two vector-like fermions besides the related gauge and Higgs fields. There is also a very heavy messenger scalar having the Yukawa couplings with the dark fermions and the heavy leptons. The heavier dark fermion can have a very long lifetime although it mostly

decays into the lighter dark fermion with a photon. The dark fermions can obtain the desired dark matter relic density through their annihilations into the dark gauge and Higgs fields. As the $U(1)_X$ and $U(1)_{B-L}$ gauge fields are allowed to have a kinetic mixing, the dark matter fermions can be verified by the ongoing and forthcoming dark matter direct detection experiments.

II. THE MODEL

For simplicity, we will not give the full Lagrangian. Instead, we only show the kinetic, mass and Yukawa terms relevant to our demonstration,

$$\begin{aligned} \mathcal{L} \supset & -\frac{1}{4}W_{L\mu\nu}^a W_{L\mu\nu}^a - \frac{1}{4}W_{R\mu\nu}^a W_{R\mu\nu}^a - \frac{1}{4}B_{\mu\nu}B_{\mu\nu} \\ & -\frac{1}{4}C_{\mu\nu}C^{\mu\nu} - \frac{\epsilon}{2}B_{\mu\nu}C^{\mu\nu} + (D_\mu\phi_L)^\dagger D^\mu\phi_L \\ & + (D_\mu\phi_R)^\dagger D^\mu\phi_R + (D_\mu\sigma)^\dagger D^\mu\sigma + (D_\mu\delta)^\dagger D^\mu\delta \\ & + i\bar{q}_L \not{D} q_L + i\bar{q}_R \not{D} q_R + i\bar{l}_L \not{D} l_L + i\bar{l}_R \not{D} l_R \\ & + i\bar{D}_L \not{D} D_L + i\bar{D}_R \not{D} D_R + i\bar{U}_L \not{D} U_L + i\bar{U}_R \not{D} U_R \\ & + i\bar{E}_L \not{D} E_L + i\bar{E}_R \not{D} E_R + i\bar{\chi}_L \not{D} \chi_L + i\bar{\chi}_R \not{D} \chi_R \\ & - M_\delta^2 \delta^\dagger \delta - \bar{D}_L M_D D_R - \bar{U}_L M_U U_R - \bar{E}_L M_E E_R \\ & - \bar{\chi}_L m_\chi \chi_R - \bar{q}_L \phi_L y_D^L D_R - \bar{q}_R \phi_R y_D^R D_L \\ & - \bar{q}_L \tilde{\phi}_L y_U^L U_R - \bar{q}_R \tilde{\phi}_R y_U^R U_L - \bar{l}_L \phi_L y_E^L E_R \\ & - \bar{l}_R \phi_R y_E^R E_L - \delta \bar{E}_L f_R \chi_R - \delta \bar{E}_R f_L \chi_L + \text{H.c.} \end{aligned} \quad (1)$$

Here W_L^a , W_R^a , B and C are the gauge fields associated with the $SU(2)_L$, $SU(2)_R$, $U(1)_{B-L}$ and $U(1)_X$ gauge groups, respectively. The Higgs scalars

$$\phi_L(+1, 0) = \begin{bmatrix} \phi_L^+ \\ \phi_L^0 \end{bmatrix}, \quad \phi_R(+1, 0) = \begin{bmatrix} \phi_R^+ \\ \phi_R^0 \end{bmatrix}, \quad \sigma(0, +2) \quad (2)$$

are an $SU(2)_L$ doublet, an $SU(2)_R$ doublet and an $SU(2)$ singlet, respectively. Here and thereafter the first and second numbers in parentheses are the $U(1)_{B-L}$ charges $B-L$ and the $U(1)_X$ charges X . The messenger scalar δ

*Electronic address: peihong.gu@mpi-hd.mpg.de

is an $SU(2)$ singlet and carries both of the $U(1)_{B-L}$ and $U(1)_X$ charges,

$$\delta(-2, +\frac{2}{3}). \quad (3)$$

Among the fermions, the $[SU(2)]$ -doublet quarks $q_{L,R}$, the $[SU(2)]$ -doublet leptons $l_{L,R}$, the $[SU(2)]$ -singlet quarks $D_{L,R}$ and $U_{L,R}$ as well as the $[SU(2)]$ -singlet leptons $E_{L,R}$ carry the $U(1)_{B-L}$ charges,

$$\begin{aligned} q_L(+\frac{1}{3}, 0) &= \begin{bmatrix} u_L \\ d_L \end{bmatrix}, \quad q_R(+\frac{1}{3}, 0) = \begin{bmatrix} u_R \\ d_R \end{bmatrix}, \\ l_L(-1, 0) &= \begin{bmatrix} \nu_L \\ e_L \end{bmatrix}, \quad l_R(-1, 0) = \begin{bmatrix} \nu_R \\ e_R \end{bmatrix}, \\ D_{L,R}(-\frac{2}{3}, 0), \quad U_{L,R}(+\frac{4}{3}, 0), \quad E_{L,R}(-2, 0), \end{aligned} \quad (4)$$

while the $[SU(2)]$ -singlet dark fermions $\chi_{L,R}$ carry the $U(1)_X$ charges,

$$\chi_{L,R}(0, -\frac{2}{3}). \quad (5)$$

The covariant derivatives are

$$\begin{aligned} D_\mu &= \partial_\mu - ig_X \frac{X}{2} C_\mu - ig_{B-L} \frac{B-L}{2} B_\mu - ig_{I_{3L}} W_{L\mu}^3 \\ &\quad - ig_R I_{3R} W_{R\mu}^3 + \dots, \end{aligned} \quad (6)$$

where we have only written down the diagonal components of the $SU(2)$ gauge fields (W_L^3, W_R^3) which can mix with the other $U(1)$ gauge fields (B, C).

A. Symmetry breaking

The $[SU(2)]$ -doublet Higgs scalars ϕ_R and ϕ_L are responsible for breaking the left-right symmetry down to the electroweak symmetry and then the electromagnetic symmetry, i.e.

$$\begin{aligned} &SU(2)_L \times SU(2)_R \times U(1)_{B-L} \\ &\quad \downarrow \langle \phi_R \rangle \\ &SU(2)_L \times U(1)_Y \\ &\quad \downarrow \langle \phi_L \rangle \\ &U(1)_{em}, \end{aligned} \quad (7)$$

where $\langle \phi_R \rangle$ and $\langle \phi_L \rangle$ are the vacuum expectation values (VEVs),

$$\begin{aligned} \langle \phi_L \rangle &= \langle \phi_L^0 \rangle = \frac{1}{\sqrt{2}} v_L \quad (v_L \simeq 246 \text{ GeV}), \\ \langle \phi_R \rangle &= \langle \phi_R^0 \rangle = \frac{1}{\sqrt{2}} v_R. \end{aligned} \quad (8)$$

As for the dark symmetry $U(1)_X$, it will be broken when the $[SU(2)]$ -singlet Higgs scalar σ develops its VEV,

$$\langle \sigma \rangle = \frac{1}{\sqrt{2}} v_X. \quad (9)$$

Roughly, the above symmetry breakings take place at the temperatures $T = \mathcal{O}(\langle \phi_R \rangle)$, $\mathcal{O}(\langle \phi_L \rangle)$ and $\mathcal{O}(\langle \sigma \rangle)$, respectively.

B. Fermions

The charged fermions have the masses,

$$\begin{aligned} \mathcal{L} \supset & -[\bar{d}_L \bar{D}_L] \begin{bmatrix} 0 & \frac{1}{\sqrt{2}} y_D^L v_L \\ \frac{1}{\sqrt{2}} y_D^{R\dagger} v_R & M_D \end{bmatrix} \begin{bmatrix} d_R \\ D_R \end{bmatrix} \\ & -[\bar{u}_L \bar{U}_L] \begin{bmatrix} 0 & \frac{1}{\sqrt{2}} y_U^L v_L \\ \frac{1}{\sqrt{2}} y_U^{R\dagger} v_R & M_U \end{bmatrix} \begin{bmatrix} u_R \\ U_R \end{bmatrix} \\ & -[\bar{e}_L \bar{E}_L] \begin{bmatrix} 0 & \frac{1}{\sqrt{2}} y_E^L v_L \\ \frac{1}{\sqrt{2}} y_E^{R\dagger} v_R & M_E \end{bmatrix} \begin{bmatrix} e_R \\ E_R \end{bmatrix} + \text{H.c.}, \end{aligned} \quad (10)$$

which can be block diagonalized by

$$\begin{aligned} \mathcal{L} \supset & -\bar{d}_L \left(-y_D^L \frac{v_L v_R}{2M_D} y_D^{R\dagger} \right) d_R - \bar{D}_L M_D D_R \\ & -\bar{u}_L \left(-y_U^L \frac{v_L v_R}{2M_U} y_U^{R\dagger} \right) u_R - \bar{U}_L M_U U_R \\ & -\bar{e}_L \left(-y_E^L \frac{v_L v_R}{2M_E} y_E^{R\dagger} \right) e_R - \bar{E}_L M_E E_R \\ & + \text{H.c.}, \end{aligned} \quad (11)$$

Remarkably, we have the heavy charged fermions besides the SM charged fermions in this universal seesaw scenario.

In the following we will work in the base where the mass matrices of the dark fermions χ and the heavy charged fermions $F = D, U, E$ are diagonal and real, i.e.

$$\begin{aligned} m_\chi &= \text{diag}\{m_{\chi_1}, m_{\chi_2}\}, \\ M_F &= \text{diag}\{M_{F_1}, M_{F_2}, M_{F_3}\}, \end{aligned} \quad (12)$$

and then define the vector-like fermions:

$$\chi_i = \chi_{Li} + \chi_{Ri}, \quad F_a = F_{La} + F_{Ra}. \quad (13)$$

Moreover, we will assume $m_{\chi_1} \leq m_{\chi_2}$ and $M_{F_1} \leq M_{F_2} \leq M_{F_3}$ without loss of generality. Note the left-right symmetry breaking scale and the heavy fermion masses should be large enough to escape from the experimental constraint. For example, the right-handed charged gauge boson should be heavier than a few TeV [70]. In the

present work, we will take $v_R \sim M_{F_{1,2,3}} = \mathcal{O}(10^3 \text{ TeV})$ to give a numerical example.

The present model can accommodate a discrete parity symmetry to solve the strong CP problem without an axion [59, 71]. Furthermore, the neutral neutrinos can have a two-loop induced Dirac mass matrix proportional to the SM charged lepton mass matrix [72]. One can introduce more fermions or scalars to generate the desired neutrino masses and mixing.

C. Gauge fields

We can remove the kinetic mixing between the $U(1)_X$ and $U(1)_{B-L}$ gauge fields by making a non-unitary transformation [73],

$$\begin{aligned} B_\mu &= \tilde{B}_\mu - \frac{\epsilon}{\sqrt{1-\epsilon^2}} \tilde{C}_\mu = \tilde{B}_\mu - \xi \tilde{C}_\mu, \\ C_\mu &= \frac{1}{\sqrt{1-\epsilon^2}} \tilde{C}_\mu, \end{aligned} \quad (14)$$

and then define the orthogonal fields,

$$A_\mu = W_{L\mu}^3 s_W + (W_{R\mu}^3 s_R + \tilde{B}_\mu c_R) c_W, \quad (15a)$$

$$Z_{L\mu} = W_{L\mu}^3 c_W - (W_{R\mu}^3 s_R + \tilde{B}_\mu c_R) s_W, \quad (15b)$$

$$Z_{R\mu} = W_{R\mu}^3 c_R - \tilde{B}_\mu s_R, \quad (15c)$$

$$Z_{X\mu} = \tilde{C}_\mu, \quad (15d)$$

with

$$s_R = \sin \theta_R, \quad c_R = \cos \theta_R \quad \text{for} \quad t_R = \tan \theta_R = \frac{g_{B-L}}{g_R},$$

$$s_W = \sin \theta_W, \quad c_W = \cos \theta_W \quad \text{for}$$

$$t_W = \tan \theta_W = \frac{g_{B-L} g_R / g}{\sqrt{g_{B-L}^2 + g_R^2}} = \frac{g'}{g}. \quad (16)$$

Here g and g' are the SM gauge couplings with $g \simeq 0.653$ and $g' \simeq 0.358$ while θ_W is the Weinberg angle $s_W^2 \simeq 0.231$. If a parity symmetry is imposed, we can determine the unknown gauge couplings g_R and g_{B-L} by

$$g_R = g \simeq 0.653, \quad g_{B-L} = \frac{gg'}{\sqrt{g^2 - g'^2}} \simeq 0.428. \quad (17)$$

Among the orthogonal fields (15), A is the massless photon γ , while Z_L , Z_R and Z_X have the mass terms as below,

$$\begin{aligned} \mathcal{L} \supset & \frac{1}{2} m_{Z_R}^2 (Z_{R\mu} + \xi Z_{X\mu} s_R) (Z_R^\mu + \xi Z_X^\mu s_R) \\ & + \frac{1}{2} m_{Z_L}^2 (Z_{L\mu} + Z_{R\mu} t_R s_W + \xi Z_{X\mu} t_R s_W) \\ & \times (Z_L^\mu + Z_R^\mu t_R s_W + \xi Z_X^\mu t_R s_W) + \frac{1}{2} m_{Z_X}^2 Z_{X\mu} Z_X^\mu \end{aligned}$$

with $\begin{cases} m_{Z_R} = \frac{1}{2 \cos \theta_R} g_R v_R, \\ m_{Z_L} = \frac{1}{2 \cos \theta_W} g v_L \simeq 91 \text{ GeV}, \\ m_{Z_X} = \frac{1}{\sqrt{1-\epsilon^2}} g_X v_X. \end{cases} \quad (18)$

In the following we will focus on the case that the orthogonal fields Z_R , Z_L and Z_X approximate to the mass eigenstates for $m_{Z_R}^2 \gg m_{Z_L}^2 \gg m_{Z_X}^2$ and $\epsilon \ll 1$. In this case, the quasi-mass-eigenstate Z_L is identified to the SM Z boson.

III. DARK GAUGE BOSON DECAY

The dark gauge field C which is mostly the quasi-mass-eigenstate Z_X can couple to the SM fermions besides the dark fermions,

$$\begin{aligned} \mathcal{L} \supset & -\frac{1}{6} \xi g_{B-L} \bar{d} \gamma^\mu d Z_{X\mu} - \frac{1}{6} \xi g_{B-L} \bar{u} \gamma^\mu u Z_{X\mu} \\ & + \frac{1}{2} \xi g_{B-L} \bar{e} \gamma^\mu e Z_{X\mu} + \frac{1}{2} \xi g_{B-L} \bar{\nu} \gamma^\mu \nu Z_{X\mu} \\ & - \frac{1}{3 \sqrt{1-\epsilon^2}} g_X \bar{\chi} \gamma^\mu \chi Z_{X\mu}. \end{aligned} \quad (19)$$

Therefore, if the dark gauge boson Z_X is heavy enough, it can decay into the SM fermion pairs $f\bar{f}$. For $2m_f < m_{Z_X} < 2m_{\chi_1}$, the decay width should be

$$\begin{aligned} \Gamma_{Z_X} &= \sum_f \Gamma_{Z_X \rightarrow f\bar{f}} \\ &\simeq \sum_f N_c \frac{\epsilon^2 g_{B-L}^2}{12\pi} \left(\frac{B-L}{2} \right)^2 m_{Z_X} \left(1 + \frac{m_f^2}{m_{Z_X}^2} \right) \\ &\quad \times \sqrt{1 - 4 \frac{m_f^2}{m_{Z_X}^2}}, \end{aligned} \quad (20)$$

where $(N_c, B-L) = (3, \frac{1}{3})$ for a quark and $(N_c, B-L) = (1, -1)$ for a lepton. We then find

$$\tau_{Z_X} \simeq \left(\frac{1.34 \times 10^{-11}}{\epsilon} \right)^2 \left(\frac{0.428}{g_{B-L}} \right)^2 \left(\frac{500 \text{ MeV}}{m_{Z_X}} \right) \text{ sec} \quad (21)$$

So, the dark gauge boson Z_X with a mass $m_{Z_X} = 500 \text{ MeV}$ can have a lifetime shorter than 1 second if we take $\epsilon > 1.34 \times 10^{-11}$. Currently, the measurement on the muon magnetic moment constrains $\epsilon^2 c_R^2 c_W^2 < 2 \times 10^{-4}$ for $m_{Z_X} = 500 \text{ MeV}$ [74].

IV. DARK MATTER RELIC DENSITY

The dark fermions $\chi_{1,2}$ can annihilate into the dark gauge field C and the dark Higgs field σ . The heavier dark fermion χ_2 can also annihilate into the lighter dark fermion χ_1 . The relevant diagrams are shown in Fig. 1. If the annihilations freeze out before the $U(1)_X$ symmetry breaking at the temperature $T = \mathcal{O}(\langle \sigma \rangle)$, the




FIG. 1: The dark fermion annihilations. Here $\chi_{i,j}$ ($i \neq j$) denotes the dark matter fermions, C is the $U(1)_X$ gauge field, while σ is the Higgs scalar for breaking the $U(1)_X$ symmetry.

annihilation cross sections should be dominated by

$$\begin{aligned} \sigma_{\chi_1} &= \langle \sigma_{\chi_1 \bar{\chi}_1 \rightarrow CC} v_{\text{rel}} \rangle + \langle \sigma_{\chi_1 \bar{\chi}_1 \rightarrow \sigma \bar{\sigma}} v_{\text{rel}} \rangle \\ &\simeq \frac{13}{5184\pi} \frac{g_X^4}{m_{\chi_1}^2}, \\ \sigma_{\chi_2} &= \langle \sigma_{\chi_2 \bar{\chi}_2 \rightarrow CC} v_{\text{rel}} \rangle + \langle \sigma_{\chi_2 \bar{\chi}_2 \rightarrow \sigma \bar{\sigma}} v_{\text{rel}} \rangle \\ &\quad + \langle \sigma_{\chi_2 \bar{\chi}_2 \rightarrow \chi_1 \bar{\chi}_1} v_{\text{rel}} \rangle \\ &\simeq \frac{13}{5184\pi} \frac{g_X^4}{m_{\chi_2}^2} \left[1 + \frac{4}{13} \left(1 + \frac{m_{\chi_1}^2}{2m_{\chi_2}^2} \right) \sqrt{1 - \frac{m_{\chi_1}^2}{m_{\chi_2}^2}} \right]. \end{aligned} \quad (22)$$

Here v_{rel} is the relative velocity between the two annihilating particles in their center-of-mass frame.

As we will clarify later, the heavier dark fermion χ_2 has a very long lifetime and hence contributes to the dark matter relic density together with the lighter and stable dark fermion χ_1 . The relic density of the dark fermion χ_i can be calculated by [75]

$$\Omega_{\chi_i + \bar{\chi}_i} h^2 = \Omega_{\chi_i} h^2 + \Omega_{\bar{\chi}_i} h^2 = \frac{1.07 \times 10^9 m_{\chi_i}}{\sqrt{g_*} M_{\text{Pl}} \sigma_{\chi_i} T_i (\text{GeV})}, \quad (24)$$

where $M_{\text{Pl}} = 1.22 \times 10^{19} \text{ GeV}$ is the Planck mass, T_i is the freeze-out temperature determined by [75]

$$\begin{aligned} \frac{m_{\chi_i}}{T_i} &= \ln(2 \times 0.038 M_{\text{Pl}} m_{\chi_i} \sigma_{\chi_i} / \sqrt{g_*}) \\ &\quad - \frac{1}{2} \ln[\ln(2 \times 0.038 M_{\text{Pl}} m_{\chi_i} \sigma_{\chi_i} / \sqrt{g_*})], \end{aligned} \quad (25)$$

while $g_* = g_*(T_i)$ is the number of relativistic degrees of freedom. The dark fermions $\chi_{1,2}$ can dominate the dark matter relic density for a proper parameter choice. For example, we input

$$\begin{aligned} g_X &= 0.592, \quad v_X = 845 \text{ MeV}, \quad m_{\chi_1} = 20 \text{ GeV}, \\ m_{\chi_2} &= 262 \text{ GeV}, \quad g_*(T_1) = 84, \quad g_*(T_2) = 94, \end{aligned} \quad (26)$$

to obtain

$$\begin{aligned} \sigma_{\chi_1} &\simeq 2.45 \times 10^{-7} \text{ GeV}^{-2}, \quad T_1 \simeq m_{\chi_1}/25.3, \\ \Omega_{\chi_1 + \bar{\chi}_1} h^2 &\simeq 0.001, \end{aligned} \quad (27)$$

and

$$\begin{aligned} \sigma_{\chi_2} &\simeq 1.87 \times 10^{-9} \text{ GeV}^{-2}, \quad T_2 \simeq m_{\chi_2}/23.0, \\ \Omega_{\chi_2 + \bar{\chi}_2} h^2 &\simeq 0.111. \end{aligned} \quad (28)$$

The total dark matter relic density thus should be

$$\Omega_{\text{DM}} h^2 = \Omega_{\chi_1 + \bar{\chi}_1} h^2 + \Omega_{\chi_2 + \bar{\chi}_2} h^2 \simeq 0.112, \quad (29)$$

which is well consistent with the observations [76]. In the above numerical estimation, the heavier dark fermion χ_2 rather than the lighter dark fermion χ_1 dominates the dark matter relic density, i.e.

$$\frac{\Omega_{\chi_1 + \bar{\chi}_1} h^2}{\Omega_{\text{DM}} h^2} \simeq 0.9\%, \quad \frac{\Omega_{\chi_2 + \bar{\chi}_2} h^2}{\Omega_{\text{DM}} h^2} \simeq 99.1\%, \quad (30)$$

as a result of their hierarchial masses $m_{\chi_2}^2 \gg m_{\chi_1}^2$. Actually, it is easy to see

$$\frac{\Omega_{\chi_1 + \bar{\chi}_1} h^2}{\Omega_{\text{DM}} h^2} \leq 50\% \leq \frac{\Omega_{\chi_2 + \bar{\chi}_2} h^2}{\Omega_{\text{DM}} h^2} \quad \text{for } m_{\chi_1} \leq m_{\chi_2}. \quad (31)$$

V. DARK MATTER DECAY

The dark matter fermions $\chi_{1,2}$ can couple to the $U(1)_{B-L}$ and $U(1)_X$ field strength tensors ($B_{\mu\nu}, C_{\mu\nu}$) at one-loop level. We show the relevant diagrams in Fig. 2. For $m_{\chi_{1,2}} \ll M_{E_a}, M_{\delta}$, the effective interactions should




FIG. 2: The dark matter fermions $\chi_{i,j}$ couple to the $U(1)_{B-L}$ and $U(1)_X$ field strength tensors $(B_{\mu\nu}, C_{\mu\nu})$ at one-loop level.

be

$$\begin{aligned} \mathcal{L} \supset & -\lambda_{ij} \bar{\chi}_{Li} \sigma^{\mu\nu} \chi_{Rj} B_{\mu\nu} - \kappa_{ij} \bar{\chi}_{Li} \sigma^{\mu\nu} \chi_{Rj} C_{\mu\nu} + \text{H.c.} \\ = & -\lambda_{ij} \bar{\chi}_{Li} \sigma^{\mu\nu} \chi_{Rj} (A_{\mu\nu} c_W c_R - Z_{L\mu\nu} s_W c_R \\ & - Z_{R\mu\nu} s_R - \xi Z_{X\mu\nu}) - \frac{1}{\sqrt{1-\epsilon^2}} \kappa_{ij} \bar{\chi}_{Li} \sigma^{\mu\nu} \chi_{Rj} Z_{X\mu\nu} \\ & + \text{H.c.}, \end{aligned} \quad (32)$$

where the couplings λ_{ij} and κ_{ij} are given by

$$\begin{aligned} \lambda_{ij} &= \frac{g_{B-L}}{32\pi^2} (f_L^\dagger)_{ia} (f_R)_{aj} \frac{1}{M_{E_a}} F_B \left(\frac{M_\delta^2}{M_{E_a}^2} \right) \quad \text{with} \\ F_B(x) &= \frac{1}{1-x} + \frac{x}{(1-x)^2} \ln x, \\ \kappa_{ij} &= \frac{g_X}{96\pi^2} (f_L^\dagger)_{ia} (f_R)_{aj} \frac{1}{M_{E_a}} F_C \left(\frac{M_\delta^2}{M_{E_a}^2} \right) \quad \text{with} \\ F_C(x) &= -\frac{1+x}{2(1-x)^2} - \frac{x}{(1-x)^3} \ln x. \end{aligned} \quad (33)$$

Therefore, the heavier dark matter fermion χ_2 can decay into the lighter dark matter fermion χ_1 and a gauge boson (γ , Z , Z_X or Z_R) as long as the kinematics is allowed. Provided that

$$m_{Z_R} \gg m_{\chi_2} - m_{\chi_1} > m_Z, m_{Z_X}, \quad (35)$$

we can obtain the decay widths,

$$\begin{aligned} \Gamma_{\chi_2 \rightarrow \chi_1 \gamma} &= \Gamma_{\bar{\chi}_2 \rightarrow \bar{\chi}_1 \gamma} \\ &\simeq \frac{c_R^2 c_W^2}{2\pi} |\lambda_{12}|^2 m_{\chi_2}^3 (1 - m_{\chi_1}^2/m_{\chi_2}^2)^3, \end{aligned} \quad (36)$$




FIG. 3: The heavier dark matter fermion χ_j ($\bar{\chi}_i$) decays into the lighter dark matter fermion χ_i ($\bar{\chi}_j$) and a charged lepton pair $e_\alpha^+ e_\beta^-$.

and

$$\begin{aligned} \Gamma_{\chi_2 \rightarrow \chi_1 Z} &= \Gamma_{\bar{\chi}_2 \rightarrow \bar{\chi}_1 Z} \\ &\simeq \frac{c_R^2 s_W^2}{2\pi} |\lambda_{12}|^2 m_{\chi_2}^3 F_2 \left(\frac{m_{\chi_1}^2}{m_{\chi_2}^2}, \frac{m_Z^2}{m_{\chi_2}^2} \right), \\ \Gamma_{\chi_2 \rightarrow \chi_1 Z_X} &= \Gamma_{\bar{\chi}_2 \rightarrow \bar{\chi}_1 Z_X} \\ &\simeq \frac{1}{2\pi} |\kappa_{12}|^2 m_{\chi_2}^3 F_2 \left(\frac{m_{\chi_1}^2}{m_{\chi_2}^2}, \frac{m_{Z_X}^2}{m_{\chi_2}^2} \right) \quad \text{with} \\ F_2(x, y) &= \left[(1-x)^2 - \frac{1}{2}y(1+x+y) \right] \\ &\quad \times \sqrt{(1-x-y)^2 - 4xy}. \end{aligned} \quad (37)$$

In Fig. 3, we see the heavier dark matter fermion χ_2 can also decay into the lighter dark matter fermion χ_1 and a pair of the SM charged leptons $e_\alpha^+ e_\beta^-$ at tree level. These three-body decay modes have the decay widths as

below,

$$\begin{aligned}\Gamma_{\chi_2 \rightarrow \chi_1 l^+ l^-} &= \Gamma_{\bar{\chi}_2 \rightarrow \bar{\chi}_1 l^+ l^-} = \sum_{\alpha, \beta} \Gamma_{\chi_2 \rightarrow \chi_1 e_\alpha^+ e_\beta^-} \\ &\simeq \frac{1}{29\pi^3} (R_{11} R_{22} + R_{11} L_{22} + R_{22} L_{11} \\ &\quad + L_{11} L_{22}) \frac{m_\delta^5}{M_\delta^4} F_3 \left(\frac{m_{\chi_1}^2}{m_{\chi_2}^2} \right),\end{aligned}\quad (38)$$

with

$$\begin{aligned}R_{ii} &= \left(f_L^\dagger \frac{v_R}{\sqrt{2}M_E} y_E^{R\dagger} y_E^R \frac{v_R}{\sqrt{2}M_E} f_L \right)_{ii}, \\ L_{ii} &= \left(f_R^\dagger \frac{v_L}{\sqrt{2}M_E} y_E^{L\dagger} y_E^L \frac{v_L}{\sqrt{2}M_E} f_R \right)_{ii},\end{aligned}\quad (39)$$

and

$$F_3(x) = \frac{1}{12} (1 - 8x + 8x^3 - x^4) - x^2 \ln x. \quad (40)$$

The heavier dark matter fermion χ_2 can have a very long lifetime. For example, by inputting

$$\begin{aligned}g_{B-L} &= 0.428, \quad g_X = 0.592, \quad \epsilon = 10^{-7}, \\ f_L = f_R &= \text{diag} \left\{ \sqrt{\frac{2m_e}{v_L}}, \sqrt{\frac{2m_\mu}{v_L}}, \sqrt{\frac{2m_\tau}{v_L}} \right\}, \\ M_{E_{1,2,3}} &= \frac{1}{\sqrt{2}} v_R = 900 \text{ TeV}, \quad M_\delta = 10^{16} \text{ GeV}, \\ m_{\chi_2} &= 262 \text{ GeV}, \quad m_{\chi_1} = 20 \text{ GeV},\end{aligned}\quad (41)$$

we can obtain

$$\begin{aligned}\Gamma_{\chi_2 \rightarrow \chi_1 \gamma} &= \Gamma_{\bar{\chi}_2 \rightarrow \bar{\chi}_1 \gamma} = 5.03 \times 10^{-53} \text{ GeV}, \\ \Gamma_{\chi_2 \rightarrow \chi_1 Z} &= \Gamma_{\bar{\chi}_2 \rightarrow \bar{\chi}_1 Z} = 1.23 \times 10^{-53} \text{ GeV}, \\ \Gamma_{\chi_2 \rightarrow \chi_1 Z_X} &= \Gamma_{\bar{\chi}_2 \rightarrow \bar{\chi}_1 Z_X} = 2.22 \times 10^{-57} \text{ GeV}, \\ \Gamma_{\chi_2 \rightarrow \chi_1 l^+ l^-} &= \Gamma_{\bar{\chi}_2 \rightarrow \bar{\chi}_1 l^+ l^-} \\ &= 6.20 \times 10^{-66} \text{ GeV},\end{aligned}\quad (42)$$

and then

$$\begin{aligned}\text{Br}_{\chi_1 \gamma} &= \text{Br}(\chi_2 \rightarrow \chi_1 \gamma) + \text{Br}(\bar{\chi}_2 \rightarrow \bar{\chi}_1 \gamma) \\ &= \frac{\Gamma_{\chi_2 \rightarrow \chi_1 \gamma} + \Gamma_{\bar{\chi}_2 \rightarrow \bar{\chi}_1 \gamma}}{\Gamma_{\chi_2} + \Gamma_{\bar{\chi}_2}} \simeq 80.3\%, \\ \tau_{\chi_2} &= \tau_{\bar{\chi}_2} = \frac{1}{\Gamma_{\chi_2}} \simeq 1.05 \times 10^{28} \text{ sec},\end{aligned}\quad (43)$$

with Γ_{χ_2} and $\Gamma_{\bar{\chi}_2}$ being the total decay width,

$$\begin{aligned}\Gamma_{\chi_2} = \Gamma_{\bar{\chi}_2} &= \Gamma_{\chi_2 \rightarrow \chi_1 \gamma} + \Gamma_{\chi_2 \rightarrow \chi_1 Z} + \Gamma_{\chi_2 \rightarrow \chi_1 Z_X} \\ &\quad + \Gamma_{\chi_2 \rightarrow \chi_1 l^+ l^-} \\ &\simeq \Gamma_{\chi_2 \rightarrow \chi_1 \gamma} + \Gamma_{\chi_2 \rightarrow \chi_1 Z} \\ &= 6.26 \times 10^{-53} \text{ GeV},\end{aligned}\quad (44)$$

Furthermore, the photons from the decays $\chi_2 \rightarrow \chi_1 \gamma$ and $\bar{\chi}_2 \rightarrow \bar{\chi}_1 \gamma$ have the determined energy

$$E_\gamma = \frac{m_{\chi_2}^2 - m_{\chi_1}^2}{2m_{\chi_2}} \simeq 130 \text{ GeV}. \quad (45)$$

The gamma-ray flux from the dark matter decays $\chi_2 \rightarrow \chi_1 \gamma$ and $\bar{\chi}_2 \rightarrow \bar{\chi}_1 \gamma$ can be written as

$$\frac{d\Phi}{dE d\Omega} = \frac{\Omega_{\chi_2 + \bar{\chi}_2} h^2}{\Omega_{\text{DM}} h^2} \frac{\text{Br}_{\chi_1 \gamma}}{4\pi m_{\chi_2} \tau_{\chi_2}} \int_{l.o.s} ds \rho[r(s, \psi)] \frac{dN}{dE}, \quad (46)$$

where dN/dE is the differential gamma spectrum per dark matter decay with E being the gamma-ray energy, $r(s, \psi) = (r_\odot^2 + s^2 - 2r_\odot s \cos \psi)^{1/2}$ is the coordinate centered on the GC with s being the distance from the Sun along the line-of-sight (l.o.s), $r_\odot$ being the distance from the Sun to the GC and ψ being the angle between the direction of observation in the sky and the GC, $\rho[r(s, \psi)]$ is the dark matter density profile. In Ref. [27], the authors have shown that the decay of a single-component dark matter fermion into a neutrino and a photon can explain the gamma-ray line in the Fermi-LAT data [27], i.e.

$$\frac{d\Phi}{dE d\Omega} = \frac{\text{Br}_{\nu \gamma}}{4\pi m_\chi \tau_\chi} \int_{l.o.s} ds \rho[r(s, \psi)] \frac{dN}{dE}. \quad (47)$$

Here

$$m_\chi \simeq 2E_\gamma \simeq 260 \text{ GeV}, \quad (48)$$

is the dark matter mass, τ_χ is the dark matter lifetime, $\text{Br}_{\nu \gamma}$ is the branching ratio of the decay modes $\chi \rightarrow \nu \gamma$. By comparing Eqs. (45-46) and (47-48), we can take

$$\begin{aligned}\frac{m_{\chi_2}^2 - m_{\chi_1}^2}{2m_{\chi_2}} &= m_\chi, \quad \text{Br}_{\chi_1 \gamma} = \text{Br}_{\nu \gamma}, \\ \tau_{\chi_2} &= \frac{\Omega_{\chi_2 + \bar{\chi}_2} h^2}{\Omega_{\text{DM}} h^2} \frac{m_\chi}{m_{\chi_2}} \tau_\chi,\end{aligned}\quad (49)$$

to account for the fitting results [27]. For example, we can obtain the Fermi-LAT gamma-ray line by inputting

$$\begin{aligned}m_{\chi_2} &= 262 \text{ GeV}, \quad m_{\chi_1} = 20 \text{ GeV}, \\ \frac{\Omega_{\chi_2 + \bar{\chi}_2} h^2}{\Omega_{\text{DM}} h^2} &= 99.1\%, \quad \text{Br}_{\chi_1 \gamma} = 80.3\%, \\ \tau_{\chi_2} &= 1.05 \times 10^{28} \text{ sec}.\end{aligned}\quad (50)$$

VI. DARK MATTER SCATTERING AND SELF-INTERACTION

As shown in Fig. 4, the dark matter fermions $\chi_{1,2}$ can scatter off the nucleon N through the kinetic mixing between the $U(1)_X$ and $U(1)_{B-L}$ gauge fields. The elastic




FIG. 4: The dark matter scattering $\chi_i N \rightarrow \chi_i N$. Here χ_i and N denote the dark matter fermions and the ordinary nucleons while B and C are the $U(1)_{B-L}$ and $U(1)_X$ gauge fields. For simplicity, we don't show the scattering $\bar{\chi}_i N \rightarrow \bar{\chi}_i N$.

scattering cross section is computed by

$$\begin{aligned} \sigma_{\chi_i N} &= \sigma_{\chi_i N \rightarrow \chi_i N} = \sigma_{\bar{\chi}_i N \rightarrow \bar{\chi}_i N} \simeq \frac{\epsilon^2 g_X^2 g_{B-L}^2}{324\pi} \frac{\mu_r^2}{m_{Z_X}^4} \\ &= 3.9 \times 10^{-45} \text{ cm}^2 \left(\frac{\epsilon}{10^{-7}} \right)^2 \left(\frac{g_X}{0.592} \right)^2 \\ &\quad \times \left(\frac{g_{B-L}}{0.428} \right)^2 \left(\frac{500 \text{ MeV}}{m_{Z_X}} \right)^4 \left(\frac{\mu_r}{1 \text{ GeV}} \right)^2. \end{aligned} \quad (51)$$

Here we have defined the reduced mass,

$$\mu_r = \frac{m_{\chi_i} m_N}{m_{\chi_i} + m_N} \simeq m_N \simeq 1 \text{ GeV} \quad \text{for } m_{\chi_i} \gg m_N. \quad (52)$$

Such scattering can be measured by the dark matter direct detection experiments. The event rate per unit time per nucleon should be

$$R_{\chi_i} \approx \frac{\Omega_{\chi_i + \bar{\chi}_i} h^2}{\Omega_{\text{DM}} h^2} \frac{\rho_\odot}{m_{\chi_i}} \sigma_{\chi_i N} \quad (53)$$

with $\rho_\odot$ being the local dark matter density. Currently, the measured experimental rate is given in the single-component dark matter hypothesis,

$$R_\chi \approx \frac{\rho_\odot}{m_\chi} \sigma_{\chi N} \quad (54)$$

where $\sigma_{\chi N}$ denotes the cross section of a single-component dark matter χ with the mass m_χ scattering off the nucleon N . The XENON10 and XENON100 experiments [77, 78] have stringently put an upper bound $\sigma_{\chi N}^{\text{exp}}$ on the cross section $\sigma_{\chi N}$ for a given mass m_χ . Therefore, we should constrain

$$\sigma_{\chi_i N} < \frac{\Omega_{\text{DM}} h^2}{\Omega_{\chi_i + \bar{\chi}_i} h^2} \sigma_{\chi N}^{\text{exp}}. \quad (55)$$

For a proper parameter choice such as

$$\begin{aligned} m_{\chi_1} &= 20 \text{ GeV}, \quad m_{\chi_2} = 262 \text{ GeV}, \\ \sigma_{\chi_1 N} &= \sigma_{\chi_2 N} = 3.9 \times 10^{-45} \text{ cm}^2, \\ \frac{\Omega_{\chi_1 + \bar{\chi}_1} h^2}{\Omega_{\text{DM}} h^2} &\simeq 0.9\%, \quad \frac{\Omega_{\chi_2 + \bar{\chi}_2} h^2}{\Omega_{\text{DM}} h^2} \simeq 99.1\%, \end{aligned} \quad (56)$$

the dark matter fermions $\chi_{1,2}$ can be verified by the on-going and forthcoming dark matter direct detection experiments.

Furthermore, the dark matter fermions $\chi_{1,2}$ can have a self-interaction as shown in Fig. 5. For $m_{\chi_i} \gg m_{Z_X}$, the self-interacting cross section should be

$$\begin{aligned} \frac{\sigma_{\chi_i \chi_i \rightarrow \chi_i \chi_i}}{m_{\chi_i}} &= \frac{\sigma_{\bar{\chi}_i \bar{\chi}_i \rightarrow \bar{\chi}_i \bar{\chi}_i}}{m_{\chi_i}} \simeq \frac{g_X^4}{628\pi} \frac{m_{\chi_i}}{m_{Z_X}^4}, \\ \frac{\sigma_{\chi_i \bar{\chi}_i \rightarrow \chi_i \bar{\chi}_i}}{m_{\chi_i}} &\simeq \frac{g_X^4}{324\pi} \frac{m_{\chi_i}}{m_{Z_X}^4}, \end{aligned} \quad (57)$$

from which we read

$$\begin{aligned} \frac{\sigma_{\chi_1 \bar{\chi}_1 \rightarrow \chi_1 \bar{\chi}_1}}{m_{\chi_1}} &\simeq 2 \frac{\sigma_{\chi_1 \chi_1 \rightarrow \chi_1 \chi_1}}{m_{\chi_1}} = 2 \frac{\sigma_{\bar{\chi}_1 \bar{\chi}_1 \rightarrow \bar{\chi}_1 \bar{\chi}_1}}{m_{\chi_1}} \\ &\simeq 3.0 \times 10^{-43} \text{ cm}^3 \left(\frac{g_X}{0.592} \right)^4 \\ &\quad \times \left(\frac{m_{\chi_1}}{20 \text{ GeV}} \right) \left(\frac{500 \text{ MeV}}{m_{Z_X}} \right)^4, \\ \frac{\sigma_{\chi_2 \bar{\chi}_2 \rightarrow \chi_2 \bar{\chi}_2}}{m_{\chi_2}} &\simeq 2 \frac{\sigma_{\chi_2 \chi_2 \rightarrow \chi_2 \chi_2}}{m_{\chi_2}} = 2 \frac{\sigma_{\bar{\chi}_2 \bar{\chi}_2 \rightarrow \bar{\chi}_2 \bar{\chi}_2}}{m_{\chi_2}} \\ &\simeq 3.9 \times 10^{-42} \text{ cm}^3 \left(\frac{g_X}{0.592} \right)^4 \\ &\quad \times \left(\frac{m_{\chi_2}}{262 \text{ GeV}} \right) \left(\frac{500 \text{ MeV}}{m_{Z_X}} \right)^4. \end{aligned} \quad (58)$$

For a single-component dark matter with the mass m , its self-interacting cross section σ has an upper bound $\sigma/m < 4.4 \times 10^{-42} \text{ cm}^3$ [79]. We hence should require

$$\begin{aligned} \frac{\Omega_{\chi_i} h^2}{\Omega_{\text{DM}} h^2} \frac{\sigma_{\chi_i \chi_i \rightarrow \chi_i \chi_i}}{m_{\chi_i}} &= \frac{\Omega_{\chi_i + \bar{\chi}_i} h^2}{2\Omega_{\text{DM}} h^2} \frac{\sigma_{\chi_i \chi_i \rightarrow \chi_i \chi_i}}{m_{\chi_i}}, \\ \frac{\Omega_{\bar{\chi}_i} h^2}{\Omega_{\text{DM}} h^2} \frac{\sigma_{\bar{\chi}_i \bar{\chi}_i \rightarrow \bar{\chi}_i \bar{\chi}_i}}{m_{\chi_i}} &= \frac{\Omega_{\chi_i + \bar{\chi}_i} h^2}{2\Omega_{\text{DM}} h^2} \frac{\sigma_{\bar{\chi}_i \bar{\chi}_i \rightarrow \bar{\chi}_i \bar{\chi}_i}}{m_{\chi_i}}, \\ \frac{\Omega_{\chi_i + \bar{\chi}_i} h^2}{\Omega_{\text{DM}} h^2} \frac{\sigma_{\chi_i \bar{\chi}_i \rightarrow \chi_i \bar{\chi}_i}}{m_{\chi_i}} &< 4.4 \times 10^{-42} \text{ cm}^3. \end{aligned} \quad (59)$$

Clearly, the parameter choice (58) can satisfy the limit (59) since we have $\frac{\Omega_{\chi_i + \bar{\chi}_i} h^2}{\Omega_{\text{DM}} h^2} < 1$.




FIG. 5: The dark matter self-interactions $\chi_i \chi_i \rightarrow \chi_i \chi_i$ and $\chi_i \bar{\chi}_i \rightarrow \chi_i \bar{\chi}_i$. Here χ_i denotes the dark matter fermions while C is the $U(1)_X$ gauge field. For simplicity, we don't show the processes $\bar{\chi}_i \bar{\chi}_i \rightarrow \bar{\chi}_i \bar{\chi}_i$.

VII. SUMMARY

It was suggested that the 130 GeV gamma-ray line hinted by the Fermi-LAT data could be understood by the dark matter annihilation or decay into monochromatic photons. We hence propose a multi-component dark matter model, where a heavier dark matter fermion mostly decays into a lighter dark matter fermion and a photon, to explain the Fermi-LAT gamma-ray line. In our model, the neutral dark matter fermions have a highly suppressed magnetic moment at one-loop level because of their Yukawa couplings to a charged scalar and three non-SM leptons. The new scalar besides the

dark matter fermions is gauged by a $U(1)_X$ symmetry which will be spontaneously broken below the GeV scale. As for the non-SM leptons, they play an essential role for generating the SM lepton masses in the $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ left-right symmetric models for the universal seesaw scenario where the strong CP problem can be solved without an axion. The dark matter fermions can obtain a thermally produced relic density through their annihilations into the $U(1)_X$ gauge and Higgs fields. The kinetic mixing between the $U(1)_X$ and $U(1)_{B-L}$ gauge fields can result in a testable dark matter scattering.

-
- [1] T. Bringmann, X. Huang, A. Ibarra, S. Vogl, and C. Weniger, JCAP **1207**, 054 (2012).
 - [2] C. Weniger, JCAP **1208**, 007 (2012).
 - [3] E. Tempel, A. Hektor, and M. Raidal, JCAP **1209**, 032 (2012).
 - [4] A. Geringer-Sameth and S.M. Koushiappas, Phys. Rev. D **86**, 021302(R) (2012).
 - [5] M. Su and D.P. Finkbeiner, arXiv:1206.1616 [astro-ph.HE].
 - [6] Y. Li and Q. Yuan, Phys. Lett. B **715**, 35 (2012).
 - [7] T. Cohen, M. Lisanti, T.R. Slatyer, and J.G. Wacker, arXiv:1207.0800 [hep-ph].
 - [8] I. Cholis, M. Tavakoli, and P. Ullio, Phys. Rev. D **86**, 083525 (2012).
 - [9] R.Z. Yang, Q. Yuan, L. Feng, Y.Z. Fan, and J. Chang, Phys. Lett. B **715**, 285 (2012).
 - [10] A. Hektor, M. Raidal, and E. Tempel, arXiv:1207.4466 [astro-ph.HE]; arXiv:1209.4548 [astro-ph.HE].
 - [11] D.P. Finkbeiner, M. Su, and C. Weniger, arXiv:1209.4562 [astro-ph.HE].
 - [12] L. Bergström, G. Bertone, J. Conrad, C. Farnier, and C. Weniger, arXiv:1207.6773 [hep-ph].
 - [13] X. Huang, Q. Yuan, P.F. Yin, X.J. Bi, and X. Chen, JCAP **1211**, 048 (2012).
 - [14] D. Hooper and T. Linden, arXiv:1208.0828 [hep-ph].
 - [15] D. Whiteson, JCAP **1211**, 008 (2012).
 - [16] A. Hektor, M. Raidal, and E. Tempel, arXiv:1209.4548 [astro-ph.HE].
 - [17] E. Dudas, Y. Mambrini, S. Pokorski, and A. Romagnoni, arXiv:1205.1520 [hep-ph].
 - [18] J.M. Cline, arXiv:1205.2688 [hep-ph].
 - [19] K.Y. Choi and O. Seto, arXiv:1205.3276 [hep-ph].
 - [20] A. Rajaraman, T.M.P. Tait, and D. Whiteson, JCAP **1209**, 003 (2012).
 - [21] B. Kyae and J.C. Park, Phys. Lett. B **718**, 1425 (2013).
 - [22] H.M. Lee, M. Park, and W.I. Park, arXiv:1205.4675 [hep-ph].
 - [23] M.R. Buckley, D. Hooper, arXiv:1205.6811 [hep-ph].
 - [24] X. Chu, T. Hambye, T. Scarna, and M.H.G. Tytgat, arXiv:1206.2279 [hep-ph].
 - [25] D. Das, U. Ellwanger, and P. Mitropoulos, arXiv:1206.2639 [hep-ph].
 - [26] Z. Kang, T. Li, J. Li, and Y. Liu, arXiv:1206.2863 [hep-ph].
 - [27] W. Buchmüller and M. Garny, arXiv:1206.7056 [hep-ph].
 - [28] J.C. Park and S.C. Park, arXiv:1207.4981 [hep-ph].
 - [29] M.T. Frandsen, U. Haisch, F. Kahlhoefer, P. Mertsch, and K. Schmidt-Hoberg, JCAP **1210**, 033 (2012).

- [30] S. Tulin, H.B. Yu, and K.M. Zurek, arXiv:1208.0009 [hep-ph].
- [31] J.M. Cline, A.R. Frey, and G.D. Moore, arXiv:1208.2685 [hep-ph].
- [32] Y. Bai and J. Shelton, arXiv:1208.4100 [hep-ph].
- [33] M. Kuhlen, J. Guedes, A. Pillepich, P. Madau, and L. Mayer, arXiv:1208.4844 [astro-ph.GA].
- [34] R. Laha, K.C.Y. Ng, B. Dasgupta, and S. Horiuchi, arXiv:1208.5488 [astro-ph.CO].
- [35] L. Bergström, arXiv:1208.6082 [hep-ph].
- [36] L. Wang and X.F. Han, arXiv:1209.0376 [hep-ph].
- [37] J. Fan and M. Reece, arXiv:1209.1097 [hep-ph].
- [38] N. Weiner and I. Yavin, arXiv:1209.1093 [hep-ph].
- [39] H.M. Lee, M. Park, and W.I. Park, arXiv:1209.1955 [hep-ph].
- [40] B. Shakya, arXiv:1209.2427 [hep-ph].
- [41] S. Baek, P. Ko, and E. Senaha, arXiv:1209.1685 [hep-ph].
- [42] M. Cvetič, J. Halverson, and H. Piragua, arXiv:1210.5245 [hep-ph].
- [43] F. D'Eramo, M. McCullough, and J. Thaler, arXiv:1210.7817 [hep-ph].
- [44] K. Schmidt-Hoberg, F. Staub, and M.W. Winkler, arXiv:1211.2835 [hep-ph].
- [45] Y. Farzan and A. Rezaei Akbarieh, arXiv:1211.4685 [hep-ph].
- [46] G. Chalons, M.J. Dolan, and C. McCabe, arXiv:1211.5154 [hep-ph].
- [47] A. Rajaraman, T.M.P. Tait, and A.M. Wijangco, arXiv:1211.7061 [hep-ph].
- [48] Y. Bai, M. Su, and Y. Zhao, arXiv:1212.0864 [hep-ph].
- [49] Y. Zhang, arXiv:1212.2730 [hep-ph].
- [50] Y. Bai, V. Barger, L.L. Everett, and G. Shaughnessy, arXiv:1212.5604 [hep-ph].
- [51] H.M. Lee, M. Park, and V. Sanz, arXiv:1212.5647 [hep-ph].
- [52] J. Kopp, E.T. Neil, R. Primulando, and J. Zupan, arXiv:1301.1683 [hep-ph].
- [53] A. Biswas, D. Majumdar, A. Sil, and P. Bhattacharjee, arXiv:1301.3668 [hep-ph].
- [54] S.D. McDermott, H.B. Yu, and K.M. Zurek, Phys. Rev. D **85**, 023519 (2012).
- [55] J.C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974); R.N. Mohapatra and J.C. Pati, *ibid.* **11**, 566 (1975); R.N. Mohapatra and J.C. Pati, *ibid.* **11**, 2558 (1975); R.N. Mohapatra and G. Senjanović, *ibid.* **12**, 1502 (1975).
- [56] Z.G. Berezhiani, Phys. Lett. B **129**, 99 (1983); D. Chang and R.N. Mohapatra, Phys. Rev. Lett. **58**, 1600 (1987); S. Rajpoot, Phys. Lett. B **191**, 122 (1987); A. Davidson and K.C. Wali, Phys. Rev. Lett. **59**, 393 (1987).
- [57] P. Minkowski, Phys. Lett. B **67**, 421 (1977); T. Yanagida, in *Proceedings of the Workshop on Unified Theory and the Baryon Number of the Universe*, edited by O. Sawada and A. Sugamoto (KEK, Tsukuba, 1979), p. 95; M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by F. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979), p. 315; S.L. Glashow, in *Quarks and Leptons*, edited by M. Lévy *et al.* (Plenum, New York, 1980), p. 707; R.N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).
- [58] M. Magg and C. Wetterich, Phys. Lett. B **94**, 61 (1980); J. Schechter and J.W.F. Valle, Phys. Rev. D **22**, 2227 (1980); T.P. Cheng and L.F. Li, Phys. Rev. D **22**, 2860 (1980); G. Lazarides, Q. Shafi, and C. Wetterich, Nucl. Phys. B **181**, 287 (1981); R.N. Mohapatra and G. Senjanović, Phys. Rev. D **23**, 165 (1981).
- [59] K.S. Babu and R.N. Mohapatra, Phys. Rev. Lett. **62**, 1079 (1989).
- [60] Z.G. Berezhiani and M.Yu. Khlopov, Z. Phys. C **49**, 73 (1991).
- [61] C. Boehm, P. Fayet, and J. Silk, Phys. Rev. D **69**, 101302 (2004).
- [62] E. Ma, Annales Fond. Broglie **31**, 285 (2006).
- [63] P.H. Gu, Phys. Lett. B **661**, 290 (2008).
- [64] T. Hur, H.S. Lee, and S. Nasri, Phys. Rev. D **77**, 015008 (2007).
- [65] Q.H. Cao, E. Ma, J. Wudka, and C.P. Yuan, arXiv:0711.3881 [hep-ph].
- [66] T. Hambye, JHEP **0901**, 028 (2009).
- [67] P.H. Gu, H.J. He, U. Sarkar, and X. Zhang, Phys. Rev. D **80**, 053004 (2009).
- [68] M. Aoki, M. Duerr, J. Kubo, and H. Takano, arXiv:1207.3318 [hep-ph].
- [69] J. Heeck and H. Zhang, arXiv:1211.0538 [hep-ph].
- [70] Y. Zhang, H. An, X. Ji, and R.N. Mohapatra, Nucl. Phys. B **802**, 247 (2008).
- [71] S.M. Barr, D. Chang, and G. Senjanović, Phys. Rev. Lett. **67**, 2765 (1991); L. Lavoura, Phys. Lett. B **400**, 152 (1997).
- [72] R.N. Mohapatra, Phys. Lett. B **210**, 517 (1988); K.S. Babu and X.G. He, Mod. Phys. Lett. A **4**, 61 (1989).
- [73] R. Foot and X.G. He, Phys. Lett. B **267**, 509 (1991).
- [74] M. Pospelov, Phys. Rev. D **80**, 095002 (2009).
- [75] E.W. Kolb and M.S. Turner, *The Early Universe*, Addison-Wesley, 1990.
- [76] D. Larson *et al.*, Astrophys. J. Suppl. **192**, 16 (2011).
- [77] J. Angle *et al.*, (XENON10 Collaboration), Phys. Rev. Lett. **107**, 051301 (2011).
- [78] J. Aprile *et al.*, (XENON100 Collaboration), arXiv:1207.5988 [astro-ph.CO].
- [79] S.W. Randall, M. Markevitch, D. Clowe, A.H. Gonzalez, and M. Bradac, arXiv:0704.0261 [astro-ph].