

Neutrinoless double beta decay and pseudo-Dirac neutrino mass predictions through inverse seesaw mechanism

Ram Lal Awasthi,^δ M. K. Parida[†] and Sudhanwa Patra[†]

[†]*Center of Excellence in Theoretical and Mathematical Sciences,
Siksha 'O' Anusandhan University, Bhubaneswar-751030, India.*

^δ*Harish-Chandra Research Institute, Chhatnag Road, Jhusi, Allahabad 211019, India.**

In the inverse seesaw extension of the standard model, supersymmetric or non-supersymmetric, while the light left-handed neutrinos are Majorana, the heavy right-handed neutrinos are pseudo-Dirac fermions. We show how one of these latter category of particles can contribute quite significantly to neutrinoless double beta decay. The neutrino virtuality momentum is found to play a crucial role in the non-standard contributions leading to the prediction of the pseudo-Dirac fermion mass in the range of 120 MeV – 500 MeV. When the Dirac neutrino mass matrix in the inverse seesaw formula is similar to the up-quark mass matrix, characteristic of high scale quark-lepton symmetric origin, the predicted branching ratios for lepton flavor violating decays are also found to be closer to the accessible range of ongoing experiments.

I. INTRODUCTION: The standard gauge theory of strong, weak, and electromagnetic interactions has confronted numerous experimental tests while the last piece of evidence on the Higgs boson is currently under rigorous scrutiny at the Large Hadron Collider (LHC). In spite of these, neutrino oscillation data uncovering tiny masses of left-handed (LH) neutrinos call for physics beyond the standard model (SM) which is most simply achieved via canonical seesaw mechanism [1, 2] that requires the addition of one heavy right-handed (RH) neutrino per generation provided both LH and RH neutrinos are Majorana fermions [3]. Several other forms of seesaw mechanism [5–7] also require Majorana fermions. Quite interestingly, ongoing experiments on neutrinoless double beta decay ($0\nu\beta\beta$) [8] is expected to resolve the issue between Majorana [3] or Dirac [4] nature of the neutrino¹. In contrast to the predicted small contribution to the $0\nu\beta\beta$ decay rate in the SM, there has been quite significant, or even more dominant predictions if, at the TeV scale, there is left-right (LR) gauge theory [10, 11]. Even, attempts have been made to predict nonstandard contributions to $0\nu\beta\beta$ decay rate due to the mediation of pseudo-Dirac neutrinos where each of them is considered to be a pair of Majorana neutrinos [11, 12]. While the possibility of left-handed neutrinos being pseudo-Dirac has been shown to be highly challenging [13], contribution of a fourth generation heavy pseudo-Dirac neutrino to $0\nu\beta\beta$ has been explored with the condition that its mass should be greater than $M_Z/2$ [14]. If the Dirac neutrino mass matrix occurring in seesaw formulas has its left-right symmetric or quark-lepton symmetric origin, descending from Pati-Salam symmetry [15] or SO(10) grand unified theory [16] at high scales, then the canonical seesaw scale is too large to be experimentally tested by high energy accelerators including LHC. Alternatively, the inverse

seesaw mechanism [17, 18], which requires one RH neutrino as well as an additional sterile fermion per generation, operates at TeV scale and is, therefore, experimentally verifiable. In this framework while the LH light neutrinos are Majorana fermions, the RH neutrinos are pseudo-Dirac by nature having heavier masses.

In this letter we show that the inverse seesaw formula explaining the light neutrino masses and mixings permits the lightest of the three pseudo-Dirac neutrinos in the mass range (120 – 500) MeV leading to new contributions to $0\nu\beta\beta$ decay comparable to, or much more than, those due to the exchanges of the light left-handed neutrinos. The neutrino virtuality momentum [19, 20], $|p| \sim 190$ MeV, is noted to play a crucial role in such new contributions. The origin of Dirac neutrino mass matrix is also found to be important in our estimations in predicting lepton flavor violating decays accessible to ongoing experimental searches. As our results are also applicable in the inverse seesaw extension of the minimal supersymmetric standard model (MSSM), they are consistent with gauge coupling unification at the MSSM-GUT scale, $M_U \simeq 2 \times 10^{16}$ GeV.

II. THE INVERSE SEESAW EXTENSION: As is customary to the implementation of inverse seesaw mechanism, we add two fermion singlets to each generation of the SM, with or without supersymmetry. While we call the first type of singlet a RH neutrino (N_R), the second type of singlet is named as a sterile neutrino (S_L) and, in the (ν_L, N_R^c, S_L) basis, the 9×9 neutrino mass matrix is [18]

$$\mathcal{M}_\nu = \begin{pmatrix} 0 & M_D & 0 \\ M_D^T & 0 & M^T \\ 0 & M & \mu_S \end{pmatrix}, \quad (1)$$

where M_D is the Dirac mass term of the neutrino, and M is the heavy Dirac mass matrix relating N_R and S_L . The matrices M_D and M are in general 3×3 complex in flavor space whereas the μ_S is 3×3 complex symmetric matrix.

Transformation from flavor to mass basis and diagonaliza-

¹ Besides the two distinct possibilities, Dirac or Majorana, very recently a new hypothesis has been advanced in which neutrinos could be schizophrenic [9].

tion are achieved through

$$|\nu\rangle_f = \mathcal{V}^* |\nu\rangle_m, \quad (2)$$

$$\mathcal{V}^\dagger \mathcal{M}_\nu \mathcal{V}^* = \hat{\mathcal{M}}_\nu = \text{Diag}\{m_{\nu_i}; M_{\zeta_j}\}, \quad (3)$$

where $|\nu\rangle_m = (\tilde{\nu}_i, \zeta_j)^T$ represents the three light and six heavy mass states, and i and j run over the light and heavy mass eigenstates, respectively. With $\mu_S, M_D \ll M$, the matrix \mathcal{M}_ν can be block diagonalized to light and heavy sectors

$$m_\nu \simeq \left(\frac{M_D}{M}\right) \mu_S \left(\frac{M_D}{M}\right)^T, \\ M_H \simeq \begin{pmatrix} 0 & M^T \\ M & \mu_S \end{pmatrix}. \quad (4)$$

where m_ν has the well known inverse seesaw formula [18] and M_H is the mass matrix for heavy pseudo-Dirac pairs of comparable masses with splitting of the order of μ_S . The μ_S term in the Lagrangian breaks the leptonic global symmetry, $U(1)_L$, which is otherwise preserved in the standard model in the limit $\mu_S \rightarrow 0$ rendering all the LH neutrinos to be massless. Hence the small μ_S should be a natural parameter in this theory in the 't Hooft sense [21]. The above block diagonalized matrices are further diagonalized through the PMNS matrix, U_ν , and a 6×6 unitary matrix U_H , respectively, so that

$$\mathcal{V} \simeq \begin{pmatrix} 1 - \frac{1}{2} B^* B^T & B^* \\ -B^T & 1 - \frac{1}{2} B^T B^* \end{pmatrix} \begin{pmatrix} U_\nu & 0 \\ 0 & U_H \end{pmatrix}, \quad (5)$$

where

$$B^T \simeq \begin{pmatrix} -M^{*-1} \mu_S^* (M_D M^{-1})^\dagger \\ (M_D M^{-1})^\dagger \end{pmatrix} \simeq \begin{pmatrix} 0 \\ X^\dagger \end{pmatrix}. \quad (6)$$

Hence, in the leading order approximation, \mathcal{V} can be written as

$$\mathcal{V} \simeq \begin{pmatrix} 1 - \frac{1}{2} X X^\dagger & 0 & X \\ 0 & 1 & 0 \\ -X^\dagger & 0 & 1 - \frac{1}{2} X^\dagger X \end{pmatrix} \begin{pmatrix} U_\nu & 0 \\ 0 & U_H \end{pmatrix}, \quad (7)$$

where $X = (M_D M^{-1})$, and all the elements in the first block are 3×3 matrices.

(II. A) μ_S from neutrino oscillation data: The inverse seesaw formula in eqn. (4) predicts light neutrino mass matrix in terms of three other matrices, M_D , M , and μ_S . At first we take $M_D \simeq M_\ell$, the charged lepton mass matrix, which may arise if the SM originates from high scale left-right gauge symmetry, $SU(2)_L \times SU(2)_R \times U(1)_{B-L} \times SU(3)_C \xrightarrow{M_R} SM$, where $M_R \gg M_W$. Assuming the matrix M to be diagonal for the sake of simplicity and using $M_D = \text{diag}\{m_e, m_\mu, m_\tau\} = \{0.0005, 0.1, 1.7\}$ GeV, we obtain μ_S from global fits to the neutrino oscillation data [22] given in TABLE I

$$\mu_S \text{ (GeV)} = X^{-1} \mathcal{N} \hat{m}_\nu \mathcal{N}^T X^{T-1} \quad (8) \\ = \begin{pmatrix} 6.71 \times 10^{-7} + 1.96 \times 10^{-7} \mathbf{i} & -1.17 \times 10^{-8} - 3.22 \times 10^{-8} \mathbf{i} & -3.71 \times 10^{-8} - 2.03 \times 10^{-8} \mathbf{i} \\ -1.17 \times 10^{-8} - 3.22 \times 10^{-8} \mathbf{i} & 1.53 \times 10^{-8} - 2.22 \times 10^{-10} \mathbf{i} & 7.0 \times 10^{-9} - 2.83 \times 10^{-9} \mathbf{i} \\ -3.71 \times 10^{-8} - 2.03 \times 10^{-8} \mathbf{i} & 7.0 \times 10^{-9} - 2.83 \times 10^{-9} \mathbf{i} & -5.50 \times 10^{-9} + 5.26 \times 10^{-11} \mathbf{i} \end{pmatrix}, \quad (9)$$

where $\mathcal{N} = (1 - \eta) U_\nu$ and $\eta = \frac{1}{2} X X^\dagger$ is a measure of unitarity violation. This particular structure of μ_S has been derived using, as an example, the normal hierarchical (NH) light neutrino masses $\hat{m}_\nu^{\text{diag}} = \text{diag}(0.00127 \text{ eV}, 0.00885 \text{ eV}, 0.0495 \text{ eV})$ and non-degenerate eigenvalues of $M = \text{diag}\{0.2, 2.6, 23.7\}$ GeV. Similar analysis predicts somewhat different structures of μ_S for inverted hierarchical (IH) and quasi-degenerate (QD) pattern of the light neutrinos and can further be easily obtained for degenerate $M_1 = M_2 = M_3$ or, partially-degenerate $M_1 = M_2 \ll M_3$ after taking care of the phenomenological bounds $|\eta_{ee}| < 2.0 \times 10^{-3}$, $|\eta_{\mu\mu}| < 8.0 \times 10^{-4}$, and $|\eta_{\tau\tau}| < 2.7 \times 10^{-3}$. Our ansatz with $M = \text{diag}(M_1, M_2, M_3)$

Neutrino oscillation parameters	Globally fitted values
$\Delta m_{\text{sol}}^2 [\text{eV}^2]$	7.58×10^{-5}
$ \Delta m_{\text{atm}}^2 [\text{eV}^2]$	2.35×10^{-3}
$\sin^2 \theta_{12}$	0.320
$\sin^2 \theta_{23}$	0.427
$\sin^2 \theta_{13}$	0.0246
δ_{CP}	0.8π

TABLE I: Mass squared differences, mixing angles, and CP -phase from global fits to neutrino oscillation data [22].

gives

$$\eta = \text{diag} \left(\frac{1.25 \times 10^{-7}}{M_1^2}, \frac{0.005}{M_2^2}, \frac{1.35}{M_3^2} \right),$$

where all masses on the right hand side are in GeV.

III. NEUTRINOLESS DOUBLE BETA DECAY PREDICTIONS

Two separate contributions due to light and heavy neutrino exchanges to $0\nu\beta\beta$ transition become transparent by writing the flavor eigenstates as linear combination of light and heavy mass eigenstates

$$\nu_\alpha = \mathcal{N}_{\alpha i} \nu_i + \mathcal{U}_{\alpha j} \zeta_j,$$

where $\mathcal{U} \simeq (0, X)U_H$ is a 3×6 matrix. Then the weak charge-current Lagrangian can be expressed as

$$\begin{aligned} \mathcal{L}_{CC} &= \frac{g}{\sqrt{2}} W_L^\mu \bar{\ell}_\alpha \gamma^\mu P_L \nu_\alpha + \text{h.c.} \\ &= \frac{g}{\sqrt{2}} W_L^\mu \bar{\ell}_\alpha \gamma^\mu P_L (\mathcal{N}_{\alpha i} \nu_i + \mathcal{U}_{\alpha j} \zeta_j) + \text{h.c.}, \end{aligned} \quad (10)$$

resulting in two different categories of Feynman amplitudes:

- \mathcal{A}_{LL}^ν which arises from the Feynman diagram of Fig. 1(a) due to only light neutrino exchanges

$$\mathcal{A}_{LL}^\nu = G_F^2 \mathcal{N}_{ei}^2 \frac{m_{\nu i}}{p^2}, \quad (11)$$

where $\langle p \rangle \simeq 190$ MeV represents neutrino virtuality momentum [19, 20].

- \mathcal{A}_{LL}^ζ which arises from the Feynman diagram of Fig. 1(b) due to heavy pseudo-Dirac neutrinos,

$$\mathcal{A}_{LL}^\zeta = G_F^2 (\mathcal{U}_{ej})^2 \frac{M_{\zeta j}}{p^2 - M_{\zeta j}^2}. \quad (12)$$




FIG. 1: Feynman diagrams contributing to neutrinoless double beta decay due to light neutrino exchanges (left-panel) and heavy pseudo-Dirac neutrino exchanges (right-panel).

The mass eigenstates of heavy pseudo-Dirac neutrinos are $(\zeta_1^+, \zeta_2^+, \zeta_3^+; \zeta_1^-, \zeta_2^-, \zeta_3^-)$ with almost degenerate pairs $(\zeta_k^+, \zeta_k^-; k=1,2,3)$ but having small mass difference μ_S between the members of the pair and the flavor states are $(N_1, N_2, N_3; S_1, S_2, S_3)$. The mixing matrix for these

pseudo-Dirac neutrinos have been discussed in Sec-II. The half-life of $0\nu\beta\beta$ transition is then found to be

$$\left[T_{1/2}^{0\nu\beta\beta} \right]^{-1} = \mathcal{K}_{0\nu} \left[|m_{\nu,LL}^{ee} + M_{\zeta,LL}^{ee}|^2 \right], \quad (13)$$

where $\mathcal{K}_{0\nu}$ contains phase space factors plus nuclear matrix elements and $m_{\nu,LL}^{ee}$ ($M_{\zeta,LL}^{ee}$) represents the effective neutrino mass derived from light neutrino (heavy pseudo-Dirac neutrino) exchanges in the mass basis. The analytic forms of the two effective masses have been estimated for this model as shown in TABLE. II:

Effective mass	Analytical expression
$m_{\nu,LL}^{ee}$	$\mathcal{N}_{ei}^2 m_{\nu i}$
$M_{\zeta,LL}^{ee}$	$(\mathcal{U}_{ej})^2 \frac{M_{\zeta j}}{p^2 - M_{\zeta j}^2} \langle p \rangle ^2$

TABLE II: Effective mass parameter for standard (non-standard) contributions due to light (heavy pseudo-Dirac) neutrino exchanges for $0\nu\beta\beta$ decay.

We discuss below three different cases:

(III. A) The standard contribution. It is well known that the standard contributions due to light neutrino exchanges are dependent on their allowed mass patterns; *normal hierarchical* (NH), *inverted hierarchical* (IH), or *quasi-degenerate* (QD),

$$\begin{aligned} m_{ee,LL}^\nu &\simeq U_{e1}^2 m_{\nu 1} + U_{e2}^2 e^{2i\alpha} m_2 + U_{e3}^2 e^{2i\beta} m_3 \\ \Rightarrow |m_{ee,LL}^\nu| &\simeq \begin{cases} 0.004 \text{ eV} & \text{NH,} \\ 0.048 \text{ eV} & \text{IH,} \\ 0.1 \text{ eV} & \text{QD.} \end{cases} \end{aligned} \quad (14)$$

In our case, $\mathcal{N}_{ei} \simeq U_{ei}$ and light neutrino exchanges in the mass basis gives almost the same contributions which are presented by solid lines shown in Fig. 2, Fig. 3, and Fig. 4.

(III. B) $M_{\zeta j} \gg |p|$: In the inverse seesaw extension under study, in addition to the standard effective mass parameter, the additional effective mass parameter for $|M_{\zeta j}| \gg |p|$ satisfies $\langle M_{\zeta \pm} \rangle^{-1} = (\mathcal{U}^\pm)^2 \frac{-1}{M_{\zeta \pm}}$. This results in new contribution to $0\nu\beta\beta$ transition half-life

$$\begin{aligned} \left[T_{1/2}^{0\nu\beta\beta} \right]^{-1} &= \mathcal{K}_{0\nu} \left| |\langle p \rangle|^2 \left(\frac{1}{\langle M_{\zeta_+} \rangle} - \frac{1}{\langle M_{\zeta_-} \rangle} \right) \right|^2 \\ &\simeq \mathcal{K}_{0\nu} \left| |\langle p \rangle|^2 (\mathcal{U}^\pm)^2_{ek} \frac{\mu_{S_{kk}}}{M_{kk}^2} \right|^2, \end{aligned} \quad (15)$$

where $\mu_{S_{kk}}$ and M_{kk} are the eigenvalues of μ_S and M , respectively. One example of this case has been shown in Fig.




FIG. 2: Predictions of effective mass $|M_{ee}|$ in $0\nu\beta\beta$ decay with Dirac neutrino mass $M_D =$ charged lepton mass M_l and the diagonal structure of M for NH (IH) pattern of light LH neutrino masses as shown in the *left (right) panel*. The standard contribution is shown by solid line and nonstandard contributions with pseudo-Dirac neutrino exchanges of different masses expressed in MeV are shown by other lines.

2 for $M_{\zeta_1} = 0.5$ GeV where the predicted effective mass parameter is nearly 3/2 (4) times larger than the standard prediction for NH (IH) case.

(III. C) $M_{\zeta_j} \simeq |p|$: In this region where different allowed values of M_{ζ_j} are of the order of neutrino virtuality momentum $|p| \simeq 190$ MeV, the new contribution to neutrinoless double beta decay due to heavy pseudo-Dirac neutrino exchange is found to be more dominant than the standard contribution and the $0\nu\beta\beta$ transition half-life is given below

$$\left[T_{1/2}^{0\nu\beta\beta} \right]_{\text{pseudo-Dirac}}^{-1} = \mathcal{K}_{0\nu} \left| \langle p \rangle^2 \mathcal{U}^{\pm} \left(\frac{M_{\zeta_+}}{p^2 - M_{\zeta_+}^2} - \frac{M_{\zeta_-}}{p^2 - M_{\zeta_-}^2} \right) \right|^2 \simeq \mathcal{K}_{0\nu} \left| \langle p \rangle^2 (\mathcal{U}^{\pm})_{ek}^2 \frac{\mu_{Skk}}{p^2 - M_{kk}^2} \right|^2. \quad (16)$$

The predicted new values of the effective mass parameters arising solely due to pseudo-Dirac neutrino exchanges have been shown in Fig. 2 in the *left-pannel (right-pannel)* for NH (IH) patterns of the light neutrino masses, respectively, where $M_{\zeta_1} = (0.15 - 0.5)$ GeV. It is quite clear from the plots that even for $M_{\zeta_1} = 0.25$ GeV or, 0.5 GeV, the new contributions are 3-6 times larger than the standard ones. While for the value of $M_{\zeta_1} = 0.18$ GeV, the contribution is nearly 100 times larger shown in Fig. 2. This large enhancement occurs as M_{ζ_1} approaches the vicinity of the neutrino virtuality momentum, $|p| \simeq 190$ MeV. We point out that such important effects of pseudo-Dirac neutrino masses are found for the first time in this work.

IV. LEPTON FLAVOR VIOLATION WITH DOMINANT $0\nu\beta\beta$ DECAY RATE: We have clearly shown that the predicted non-standard contributions to neutrinoless double beta decay rate are dominant for the lightest allowed pseudo-Dirac neutrino mass $M_{\zeta_1} \simeq (0.15 - 0.5)$ GeV. However, because of the diagonal nature of M_D and assumed structure of M , the branching ratios for lepton flavor violating (LFV) decays, $\mu \rightarrow e + \gamma$, $\tau \rightarrow e + \gamma$, and $\tau \rightarrow \mu + \gamma$ are as small as the SM predictions.

In the next two examples we adopt plausible parametrization predicting significantly larger contribution to these branching ratios while retaining the dominant contributions to $0\nu\beta\beta$ transition.

(IV. A) $M_D \simeq M_l$ with non-diagonal M : We generate non-diagonal matrix M to satisfy the existing phenomenological bounds on the elements of η [23],

$$\begin{aligned} |\eta_{ee}| &< 2.0 \times 10^{-3}, & |\eta_{e\mu}| &< 3.5 \times 10^{-5}, \\ |\eta_{e\tau}| &< 8.0 \times 10^{-3}, & |\eta_{\mu\mu}| &< 8.0 \times 10^{-4}, \\ |\eta_{\mu\tau}| &< 5.1 \times 10^{-3}, & |\eta_{\tau\tau}| &< 2.7 \times 10^{-3}. \end{aligned} \quad (17)$$

Using the parametrization of the type used in ref. [24], M can be expressed as

$$\left[M \left(\frac{1}{\sqrt{2}\eta} O^T M_D \right)^{-1} \right]^\dagger \left[M \left(\frac{1}{\sqrt{2}\eta} O^T M_D \right)^{-1} \right] = \mathbf{1}_3 = V^\dagger V \quad (18)$$

where O is the matrix diagonalizing $|\eta|$ and V is an arbitrary unitary matrix. Choosing, for the sake of simplicity, $V = \mathbf{1}_3$ and we note that that a lightest pair with $M_{\zeta_1} \simeq 0.16$ GeV, in the vicinity of neutrino virtuality momentum, can be achieved by suitable rescaling, *e.g.* $\eta_{\alpha\beta} \rightarrow \eta_{\alpha\beta}/(1500)$. After this scaling we find

$$\begin{aligned} M(\text{GeV}) &= \left(\frac{1}{\sqrt{2}\eta} O^T M_D \right) \\ &= \begin{pmatrix} 0.092i & 14.08i & -383.7i \\ 0.217 & -70.36 & -80.39 \\ 0.074 & 8.866 & 320.5 \end{pmatrix}, \end{aligned} \quad (19)$$

where

$$O = \begin{pmatrix} 0.5874i & 0.5446 & 0.5987 \\ 0.4284i & -0.8368 & 0.3409 \\ -0.6866i & -0.0562 & 0.7248 \end{pmatrix}.$$




FIG. 3: Same as Fig. 2 but now with non-diagonal structure of M and reduced values of nonunitarity matrices η as described in the text.

With the allowed mass eigenvalues for the heavy pseudo-Dirac neutrinos, $M_\zeta = \text{diag}\{0.159, 72.0, 506.4\}$ GeV, the predicted branching ratios for lepton flavor violating decays are [25]

$$\begin{aligned} \text{Br}(\mu \rightarrow e + \gamma) &= 1.56 \times 10^{-26}, \\ \text{Br}(\tau \rightarrow e + \gamma) &= 5.79 \times 10^{-27}, \\ \text{Br}(\tau \rightarrow \mu + \gamma) &= 1.10 \times 10^{-18}. \end{aligned} \quad (20)$$

Although all the three branching ratios are much smaller than their corresponding experimental upper limits [27], they are considerably larger than the SM predictions. However we note below that with M_D similar to M_u , the up-quark mass matrix, a phenomenon underlying the possible origin of SM from Pati-Salam [15] or SO(10) model, LFV decays have much larger predicted values, accessible to ongoing experimental searches, while similar predictions on dominant $0\nu\beta\beta$ decay are maintained.

(IV. B) $M_D \simeq M_u$ and GUT connection: In this case Dirac neutrino mass matrix is approximated to be up-quark mass matrix, which originates if the high scale symmetry is Pati-Salam or SO(10) GUT, $SU(2)_L \times SU(2)_R \times SU(4)_C$ or SO(10) $\xrightarrow{M_R}$ SM. Using running masses $(m_u, m_c, m_t) = (0.00233, 1.275, 160)$ GeV and Cabbibo-Kobayashi-Maskawa mixing matrix, V_{CKM} [28],

$$M_D(\text{GeV}) \simeq M_u = V_{CKM} \hat{M}_u V_{CKM}^T = \begin{pmatrix} 0.067 - 0.004i & 0.302 - 0.022i & 0.55 - 0.53i \\ 0.302 - 0.022i & 1.48 - 0.0i & 6.534 - 0.001i \\ 0.55 - 0.53i & 6.534 - 0.0009i & 159.72 + 0.0i \end{pmatrix}. \quad (21)$$

At first using the phenomenological bounds from eqn. (17) and saturating our ansatz for $\eta_{\alpha\beta} = |\eta|_{\max}/n$, $n = 1, \dots, 5$, we search for matrix M through eqn. (18) which gives μ_S from

eqn. (8). We obtain for $n = 4$

$$M(\text{GeV}) = \begin{pmatrix} -5.9 - 3.45i & 0.2 - 60.72i & 5.15 - 1760i \\ -10.44 + 2.13i & -60.9 - 0.5i & -598.0 - 12.16i \\ 7.08 - 5.09i & 70.86 - 0.18i & 1547 - 4.15i \end{pmatrix}, \quad (22)$$

$\mu_S(\text{eV})$

$$= \begin{pmatrix} -3.42 - 0.51i & -1.92 + 5.55i & 0.28 - 1.92i \\ -1.92 + 5.55i & 39.1 + 5.68i & -12.1 + 0.11i \\ 0.28 - 1.92i & -12.1 + 0.11i & 4.10 - 0.68i \end{pmatrix}. \quad (23)$$

Our predictions on numerical values of the effective mass parameter for $0\nu\beta\beta$ are shown in Fig. 4 for NH, IH and QD cases. For NH light neutrinos we find that the predicted value of $|M_{ee}|$ is increased by a factor 3 for $\eta = |\eta|_{\max}/3$, corresponding to lightest pair $M_{\zeta_1} = 131$ MeV, while the increment is 10 times for $\eta = |\eta|_{\max}/4$ and $M_{\zeta_1} = 152$ MeV, and 30 times for $\eta = |\eta|_{\max}/5$ and $M_{\zeta_1} = 169$ MeV. We find that the enhancement survives as long as lightest pair $M_{\zeta_1} \simeq 120 - 350$ MeV. For the IH light neutrino masses the results are similar as shown on the right panel of Fig. 4. The branching ratios for lepton flavor violating decays predicted in this scenario with $M_\zeta = (0.152, 39.5, 2426)$ GeV are

$$\begin{aligned} \text{Br}(\mu \rightarrow e + \gamma) &= 3.6 \times 10^{-13}, \\ \text{Br}(\tau \rightarrow e + \gamma) &= 4.2 \times 10^{-14}, \\ \text{Br}(\tau \rightarrow \mu + \gamma) &= 3.3 \times 10^{-12}, \end{aligned} \quad (24)$$

while the present experimental limits at 90% C.L. on these branching ratios are $\text{Br}(\mu \rightarrow e + \gamma) \leq 1.2 \times 10^{-11}$, $\text{Br}(\tau \rightarrow e + \gamma) \leq 3.3 \times 10^{-8}$, and $\text{Br}(\tau \rightarrow \mu + \gamma) \leq 4.4 \times 10^{-8}$ [27]. The projected reach of sensitivity in the future is $\text{Br}(\tau \rightarrow e + \gamma)$, $\text{Br}(\tau \rightarrow \mu + \gamma) \leq 10^{-9}$ and specifically $\text{Br}(\mu \rightarrow e + \gamma) \leq 10^{-19}$ [27].

The predicted nonstandard contributions to $0\nu\beta\beta$ transition are shown in the *left-panel* for NH and in the *right-panel*




FIG. 4: The effective mass parameter, $|M_{ee}|$, prediction for $0\nu\beta\beta$ decay due to light (Solid line) and pseudo-Dirac (dashed lines) neutrino exchange where $M_D \simeq M_u$ and $\eta = |\eta_{max}|/n$, $n = 1, 3, 4, 5$ as discussed in the text.

for IH case of Fig. 4. In view of the M_{ζ_1} -dependent enhancements of $0\nu\beta\beta$ decay rates discussed above it is tempting to search for the possibility of the lightest pseudo-Dirac neutrino mass which we perform by the replacement $M_{\zeta_1}^2 \rightarrow M_{\zeta_1}^2 + iM_{\zeta_1}\Gamma_1$, where Γ_1 corresponds to plausible value of width of the particle. Using, for example, $\Gamma_1 \simeq 0.1$ keV, our predictions are presented by solid curve in Fig. 5 for NH light neutrino masses where the resonant behavior is clearly exhibited around $M_{\zeta_1} = 190$ MeV.




FIG. 5: Variation of effective mass $|M_{ee}|$ prediction as a function of lightest pseudo-Dirac neutrino mass M_{ζ_1} where μ_S matrix has been determined for NH light neutrino masses. The resonance peak is at $M_{\zeta_1} \simeq 190$ MeV as shown by the solid line. For comparison, the standard contribution with NH masses is shown by the dashed horizontal line.

V. DISCUSSIONS AND CONCLUSION: In this letter we have shown that in the inverse seesaw framework of the standard model, the lightest of the pseudo-Dirac neutrino could be of $\mathcal{O}(100)$ MeV in concordance with tiny left handed neutrino masses and the oscillation data. This pseudo-Dirac neutrino mass being in the vicinity of the neutrino virtuality momentum $|p| \simeq 190$ MeV, gives very significant non-standard con-

tributions to $0\nu\beta\beta$ decay rates, even far exceeding the standard contributions. The Dirac neutrino mass possibly originating from high scale Pati-Salam symmetry or SO(10) grand unification, plays a crucial role in determining dominant contributions to $0\nu\beta\beta$ decay rates simultaneously with LFV decays with predicted branching ratios accessible to on going search experiments. The underlying mechanism provides three distinct platforms for its falsifiability (i) $0\nu\beta\beta$ decay rates, (ii) determination of light pseudo-Dirac neutrino mass $M_{\zeta_1} \simeq 120 - 500$ MeV, and (iii) the three predicted branching ratios of eqn. (24). As all our results are applicable in the case of inverse seesaw extended supersymmetric standard model, they are also consistent with gauge coupling unification at the MSSM-GUT scale, $M_U \simeq 2 \times 10^{16}$ GeV. The Pati-Salam or SO(10) completion of the model discussed in Sec. IV.B will be reported elsewhere in future publication [26].

ACKNOWLEDGEMENT: Ram Lal Awasthi acknowledges the hospitality at Center of Excellence in Theoretical and Mathematical Sciences, SOA University where the present work has been completed.

* Electronic address: ramlal@hri.res.in, sudha.astro@gmail.com, paridamk@s

- [1] P. Minkowski, Phys. Lett. B **67**, 421 (1977); T. Yanagida, proceedings of the *Workshop on Unified Theories and Baryon Number in the Universe*, Tsukuba, 1979, eds. A. Sawada, A. Sugamoto, KEK Report No. 79-18, Tsukuba; S. Glashow, in *Quarks and Leptons, Cargèse 1979*, eds. M. Lévy. et al., (Plenum, 1980, New York); M. Gell-Mann, P. Ramond, R. Slansky, proceedings of the *Supergravity Stony Brook Workshop*, New York, 1979, eds. P. Van Nieuwenhuizen, D. Freeman (North-Holland, Amsterdam); R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [2] J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980).
- [3] E. Majorana; N. Cim. **14** (1937) 171.
- [4] P. A. M. Dirac; Proceedings of the Royal Society of London, **109**, 752 (Dec. 1, 1925), pp. 642-653.
- [5] R. N. Mohapatra and G. Senjanovic, Phys. Rev. D **23**, 165

- (1981);
- [6] G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B **181**, 287 (1981);
- [7] R. Foot, H. Lew, X. G. He and G. C. Joshi, Z. Phys. C **44**, 441 (1989); B. Bajc, G. Senjanović, JHEP **0708**, 014 (2007) [hep-ph/0612029]; E. Ma and D. P. Roy, Nucl. Phys. B **644**, 290 (2002); W. Grimus and L. Lavoura, JHEP **0011**, 042 (2000), arXiv: 0008179 [hep-ph]; M. Hirsch, H. V. Klapdor-Kleingrothaus and O. Panella, Phys. Lett. B **374**, 7 (1996), arXiv: 9602306 [hep-ph]; B. Bajc, M. Nemevšek, G. Senjanović, Phys. Rev. **D76** (2007) 055011 [hep-ph/0703080];
- [8] H. V. Klapdor-Kleingrothaus, A. Dietz, L. Baudis, G. Heusser, I. V. Krivosheina, S. Kolb, B. Majorovits, H. Pas et al., Eur. Phys. J. A12 (2001) 147-154; C. Arnaboldi et al. [CUORICINO Collaboration], Phys. Rev. C78, 035502 (2008); C. E. Aalseth et al. [IGEX Collaboration], Phys. Rev. D65 (2002) 092007; J. Argyriades et al. [NEMO Collaboration], Phys. Rev. **C80**, 032501 (2009); I. Abt, M. F. Altmann, A. Bakalyarov, I. Barabanov, C. Bauer, E. Bellotti, S. T. Belyaev, L. B. Bezrukov et al., [hep-ex/0404039]; S. Schonert et al. [GERDA Collaboration], Nucl. Phys. Proc. Suppl. 145, 242-245 (2005); C. Arnaboldi et al. [CUORE Collaboration], Nucl. Instrum. Meth. A518, 775-798 (2004); H. V. Klapdor-Kleingrothaus, I. V. Krivosheina, A. Dietz, O. Chkvorets, Phys. Lett. B586, 198-212 (2004); H. V. Klapdor-Kleingrothaus, I. V. Krivosheina, Mod. Phys. Lett. A21, 1547-1566 (2006).
- [9] James Barry, Rabindra N. Mohapatra and Werner Rodejohann; Phys.Rev. **D 83** (2011) 113012; R. Allahverdi, B. Dutta, and R.N. Mohapatra, Phys.Lett. **B 695** (2011) 181-184.
- [10] V. Tello, M. Nemevšek, F. Nesti, G. Senjanović, F. Vissani, Phys. Rev. Lett. **106**, 151801 (2011); Joydeep Chakraborty, H.Zeen Devi, Srubabati Goswami and Sudhanwa Patra, JHEP 1208 (2012) 008. arXiv:1204.2527 [hep-ph].
- [11] M.K. Parida and Sudhanwa Patra; Phys. Lett. **B 718** (2013) 1407.
- [12] H. Zhang, and S. Zhou, Phys.Lett. **B 685** (2010) 297.
- [13] J.F. Beacom, N.F. Bell, D. Hooper, J.G. Learned, S. Pakvasa, and T.J. Weiler, Phys. Rev. Lett. 92 (2004) 011101.
- [14] A. Lenz, H. Pas, and D. Schalla, Phys.Rev. **D85** (2012) 075025.
- [15] J. C. Pati and A. Salam, Phys. Rev. D 10 (1974) 275.
- [16] H. Georgi, Particles and Fields, Proceedings of APS Division of Particles and Fields, ed C. Carlson, p 575 (1975); H. Fritzsch, P. Minkowski, Ann. Phys. 93, 193 (1975).
- [17] R. N. Mohapatra, Phys. Rev. Lett. **56**, 561-563 (1986); R. N. Mohapatra, J. W. F. Valle, Phys. Rev. **D34**, 1642 (1986).
- [18] D. Wyler, L. Wolfenstein, Nucl. Phys. **B218**, 205 (1983); E. Witten, Nucl. Phys. **B268**, 79 (1986).
- [19] R.N. Mohapatra; Phys. Rev. D 34 (1986) 909; M. Doi and T. Kotani, Prog. Theor. Phys. 89 (1993) 139.
- [20] K. Muto, I. Blender, and H. V. Klapdor-Kleingrothaus, Z. Phys. **A 334** (1989) 177; M. Hirsch, K. Muto, T. Oda, and H. V. Klapdor-Kleingrothaus, Z. Phys. **A 347** (1994) 151; J.J. Gomez-Cadenas, J. Martin-Albo, M. Mezzetto, F. Monrabal, and M. Sorel, Riv.Nuovo Cim. 35 (2012) 29-98, e-Print: arXiv:1109.5515 [hep-ex].
- [21] G. t Hooft, in Proceedings of the 1979 Cargese Summer Institute on Recent Developments in Gauge Theories, edited by G. t Hooft et al. (Plenum Press, New York, 1980).
- [22] F. P. An *et. al.* [DAYA-BAY Collaboration], arXiv:1203.1669 [hep-ex]; J. K. Ahn *et. al.* [Soo-Bong Kim for RENO collaboration], arXiv:1204.0626v1 [hep-ex]; G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. M. Rotunno, Phys. Rev. **D 84**, 053007 (2011), arXiv:1106.6028 [hep-ph]; D. V. Forero, M. Tortola and J. W. F. Valle, arXiv:1205.4018 [hep-ph].
- [23] S. Antusch, J. P. Baumann, and E. Fernandez-Martinez, Nucl. Phys. **B810**, 369 (2009); Ram Lal Awasthi and Mina K. Parida, Phys.Rev. **D 86** (2012) 093004.
- [24] J. A. Casas and A. Ibarra, Nucl. Phys. **B 618**, 171 (2001).
- [25] A. Ilakovac, A. Pilaftsis, Nucl. Phys. **B437**, 491 (1995) [hep-ph/9403398]; F. Deppisch, J. W. F. Valle, Phys. Rev. **D72**, 036001 (2005) [hep-ph/0406040]; C. Arina, F. Bazzocchi, N. Fornengo, J. C. Romao, J. W. F. Valle, Phys. Rev. Lett. **101**, 161802 (2008) [arXiv:0806.3225 [hep-ph]]; M. Malinsky, T. Ohlsson, Z. -z. Xing, H. Zhang, Phys. Lett. **B679**, 242-248 (2009) [arXiv:0905.2889 [hep-ph]]; M. Hirsch, T. Kernreiter, J. C. Romao, A. Villanova del Moral, JHEP **1001**, 103 (2010) [arXiv:0910.2435 [hep-ph]]; F. Deppisch, T. S. Kosmas, J. W. F. Valle, Nucl. Phys. **B752**, 80-92 (2006) [arXiv:0910.3924 [hep-ph]].
- [26] Ram Lal Awasthi, M. K. Parida, and Sudhanwa Patra (*under preparation*).
- [27] M. L. Brooks et al. [MEGA Collaboration], Phys. Rev. Lett. 83, 1521 (1999); B. Aubert [The BABAR Collaboration], arXiv:0908.2381 [hep-ex]; Y. Kuno (PRIME Working Group), Nucl. Phys. B. Proc. Suppl. 149, 376 (2005). For a review see F. R. Joaquim, A. Rossi, Nucl. Phys. B 765, 71 (2007).
- [28] C. Amsler et al., Phys. Lett. **B 667**, 1 (2008).