

# QCD sum rule study on the $f_0(980)$ structure as a pure $K\bar{K}$ bound state

Hee-Jung Lee,<sup>1,\*</sup> N. I. Kochelev,<sup>2,3,†</sup> and Yongseok Oh<sup>2,4,‡</sup>

<sup>1</sup>*Department of Physics Education, Chungbuk National University, Cheongju, Chungbuk 361-763, Korea*

<sup>2</sup>*Department of Physics, Kyungpook National University, Daegu 702-701, Korea*

<sup>3</sup>*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna, Moscow region, 141980 Russia*

<sup>4</sup>*Asia Pacific Center for Theoretical Physics, Pohang, Gyeongbuk 790-784, Korea*

(Dated: April 3, 2024)

We perform a QCD sum rule analysis for the scalar  $f_0(980)$  meson to investigate whether it can be described as a pure bound state of  $K$  and  $\bar{K}$  mesons. Based on the QCD sum rule with the operators of up to dimension 10 within the operator product expansion, we found that it is hard to treat the  $f_0(980)$  as a simple  $K\bar{K}$  bound state, which implies that the  $f_0(980)$  scalar meson has more complicated structure being mixed states of various configurations.

PACS numbers: 11.55.Hx, 12.38.Lg, 14.40.Be

The structure of the scalar meson nonet has been a long-standing puzzle in hadron physics. It is now widely accepted that the simplest picture, where the scalar mesons are described as orbital excitations of quark-antiquark pairs, is not compatible with the experimental observations on the decay modes and mass spectra [1]. This led to the idea that these scalar mesons are crypto-exotic tetraquark states [2], and there have been a lot of studies along this direction. Depending on the details of the structure of the tetraquark states, the scalar mesons are considered as diquark-antidiquark bound states [3–8], two-meson molecular states [9–14], or hybrid states [15]. (See also Ref. [16].)

Among the low-lying scalar mesons, the  $f_0(980)$  attracts much interests since the seminal work of Weinstein and Isgur, which investigated the  $f_0(980)$  as a  $K\bar{K}$  molecular state [9]. In a recent work [13], for example, the properties of the  $f_0(980)$  were reanalyzed in a phenomenological Lagrangian approach assuming a pure  $K\bar{K}$  bound state, and the calculated decay widths for  $f_0(980) \rightarrow \pi\pi$  and  $f_0(980) \rightarrow \gamma\gamma$  were claimed to be consistent with the available data. In a recent work [14], however, the scalar and isoscalar meson resonances are investigated in various channels of  $\pi\pi$  scattering, which raised the possibility of the  $f_0(980)$  as a pure  $\eta\eta$  bound state rejecting the pure  $K\bar{K}$  structure. All these ambiguities show that the structure of scalar mesons is non-trivial and more QCD-based approaches are required for

understanding the structure of scalar mesons.

The QCD sum rule (QCDSR) approach is known to be one of the ways to investigate the hadron properties from QCD in a direct way [17]. This approach was used to study the diquark picture of scalar mesons [6], and it was recently shown by one of us that the QCDSR does not support the picture of the  $f_0(980)$  as a pure  $\eta\eta$  bound state [18]. In the present work, we construct the QCDSR for the  $f_0(980)$  to test whether it can be described as a pure  $K\bar{K}$  bound state. To this end, we obtain the QCDSR up to dimension  $d = 10$  operators within the operator product expansion (OPE).

The wave function of the  $f_0(980)$  meson as a pure  $K\bar{K}$  bound state is written generally as

$$|f_0(980)\rangle = \alpha|K^+K^- \rangle + \beta|K^0\bar{K}^0 \rangle. \quad (1)$$

With the following  $K$  meson interpolating currents,

$$\begin{aligned} J_{K^+} &= i\bar{s}\gamma_5 u, & J_{K^-} &= i\bar{u}\gamma_5 s, \\ J_{K^0} &= i\bar{s}\gamma_5 d, & J_{\bar{K}^0} &= i\bar{d}\gamma_5 s, \end{aligned} \quad (2)$$

the interpolating current for the  $f_0(980)$  in QCDSR approach becomes

$$\begin{aligned} J_{f_0} &= \alpha J_{K^+} J_{K^-} + \beta J_{K^0} J_{\bar{K}^0} \\ &= -[\alpha(\bar{s}\gamma_5 u)(\bar{u}\gamma_5 s) + \beta(\bar{s}\gamma_5 d)(\bar{d}\gamma_5 s)]. \end{aligned} \quad (3)$$

Then the vacuum expectation value of the time ordered product of the currents reads

$$\begin{aligned} \langle 0|T J_{f_0}(x) J_{f_0}^\dagger(0)|0\rangle &= \langle 0|T \left\{ \alpha^2 [\bar{s}(x)\gamma_5 u(x)][\bar{u}(x)\gamma_5 s(x)][\bar{s}(0)\gamma_5 u(0)][\bar{u}(0)\gamma_5 s(0)] \right. \\ &\quad + \alpha\beta [\bar{s}(x)\gamma_5 u(x)][\bar{u}(x)\gamma_5 s(x)][\bar{s}(0)\gamma_5 d(0)][\bar{d}(0)\gamma_5 s(0)] + \alpha\beta [\bar{s}(x)\gamma_5 d(x)][\bar{d}(x)\gamma_5 s(x)][\bar{s}(0)\gamma_5 u(0)][\bar{u}(0)\gamma_5 s(0)] \\ &\quad \left. + \beta^2 [\bar{s}(x)\gamma_5 d(x)][\bar{d}(x)\gamma_5 s(x)][\bar{s}(0)\gamma_5 d(0)][\bar{d}(0)\gamma_5 s(0)] \right\} |0\rangle. \end{aligned} \quad (4)$$

\* hjl@cbnu.ac.kr

† kochelev@theor.jinr.ru

‡ yohphy@knu.ac.kr

Since the disconnected terms do not contribute to the QCSSR, here we present only the connected terms. Then the first term can be transformed as

$$\langle 0 | T[\bar{s}(x)\gamma_5 u(x)][\bar{u}(x)\gamma_5 s(x)][\bar{s}(0)\gamma_5 u(0)][\bar{u}(0)\gamma_5 s(0)] | 0 \rangle = \text{Tr}[S_s^{ba'}(x, 0)\gamma_5 S_u^{a'b}(0, x)\gamma_5] \text{Tr}[S_s^{b'a}(0, x)\gamma_5 S_u^{ab'}(x, 0)\gamma_5], \quad (5)$$

in terms of the quark propagator  $S_q^{ab}(x, y)$  with the color indexes  $a, b$ . One can easily verify that, in Eq. (4), replacing the quark flavor  $u$  in the first two terms by  $d$  yields the last two terms. Since we are working in the chiral limit  $m_u = m_d = 0$ , the first two terms and the last two terms give the same contribution. Furthermore, the second and the third terms have disconnected diagrams only, which leads to the overall factor  $\alpha^2 + \beta^2$  in

Eq. (4).

The correlator  $\Pi(q^2) = i \int d^4x e^{iq \cdot x} \langle 0 | T J_{f_0}(x) J_{f_0}^\dagger(0) | 0 \rangle$  can then be calculated within the OPE up to  $O(m_s)$  and  $O(g_c^2)$  keeping the operators of dimension up to 10. By making use of the quark propagator of Ref. [19], the imaginary part of the correlator is obtained as

$$\begin{aligned} \frac{1}{\pi} \text{Im} \Pi^{\text{OPE}}(q^2) = & (\alpha^2 + \beta^2) \left[ \frac{1}{2^{14} (5\pi^6)} (q^2)^4 + \frac{g_c^2 \langle G^2 \rangle}{2^{12} \pi^6} (q^2)^2 + \frac{m_s}{2^8 \pi^4} (\langle \bar{s}s \rangle - 2\langle \bar{u}u \rangle) (q^2)^2 \right. \\ & + \frac{m_s}{2^8 \pi^4} \{ 2ig_c \langle \bar{s}\sigma \cdot Gs \rangle + 3ig_c \langle \bar{u}\sigma \cdot Gu \rangle \} q^2 \\ & + \frac{m_s ig_c \langle \bar{u}\sigma \cdot Gu \rangle}{2^7 \pi^4} q^2 \{ -2 \ln(q^2/\Lambda^2) + \ln \pi + \psi(3) + \psi(2) + 2\gamma_{\text{EM}} \} + \frac{\langle \bar{u}u \rangle \langle \bar{s}s \rangle}{2^4 \pi^2} q^2 \\ & + \frac{m_s g_c^2 \langle G^2 \rangle}{2^9 \pi^4} \left\{ \langle \bar{s}s \rangle - \frac{2^3}{3} \langle \bar{u}u \rangle \right\} - \frac{m_s g_c^2 \langle G^2 \rangle \langle \bar{u}u \rangle}{2^8 \pi^4} \left\{ -2 \ln(q^2/\Lambda^2) + \ln \pi + \psi(2) + \psi(1) + 2\gamma_{\text{EM}} - \frac{2}{3} \right\} \\ & - \frac{1}{2^5 \pi^2} (\langle \bar{u}u \rangle ig_c \langle \bar{s}\sigma \cdot Gs \rangle + \langle \bar{s}s \rangle ig_c \langle \bar{u}\sigma \cdot Gu \rangle) - \frac{m_s \langle \bar{u}u \rangle \langle \bar{s}s \rangle}{2 \cdot 3} \left( \langle \bar{u}u \rangle - \frac{1}{2} \langle \bar{s}s \rangle \right) \delta(q^2) \\ & \left. + \frac{5g_c^2 \langle G^2 \rangle \langle \bar{u}u \rangle \langle \bar{s}s \rangle}{2^6 \cdot 3^2 \pi^2} \delta(q^2) + \frac{13ig_c \langle \bar{u}\sigma \cdot Gu \rangle ig_c \langle \bar{s}\sigma \cdot Gs \rangle}{2^9 \cdot 3 \pi^2} \delta(q^2) \right], \quad (6) \end{aligned}$$

where  $g_c$  is the strong coupling constant and  $\psi(n) = 1 + 1/2 + \dots + 1/(n-1) - \gamma_{\text{EM}}$  with the Euler-Mascheroni constant  $\gamma_{\text{EM}}$ . Here, we have used the factorization hypothesis in calculating the condensates of the operators of dimension higher than 6. The diagrammatic representation of each term is shown in Fig. 1.

Decomposing the spectral sum, which is generated from the dispersion relation of the correlator, into a narrow single resonance and the continuum, and applying the hadron-quark duality hypothesis with the Borel

transform as well, we have the following sum rule,

$$\frac{1}{\pi} \int_0^{s_0^2} ds^2 e^{-s^2/M^2} \text{Im} \Pi^{\text{OPE}}(s^2) = 2f_{f_0}^2 m_{f_0}^8 e^{-m_{f_0}^2/M^2}, \quad (7)$$

with the convention  $\langle 0 | J_{f_0}(0) | f_0(980) \rangle = \sqrt{2} f_{f_0} m_{f_0}^4$ . Here,  $s_0$  and  $M$  denote the threshold for the continuum and the Borel mass, respectively. The imaginary part of the correlator in Eq. (6) gives the explicit QCDSR for the  $f_0(980)$  as

$$\begin{aligned} 2f_{f_0}^2 m_{f_0}^8 e^{-m_{f_0}^2/M^2} = & (\alpha^2 + \beta^2) \left[ \frac{3}{2^{11} (5\pi^6)} M^{10} E_4(M^2) + \frac{g_c^2 \langle G^2 \rangle}{2^{11} \pi^6} M^6 E_2(M^2) + \frac{m_s}{2^7 \pi^4} \left\{ \langle \bar{s}s \rangle - 2\langle \bar{u}u \rangle \right\} M^6 E_2(M^2) \right. \\ & + \frac{m_s}{2^8 \pi^4} \left\{ 2ig_c \langle \bar{s}\sigma \cdot Gs \rangle + 3ig_c \langle \bar{u}\sigma \cdot Gu \rangle \right\} M^4 E_1(M^2) + \frac{m_s ig_c \langle \bar{u}\sigma \cdot Gu \rangle}{2^7 \pi^4} M^4 \bar{W}_1(M^2) \\ & + \frac{\langle \bar{u}u \rangle \langle \bar{s}s \rangle}{2^4 \pi^2} M^4 E_1(M^2) + \frac{m_s g_c^2 \langle G^2 \rangle}{2^9 \pi^4} \left\{ \langle \bar{s}s \rangle - \frac{2^3}{3} \langle \bar{u}u \rangle \right\} M^2 E_0(M^2) - \frac{m_s g_c^2 \langle G^2 \rangle \langle \bar{u}u \rangle}{2^8 \pi^4} M^2 W_0(M^2) \\ & - \frac{1}{2^5 \pi^2} \left\{ \langle \bar{u}u \rangle ig_c \langle \bar{s}\sigma \cdot Gs \rangle + \langle \bar{s}s \rangle ig_c \langle \bar{u}\sigma \cdot Gu \rangle \right\} M^2 E_0(M^2) - \frac{m_s \langle \bar{u}u \rangle \langle \bar{s}s \rangle}{2^2 \cdot 3} \left\{ \langle \bar{u}u \rangle - \frac{1}{2} \langle \bar{s}s \rangle \right\} \\ & \left. + \frac{5g_c^2 \langle G^2 \rangle \langle \bar{u}u \rangle \langle \bar{s}s \rangle}{2^7 \cdot 3^2 \pi^2} + \frac{13ig_c \langle \bar{u}\sigma \cdot Gu \rangle ig_c \langle \bar{s}\sigma \cdot Gs \rangle}{2^{10} \cdot 3 \pi^2} \right], \quad (8) \end{aligned}$$




FIG. 1. Diagrammatic representations of the terms in Eq. (6). Upper two lines correspond to the  $s$  quark and lower two lines to the  $u$  quark. The symbol  $\times$  denotes the strange quark mass  $m_s$ . Here only the nonvanishing diagrams are shown.

where we have used  $\langle \bar{u}u \rangle = \langle \bar{d}d \rangle$  in the chiral limit and

$$E_n(M^2) = \frac{1}{\Gamma(n+1)M^{2n+2}} \int_0^{s_0^2} ds^2 e^{-s^2/M^2} (s^2)^n,$$

$$\overline{W}_n(M^2) = \frac{1}{\Gamma(n+1)M^{2n+2}} \int_0^{s_0^2} ds^2 e^{-s^2/M^2} (s^2)^n \{-2 \ln(s^2/\Lambda^2) + \ln \pi + \psi(n+1) + \psi(n+2) + 2\gamma_{\text{EM}}\}, \quad (9)$$

with  $W_n(M^2) = \overline{W}_n(M^2) - \frac{2}{3}E_n(M^2)$ .

For numerical analysis, we use the standard values and relations for  $m_s$  and the condensates as

$$\begin{aligned} \langle \bar{u}u \rangle &= -(0.25)^3 \text{ GeV}^3, & \langle \bar{s}s \rangle &= f_s \langle \bar{u}u \rangle, \\ \langle g_c^2 G^2 \rangle &= 0.5 \text{ GeV}^4, & m_s &= 0.15 \text{ GeV}, \\ ig_c \langle \bar{u}\sigma \cdot Gu \rangle &= 0.8 \text{ GeV}^2 \langle \bar{u}u \rangle, \\ ig_c \langle \bar{s}\sigma \cdot Gs \rangle &= f_s ig_c \langle \bar{u}\sigma \cdot Gu \rangle \end{aligned} \quad (10)$$

with  $f_s = 0.8$  and  $\Lambda = 0.5 \text{ GeV}$ . Since the QCDSR is proportional to  $\alpha^2 + \beta^2$ , the results are independent of the choice on  $\alpha$  and  $\beta$ .

Defining the right hand side of the sum rule in Eq. (8) by  $L^{\text{OPE}}(M)$ , we analyze its behavior as a function of the Borel mass  $M$ . Shown in Fig. 2 is  $L^{\text{OPE}}(M)$  for the threshold  $s_0 = 1.2 \text{ GeV}$  and  $1.5 \text{ GeV}$ . Here, the dashed, dot-dashed, and solid lines correspond to  $L^{\text{OPE}}(M)$  with the operators of  $d \leq 6$ ,  $d \leq 8$ , and  $d \leq 10$ , respectively. This shows that the contribution from the operators of dimension 8 to the QCDSR is large and negative for both cases. For  $s_0 = 1.2 \text{ GeV}$ , in contradiction with a positive definite value of the left hand side of Eq. (8), the large negative contribution from the operators of dimension 8 makes the full  $L^{\text{OPE}}(M)$  have a definite negative value in the physical Borel region less than the threshold. This is similar to the result found in Ref. [20], where the

QCDSR for the light scalar meson nonet was analyzed by assuming the scalar diquark-antidiquark structure. For  $s_0 = 1.5 \text{ GeV}$ , the contributions from the operators of dimension 6 and 10 are large enough to overcome the negative contribution from the dimension 8 operators in the Borel region  $M \geq 1 \text{ GeV}$ . However, as shown in Fig. 3, it is difficult to find the Borel window, where the fitted mass does not have strong dependence on  $M$ . Furthermore, the fact that the fitted mass is larger than the value of the threshold is in contradiction with the basic concept of the QCDSR. In addition, the ratio of the pole to continuum contributions is found to be very small ( $\sim 0.03$ ), which violates one of the main requirements to have a reliable QCDSR as discussed in Ref. [21].

We have also tested the sum rule with  $s_0 > 1.5 \text{ GeV}$  to find that the Borel region of positive  $L^{\text{OPE}}(M)$  becomes wider. However, the fitted mass is very high (about  $1.8 \text{ GeV}$  for  $s_0 = 2.0 \text{ GeV}$ , for example) compared to the  $f_0(980)$  mass. These observations lead us to conclude that it is hard to consider the  $f_0(980)$  as a pure  $K\bar{K}$  bound state. We also point out that the possible strong deviations of the values of the condensates of dimension 6 and 8 from the factorization hypothesis in the level presented in Ref. [22] does not change our main




FIG. 2.  $L^{\text{OPE}}$  as a function of  $M$  for (a)  $s_0 = 1.2$  GeV and (b)  $s_0 = 1.5$  GeV. Dashed and dot-dashed lines are  $L^{\text{OPE}}$  obtained with the operators of up to dimension 6 and 8, respectively. The solid lines show the full calculation of up to dimension 10.




FIG. 3. The fitted mass from the sum rule (8) as a function of  $M$  for  $s_0 = 1.5$  GeV.

conclusion.

In summary, we have constructed and analyzed the QCDSR within the OPE with the operators of  $d \leq 10$  by assuming the pure  $K\bar{K}$  structure for the  $f_0(980)$ . Our

analyses show that there is no value of the threshold which guarantees the positivity of  $L^{\text{OPE}}$  and weak dependency of the fitted mass for the  $f_0(980)$  on the Borel mass simultaneously. This leads to the conclusion that the  $f_0(980)$  has a very complicated structure other than a pure  $K\bar{K}$  state. Therefore, it would be interesting to investigate an admixture of four quark configurations and two quark configuration for the internal structure of the  $f_0(980)$ .

We are grateful to S. B. Gerasimov for fruitful discussions on various aspects of this research. Y.O. thanks H. Kim for valuable comments. This work was supported in part by the National Research Foundation funded by the Korean Government (Grant No. NRF-2011-220-C00011 and Grant No. NRF-2010-0009381). The work of N.K. was supported in part by the MEST of the Korean Government (Brain Pool Program No. 121S-1-3-0318). We also acknowledge that this work was initiated through the series of APCTP-BLTP JINR Joint Workshop.

- 
- [1] C. Amsler and N. A. Törnqvist, Phys. Rep. **389**, 61 (2004).
  - [2] R. L. Jaffe, Phys. Rev. D **15**, 267 (1977).
  - [3] R. L. Jaffe, Phys. Rep. **409**, 1 (2005).
  - [4] L. Maiani, F. Piccinini, A. D. Polosa, and V. Riquer, Phys. Rev. Lett. **93**, 212002 (2004).
  - [5] T. V. Brito, F. S. Navarra, M. Nielsen, and M. E. Bracco, Phys. Lett. B **608**, 69 (2005).
  - [6] Z.-G. Wang and W.-M. Yang, Eur. Phys. J. C **42**, 89 (2005).
  - [7] H.-X. Chen, A. Hosaka, and S.-L. Zhu, Phys. Rev. D **74**, 054001 (2006); Phys. Lett. B **650**, 369 (2007); Phys. Rev. D **76**, 094025 (2007); H.-X. Chen, A. Hosaka, H. Toki, and S.-L. Zhu, Phys. Rev. D **81**, 114034 (2010).
  - [8] N. Mathur et al., Phys. Rev. D **76**, 114505 (2007).
  - [9] J. Weinstein and N. Isgur, Phys. Rev. Lett. **48**, 659 (1982); Phys. Rev. D **27**, 588 (1983); **41**, 2236 (1990).
  - [10] F. E. Close, N. Isgur, and S. Kumano, Nucl. Phys. **B389**, 513 (1993).
  - [11] N. N. Achasov, V. V. Gubin, and V. I. Shevchenko, Phys. Rev. D **56**, 203 (1997).
  - [12] V. Baru et al., Phys. Lett. B **586**, 53 (2004).
  - [13] T. Branz, T. Gutsche, and V. E. Lyubovitskij, Eur. Phys. J. A **37**, 303 (2008).
  - [14] Yu. S. Surovtsev, P. Bydzovsky, and V. E. Lyubovitskij, Phys. Rev. D **85**, 036002 (2012).
  - [15] A. V. Anisovich et al., hep-ph/0508260.
  - [16] D. Black, M. Harada, and J. Schechter, Phys. Rev. Lett. **88**, 181603 (2002); Phys. Rev. D **73**, 054017 (2006); D. Harnett et al., Nucl. Phys. A **850**, 110 (2011).
  - [17] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. **B147**, 385 (1979).

- [18] H.-J. Lee, New Physics (Sae Mulli) **61**, 586 (2011).
- [19] H.-J. Lee, N. I. Kochelev, and V. Vento, Phys. Rev. D **73**, 014010 (2006).
- [20] H.-J. Lee, Eur. Phys. J. A **30**, 423 (2006).
- [21] M. E. Bracco, M. Chiapparini, F. S. Navarra, and M. Nielsen, Prog. Part. Nucl. Phys. **67**, 1019 (2012).
- [22] S. Narison, Phys. Lett. B **624**, 223 (2005).