

A 125 GeV Scalar Boson and $SU(N_{TC}) \otimes SU(3)_L \otimes U(1)_X$ models

A. Doff¹ and A. A. Natale^{2,3}

¹*Universidade Tecnológica Federal do Paraná - UTFPR - DAFIS,
Av. Monteiro Lobato Km 04, 84016-210, Ponta Grossa - PR, Brazil*

²*Centro de Ciências Naturais e Humanas,
Universidade Federal do ABC, 09210-170, Santo André - SP, Brazil*

³*Instituto de Física Teórica, UNESP,
Rua Dr. Bento T. Ferraz, 271, Bloco II,
01140-070, São Paulo - SP, Brazil*

(Dated: February 7, 2022)

Abstract

We verify that $SU(N)_{TC} \otimes SU(3)_L \otimes U(1)_X$ models, where the gauge symmetry breaking is totally dynamical and promoted by the non-Abelian technicolor (TC) group and the strong Abelian interactions, are quite constrained by the LHC data. The theory contains a T quark self-energy involving the mixing between the neutral gauge bosons, which introduces the coupling between the light and heavy composite scalar bosons of the model. We determine the lightest scalar boson mass for these models from an effective action for composite operators, assuming details about the dynamics of the strong interaction theories. Comparing the value of this mass with the ATLAS and CMS observation of a new boson with a mass $M_\phi \sim 125\text{GeV}$ and considering the lower bound determined by the LHC Collaborations on the heavy neutral gauge boson (Z') present in these models, we can establish constraints on the possible models. For example, if $SU(N)_{TC} \equiv SU(2)_{TC}$, with technifermions in the fundamental representation, the model barely survives the confrontation with the LHC data.

I. INTRODUCTION

The Standard Model (SM) of electroweak interactions is in excellent agreement with the experimental data and has explained many features of particle physics throughout the years. Despite its success there are some points in the model as, for instance, the enormous range of masses between the lightest and heaviest fermions and other peculiarities that could be better explained at a deeper level assuming the introduction of new fields or symmetries.

Recently the ATLAS and CMS collaborations reported the observation of a new boson with a mass $M_\phi \sim 125\text{GeV}$ which is suspected to be the SM Higgs boson. The current data on this boson diphoton event rate exhibit the signal strength about (1.5 - 2) larger than the one expected for the standard model Higgs boson. There are already many SM extensions trying to explain this possible enhancement of the $\gamma\gamma$ decay. In particular, the increase of this decay rate is natural in the context of a 3-3-1 model [2, 3] and its alternative version with exotic leptons[4], due to the presence of an extra charged vector boson and a doubly charged one as discussed in Ref.[5].

This class of models predicts interesting new physics at TeV scale [6] and addresses some fundamental questions that cannot be explained in the framework of the Standard Model[7, 8]. These models also contain a set of fundamental scalar bosons, with many parameters and clearly suffering from the problems of naturalness and hierarchy [9, 10]. However in Refs.[11, 12] it was suggested that the gauge symmetry breaking in some versions of the 3-3-1 model [4] could be promoted dynamically, because at the scale of a few TeVs the $U(1)_X$ coupling constant becomes strong and the exotic quark T that appears in the model forms a condensate breaking $SU(3)_L \otimes U(1)_X$ to the SM electroweak symmetry. This is a very interesting feature and peculiar to this class of models. Unfortunately the SM gauge symmetry still remains intact, and the nice characteristics of the model could be missed with the introduction of a elementary scalar field in order to break the electroweak gauge symmetry, leading to an unpleasant system of composite and elementary fields responsible for the gauge symmetry breaking.

In Ref.[13] the full realization of the dynamical symmetry breaking of a $SU(3)_L \otimes U(1)_X$ extension of the SM [4] was explored. This was accomplished assuming the gauge symmetry $SU(2)_{TC} \otimes SU(3)_L \otimes U(1)_X$, where the electroweak symmetry is broken dynamically by a technifermion condensate generated by the $SU(2)_{TC}$ Technicolor (TC) gauge group, i.e.

besides the exotic T quark condensate and respective composite scalar, we do now have another composite scalar boson formed by $SU(2)_{TC}$ technifermions. This symmetry breaking also occurs when we exchange the $SU(2)_{TC}$ group by the $SU(N)_{TC}$ group, as well as when we deal with different technifermion representations [14].

In 3-3-1 models where the gauge symmetry breaking is promoted by elementary scalar fields, the many parameters in the scalar potential can be varied in a large range leaving space to scape, up to now, to the LHC experimental constraints. However, in the case where the gauge symmetry breaking is totally dynamical, once we describe the possible dynamics of the theory, we may already have some limitation on the possible models. The study of possible constraints in this class of models is the main motivation of this work. We compute the effective potential for composite operators of a class of 3-3-1 models where the gauge symmetry is dynamically broken, with the main purpose of determining the composite scalar masses. If, for instance, we consider $SU(N)_{TC} \equiv SU(2)_{TC}$, we verify that it is quite difficult to generate a scalar boson mass of 125 GeV, assuming that it is a composite scalar that has been observed at the LHC, obeying, at the same time, the lower limit on the Z' .

The composite scalar system of these 3-3-1 models have a mixing related to the Z and Z' mixing, which is present in the exotic T quark self-energy. This mixing will appear in the calculation of the effective potential for composite operators [16], which, when minimized, supply the physical scalar masses, and it is important to know its amount because the scalar masses may be modified by this effect. There are other possible contributions to this mixing, that would appear in an extended theory necessary to explain the fermion masses, whose spectrum has no explanation in any dynamical symmetry breaking model up to now. However these corrections are expected to be small compared to the ones that we discuss here, and it will become clear that already at this level some versions of these models may be excluded by the recent LHC data.

The distribution of our paper is the following: In Section II we present the main aspects of the $SU(N_{TC}) \otimes SU(3)_L \otimes U(1)_X$ models. In Section III we discuss the mixing of the neutral gauge boson system, its relation to the T quark dynamical mass and present the self-energies that will be used in the effective action calculation. The mixing in the T quark self-energy will be responsible by the coupling between the composite scalars (ϕ_T, ϕ) , associated respectively with the breaking of symmetries $SU(3)_L \otimes U(1)_X$ and $SU(2)_L \otimes U(1)_Y$ in the effective potential. Section IV contains the calculation of the effective action and in

Section V we compute numerically the scalar boson masses for different TC groups. Finally, in Section VI we draw our conclusions.

II. $SU(N)_{TC} \otimes SU(3)_L \otimes U(1)_X$ MODELS

On the lines below we describe the main features of the models, which are similar to that proposed in Ref.[13], the fermionic content has the following form

$$Q_{3L} = \begin{pmatrix} t \\ b \\ T \end{pmatrix}_L \sim (\mathbf{1}, \mathbf{3}, 2/3)$$

$$t_R \sim (\mathbf{1}, \mathbf{1}, 2/3), b_R \sim (\mathbf{1}, \mathbf{1}, -1/3)$$

$$T_R \sim (\mathbf{1}, \mathbf{1}, 5/3)$$

$$Q_{\alpha L} = \begin{pmatrix} D \\ u \\ d \end{pmatrix}_{\alpha L} \sim (\mathbf{1}, \mathbf{3}^*, -1/3)$$

$$u_{\alpha R} \sim (\mathbf{1}, \mathbf{1}, 2/3), d_{\alpha R} \sim (\mathbf{1}, \mathbf{1}, -1/3)$$

$$D_{\alpha R} \sim (\mathbf{1}, \mathbf{1}, -4/3) \tag{1}$$

where $\alpha = 1, 2$ is the family index and we represent the third quark family by Q_{3L} . In these expressions $(\mathbf{1}, \mathbf{3}, X)$, $(\mathbf{1}, \mathbf{3}^*, X)$ or $(\mathbf{1}, \mathbf{1}, X)$ denote the transformation properties under $SU(N_{TC}) \otimes SU(3)_L \otimes U(1)_X$ and X is the corresponding $U(1)_X$ charge. The leptonic sector includes besides the conventional charged leptons and their respective neutrinos the charged heavy leptons E_a [4].

$$l_{aL} = \begin{pmatrix} \nu_a \\ l_a \\ E_a^c \end{pmatrix}_L \sim (\mathbf{1}, \mathbf{3}, 0) \tag{2}$$

where $a = 1, 2, 3$ is the family index and l_{aL} transforms as triplets under $SU(3)_L$. Moreover, we have to add the corresponding right-handed components, $l_{aR} \sim (\mathbf{1}, \mathbf{1}, -1)$ and $E_{aR}^c \sim (\mathbf{1}, \mathbf{1}, +1)$.

The fermionic content associated with the TC sector has the form

$$\begin{aligned}\Psi_{1L} &= \begin{pmatrix} U_1 \\ D_1 \\ U' \end{pmatrix}_L \sim (\mathbf{N}_{\mathbf{TC}}, \mathbf{3}, 1/2) \\ U_{1R} &\sim (\mathbf{N}_{\mathbf{TC}}, \mathbf{1}, 1/2), \quad D_{1R} \sim (\mathbf{N}_{\mathbf{TC}}, \mathbf{1}, -1/2) \\ U'_R &\sim (\mathbf{N}_{\mathbf{TC}}, \mathbf{1}, 3/2), \\ \Psi_{2L} &= \begin{pmatrix} D' \\ U_2 \\ D_2 \end{pmatrix}_L \sim (\mathbf{N}_{\mathbf{TC}}, \mathbf{3}^*, -1/2) \\ U_{2R} &\sim (\mathbf{N}_{\mathbf{TC}}, \mathbf{1}, 1/2), \quad D_{2R} \sim (\mathbf{N}_{\mathbf{TC}}, \mathbf{1}, -1/2) \\ D'_R &\sim (\mathbf{N}_{\mathbf{TC}}, \mathbf{1}, -3/2).\end{aligned}\tag{3}$$

where 1 and 2 label the first and second techniquark families, U' and D' can be considered as exotic techniquarks making an analogy with quarks T and D that appear in the ordinary fermionic content of the model. The model is anomaly free if we have equal numbers of triplets and antitriplets, counting the color of $SU(3)_c$. Therefore, in order to make the model anomaly free two of the three quark generations transform as $\mathbf{3}^*$, the third quark family and the three leptons generations transform as $\mathbf{3}$. It is easy to check that all gauge anomalies cancel out in this model, in the TC sector the triangular anomaly cancels between the two generations of technifermions. In the present version of the model the technifermions are singlets of $SU(3)_c$.

As pointed out in Refs.[11, 12], one interesting feature of the versions[2–4] of 3-3-1 models is the following relationship among the coupling constants g and g' associated to the gauge group $SU(3)_L \otimes U(1)_X$

$$t^2 \equiv \frac{\alpha'}{\alpha} = \frac{\alpha_X}{\alpha} = \frac{\sin^2 \theta_w(\mu)}{1 - 4 \sin^2 \theta_w(\mu)}\tag{4}$$

where $\alpha = g^2/4\pi$, $\alpha' = g'^2/4\pi$ and θ_w is the electroweak mixing angle. According to the discussion presented in[11, 12], it is precisely this feature of the model that allows the gauge

symmetry breaking of this version of the $SU(3)_L \otimes U(1)_X$ model to the SM symmetry, because at the scale of a few TeVs the $U(1)_X$ coupling constant becomes strong as we approach the peak existent in Eq.(4). Therefore, in the model described in this section the exotic quark T will form a condensate breaking $SU(3)_L \otimes U(1)_X$ to the electroweak gauge symmetry, while the SM gauge symmetry will be broken dynamically by a technifermion condensate.

In order to compute the effective action generated for the composite scalar bosons resulting from the two symmetry breaking stages, in the next section we discuss the T quark self-energy, that is related to the mixing between the standard model neutral gauge boson Z with the Z' boson.

III. THE (Z' - Z) MIXING AND SELF-ENERGIES

In the models that we consider here there is a mixing between the standard model neutral gauge boson Z with the Z' boson, the mass eigenstates are[17]

$$Z_1 = Z \cos \theta - Z' \sin \theta \quad (5)$$

$$Z_2 = Z' \cos \theta + Z \sin \theta \quad (6)$$

where the mixing angle(θ) is given by[17]

$$\tan \theta = \frac{M_Z^2 - M_{Z_1}^2}{M_{Z_2}^2 - M_Z^2}, \quad (7)$$

in this case (Z_1) represents the SM neutral boson and (Z_2) corresponds to the additional 3-3-1 heavy neutral boson. Therefore, assuming this mixing we can write in the Euclidean space the following linearized gap equation ($\Sigma_T(p^2)$) for the T quark:

$$\begin{aligned} \Sigma_T(p^2) = & a \cos \theta \int dk^2 k^2 \frac{\Sigma_T(k^2)}{[k^2 + \mu_X^2]} \frac{1}{[(p-k)^2 + M_{Z'}^2]} + \\ & a \sin \theta \int dk^2 k^2 \frac{\Sigma_T(k^2)}{[k^2 + \mu_X^2]} \frac{1}{[(p-k)^2 + M_Z^2]} \end{aligned} \quad (8)$$

where

$$a = \frac{3g_X^2 X_L X_R}{16\pi^2},$$

μ_X is the energy scale where the $U(1)_X$ interaction becomes sufficiently strong to break dynamically the $SU(3)_L \otimes U(1)_X$ to $SU(2)_L \otimes U(1)_Y$, g_X^2 is the $U(1)_X$ coupling constant,

X_L and X_R are respectively $U(1)_X$ charges attributed to the chiral components of the exotic quark T .

Besides the condensate and composite states (scalar and pseudoscalar) associated to Eq.(8), we have similar entities due to the $SU(N)_{TC}$ group condensation at the scale μ_{TC} , generated by a non-trivial technifermion self-energy ($\Sigma_{TC}^2(p^2)$). As discussed in Ref.[13] the technifermions multiplets, Ψ_1 and Ψ_2 described in Eq.(3), lead to the formation of composite scalar bosons (ϕ_1 and ϕ_2) that are equivalents to the set of fundamental scalar fields, ρ and η [2, 4], so, in order to obtain a structure of the scalar potential similar the described in[2, 4], we will assume that

$$\phi = \frac{\phi_1 + \phi_2}{\sqrt{2}}, \quad (9)$$

this normalization results from the fact that $f_\pi^2 \propto \phi^2$ (as we shall describe in Eq.(29)), and from this it is possible to verify that

$$\phi^2 = \phi_1^2 + \phi_2^2, \quad (10)$$

once $f_\pi^2 = f_{\pi_1}^2 + f_{\pi_2}^2$, and $\langle \phi_1 \rangle = \langle \phi_2 \rangle$, typical of the two technifermion generations of Eq.(3). The f_{π_i} are the technipions decay constants that can be computed through the linearized Pagels and Stokar relation[15]

$$f_\pi^2 \approx \frac{N_{TC}}{4\pi^2} \int \frac{dp^2 p^2 \Sigma_{TC}^2(p^2)}{(p^2 + \mu_{TC}^2)^2} \quad (11)$$

whereas the pseudoscalar decay constant associated to T quark self-energy, $\Sigma'_T(p^2)$ will be written as

$$F_\pi^2 \approx \frac{1}{4\pi^2} \int \frac{dp^2 p^2 \Sigma_T^2(p^2)}{(p^2 + \mu_X^2)^2}. \quad (12)$$

To compute the effective potential for composite operators [16] we need to know the self-energies of the strongly interacting fermions: The T quark and the fermions with $SU(N)_{TC}$ charges. Eq.(8) has two possible solutions, and in the program developed in Refs.[18, 19] it was verified that the solution falling slowly with the momentum is the dominating one if suitable new interactions are assumed to be relevant at the scale of the (UV) cutoff. In their case the gauge boson mass integrals receive significant contributions from a very large range of loop momenta and the SM gauge boson masses M_W and M_Z turn out to be of similar magnitude when compared to the top quark mass. This is exactly the situation that we have in the approach proposed to promote the gauge symmetry breaking of $SU(3)_L \times U(1)_X$ to the

electroweak symmetry, where at the scale of a few TeVs the $U(1)_x$ coupling constant becomes strong as we approach the peak existent in Eq.(4). In the Ref.[11] after the numerical calculation of M_T , it was found that the magnitudes of $M_{Z'}$ and M_T are the same order. Therefore, considering the above comments we will assume that the solution of Eq.(8) is giving by

$$\Sigma_T(p^2) \approx \Sigma'_T(p^2) \left(1 - h(\omega) \ln \left(\frac{p^2}{\mu_X^2} \right) \tan \theta \right), \quad (13)$$

and is the one that will be used for the T quark self-energy to determine the effective potential (Ω_T). To write Eq.(13) we defined the following quantities:

$$\Sigma'_T(p^2) = \mu_X \left(\frac{p^2}{\mu_X^2} \right)^{-\left(\frac{1-\omega}{2}\right)},$$

where $\omega = \sqrt{1-4A}$, $A = a \cos \theta$ and

$$h(\omega) = \left(1 - \frac{1}{\omega} + \frac{A}{\omega} \right). \quad (14)$$

In the case of fermions with $SU(N)_{TC}$ charges the self-energy will be given by

$$\Sigma(p^2) \sim \mu_{TC} [1 + bg^2 \ln(p^2/\mu_{TC}^2)]^{-\gamma}, \quad (15)$$

where μ_{TC} is the $SU(N)_{TC}$ characteristic scale of mass generation,

$$\gamma = 3c/16\pi^2 b \quad (16)$$

and $c = \frac{1}{2} [C_2(R_1) + C_2(R_2) - C_2(R_3)]$ where $C_2(R_i)$ are the Casimir operators for fermions in the representations R_1 and R_2 that condense in the representation R_3 , $b = (11N - 2N_f)/48\pi^2$ for the $SU(N)_{TC}$ group with N_f flavors, g^2 is the coupling constant for which we assume the expression

$$g^2(k^2) = \frac{1}{b \ln[(k^2 + 4m_g^2)/\Lambda^2]}, \quad (17)$$

where, m_g is an infrared dynamical gauge boson mass, whose phenomenologically preferred value is $m_g \approx 2\Lambda$ [20, 21], and we will set $\Lambda = \mu_{TC}$. Note that, using the above coupling constant, we are assuming that non-Abelian gauge theories generate dynamical masses for their gauge bosons [20, 22]. As a consequence it is expected that confinement should be necessary to generate non-trivial fermionic self-energies [23, 24], and the expression for the

self-energy is the one of Eq.(15), as discussed at length in Ref.[25]. The main features of Eq.(15), is that it causes the decoupling of heavier degrees of freedom in models where there is an interaction connecting different fermionic families [25], it leads to the deepest minimum of energy, with a vacuum expectation value proportional to $1/g^2$ [26–28], it is the only self-energy able to naturally explain fermion masses as heavy as the top quark [29], as well as the unique possible form of solution that may generate a light composite scalar boson [30, 31].

In the self-energies that we discussed above the characteristic scales μ_X and μ_{TC} have not been determined up to now. However they should be constrained by the value of the $SU(3)_L \otimes U(1)_X$ gauge boson masses. In order to do so we notice that for $M_{Z'} \gg M_Z$, it is possible to show that[17]

$$\tan \theta \approx \frac{1}{2\sqrt{3}t^2} \frac{M_Z^2}{M_{Z'}^2}, \quad (18)$$

assuming the result described in Ref.[13] we obtain the masses

$$\begin{aligned} M_Z^2 &= \frac{g^2}{4} (f_{\pi_1}^2 + f_{\pi_2}^2) \left[\frac{1 + 4t^2}{1 + 3t^2} \right] \\ &= \frac{g^2}{4} f_\pi^2 \left[\frac{1 + 4t^2}{1 + 3t^2} \right], \end{aligned} \quad (19)$$

$$M_{Z'}^2 = \frac{g^2}{4} F_\Pi^2 \left[\frac{4}{3} + 4t^2 \right] .. \quad (20)$$

With these masses and Eq.(18), we can write Eq.(13) in the form

$$\Sigma_T(p^2) \approx \Sigma'_T(p^2) \left(1 + A(\omega) \frac{f_\pi^2}{F_\Pi^2} \ln \left(\frac{p^2}{\mu_X^2} \right) \right), \quad (21)$$

where for $SU(N)_{TC}$

$$A(\omega) = -\frac{\sqrt{3}N_{TC}}{16t^2} \frac{(1 + 4t^2)}{(1 + 3t^2)^2} h(\omega).$$

The f_π decay constant is related to the SM vacuum expectation value(vev) through

$$f_\pi^2 = (f_{\pi_1}^2 + f_{\pi_2}^2) = v^2 = \frac{4M_W^2}{g^2} = (246 GeV)^2. \quad (22)$$

and in the case of T quark self-energy, since there are no evidences for the Z' boson, we just assume $F_\Pi \sim O(\mu_X) \sim O(TeV)$. Equations (15) and (21) are the main ingredients to compute the effective action for the model described in section II.

IV. THE EFFECTIVE ACTION FOR COMPOSITE SCALAR BOSONS OF THE $SU(N_{TC}) \otimes SU(3)_L \otimes U(1)_X$ MODEL

The effective potential for composite operators [16] is a function of the Green's functions of the theory, in particular it can be written as a function of the complete fermion (S) and gauge boson (D) propagators as

$$V(S, D) = -i \int \frac{d^4 p}{(2\pi)^4} \text{Tr} (\ln S_0^{-1} S - S_0^{-1} S + 1) + V_2(S, D), \quad (23)$$

where S_0 (and D_0) stands for the bare fermion (gauge boson) propagator and $V_2(S, D)$ is the sum of all two-particle irreducible vacuum diagrams. The physically meaningful quantity that we must compute is the vacuum energy density given by

$$\Omega_V = V(S, D) - V(S_0, D_0), \quad (24)$$

where we are subtracting the symmetric part of the potential from the potential that admits condensation in the scalar channel, that is denoted by $V(S_0, D_0)$ and is a function of the perturbative propagators (S_0 and D_0).

The vacuum energy density, if we remove all indices and integrations, can be written as [16, 26]

$$\begin{aligned} \Omega_V = & -i \text{Tr} (\ln S_0^{-1} S - S_0^{-1} S + 1) + i \text{Tr} \Sigma (S - S_0) \\ & + \frac{1}{2} i \text{Tr} (\Gamma S \Gamma S - \Gamma S_0 \Gamma S_0) D. \end{aligned} \quad (25)$$

The self-energies described by Eqs.(15) and (21) enter into the definition of S .

How the effective action for composite scalar bosons emerge from Eq.(25) in the case of a dynamically broken gauge theory including the kinetic term has been detailed by Cornwall and Shellard [32], and is also discussed in Refs.[26, 33]. Here we will skip lengthy details and follow closely the work of Ref.[33], where it was shown that the effective action generated for a TC model could be written in the following way

$$\Omega_{TC} = \int d^4 x \left[\frac{1}{2Z_{TC}} \partial_\mu \phi \partial^\mu \phi - \frac{\lambda_{4TC}}{4} \phi^4 - \frac{\lambda_{6TC}}{6} \phi^6 - \dots \right]. \quad (26)$$

The effective scalar field $\phi(x)$ acts like a dynamical effective scalar field with anomalous dimension 2γ , is related to the bilinear self-energy $\Sigma(k, p) \propto \phi(k)\Sigma(p)$, it is seen as a

variational parameter, and the kinetic term for our effective theory is obtained inserting $\phi(k)$ in the effective action and expanding around $k = 0$ [26, 32, 33]. In the above equation

$$Z_{TC} \approx \frac{4\pi^2\beta(2\gamma - 1)}{N_{TC}N_f}, \quad (27)$$

and

$$\begin{aligned} \lambda_{4TC} &= \frac{N_{TC}N_f}{4\pi^2} \left(\frac{1}{\beta(4\delta - 1)} + \frac{1}{2} \right) \\ \lambda_{6TC} &= -\frac{N_{TC}N_f}{4\pi^2} \left[\frac{1}{\mu_{TC}^2} \right]. \end{aligned} \quad (28)$$

In this expressions, $\beta = bg^2$, N_f denote the number of technifermions, γ has been defined in Eq.(16) and is calculated for the respective TC representations.

In Eq.(26) a term proportional to ϕ^2 does not appear because we assume that the self-energies are exact solutions of the linearized gap equations [16], also odd terms in ϕ do not appear because we do not have current fermion masses. The constant Z_{TC} arises when the contribution of the kinetic term is included in the calculation of the effective action[26, 32, 33], and this acts as a normalization constant. This contribution is important in our calculation because it will give the correct normalization of the effective fields, ϕ and ϕ_T , as discussed in the Refs.[26, 32, 33]. In terms of these fields we can also write the decay constants for the TC and T fermions as

$$f_\pi^2 = \frac{N_{TC}}{4\pi^2} \frac{\phi^2}{\beta} \frac{1}{(2\gamma - 1)}. \quad (29)$$

While for the self-energy, $\Sigma'_T(p^2)$, we obtain the following relation

$$F_\Pi^2 = \frac{\phi_T^2}{4\pi^2} \frac{1}{2a}, \quad (30)$$

and in this case $\langle \phi_T \rangle \simeq \mu_x$.

We can now present the contribution to the effective action due to the composite scalar boson formed by the strong interaction of the exotic T quark. Below we show the ϕ_T^4 and ϕ_T^6 terms of the effective potential Ω_T and the corrections $\Delta\Omega_T$ assuming the mixture in Eq.(13) and the comments leading to Eq.(21):

$$\begin{aligned} \Omega_T &= \int d^4x \left[\frac{1}{2Z_T} \partial_\mu \phi_T \partial^\mu \phi_T - \frac{\lambda_{4T}}{4} \phi_T^4 - \frac{\lambda_{6T}}{6} \phi_T^6 + \right. \\ &\quad \left. - \frac{\Delta\lambda_{4T}}{4} \phi_T^2 \phi^2 - \frac{\Delta\lambda_{6T}}{6} \phi_T^4 \phi^2 + \dots \right] \end{aligned} \quad (31)$$

where we identify

$$\begin{aligned} Z_T &\approx 8\pi^2 a \quad , \quad \lambda_{4T} \approx \frac{1}{4\pi^2} \left(\frac{1}{4a} + \frac{1}{4} \right) \\ \lambda_{6T} &\approx -\frac{1}{4\pi^2} \frac{1}{\mu_X^2} \quad , \quad \Delta\lambda_{4T} = \frac{a\Delta\lambda_4}{\pi^2} \left(\frac{1}{4a} + \frac{1}{4} \right) \end{aligned} \quad (32)$$

$$\begin{aligned} \Delta\lambda_{6T} &= -\frac{\Delta\lambda_4}{4\pi^2} \frac{24a^2}{1+2a} \frac{1}{\mu_X^2} \\ \Delta\lambda_4 &= \frac{\sqrt{3}N_{TC}}{16\beta t^2} \left(\frac{1}{(2\gamma-1)(1-2a)} \right) \\ &\quad \times \frac{(1+4t^2)}{(1+3t^2)^2}, \end{aligned} \quad (33)$$

In these expressions we assumed the existence of just one exotic quark that condenses in the most attractive channel(MAC) [34].

In order to reproduce a standard scalar effective field theory we introduce in our effective Lagrangian the normalized fields

$$\Phi(x) = Z_{TC}^{-1/2} \phi(x) \quad , \quad (34)$$

$$\Phi_T(x) = Z_T^{-1/2} \phi_T(x) \quad . \quad (35)$$

Now, considering the Eqs.(9), where we see that the field ϕ actually represent two fields (ϕ_1 and ϕ_2), and adding Eq.(26) to Eq.(31) we can write down the full effective action in terms of the normalized fields Φ_T and $\Phi(x)$ (composed by Φ_1 and Φ_2):

$$\begin{aligned} \Omega(\Phi_T, \Phi) = \int d^4x \left[\frac{1}{2} \partial_\mu \Phi_1 \partial^\mu \Phi_1 + \frac{1}{2} \partial_\mu \Phi_2 \partial^\mu \Phi_2 + \frac{1}{2} \partial_\mu \Phi_T \partial^\mu \Phi_T - \frac{\lambda_{4T}^R}{4} \Phi_T^4 - \frac{\lambda_{4TC}^{R(a)}}{4} \Phi_1^4 - \frac{\lambda_{4TC}^{R(a)}}{4} \Phi_2^4 \right. \\ - \frac{\lambda_{4TC}^{R(a)}}{4} \Phi_1^2 \Phi_2^2 - \frac{\lambda_{4TC}^{R(a)}}{4} \Phi_2^2 \Phi_1^2 - \frac{\lambda_{4TC}^{R(b)}}{4} \Phi_T^2 \Phi_1^2 - \frac{\lambda_{4TC}^{R(b)}}{4} \Phi_T^2 \Phi_2^2 - \frac{\lambda_{6T}^R}{6} \Phi_T^6 - \frac{\lambda_{6TC}^{R(a)}}{4} \Phi_1^6 - \frac{\lambda_{6TC}^{R(a)}}{4} \Phi_2^6 \\ - \frac{\lambda_{6TC}^{R(a)}}{4} \Phi_1^2 \Phi_2^4 - \frac{\lambda_{6TC}^{R(a)}}{4} \Phi_2^2 \Phi_1^4 - \frac{\lambda_{6TC}^{R(a)}}{4} \Phi_1^2 \Phi_2^2 \Phi_1^2 - \frac{\lambda_{6TC}^{R(a)}}{4} \Phi_2^2 \Phi_1^2 \Phi_2^2 - \frac{\lambda_{6TC}^{R(b)}}{4} \Phi_T^4 \Phi_1^2 - \frac{\lambda_{6TC}^{R(b)}}{4} \Phi_T^4 \Phi_2^2 \\ \left. - \frac{\lambda_{6TC}^{R(b)}}{4} \Phi_T^4 \Phi_1 \Phi_2 - \frac{\lambda_{6TC}^{R(b)}}{4} \Phi_T^4 \Phi_2 \Phi_1 \right] \end{aligned} \quad (36)$$

where the normalized couplings for the composite fields, ϕ_T , Φ_1 and Φ_2 , are given respectively by

$$\begin{aligned}
\lambda_{4T}^R &= \frac{Z_T^2}{4\pi^2} \left(\frac{1}{4a} + \frac{1}{4} \right), \quad \lambda_{4TC}^{R(a)} = \frac{N_{TC} N_f Z_{TC}^2}{4\pi^2} \left(\frac{1}{\beta(4\gamma - 1)} + \frac{1}{2} \right) \\
\lambda_{4TC}^{R(b)} &= \frac{a\Delta\lambda_4 Z_T Z_{TC}}{\pi^2} \left(\frac{1}{4a} + \frac{1}{4} \right), \quad \lambda_{6T}^R = -\frac{Z_T^3}{4\pi^2} \frac{1}{\mu_X^2} \\
\lambda_{6TC}^{R(a)} &= -\frac{N_{TC} N_f Z_{TC}^3}{4\pi^2} \frac{1}{\mu_{TC}^2}, \quad \lambda_{6TC}^{R(b)} = -\frac{\Delta\lambda_4 Z_T^2 Z_{TC}}{\pi^2} \frac{6a^2}{1 + 2a} \frac{1}{\mu_X^2}.
\end{aligned} \tag{37}$$

V. THE 125 GEV SCALAR BOSON AND THE EFFECTIVE ACTION FOR THE $SU(N_{TC}) \otimes SU(3)_L \otimes U(1)_X$ MODEL

To compute the scalar masses from the effective Lagrangian described in Eq.(36) we use

$$M_{\Phi_i}^2 = \frac{\partial^2 \Omega(\Phi_T, \Phi)}{\partial \Phi_i^2} \Big|_{\Phi_i = \Phi_{i_{min}}}. \tag{38}$$

where $i = 1, 2$, and $(1, 2)$ label the first and second techniquark families. After neglecting terms of higher order when substituting the minimum value in the potential we obtain

$$M_{\Phi_i}^2 \approx 2\lambda_{4TC}^{R(a)} \left(\frac{\lambda_{4TC}^{R(a)}}{\lambda_{6TC}^{R(a)}} \right) + 2\lambda_{4TC}^{R(b)} \left(\frac{\lambda_{4TC}^{R(b)}}{\lambda_{6TC}^{R(b)}} \right). \tag{39}$$

Assuming the above equation, with couplings $\lambda_{4TC}^{R(a)}$, $\lambda_{6TC}^{R(a)}$, $\lambda_{4TC}^{R(b)}$ and $\lambda_{6TC}^{R(b)}$ defined by Eq.(37), we finally can make a scan in the parameter space $[\mu_{TC} \times \mu_X]$ limiting the Higgs mass to the range $120\text{GeV} < M_H < 130\text{GeV}$, in order to exclude possible candidates for TC models. Note that for this purpose we assume that the ATLAS and CMS are indeed observing a new scalar boson with a mass $M_\phi \sim 125\text{GeV}$, and consider also the strong limit on the Z' mass announced by these collaborations[35].

The extra Z' boson is predicted in many extensions of the Standard Model at the TeV mass scale, as in the Sequential Standard Model (Z_{ssm})[36], with SM like couplings. With LHC data at $(\sqrt{s} = 8\text{TeV})$, recently the ATLAS and CMS collaborations placed strong constraints on the mass of these particles[35]. These constraints depend on the knowledge of the coupling of this boson with SM fermions. In the case of the Z_{ssm} model with SM like couplings the Z' mass can be excluded below 2.49 TeV. This limit can also be taken as a lower limit on the Z' mass of the models discussed here. In this particular case, if $M_{Z'} > 2.49\text{TeV}$ the energy scale μ_X should be limited to $\mu_X > 1.1\text{TeV}$. In Fig.1 we present the allowed region of parameters for the $SU(2)_{TC}$, $SU(3)_{TC}$ and $SU(4)_{TC}$ cases, with a scalar

composite “Higgs” mass range $120\text{GeV} < M_H < 130\text{GeV}$. The solid black line corresponds to the lower limit on μ_X , and from this figure we verify that if $SU(N)_{TC} \equiv SU(2)_{TC}$ the model barely survives the confrontation with the LHC data.




FIG. 1: The region of parameters, μ_{TC} and μ_X , for a composite scalar mass range $120\text{GeV} < M_H < 130\text{GeV}$ in the case of $SU(2)_{TC}$, $SU(3)_{TC}$ and $SU(4)_{TC}$ TC groups. The region in red corresponds to $SU(2)_{TC}$, the blue region to $SU(3)_{TC}$ and the yellow region to $SU(4)_{TC}$. In this figure we assumed $N_f = 6$ for all groups. The solid line corresponds to the lower limit on μ_X (i.e. $M_{Z'}$).

In order to complement the analysis, in the Fig.2 we show the interval of parameters in the $SU(3)_{TC}$ case assuming $N_f = 8$ and $N_f = 10$. As can be noticed in the Fig.2, the range of values that the parameters μ_{TC} and μ_X can assume decreases when the number of technifermions is increased. The N_f dependence on the kinetic term is important and this one is transferred to the λ'_6 s couplings and results in a decrease of the scalar mass.

We can write the Eq.(39) in the following approximate form

$$M_{\Phi_i} \propto \delta_1 \sqrt{\beta(2\gamma - 1)} \mu_{TC} + \delta_2 \sqrt{\beta(2\gamma - 1)} \mu_X \quad (40)$$

where δ_1 and δ_2 are constants. The increase of the number of fermions implies in an increase of $\sqrt{\beta(2\gamma - 1)}$, then to keep the interval to the Higgs mass $120\text{GeV} < M_H < 130\text{GeV}$ we

observe a decrease in the area $[\mu_{TC} \times \mu_X]$ in the parameter space. This effect is the same that appears in the normalization of the Bethe-Salpeter wave function discussed in Ref.[30].




FIG. 2: The region of parameters, μ_{TC} and μ_X , for a composite scalar mass range $120\text{GeV} < M_H < 130\text{GeV}$ in the $SU(3)_{TC}$ case: In this figure the blue region corresponds to $N_f = 6$, the light blue to $N_f = 8$ and magenta to $N_f = 10$.

In the Fig.3 we show the interval of parameters in the $SU(2)_{TC}$ case assuming $N_f = 7$ (in Green), and again we verify that $SU(2)_{TC}$ case with this number of fermions can be ruled out in confrontation with the LHC data.

VI. CONCLUSIONS

In this work we computed an effective action for the composite scalar boson system, ϕ_T , ϕ_1 and ϕ_2 , formed by the fermions and technifermions Q_3 , Ψ_1 and Ψ_2 described in the Eqs.(1) and (3). We include in this calculation the kinetic term of the effective theory. This term is important because it provides a normalization factor for the effective scalar boson Lagrangian. The effective Lagrangian is then normalized in order to reproduce a standard scalar effective field theory, leading to a non-trivial set of scalar self-couplings. From this




FIG. 3: The region of parameters, μ_{TC} and μ_X , for a composite scalar mass range $120\text{GeV} < M_H < 130\text{GeV}$ in the $SU(2)_{TC}$ case: The small Green region corresponds to $N_f = 7$, and the solid line corresponds to the lower limit on μ_X .

Lagrangian we can determine the scalar boson masses of the theory.

To compute the effective action for the model described in the Section II, we firstly determined the correction to the T quark self-energy that result from the mixing between the standard model neutral gauge boson Z with the Z' boson. We show that this correction is the responsible by introduce the coupling between the composite scalars, ϕ_T and $\phi_{1,2}$ associated respectively to the T quark and to the technifermions. In Section III we discussed the self-energies used to compute the effective potential (Ω_T), where we assumed that the interaction $U(1)_x$ plays a role analogous to the ultraviolet dynamical symmetry breaking program proposed in Ref.[18, 19]. We also assume a TC self-energy that decays slowly with the momentum whose origin is due to the introduction of confinement in the gap equation, as discussed in Ref.[25], that is typical of the gauged Nambu-Jona-Lasinio (NJL) type of models where the anomalous dimension is $\gamma_m \approx 2$. Note that it is hardly possible to generate light scalar composite bosons without this particular choice [30, 31]. We finally determined an effective scalar boson Lagrangian and from it we obtained the scalar boson masses associated

to these models.

As already discussed in the introduction, the models that we consider here are interesting due to the particular form of their anomaly cancellation, due to the fact that it is one example of model where we have a naturally strong Abelian theory at the TeV scale, capable of producing a dynamical symmetry breaking of the model to the SM symmetry, whereas the SM gauge symmetry is broken by a TC condensate. Within this class of models we can also explain the larger decay rate of the 125 GeV boson into photons that is observed by the LHC experimental groups, caused by the presence of extra charged vector boson(V^+ , U^{++}) [5]. The only ingredients of our calculation are the self-energies of the strongly interacting theories. From these ones we can compute the scalar masses and the neutral gauge boson masses. Assuming that the 125 GeV observed at the LHC is a scalar boson and using their lower limit in the Z' mass, we compare the experimental results with the mass values that we obtained for these quantities. We verified that the models are strongly constrained, showing that it is rather difficult to have light scalar composites and at the same time to generate masses for neutral gauge bosons where one of them (the Z') is quite heavy. It is possible that in models with the presence of fundamental scalar bosons such difficulty is not present due to the many parameters that can be adjusted in the scalar potential, however, in this case, we may also foresee that this adjustment may lead to unnatural values of the coupling constants, unless some discrete symmetries are introduced by hand in order to avoid undesirable terms in the scalar potential.

Acknowledgments

This research was partially supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

-
- [1] ATLAS Collaboration, G. Aad *et al.*, Phys. Lett. B **716**, 1 (2012); CMS Collaboration, S. Chatrchyan *et al.*, Phys. Lett. B **716**, 30 (2012).
 - [2] F. Pisano and V. Pleitez, Phys. Rev. D**46**, 410 (1992).
 - [3] P. H. Frampton, Phys. Rev. Lett. **69**, 2889 (1992).
 - [4] V. Pleitez and M.D. Tonasse, Phys. Rev. D**48**, 2353 (1993).

- [5] A. Alves, E. Ramirez Barreto, A.G. Dias, C.A. de S.Pires, F.S. Queiroz and P.S. Rodrigues da Silva, Phys. Rev. D **84**, 115004 (2011); idem, hep-ph/1207.3699v2.
- [6] Alex G. Dias, C.A. de S.Pires, V. Pleitez and P.S. Rodrigues da Silva, Phys. Lett. **B621**, 151 (2005); Alex G. Dias, A. Doff, C. A. de S. Pires and P.S. Rodrigues da Silva, Phys. Rev. **D72**, 035006 (2005); Alex Gomes Dias, Phys. Rev. **D71**, 015009 (2005); Alex G. Dias, J.C. Montero and V. Pleitez, Phys. Lett. **B637**, 85 (2006); Alex G. Dias and V. Pleitez, Phys. Rev. **D73**, 017701 (2006), A. Doff, C. A. de S. Pires and P. S. Rodrigues da Silva, Phys. Rev. **D74**, 015014 (2006).
- [7] A. Doff and F. Pisano, Mod. Phys. Lett. **A15**, 1471 (2000).
- [8] C.A de S. Pires and O. P. Ravinez, Phys. Rev. **D58**, 035008 (1998); A. Doff and F. Pisano, Mod. Phys. Lett. **A14**, 1133 (1999); A. Doff and F. Pisano, Phys. Rev. **D63**, 097903 (2001).
- [9] L. Susskind, Phys. Rev. D **20**, 2619 (1979)
- [10] G. 't Hooft, "Naturalness, chiral symmetry and spontaneous chiral symmetry breaking", *NATO Adv. Study Inst. Ser. B Phys.* **59**, 135 (1980).
- [11] Prasanta Das and Pankaj Jain, Phys. Rev. **D 62**, 075001 (2000).
- [12] A. Doff, Phys. Rev. **D 76**, 037701 (2007).
- [13] A. Doff, Phys. Rev. **D 81**, 117702 (2010).
- [14] A. Doff and A. A. Natale, Int. J. Mod. Phys. A **27**, 1250156 (2012).
- [15] H. Pagels and S. Stokar, Phys. Rev. **D20**, 2947 (1979).
- [16] J. M. Cornwall, R. Jackiw and E. Tomboulis, Phys. Rev. **D10**, 2428 (1974).
- [17] Daniel Ng, *Phys. Rev. D* **49**, 4805 (1994); hep-ph/9212284.
- [18] J. Carpenter, R. Norton, S. Siegemund-Broka and A. Soni, Phys. Rev. Lett. **65**, 153 (1990).
- [19] J. D. Carpenter, R. E. Norton and A. Soni, Phys. Lett. **B212**, 63 (1988).
- [20] J. M. Cornwall, Phys. Rev. D **26**, 1453 (1982).
- [21] A. A. Natale, PoS QCD-TNT **09**, 031 (2009).
- [22] A. C. Aguilar, D. Binosi and J. Papavassiliou, Phys. Rev. D **78**, 025010 (2008).
- [23] J. M. Cornwall, Phys. Rev. D **83** (2011) 076001.
- [24] A. Doff, F. A. Machado and A. A. Natale, Annals of Physics **327** 1030 (2012).
- [25] A. Doff, F. A. Machado and A. A. Natale, New. J. Phys. **14**, 103043 (2012).
- [26] A. A. Natale, Nucl. Phys. **B226**, 365 (1983).
- [27] J. C. Montero, A. A. Natale, V. Pleitez and S. F. Novaes, Phys. Lett. B **161**, 151 (1985).

- [28] A. Doff and A. A. Natale, Phys. Lett. B **537**, 275 (2002).
- [29] A. Doff and A. A. Natale, Phys. Rev. D **68**, 077702 (2003), A. Doff and A. A. Natale, Eur. Phys. J. **C32**, 417 (2003).
- [30] A. Doff, A. A. Natale and P. S. Rodrigues da Silva, Phys. Rev. D **80**, 055005 (2009).
- [31] A. Doff, E. G. S. Luna and A. A. Natale, hep-ph/1303.3248.
- [32] J. M. Cornwall and R. C. Shellard, Phys. Rev. D **18**, 1216 (1978).
- [33] A. Doff, A. A. Natale and P. S. Rodrigues da Silva, Phys. Rev. D **77**, 075012 (2008).
- [34] S. Raby, S. Dimopoulos and L. Susskind, Nucl. Phys. B **169**, 373 (1980).
- [35] ATLAS Collaboration, "Search for high-mass dilepton resonances in $6.1fb^{-1}$ of pp collisions at $\sqrt{s} = 8\text{TeV}$ with the ATLAS experiment", **ATLAS-CONF-2012-129**, 2012; CMS Collaboration, "Search for Resonances in Dilepton Mass Spectra in pp Collisions at $\sqrt{s} = 8\text{ TeV}$ ", **CMS PAS EXO-12-015**, 2012.
- [36] S. Chatrchyan et al. [CMS Collab.], JHEP **1105**, 093 (2011).