

Equation of State within Gluon Dominated QGP Model in Relativistic Hydrodynamics Approach

T. P. Djun

Graduate Study in Material Science, University of Indonesia, Kampus UI Salemba, Jakarta 10430, Indonesia

Group for Theoretical and Computational Physics, Research Center for Physics, Indonesian Institute of Sciences, Kompleks Puspiptek Serpong, Tangerang 15310, Indonesia

E-mail: tp.djun@sci.ui.ac.id

M. K. N. Patmawijaya

Departemen Fisika, FMIPA, Universitas Indonesia, Depok 16424, Indonesia

R. Utama

Departemen Fisika, FMIPA, Universitas Indonesia, Depok 16424, Indonesia

L. T. Handoko

Group for Theoretical and Computational Physics, Research Center for Physics, Indonesian Institute of Sciences, Kompleks Puspiptek Serpong, Tangerang 15310, Indonesia

Abstract. The dynamics of Quark-gluon plasma (QGP) as a lump of deconfined free quarks and gluons is elaborated. Based on the first principal we construct the Lagrangian that represents the dynamics of QGP. To induce a hydrodynamics approach, we substitute the gluon fields with flow fields. As a result, the derived equation of Motion (E.O.M) for gluon dominated QGP shows the form that similar to Euler equation, and the energy momentum tensor also represents explicitly the system of ideal fluid. Combining the E.O.M and energy momentum tensor, the pressure and energy density distribution as the equation of states are analytically derived.

1. Introduction

Recent experiments on heavy-ion collisions show a strong indication that hot dense deconfined phase of free quark and gluon, the so called quark-gluon plasma (QGP), is conjectured to exist. The study of QGP itself has been carried out through a number of different approaches in the previous works. Some results of these studies were obtained in the framework of quantum chromodynamics (QCD) theory by utilizing the lattice gauge calculation [1, 2]. Other calculations of QGP were based on the relativistic hydrodynamics approach [3, 4]. In the latter,

the QGP could be either quark [4] or gluon [3] dominated matter. For the quark dominated approach, a very small ratio of shear viscosity over entropy is required to get a good fit of the spectra of transverse momentum, energy density distribution and other physical observables that are obtained from experiments [5, 6, 7, 8, 9, 10]. On the other hand, the gluon dominated plasma motivated by the discoveries of jet quenching in the heavy-ion collision at RHIC indicates the shock waves in the form of Mach cone [11, 12]. The present paper adopts the so-called fluid QCD model [14, 15] to produce the equation of motion and energy momentum tensor for quark and gluon in a lump of QGP, and subsequently to investigate the distribution of the pressure and energy density.

This paper is organized as follows. In Section 2 the fluid QCD model is briefly revisited, and the energy momentum tensor for ideal fluid is derived. Then, it is followed by the derivation for the equation of state and the explicit expression of gluon field in Section 3. Finally, the summary and discussion will be given in Section 4. Throughout this work, we use the natural units, i.e., $\hbar \equiv c \equiv 1$.

2. Model

The Lagrangian of QGP that describes the unification of fermions and bosons from different gauge groups with preserving $SU(3)_F \otimes U(1)_G$ gauge symmetry can be written as [13, 14]

$$\mathcal{L} = i\bar{Q}\gamma^\mu\partial_\mu Q - m_Q\bar{Q}Q - \frac{1}{4}S_{\mu\nu}^a S^{a\mu\nu} - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + g_F J_{F\mu}^a U^{a\mu} + g_G J_{G\mu} A^\mu, \quad (1)$$

where Q and \bar{Q} represent the quark and anti-quark triplet, γ^μ is Dirac gamma matrices, and m_Q is the mass of quark. The factor $\frac{1}{4}S_{\mu\nu}^a S^{a\mu\nu}$ is the gauge invariant kinetic term of gluon field. The gluon field strength tensor itself is expressed as $S_{\mu\nu}^a \equiv \partial_\mu U_\nu^a - \partial_\nu U_\mu^a + g_F f^{abc} U_\mu^b U_\nu^c$. In the latter, U_μ^a indicates the gluon field, g_F is the strong coupling constant, and f^{abc} is the structure constant of $SU(3)$ gauge group. The kinetic term for gauge boson A_μ is built inside its field strength tensor, i.e., $F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu$. The last two terms of Eq. (1), $J_{F\mu}^a = \bar{Q}T_F^a\gamma_\mu Q$ and $J_{G\mu} = \bar{Q}\gamma_\mu Q$, represent the quark currents from the $SU(3)$ and $U(1)$ gauge groups, respectively, whereas g_G denotes the coupling constant from $U(1)$ gauge group.

The QCD Lagrangian in Eq. (1) is constructed with a purpose to reproduce the energy momentum tensor that has the same form as the energy momentum tensor of an ideal fluid [13]. In the succeeding step, the gluon field U_μ^a , that is designed to act as the flow field in the system is proposed to be formulated in a configuration that inherently contains the relativistic flow. It is formulated as $U_\mu^a = (U_0^a, \mathbf{U}^a) \equiv u_\mu \phi^a$ [14, 15]. Here, ϕ^a is the dimension one scalar field, and $u_\mu \equiv \gamma(1, \mathbf{v})$ is the relativistic velocity, with $\gamma = (1 - |\mathbf{v}|^2)^{-1/2}$.

This formulation provides a clue that a single gluonic field U_μ^a may behaves as a fluid at certain scale, beside its conventional point particle properties with a polarization vector ϵ_μ in the form of $U_\mu^a = \epsilon_\mu \phi^a$. One can then consider that there is a kind of “phase transition”,

$$\underbrace{\text{hadronic state}}_{\epsilon_\mu} \longleftrightarrow \underbrace{\text{QGP state}}_{u_\mu}.$$

As the gluon field behaves as a point particle, it is in a stable hadronic state and is characterized by its polarization vector. On the other hand in the pre-hadronic state (before hadronization) like hot QGP, the gluon field behaves as a highly energized flow particle and the properties are dominated by its relativistic velocity.

The field U_μ^a is actually analogous to the gauge boson from $U(1)$ gauge group in particle physics, where the polarization vector ϵ_μ from the free photon solution $\sim \epsilon_\mu \exp(-ip_\mu x^\mu)$ is replaced by the 4-velocity u_μ . Recall that the wavefunction U_μ^a for a free particle satisfies $[g^{\mu\nu}(\partial^2 + m^2) - \partial^\nu \partial^\mu]U_\mu^a = 0$ with solution $U_\mu^a \sim \epsilon_\mu \phi^a \exp(-ip_\mu x^\mu)$. For a massive vector

particle, $m \neq 0$, we have no choice but to take $\partial^\mu U_\mu^a = 0$. It is not a gauge condition like the case of massless particle. This then demands $p^\mu \epsilon_\mu = 0$. The number of independent polarization vectors is reduced from four to three in a covariant fashion. However, one can still perform another gauge transformation to the massless U_μ^a which makes finally only two degrees of freedom remain. Therefore, one should keep in mind that in the present model the spatial velocity has only two degrees of freedom, that means one component must be described by another two vector components.

Further, when Euler-Lagrange equation is applied to Eq. (1) one obtain

$$\frac{\partial}{\partial t}(\gamma \mathbf{v} \phi^a) + \nabla(\gamma \phi^a) = -g_F \oint d\mathbf{x} (J_{F0}^a + F_0^a), \quad (2)$$

where J_{F0}^a denotes the covariant current originating from the quarks that are surrounded by and interact with the gluon "fluid", while the term F_μ^a is induced by the fluid self-interaction and the interacting gauge fields A_μ^a . Equation (2) is considered as the general relativistic fluid equation for single gluon field U_μ^a , since in the non-relativistic limit, i.e. $\phi^a \sim 1$ and $\gamma \sim 1 + \frac{1}{2}|\mathbf{v}|^2$, Eq. (2) transforms to the classical equation of motion of fluid dynamics [14]

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} = -g_F \oint d\mathbf{x} (J_{F0}^a + F_0^a) |_{\text{non-rel}}. \quad (3)$$

This shows that from a certain point of view the Lagrangian describes a general relativistic fluid system interacting with another gauge field and the matter inside. The flow characteristic that appears in the equation of motion comes from the contribution of gluon field U_μ^a . This fact indicates that the system we are working with is a gluon dominated QGP. In such system the terms that do not contain gluon field may be omitted. As a consequence we have

$$\mathcal{L} = -\frac{1}{4} S_{\mu\nu}^a S^{a\mu\nu} + g_F J_{F\mu}^a U^{a\mu}. \quad (4)$$

The Lagrangian given by Eq. (4) describes the kinetics of gluons, the self-interaction of gluons, the self-interaction between small number of quark and anti-quark, and the interaction between quark with gluon "fluid". Electromagnetic interaction also exist in the system, but the scale is suppressed due to its tiny value compared with the strong interaction. From the Lagrangian given in Eq. (4) we can derive the energy momentum tensor [17]

$$\begin{aligned} \mathcal{T}_{\mu\nu} &= \frac{2}{\sqrt{-g}} \frac{\delta(\mathcal{L}\sqrt{-g})}{\delta g^{\mu\nu}} \\ &= S_{\mu\rho}^a S^{a\rho}{}_\nu - g_{\mu\nu} \mathcal{L} + 2g_F J_{F\mu}^a U_\nu^a \\ &= [2g_F g_{\mu\nu} J_{F\mu}^a U^{a\mu} + g_F^2 f^{abc} f^{ade} U_\mu^b U_\rho^c U^{d\rho} U_\nu^e] \\ &\quad - [g_F g_{\mu\nu} J_{F\mu}^a U^{a\mu} - \frac{1}{4} g_{\mu\nu} g_F^2 f^{abc} f^{ade} U_\mu^b U_\nu^c U^{d\mu} U^{e\nu}]. \end{aligned} \quad (5)$$

The factor $J_{F\mu}^a U^{a\mu}$ in Eq. (5) might be expressed in a more elementary form. To this end, we can start with the assumption that the solution of the Dirac equation for single color quark is in the form of $Q(p, x) = q(p) \exp(-ip \cdot x)$. By inserting this solution to the Dirac equation, i.e., $(i\gamma^\mu \partial_\mu - m)q(p) \exp(-ip \cdot x) = 0$, and multiplying with the anti-quark solution, we obtain $\bar{q}\gamma_\mu q = 4p_\mu$. Since $J_{F\mu}^a = \bar{Q}\gamma_\mu Q T^a = \bar{q}\gamma_\mu q T^a = 4p_\mu T^a$, and $U^{a\mu} = u^\mu \phi^a$, we finally obtain $J_{F\mu}^a U^{a\mu} = 4m_Q T^a \phi^a$. Note that we have utilized $u_\mu u^\mu = 1$ along with the assumption that all quarks and anti-quarks have the same momenta P^μ and that the velocities of all gluons and quarks are homogeneous. Thus, the energy momentum tensor in the function of field ϕ^a reads

$$\begin{aligned} \mathcal{T}_{\mu\nu} &= [8g_F m_Q T^a \phi^a + g_F^2 f^{abc} f^{ade} \phi^b \phi^c \phi^d \phi^e] u_\mu u_\nu \\ &\quad - [4g_F m_Q T^a \phi^a - \frac{1}{4} g_F^2 f^{abc} f^{ade} \phi^b \phi^c \phi^d \phi^e] g_{\mu\nu}. \end{aligned} \quad (6)$$

With this form, $\mathcal{T}_{\mu\nu}$ is obviously describing a system of perfect fluid. Further, we assume that the gluon color states are homogeneous, i.e., $\phi^a = \phi$ for all $a = 1, \dots, 8$. As the follow-on to this assumption, the generator T^a can be compactly written as $T = \sum_a T^a$ and the 2nd and 4th terms of $\mathcal{T}_{\mu\nu}$ can be omitted due to the completely anti-commutative property of the structure constant f^{abc} . The energy momentum tensor then reduces to

$$\mathcal{T}_{\mu\nu} = (8Tg_F m_Q \phi) u_\mu u_\nu - (4Tg_F m_Q \phi) g_{\mu\nu} \quad (7)$$

It reveals the collective gluons flow in the system. Therefore, we can temporarily summarize that the system represents an isotropic homogeneous perfect fluid of gluon dominated QGP. Further, to discuss the dynamics of glue lumps, it is also plausible to take the gluon as a field that independent of time, $\phi = \phi(x)$. If we compare Eq.(7) with energy momentum tensor of ideal fluid $\mathcal{T}_{\mu\nu} = (\mathcal{E} + \mathcal{P})u_\mu u_\nu - \mathcal{P}g_{\mu\nu}$, it becomes obvious that

$$\mathcal{E} = \mathcal{P} = 4Tg_F m_Q \phi. \quad (8)$$

Here \mathcal{P} and \mathcal{E} denote the isotropic pressure and energy density for fluid field, respectively. Or, in currently discussed topic, it is the energy density and pressure distribution in the lump of gluon dominated QGP.

3. Expression for distribution field

The expression for the scalar field ϕ can be obtained by deriving the solution of Eq. (3). After applying the assumption of the gluon homogeneity, the equation can be expressed as follow.

$$\delta \partial_{tr} \phi + \gamma \partial_{rr} \phi + \xi + \beta \phi \partial_t \phi + \beta \phi \partial_r \phi - \lambda \partial_t \phi - \lambda \partial_r \phi = 0, \quad (9)$$

with $\delta = \gamma|\mathbf{v}|$, $\xi = g_F \bar{Q} \gamma_\mu T Q$, $\beta = i\gamma^2 |\mathbf{v}|^2$, $\lambda = (g_G/g_F) A \gamma$, and $\gamma = (1 - v^2)^{-\frac{1}{2}}$. For a simplification but stay relevant, the pressure and energy density distribution of the lump of QGP is assumed to depend only to r , it is $\phi = \phi(r)$, and $\partial_r \rightarrow d_r$. So the term that involve the derivative of t is vanished.

$$\gamma d_{rr} \phi + \beta \phi d_r \phi - \lambda d_r \phi + \xi = 0, \quad (10)$$

The solution for this equation is

$$\begin{aligned} \phi = & \left(2\gamma \left(\frac{\exp[\frac{r\lambda}{2\gamma}] \Lambda_1[x]}{2\gamma} - \frac{\exp[\frac{r\lambda}{2\gamma}] \beta \xi \Lambda_2[x]}{2^{1/3} \gamma^2 (-\frac{\beta \xi}{\gamma^2})^{2/3}} \right. \right. \\ & \left. \left. + \left(\frac{\exp[\frac{r\lambda}{2\gamma}] \Lambda_3[x]}{2\gamma} - \frac{\exp[\frac{r\lambda}{2\gamma}] \beta \xi \Lambda_4[x]}{2^{1/3} \gamma^2 (-\frac{\beta \xi}{\gamma^2})^{2/3}} \right) C_2 \right) \right) \\ & \times 1 / \left(\beta \left(\exp[\frac{r\lambda}{2\gamma}] \Lambda_1[x] + \exp[\frac{r\lambda}{2\gamma}] \Lambda_3[x] C_2 \right) \right). \end{aligned} \quad (11)$$

When we substitute $\delta, \xi, \beta, \lambda$, and γ back to the equation, then ϕ appears as

$$\begin{aligned} \phi = & \left(\frac{2}{\sqrt{1-v^2}} \left(\frac{\exp[\frac{rgA}{2}] \Lambda_1[x]}{2/\sqrt{1-v^2}} - \frac{\exp[\frac{rgA}{2}] (iv^2 \xi)^{1/3} \Lambda_2[x]}{2^{1/3}} \right. \right. \\ & \left. \left. + \left(\frac{\exp[\frac{rgA}{2}] \Lambda_3[x]}{2/\sqrt{1-v^2}} - \frac{\exp[\frac{rgA}{2}] (iv^2 \xi)^{1/3} \Lambda_4[x]}{2^{1/3}} \right) C_2 \right) \right) \\ & \times 1 / \left(\frac{iv^2}{1-v^2} \left(\exp[\frac{rgA}{2}] \Lambda_1[x] + \exp[\frac{rgA}{2}] \Lambda_3[x] C_2 \right) \right). \end{aligned} \quad (12)$$




Figure 1. Pressure inside QGP-lump in the function of radius

Here,

$$\begin{aligned}
\Lambda_1[x] &= \frac{1}{3^{1/6}\Gamma[\frac{2}{3}]} + \frac{3^{1/6}x}{\Gamma[\frac{1}{3}]} + \frac{x^3}{6(3^{1/6}\Gamma[\frac{2}{3}])} + \frac{x^4}{4(3^{5/6}\Gamma[\frac{1}{3}])} + \mathcal{O}[x]^5, \\
\Lambda_2[x] &= \frac{3^{1/6}}{\Gamma[\frac{1}{3}]} + \frac{x^2}{2(3^{1/6}\Gamma[\frac{2}{3}])} + \frac{x^3}{3^{5/6}\Gamma[\frac{1}{3}]} + \frac{x^5}{30(3^{1/6}\Gamma[\frac{2}{3}])} + \mathcal{O}[x]^6, \\
\Lambda_3[x] &= \frac{1}{3^{2/3}\Gamma[\frac{2}{3}]} - \frac{x}{3^{1/3}\Gamma[\frac{1}{3}]} + \frac{x^3}{6(3^{2/3}\Gamma[\frac{2}{3}])} - \frac{x^4}{12(3^{2/3}\Gamma[\frac{2}{3}])} + \mathcal{O}[x]^5, \\
\Lambda_4[x] &= -\frac{1}{3^{1/3}\Gamma[\frac{1}{3}]} + \frac{x^2}{2(3^{2/3}\Gamma[\frac{2}{3}])} - \frac{x^3}{3(3^{1/3}\Gamma[\frac{1}{3}])} + \frac{x^5}{30(3^{2/3}\Gamma[\frac{2}{3}])} + \mathcal{O}[x]^6,
\end{aligned}$$

and

$$x = \frac{-i2v^2(\xi r + \frac{C_1}{\sqrt{1-v^2}}) + g^2 A^2}{2^{4/3}(-iv^2\xi)^{2/3}}$$

So far this non-trivial solution is well expressed. But to have a firm solution, some adjustment still need to be carry out at the future works. While for the rest, we will explore the expression for pressure and energy density in the system of gluon dominated quark-gluon plasma.

4. Summary and Discussion

Return to Eq. (8), $\mathcal{E} = \mathcal{P} = 4Tg_F m_Q \phi$, the energy and pressure can be expressed as follows [16]

$$\rho = P = \int \mathcal{E} d^4x = \int \mathcal{P} d^4x = 4Tg_F f_Q m_Q \int \int \phi dt dV, \quad (13)$$

where \mathcal{E} and \mathcal{P} denote the isotropic pressure and density for single fluid field, respectively. By adopting FRW geometry as the background and do integration on the spatial dimension,

Eqs. (13) read

$$\rho = P = 4\pi\lambda \int R^3 \phi dt \int \frac{r^2 dr}{\sqrt{1 - kr^2}}. \quad (14)$$

Here we have used $dV = (R^3 r^2 \sin\theta)/\sqrt{1 - kr^2} dr d\theta d\vartheta$, and $\lambda = 4T_{GF} f_Q m_Q$. Further, if we assign $\int_0^{2\pi} \int_0^\pi \int dV = 4\pi \int \frac{r^2 dr}{\sqrt{1 - kr^2}}$ as ζ , then the expression for P and ρ can be written as

$$P = \rho = \lambda\zeta \int R^3 \phi dt. \quad (15)$$

In a case when r is constant, $p = \rho \propto \phi(t)dt$, and also if $\phi(t) \propto 1/t$, Then one arrive at $P = \rho \propto \lambda\zeta R^3/t$. It indicates that the pressure and energy in the system decrease asymptotically following the increases of time.

Figure 1 shows the total pressure inside the QGP-lump as the function of radius. It is drawn for the simplest condition, where ϕ is assumed as a constant, and the scale factor $R = 1$. While $k = 1, 0, -1$ represent the space-time curvature that describe the close, flat, or open space-time that analogous to the close, flat, or open universe. One of the obvious property that appears here is the equation of state $P/\rho = 1$. Such a ratio can indicates that in this model the QGP exist within radiation state. This estimate comes from the fact that in general the prerequisite condition for the radiation state is $P \sim 1/3\rho$, and P/ρ is getting smaller along the transition from radiation state to matter state.

Finally, the study revealed observables which should be accessible through heavy ion collisions experiments at the RHIC and LHC. With currently proposed calculation results, such experiments are in future expected to become a reference for adjustments and verifications of the dynamics theory of macroscopic behavior of QGP.

Acknowledgments

TPD thanks the Group for Theoretical and Computational Physics, Research Center for Physics, Indonesian Institute of Sciences (LIPI) for the warm hospitality during the completion of this work. T.P.D. and L.T.H. are supported by Riset Kompetitif LIPI under Contract No. 11.04/SK/KPPI/II/2016.

5. References

- [1] Gottlieb S 2007 *J. Phys. Conf. Ser.* **78**, 012023
- [2] Petreczky P 2008 *Eur. Phys. J. ST* **155**, 123
- [3] Bouras I, Molnar E, Niemi H, Xu Z, El A, Fochler O, Greiner C, Rischke D H 2009 *Phys. Rev. Lett.* **103**, 032301
- [4] Romatschke P 2010 *Int. J. Mod. Phys. E* **19**, 1
- [5] Teaney D, Lauret J, Shuryak E V 2001 *Phys. Rev. Lett.* **86**, 4783
- [6] Huovinen P, Kolb P F, Heinz U W, Ruuskanen P V, Voloshin S A 2001 *Phys. Lett. B* **503**, 58
- [7] Kolb P F, Heinz U W, Huovinen P, Eskola K J, Tuominen K 2001 *Nucl. Phys. A* **696**, 197
- [8] Kolb P F, Rapp R 2003 *Phys. Rev. C* **67**, 044903
- [9] Hirano T, Tsuda K 2002 *Phys. Rev. C* **66**, 054905
- [10] Baier R, Romatschke P 2007 *Eur. Phys. J. C* **51**, 677 (2007).
- [11] Adams J *et al.* 2003 *Phys. Rev. Lett.* **91**, 172302
- [12] Adare A *et al.* 2008 *Phys. Rev. Lett.* **101**, 232301
- [13] Djun T P, Handoko L T, Soegijono B, Mart T 2015 *Int. J. Mod. Phys. A* **30** 1550077
- [14] Sulaiman A, Fajarudin A, Djun T P, Handoko L T 2009 *Int. J. Mod. Phys. A* **24**, 3630
- [15] Djun T P, Handoko L T 2010 in *Proceeding of the Conference in Honor of Murray Gell-Mann's 80th Birthday: Quantum Mechanics, Elementary Particles, Quantum Cosmology & Complexity*, edited by H. Fritzsch and K. K. Phua (Nanyang Technological University, Singapore) , p. 419 arXiv:1109.6066 [hep-ph].
- [16] Nugroho C S, Latief A O, Djun T P, Handoko L T 2012 *Grav. Cosmol.* **18**, 32
- [17] Hobson M P, Efstathiou G, and Lasenby A N (Cambridge University Press 2006) *General Relativity*

