arXiv:1304.1166v1 [hep-ph] 3 Apr 2013

Decay coupling constant sum rules for tetraquarks $T[(\bar{Q}q)(Q\bar{q})]$ with broken SU(3) symmetry

V. Gupta* and G. Sánchez-Colón†

Departamento de Física Aplicada.
Centro de Investigación y de Estudios Avanzados del IPN.
Unidad Mérida.
A.P. 73, Cordemex.
Mérida, Yucatán, 97310. MEXICO.

S. Rajpoot‡

Department of Physics & Astronomy.
California State University, Long Beach.
Long Beach, CA 90840. USA.

(Dated: February 25, 2022)

Abstract

For tetraquarks of the form $T[(\bar{Q}q)(Q\bar{q})]$ we give sum rules for their decay coupling constants, taking into account the SU(3) symmetry breaking interactions to first order.

PACS numbers: 14.40.Rt, 13.25.Jx
Keywords: tetraquarks, sum rule

*virendra@mda.cinvestav.mx
†gsanchez@mda.cinvestav.mx
‡rajpoot@csulb.edu

I. INTRODUCTION

Two hidden-bottom charged meson resonances, $Z_b(10610)$ and $Z_b(10650)$ with $J^P = 1^+$, were observed recently by the Belle Collaboration [1]. Since these decay into $\pi^\pm\Upsilon(nS)$ $(n = 1, 2, 3)$ and $\pi^\pm h_b(mP)$ $(m = 1, 2)$ they have been interpretated [2–4] as tetraquarks of the form $T(Q\bar{Q}\,q\bar{q})$. A natural decay mode for such a tetraquark would be

$$T(Q\bar{Q}\,q\bar{q}) \to H(Q\bar{Q}) + M(q\bar{q}), \tag{1}$$

where $H(Q\bar{Q})$ is a heavy meson with $Q = b$ or c and $M(q\bar{q})$ could be the pseudoscalar or vector meson SU(3) flavour octet. For such tetraquarks we recently [5] gave sum rules for tetraquark decay coupling constants taking into account the SU(3) breaking interactions to first order.

In this paper we consider tetraquarks of the form $T[(\bar{Q}q)\,(Q\bar{q})]$ where the heavy quark $Q = b$ or c and the light quarks $q = u, d,$ or s. For a given Q there will be nine such tetraquarks. The natural decay mode will be into two mesons $(\bar{Q}q)$ and $(Q\bar{q})$ subject to angular momentum and parity selection rules. For example, a $J^P = 0^+$ tetraquark would decay into two pseudoscalar mesons.

We consider the decays

$$T[(\bar{Q}q)\,(Q\bar{q})] \to M(\bar{Q}q) + M(Q\bar{q}), \tag{2}$$

in Sec. III below. We give sum rules for the nine decay couplings constants taking into account the SU(3) breaking interactions to first order.

An important difference between the tetraquarks $T(Q\bar{Q}\,q\bar{q})$ considered earlier and the tetraquarks $T[(\bar{Q}q)\,(Q\bar{q})]$ considered here is that the latter can have OZI [6–8] suppressed decays through annihilation of $\bar{Q}^\alpha Q_\beta$ through gluons into lighter quarks $\bar{q}^\alpha q_\beta$ (greek letters represent the color indices). These modes are briefly discussed in the concluding remarks.

II. NOTATION

We denote the tetraquark made of a heavy meson and its antiparticle as

$$T^i_k = T[(\bar{Q}q_k)\,(Q\bar{q}^i)], \tag{3}$$

with $i, k = 1, 2, 3$ (latin letters representing flavour indices). The heavy quark $Q = b$ or c is SU(3) flavour singlet. The light quarks q_k $(k = 1, 2, 3)$ transform as SU(3) $\mathbf{3}$ while the antiquarks $\bar{q}^i$ $(i = 1, 2, 3)$ transform as $\bar{\mathbf{3}}$. In color space, $M_k \equiv (\bar{Q}q_k)$ and $\bar{M}^i \equiv (Q\bar{q}^i)$ form color singlets and transform as $\mathbf{3}$ and $\bar{\mathbf{3}}$ of SU(3) flavour, respectively.

The natural decay mode for the tetraquark would be

$$T^i_k \to M_k + \bar{M}^i. \tag{4}$$

The tetraquark states can be represented as a 3×3 matrix T with T^i_k as matrix elements, so that,

$$T = \begin{pmatrix} T^1_1 & T^2_1 & T^3_1 \\ T^1_2 & T^2_2 & T^3_2 \\ T^1_3 & T^2_3 & T^3_3 \end{pmatrix}. \tag{5}$$

In SU(3) flavour space this transforms as $\mathbf{1}\oplus\mathbf{8}$. The mesons M_k and $\bar{M}^i$ transform as SU(3) $\mathbf{3}$ and $\bar{\mathbf{3}}$, respectively. The 3×3 matrix $\mathcal{M}$ representing the final states is

$$\mathcal{M} \;=\; \begin{pmatrix} \mathcal{M}_1^1 & \mathcal{M}_1^2 & \mathcal{M}_1^3 \\ \mathcal{M}_2^1 & \mathcal{M}_2^2 & \mathcal{M}_2^3 \\ \mathcal{M}_3^1 & \mathcal{M}_3^2 & \mathcal{M}_3^3 \end{pmatrix}, \tag{6}$$

where $\mathcal{M}_k^i$ represents $M_k+\bar{M}^i$ in the final state. In flavour space it also transforms as $\mathbf{1}\oplus\mathbf{8}$, this is clear since $(\bar{Q}q)\otimes(Q\bar{q})$ transform as $\mathbf{3}\otimes\bar{\mathbf{3}}=\mathbf{1}\oplus\mathbf{8}$.

III. SUM RULES

In unbroken SU(3) there will be a single decay coupling constant G_0; with λ_8 breaking the tetraquark octet can form two octets, symmetric (coupling constant G_D) and antisymmetric (coupling constant G_F). In matrix form we can write the decay coupling constant as

$$\begin{aligned} &G\left[T_k^i(\mathbf{8})\to M_k+\bar{M}^i\right] = \\ &\quad G_0\,\mathrm{Tr}[\widetilde{T}\mathcal{M}] + G_D\,\mathrm{Tr}[\widetilde{T}(\lambda_8\mathcal{M}+\mathcal{M}\lambda_8)] + G_F\,\mathrm{Tr}[\widetilde{T}(\lambda_8\mathcal{M}-\mathcal{M}\lambda_8)]. \end{aligned} \tag{7}$$

Explicitly, for the nine tetraquark decays

$$G(T_1^1\to M_1+\bar{M}^1) = G_0+2G_D, \tag{8}$$

$$G(T_2^2\to M_2+\bar{M}^2) = G_0+2G_D, \tag{9}$$

$$G(T_3^3\to M_3+\bar{M}^3) = G_0-4G_D, \tag{10}$$

$$G(T_1^2\to M_1+\bar{M}^2) = G_0+2G_D, \tag{11}$$

$$G(T_2^1\to M_2+\bar{M}^1) = G_0+2G_D, \tag{12}$$

$$G(T_1^3\to M_1+\bar{M}^3) = G_0-G_D+3G_F, \tag{13}$$

$$G(T_2^3\to M_2+\bar{M}^3) = G_0-G_D+3G_F, \tag{14}$$

$$G(T_3^1\to M_3+\bar{M}^1) = G_0-G_D-3G_F, \tag{15}$$

$$G(T_3^2\to M_3+\bar{M}^2) = G_0-G_D-3G_F. \tag{16}$$

Nine decays and three coupling constants. So, six sum rules or relations,

$$G(T_1^1\to M_1+\bar{M}^1) = G(T_2^2\to M_2+\bar{M}^2) = G(T_1^2\to M_1+\bar{M}^2) = G(T_2^1\to M_2+\bar{M}^1); \tag{17}$$

$$G(T_1^3 \to M_1 + \bar{M}^3) = G(T_2^3 \to M_2 + \bar{M}^3), \quad G(T_3^1 \to M_3 + \bar{M}^1) = G(T_3^2 \to M_3 + \bar{M}^2); \tag{18}$$

$$G(T_2^2 \to M_2 + \bar{M}^2) + G(T_3^3 \to M_3 + \bar{M}^3) = G(T_1^3 \to M_1 + \bar{M}^3) + G(T_3^1 \to M_3 + \bar{M}^1). \tag{19}$$

If and when such tetraquarks are observed one can extract the coupling constants from the observed decay rates. For a $J^P = 0^+$ tetraquark decaying into two pseudoscalar mesons the decay width is

$$\Gamma = \frac{1}{8\pi} \frac{k}{M^2} |A|^2, \tag{20}$$

where M is the tetraquark mass, k is the momentum of a decay particle and A is the transition amplitude. For s-wave decay ($0^+ \to 0^- + 0^-$) $A = GM$ with G the coupling constant.

IV. CONCLUDING REMARKS

To date the tetraquarks $T[(\bar{Q}q)\,(Q\bar{q})]$ have not been observed experimentally. However, their decays products, the heavy mesons (for example $D_S^+ = c\bar{s}$, $B^+ = u\bar{b}$, etc.) have been observed [9]. It is conceivable that this type of tetraquark may be seen as a resonance in the heavy quark particle-antiparticle mass spectra, for example, $D_S^+ + D_S^-$, $D^0 + \bar{D}^0$, etc. Another possibility is that they may be observed through their decay into light meson plus anti meson pairs, for example, $q_k\bar{q}^l + q_l\bar{q}^i$ (see Fig. 1). These are possible through annihilation of $\bar{Q}Q$ through gluons into $\bar{q}q$. However, such decays are likely to be OZI [6–8] suppressed.

Acknowledgments

V. Gupta and G. Sánchez-Colón would like to thank CONACyT (México) for partial support. The work of S. Rajpoot was supported by DOE Grant #: DE-FG02-10ER41693.

[1] Belle Collab. (A. Bondar *et al.*), *Phys. Rev. Lett.* **108**, 122001 (2012).
[2] T. Guo, L. Cao, M. Z. Zhou and H. Chen, arXiv: 1106.2284 [hep-ph].
[3] F. S. Navarra, M. Nielsen and J.- M. Richard, arXiv: 1108.1230 [hep-ph].
[4] For a discussion of all tetraquarks containing only c and s quarks see: V. Gupta, *Int. J. Mod. Phys. A.* **20**, 5891 (2005).
[5] V. Gupta, G. Sánchez-Colón, and S. Rajpoot, *Mod. Phys. Lett. A* **27**, 1250165 (2012).
[6] S. Okubo, *Phys. Lett.* **5**, 165 (1963).
[7] G. Zweig, CERN Report. TH 401 and 412 (1964).
[8] J. Izuka, *Prog. Theor. Phys. Suppl.* **37** and **38**, 21 (1966).
[9] Particle Data Group (J. Beringer *et al.*), *Phys. Rev. D* **86**, 010001 (2012).




FIG. 1: Tetraquarks $T[(\bar{Q}q)\,(Q\bar{q})]$ observed through their OZI suppressed decay into light meson plus anti meson pairs, $q_k\bar{q}^l + q_l\bar{q}^i$. Greek letters represent the color indices and latin letters represent flavor indices.