

The Standard Model

W-Y. Pauchy Hwang¹

*Asia Pacific Organization on Cosmology and Particle Astrophysics,
Institute of Astrophysics, Center for Theoretical Sciences,
and Department of Physics, National Taiwan University, Taipei 106, Taiwan*

(April 17, 2013; updated on August 25, 2015; August 25, 2017)

Abstract

We declare that we live in the quantum 4-dimensional Minkowski space-time with, *via the gauge principle*, the force-fields gauge-group structure, $SU_c(3) \times SU_L(2) \times U(1)$ applied to the quark world or $SU_L(2) \times U(1) \times SU_f(3)$ applied to the lepton world, built-in from the very beginning. The quark world with the triplets from the "quark" group $SU_Q(3)$ (used to be called "flavor $SU(3)$ symmetry"), which is of nuclear sizes and is protected by $SU_c(3) \times SU_L(2) \times U(1)$ (i.e. the (123) symmetry), can be seen by our world, while the lepton world, as of atomic sizes and protected by $SU_L(2) \times U(1) \times SU_f(3)$ or another (123) symmetry, can also be seen, as singlets of the quark group $SU_Q(3)$, by our world. Apart from the "ignition" term, the entire Standard Model is dimensionless and massless in the quantum 4-dimensional Minkowski space-time; that is, all couplings are dimensionless and there are no mass terms in the quantum 4-dimensional Minkowski space-time.

Therefore, there exist the well-studied $3^\circ K$ cosmic microwave background (CMB) and the now clustered neutrino halos formed from cosmic background (CB) neutrinos (of three flavors, and antineutrinos), both present in the overall background of our Universe. In theory, it yields neutrino oscillations, as some lepton-flavor-violating processes, in a natural manner.

PACS Indices: 12.60.-i (Models beyond the standard model); 98.80.Bp (Origin and formation of the Universe); 12.10.-g (Unified field theories and models).

1 The Excerpts

We declare that our world is the quantum 4-dimensional Minkowski space-time with, *via the gauge principle*, the force-fields gauge-group structure, $SU_c(3) \times SU_L(2) \times U(1)$ at the fermi scale or $SU_L(2) \times U(1) \times SU_f(3)$ at the anstron scale, implemented at the very beginning.

The *matter* of our world has triplets of the quark group $SU_Q(3)$ and singlets of the quark group $SU_Q(3)$, each in terms of one left-handed $SU_L(2)$ doublet and two right-handed $SU_L(2)$ singlets, while the triplets (quarks) of the quark group have the gauge symmetry $SU_c(3) \times SU_L(2) \times U(1)$ and the singlets (leptons) of the quark group have the gauge symmetry $SU_L(2) \times U(1) \times SU_f(3)$. The *overall background* is the quantum 4-dimensional Minkowski space-time with, *via the gauge principle*, the force-field gauge-group structure suitably built-in at the very beginning.

¹Correspondence Author; Email: wyhwang@phys.ntu.edu.tw; A completed version (Version 3) of arXiv:1304.4705v1 [hep-ph] 17 April 2013 (the original version).

The (123) gauge symmetry, under $SU_c(3) \times SU_L(2) \times U(1)$, and the other (123) gauge symmetry, under $SU_L(2) \times U(1) \times SU_f(3)$, are explained later on in this paper - c.f., Section 7 when the meanings of the notations become clear.

Mathematically, the Standard Model [1] is a group theory - a group of the elements, triplets and singlets, of the quark group $SU_Q(3)$; they interact among themselves through the gauge symmetry, either $SU_c(3) \times SU_L(2) \times U(1)$ at the fermi sizes or $SU_L(2) \times U(1) \times SU_f(3)$ at the atomic sizes. They interact on the background of the quantum 4-dimensional Minkowski space-time.

2 Introduction

According to Newton's doctrine, the motion of an object has to be specified by both its space-time point (i.e., coordinates) and its behavior at a nearby point (thus, its velocities). The force information, which controls the change of the velocity, is characterized by the gauge principle, $\partial_\mu \rightarrow D_\mu$. In modern days, the difference, $D_\mu - \partial_\mu$, is expressed by the gauge fields. Thus, we still stick to Newton's doctrine and the various gauge fields should be characterized at the very beginning of the game.

The next thing in the game is how to introduce the matter worlds, i.e., the quark world (at the fermi scale) and the lepton world (at the atomic scale). The gauge principle is expected to act on some basic unit of matter - these units of matter are expressed in terms of the entries from the quark world, or, the entries from the lepton world. As seen below, the three entries, two right-handed units of matter and one left-handed unit of matter, together are used to define the lepton world; likewise, the three similar entries, in terms of the triplets of some symmetry group $SU_Q(3)$ (to be explained later on), are used to define the quark world.

We believe that the language will be used, for centuries to come, to describe the smallest units of matter, including electrons, neutrinos, and quarks, on the basis of Einstein's relativity principle and the quantum principle.

Nowadays it is a general belief (of others so far, but of not the view of this paper) that there are three generations of quarks and three generations of leptons, at the level of the so-called "point-like Dirac particles". According to another newly established belief in Cosmology, the content of the current Universe would be 25% in the dark matter while only 5% in the visual ordinary matter, the latter described by the "minimal Standard Model" (mSM). In this language, the dark-matter particles are supposed to be described by some real Standard Model. Indeed, there is certain urgent need for a truly Standard Model, which also accommodates those phenomena which are currently classified as "beyond the Standard Model".

We could call the above statements as "the old standard wisdom" (i.e., before the year of 2017). They would appear in talks, in papers, etc., as of today (2017). The Standard Model, the name that was invented by S. Weinberg in 1970's, of this paper deviates from the above standard wisdom in that there is new family gauge group $SU_f(3)$ on top of the well-known force-fields gauge group $SU_c(3) \times SU_L(2) \times U(1)$. The three "generations" of leptons are easily explained away and neutrino oscillations among three "generations" can be easily understood. In the quark world, all the entries are the triplets of the quark group

$SU_Q(3)$, such that the old notion of "generations" also never appear.

What we are doing in this paper is to describe the Standard Model that is based on Einstein's relativity principle and the quantum principle (established in the 20th Century) and that describes the *smallest units of matter*, including electrons, neutrinos, and quarks. The Standard Model [1] is, apart from the SSB "ignition" term, both the originally massless theory and the dimensionless theory. Both the gauge principle and the principle of renormalizability are now elevated to the status of the (mathematical) principles. (SSB means "spontaneous symmetry breaking".)

Nowadays we all know that we have firmly established the validity of Einstein's relativity principle and the quantum principle, the so-called two foundation pillars established as of the 20th Century. Starting from the beginning of the 21st Century, we should take it very seriously that there *exist* the *smallest* units of matter, such as the electrons, neutrinos, quarks, etc., a finite number of them. The behaviors of these smallest units of matter are described by the Standard Model, which we try to discuss in the present paper.

We insist on the quantum principle, in the sense that electrons, neutrinos, and quarks satisfy the Dirac equations, thereby satisfying the mysterious closed anti-commuting Dirac algebra, and they are fermions satisfying Pauli's exclusion principle when stacking up. Thus, these are truly the quantum phenomena, although this used to be a rather murky area as far as the quantum principle is concerned.

Theoretically, we suggest that we live in the quantum 4-dimensional Minkowski space-time with, *via the gauge principle*, the force-fields gauge-group structure, $SU_c(3) \times SU_L(2) \times U(1)$ applied in the triple quark world and $SU_L(2) \times U(1) \times SU_f(3)$ applied in the lepton world, built-in from the very beginning. In this overall background, the triple quark world is observed, of nuclear sizes, because of the (123) symmetry (i.e., under $SU_c(3) \times SU_L(2) \times U(1)$), while the lepton world is observed, of atomic sizes, in view of the other (123) symmetry (i.e., under $SU_L(2) \times U(1) \times SU_f(3)$). This gives us "our world" or "our Universe" [1].

The gauge principle helps to link the forces with our Minkowski space-time, since it tells us how to move the coordinates x_μ to a nearby point $x_\mu + \delta x_\mu$. The substitution, $\partial_\mu \rightarrow D_\mu$ for each basic unit of matter, reflects the information of the gauge principle.

Here $SU_f(3)$ stands for the $SU_{family}(3)$ family gauge theory, rather than $SU_{flavor}(3)$ (to be named later as $SU_Q(3)$, to avoid any further confusion) which derives from the isospin symmetry - the latter *not* a gauge theory. Presumably the size consideration forces the choice of strong $SU_c(3)$ over family $SU_f(3)$ for the triple quark world while, for the lepton world, we don't understand why the strong $SU_c(3)$ is completely shut off. In the lepton world, we insist on the symmetry $SU_L(2) \times U(1) \times SU_f(3)$, for the sake of the well-behaved limits at the small distances.

Among the detailed efforts to include the $SU_f(3)$ gauge group, we should mention those of T. Yanagida [2] and of Yue-Liang Wu [3]. In our language, the force-fields background, the triple quark world, and the lepton world are separate entities to begin with - they have their own individual ranges and other characteristics. Each of them are well-behaved, mathematically and physically. Thus, it is more natural that $SU_f(3)$ covers only the lepton world - making it free of Landau ghosts and making it asymptotically free. Thus, our $SU_f(3)$ does not cover the triple quark world, unlike [2, 3]. The applicability of $SU_f(3)$ to leptons, or quarks, or both, is the issue. In fact, the gauge principle forces us to make one

choice, and only one choice, from either of $SU_c(3)$ and $SU_f(3)$.

There are various reasons why the $SU_f(3)$ should be with the lepton world. Without the $SU_f(3)$, Landau's ghosts would kill it as a consistent theory, as conceptually it would blow up as the distance goes to zero. With the $SU_f(3)$ in the lepton world, the left-handed leptons have to put together to form a $(3, 2)$ multiplet; that would call for a $(3, 2)$ family Higgs multiplet; and then to make every $SU_f(3)$ gauge boson massive we require a $(3, 1)$ purely family Higgs multiplet. It turns out that it is harmless to have the charged partner in $\Phi(3, 2)$ - they would not go through the spontaneous symmetry breaking (SSB). In terms of the degrees of freedom (DOF), it turns out to be just right. It accommodates three kinds of neutrinos to oscillate among themselves.

Besides all these, the complex scalar field $\phi(x)$ alone cannot exist owing to the self-repulsive interaction $\lambda(\phi^\dagger\phi)^2$. So, we need $\Phi(3, 2)$ and $\Phi(3, 1)$ to guarantee the existence of the Standard-Model (SM) Higgs. It is indeed amazing.

3 The Entries and the Background of the Standard Model

In Newton's doctrine, it is *not* sufficient to specify an object by just giving the space-time point x_μ - instead, a complete description also needs the velocity information in terms of ∂_μ . Thus, the lagrangian is a functional of the coordinates and their first derivatives (velocities). The gauge principle means to replace the first derivatives by the gauge-invariant derivatives D_μ , with the differences $D_\mu - \partial_\mu$ characterized linearly by the gauge fields (potentials).

Thus, the background of the Standard Model contains not only the space-time coordinates but also their changes, i.e., the velocities; the changes in velocities through the gauge principle which contains the gauge-fields force information are required. So, this defines "the background" or "the overall background".

We have two kinds of the inputs (or the entry points): One input is at the fermi or 10^{13} cm sizes while the other input is at the atomic or 10^{-8} cm sizes.

The lepton world, or the atomic world, *via the gauge principle*, sees the force fields characterized by the gauge group $SU_L(2) \times U(1) \times SU_f(3)$. The typical sizes are of $(10^{-8} \text{ cm})^3$.

The quark world with the entries from members of the "quark" group $SU_Q(3)$ (used to be called "flavor $SU(3)$ symmetry"), or the nuclear world, *via the gauge principle*, sees the force fields characterized by the gauge group $SU_c(3) \times SU_L(2) \times U(1)$. The typical sizes are of $(10^{-13} \text{ cm})^3$.

To make it clear, the lepton world is the singlet members of the quark group $SU_Q(3)$. One should reckon the supreme status of the group $SU_Q(3)$ - it should cover everything. By our Universe, the Standard Model [1] should cover everything, including its mathematical meaning as a group. We call $SU_Q(3)$ the "quark" group mainly because the entry points of the quark world are the triplets of the group $SU_Q(3)$.

To be precise, the lepton world is defined by one left-handed $SU_Q(3)$ singlet and two right-handed $SU_Q(3)$ singlets. Meanwhile, the quark world is defined by one left-handed $SU_Q(3)$ triplet and two right-handed $SU_Q(3)$ triplets. These are the inputs of the Standard Model - we are trying to use the group theory to say things.

In the mathematical language, it may be appropriate to introduce the concept of the quark group $SU_Q(3)$ - a generalization of the flavor $SU(3)$ symmetry. It should apply to the

lepton world, as well. That makes the group concept omnipotent and the members of the group have some certain common characteristics to begin with. Mathematics and physics have some common traits, after all.

We wish to remark that the meaning of the "generations" disappears altogether in this Standard Model - in the quark world the triplet members of the quark group $SU_Q(3)$ serve as the basic entries, while in the lepton world the "generations" are replaced by the fermion members of the family gauge group $SU_f(3)$. Thus, "the three generations" are replaced by the entries of the gauge group $SU_f(3)$ while "the three generations" are replaced by the triplet entries of the quark group $SU_Q(3)$ - it turns out that we no longer need to talk about the "generations".

Both the matter worlds, the lepton world at the atomic scale and the quark world at the fermi scale, are asymptotically free, and thus free of Landau ghosts. In fact, the overall consistency of the Standard Model [1] guarantees its mathematical existence. In other world, we can analyze its (mathematical) existence by studying its overall consistency.

In view of the fundamental importance, we would like to suggest to promote "the gauge principle" and "the principle of renormalizability" to the full status of the principles, its meaning more as the mathematical principles (than in the physics principles).

4 The Principle of Renormalizability

A real strange thing happens in the 4-dimensional Minkowski space-time [1] - a story which we should have known but so far didn't. A complex scalar field $\phi(x)$ should have the self-interaction $\lambda(\phi^\dagger\phi)^2$ with a dimensionless (positive) λ cannot be observed if being alone - since a positive λ means self-repulsive. It is dimensionless so it is determined by the space-time, but *not* by the field itself. Maybe $\lambda = \frac{1}{8}$ but we can't prove it so far. Thus, the "related" complex scalar fields $\Phi(1,2)$ (Standard-Model Higgs), $\Phi(3,2)$ (mixed family Higgs), and $\Phi(3,1)$ (purely family Higgs) come together to sing a chorus - an SSB chorus with the various force-fields gauge fields.

The self-repulsive term $\lambda(\phi^\dagger\phi)^2$ ($\lambda > 0$) is completely acceptable in terms of renormalizability, but it precludes the existence of the field $\phi(x)$. Thus, we need the existence of another "related" complex scalar field $\phi_2(x)$ such that a mutual-attractive term $-4\lambda(\phi_1^\dagger\phi_2) \cdot (\phi_2^\dagger\phi_1)$ exist to save the situation.

The logic is rather simple. We're living in the quantum 4-dimensional Minkowski space-time. Through the gauge principle, we (i.e., in terms of the basic units of matter) are also connected with the various force fields (gauge fields). We have complex scalar fields, to make up the gauge fields (to make them massive). The so-called "matter", in terms of Dirac fields, including electrons, neutrinos, and quarks, all exist there. The complex scalar fields have the similar characteristics as the overall background - the quantum 4-dimensional Minkowski space-time. They are self-repulsive and cannot exist alone by itself. But they are "related" - the SM Higgs $\Phi(1,2)$, the mixed family Higgs $\Phi(3,2)$, and the purely family Higgs $\Phi(3,1)$, "related" to lower their total energy and to make the SM and family gauge bosons massive. This makes up the so-called "overall background". The triple quark world (with the entries as members of the quark group $SU_Q(3)$), with the (123) gauge symmetry, is accepted by this background. Further, the lepton world, with another (123) gauge symmetry and with

a much bigger scale (sizes), is also accepted by the same background. We are talking about the origin of fields or of point-like particles, talking about the "renormalizable" Standard Model [4, 1]. We should figure out a way to think about our entire world, even if we might not be on the right track.

At the end, the Standard Model is a consistent language of mathematics - each field represents a certain particle, no more redundant field (particle). There is the "overall" background, namely, the quantum 4-dimensional Minkowski space-time (a physical system) with, *via the gauge principle*, the force-fields gauge-group structure suitably built-in from the very beginning, plus the triple quark world and the lepton world.

5 The Gauge Principle

The "*basic units of matter*", which are based on the right-handed Dirac components or left-handed Dirac components but have their group assignments *explicit* under the group $SU_c(3) \times SU_L(2) \times U(1) \times SU_f(3)$, would be more appropriate than the so-called "building blocks of matter". Moreover, each basic unit derives from one kinetic-energy term, $-\bar{\Psi}_R \gamma_\mu \partial_\mu \Psi_R$ or $-\bar{\Psi}_L \gamma_\mu \partial_\mu \Psi_L$, and from only one such term. The gauge principle tells us that ∂_μ is to be replaced by some "gauge-invariant" derivative D_μ . It is clear that this would be an economical way to write down a theory in the globally consistent manner.

In fact, the gauge group would be $SU_c(3) \times SU_L(2) \times U(1)$ or $SU_L(2) \times U(1) \times SU_f(3)$, since the two $SU(3)$, if showing up simultaneously, would contradict each other. Moreover, if we introduce the "quark" group $SU_Q(3)$, used to be called "the flavor $SU(3)$ symmetry", the entries in the theory should be the triple (d'_R, s'_R, b'_R) , etc., instead of d'_R individually, and so on - thus, the three "generations" in the quark case is a misnomer.

These *basic* units of matter turn out to be the *smallest* units of matter - so, we may use these names interchangeably. For example, an electron, being one basic unit of matter, is indivisible by any means. So are the other basic units of matter, such as neutrinos, quarks, etc.

The most important aspect is that the kinetic energy term $-\bar{\Psi}(x) \gamma_\mu \partial_\mu \Psi(x)$ can appear *once and only once* in the lagrangian - this fact was given in Newton's classic doctrine in the description of particle's motions. The gauge principle $\partial_\mu \rightarrow D_\mu$ then implies that there is *one and only one* D_μ . The difference, $D_\mu - \partial_\mu$, is a measure of the influence of the force(s). This is why, in statement of the Standard Model, the force-fields gauge-group structure has to be built in from the very beginning - it has to precede the existence of the quark world or of the lepton world.

Clearly, we may split the ordinary-matter world into the "quark" world and the "lepton" world, as the ranges of their "livings" are quite different from each other - the quark world is at the fermi scale while the lepton world at the atomic scale. At low enough temperature, the systems of quarks (and anti-quarks) are confined while systems in leptons could roam anywhere.

We know that the right-handed neutrinos do not appear in the minimal Standard Model - so, $(\nu_{\tau R}, \nu_{\mu R}, \nu_{e R})$ would make a perfect triplet under $SU_f(3)$. In a recent paper [5], we proposed to put $((\nu_\tau, \tau)_L, (\nu_\mu, \mu)_L, (\nu_e, e)_L)$ (*columns*) ($\equiv \Psi_L(3, 2)$) as the $SU_f(3)$ triplet and $SU_L(2)$ doublet. The next question is how to assign right-handed τ_R, μ_R , and e_R under

$SU_f(3)$. We could write down the mass term for the charged leptons - if they are singlets or (τ_R, μ_R, e_R) is an $SU_f(3)$ triplet? The singlet assignment can be ruled out since there are undesirable crossed terms in, e.g., $\Psi_L(\bar{3}, 2)e_R\Phi(3, 2)$. So, three of them would better form a triplet - $\Psi_R^C(3, 1)$. In fact, This completes the list of "the basic units" in the lepton world.

The lepton world is constructed from three group elements $(\nu_{\tau R}, \nu_{\mu R}, \nu_{e R})$, (τ_R, μ_R, e_R) , and $((\nu_{\tau}, \tau)_L, (\nu_{\mu}, \mu)_L, (\nu_e, e)_L)$ (columns). The group is $SU_L(2) \times U(1) \times SU_f(3)$. This should be regarded as the starting point of the lepton world.

Since we wish to propose the $SU_f(3)$ family gauge theory as a way to understand why there are three generations, it requires all additional particles, i.e., (eight) gauge bosons and (four) residual family Higgs, very massive. In the proposal of Hwang and Yan [5], we would have one Standard-Model Higgs field $\Phi(1, 2)$, one complex family Higgs triplet $\Phi(3, 1)$, and another family triplet-doublet complex scalar fields $\Phi(3, 2)$. Amazingly enough, two neutral complex triplets, $\Phi(3, 1)$ and $\Phi^0(3, 2)$ would indeed undergo the desired Higgs mechanism and the three charged scalar fields would remain massive, i.e. there is no spontaneous symmetry breaking (SSB) in the charged Higgs sector.

So far, we have decided on the basic units of matter - those left-handed and right-handed objects (quarks or leptons); the gauge group is chosen to be $SU_c(3) \times SU_L(2) \times U(1) \times SU_f(3)$. As we said, we live in the quantum 4-dimensional Minkowski space-time with, *via the gauge principle*, the force-fields gauge-group structure $SU_c(3) \times SU_L(2) \times U(1) \times SU_f(3)$ built-in from the very beginning. This is the "background" of everything, i.e., the lepton world and the quark world. For notations, we use Wu and Hwang [6].

The gauge symmetry $SU_L(2) \times U(1) \times SU_f(3)$ applies to the lepton world, since the lepton world does not recognize the strong interaction $SU_c(3)$. The gauge symmetry $SU_c(3) \times SU_L(2) \times U(1)$ applies to the quark world, and the "quark" group $SU_Q(3)$ (which is not a gauge symmetry) leaves the mixing heavily among the three "generations" of quarks. Note that one $SU(3)$ is enough but two of them would be too much, as far as the gauge principle is concerned. *Thus, the deep mystery can be resolved by introducing the "quark" group $SU_Q(3)$, used to be called "the flavor $SU(3)$ symmetry". The quark entities are members of this group.*

In the quark world, we have the "basic unit of matter", for the up-type right-handed quarks (u_R, c_R, t_R) (column), a triplet under the quark group $SU_Q(3)$,

$$D_\mu = \partial_\mu - ig_c \frac{\lambda^a}{2} G_\mu^a - i \frac{2}{3} g' B_\mu, \quad (1)$$

and, for the rotated down-type right-handed quarks (d'_R, s'_R, b'_R) (column), a triplet under the quark group $SU_Q(3)$,

$$D_\mu = \partial_\mu - ig_c \frac{\lambda^a}{2} G_\mu^a - i(-\frac{1}{3}) g' B_\mu. \quad (2)$$

On the other hand, we have the basic unit of matter, for the $SU_L(2)$ quark doublets and a triplet under the quark group $SU_Q(3)$,

$$D_\mu = \partial_\mu - ig_c \frac{\lambda^a}{2} G_\mu^a - ig \frac{\vec{\tau}}{2} \cdot \vec{A}_\mu - i \frac{1}{6} g' B_\mu. \quad (3)$$

Thus, the term of "generation" is a misnomer when the quark group $SU_Q(3)$ is introduced. Similarly, the "generation" in the lepton world becomes a misnomer after the gauge

group $SU_f(3)$ is introduced. On the quark group $SU_Q(3)$, we should recall the role of the CKM matrix [7] to ascertain that the elements of the quark group $SU_Q(3)$ should be the input entities, rather than each individual quark, in these discussions.

At this juncture, we wish to discuss the basic role of the gauge principle, a strange topic which was discussed only along one line or two in the 20th Century [6]. The reasons which we have to decide the D_μ in the substitution $\partial_\mu \rightarrow D_\mu$ for each basic unit of matter are at least two-fold.

For each basic unit matter, such as the three quark units as given in the above, there is only one kinetic-energy term in the lagrangian, and there should be only one.

The purpose of the kinetic-energy term is to tell how the basic unit of matter changes when the space-time point moves from one point to another. When we introduce the quantum 4-dimensional Minkowski space-time, the coordinates (x_μ) alone are not adequate to describe the motion of the particle.

In the above example of the quark world, the gauge principle requires the knowledge of the force fields through the gauge fields. That is why the gauge-group structure has to be implemented "from the very beginning".

Moreover, the implementation of $SU_c(3) \times SU_L(2) \times U(1)$ precludes the assignment of $SU_{family}(3)$ of the various basic units of matter in the quark world. We presume the reason might be that the $SU_f(3)$ coupling κ is much weaker than the strong $SU_c(3)$ coupling g_c . Thus, the implementation of $SU_c(3)$ precludes the contradictory assignments of $SU_f(3)$ from occurrence. But the space-time itself is $SU_c(3) \times SU_L(2) \times U(1) \times SU_f(3)$ in nature.

As for the lepton world, we introduce the family triplets, $(\nu_\tau^R, \nu_\mu^R, \nu_e^R)$ (column) ($\equiv \Psi_R(3,1)$) and (τ_R, μ_R, e_R) (column) ($\equiv \Psi_R^C(3,1)$), under $SU_f(3)$. The gauge principle for the "basic unit of matter" is, for $\Psi_R(3,1)$,

$$D_\mu = \partial_\mu - i\kappa \frac{\bar{\lambda}^a}{2} F_\mu^a. \quad (4)$$

Similarly for $\Psi_R^C(3,1)$.

And, for the left-handed $SU_f(3)$ -triplet and $SU_L(2)$ -doublet $((\nu_\tau^L, \tau^L), (\nu_\mu^L, \mu^L), (\nu_e^L, e^L))$ (all columns) ($\equiv \Psi_L(3,2)$), the basic unit is

$$D_\mu = \partial_\mu - i\kappa \frac{\bar{\lambda}^a}{2} F_\mu^a - ig \frac{\vec{\tau}}{2} \cdot \vec{A}_\mu + i \frac{1}{2} g' B_\mu. \quad (5)$$

The generation of the various quark masses is through the Standard-Model way. For the charged leptons, we have to choose $\bar{\Psi}_L(3,2) \Psi_R^C(3,1) \Phi(1,2)$. Note that, in the U-gauge, the SM Higgs looks like $(0, \frac{1}{\sqrt{2}}(v + \eta(x)))$, only one degree of freedom [6]. Thus, the neutral neutrino triplet $\Psi_R(3,1)$ cannot be used in the above coupling, because of the charge conservation.

Moreover, there is an important point in the lepton world. In the $SU_c(3) \times SU_L(2) \times U(1) \times SU_f(3)$ Standard Model [8], the neutrino mass term assumes a new form:

$$i \frac{h}{2} \bar{\Psi}_L(3,2) \times \Psi_R(3,1) \cdot \tilde{\Phi}(3,2) + h.c., \quad (6)$$

where $\Psi(3, i)$ is the neutrino triplet (with the first label for $SU_f(3)$ and the second for $SU_L(2)$). The cross-dot (curl-dot) product is somewhat new, referring to the singlet combination of three triplets in $SU(3)$. The Higgs field $\Phi(3, 2)$ is new in this effort [5], because it carries some nontrivial $SU_L(2)$ charge. The consistent definition of $\tilde{\Phi}(3, 2)$ is given below.

This off-diagonal neutrino mass term offers us a natural way to describe neutrino oscillations, since the neutral part of $\Phi(3, 2)$ could each receive vacuum expectation values.

Note that, for charged leptons, the Standard-Model choice is $\bar{\Psi}_L(\bar{3}, 2)\Psi_R^C(3, 1)\Phi(1, 2) + c.c.$, which gives three leptons an equal mass. But, in view of that if (ϕ_1, ϕ_2) is an $SU(2)$ doublet then $(\phi_2^\dagger, -\phi_1^\dagger)$ is another doublet, we could form $\tilde{\Phi}^\dagger(3, 2)$ from the doublet-triplet $\Phi(3, 2)$. So, this $\Phi(3, 2)$ is used below, in a consistent notation as $\Phi(1, 2)$, while its "related" $\tilde{\Phi}(3, 2)$ is used above.

$$i\frac{h^C}{2}\bar{\Psi}_L(3, 2) \times \Psi_R^C(3, 1) \cdot \Phi(3, 2) + h.c.. \quad (7)$$

Here vacuum expectation values of $\Phi^0(3, 2)$ give rise to the imaginary off-diagonal (hermitian) elements in the 3×3 mass matrix, so removing the equal masses of the charged leptons.

Here the couplings h^C and h are closely related to the coupling strength κ for the family gauge bosons [9]. Note that all these couplings are dimensionless in the 4-dimensional Minkowski space-time; thus, it is the intrinsic property of the 4-dimensional Minkowski space-time - in other words, it should be *not* adjustable by those living in this space-time, at will.

The triple quark world in the quantum 4-dimensional Minkowski space-time is governed by the six dimensionless couplings, three for the strengths of the interactions and another three mass-related parameters. Similarly, the lepton world in the quantum 4-dimensional Minkowski space-time is governed by six dimensionless couplings, three for the interactions and three for the mass-related parameters. The entries for the quark world are members of the quark group $SU_Q(3)$ while the entries for the lepton world are singlets under $SU_Q(3)$. Using our arguments, the dimensionless couplings are determined *globally* by the quantum 4-dimensional Minkowski space-time, the covering space of the game. All these might lead to some philosophical insights.

Mathematics-wise, the Standard Model [1] is basically a group theory, though fairly complicated. The entry point is characterized by the basic units of matter, which are expressed in terms of quarks, electrons, neutrinos, etc. The gauge principle, i.e., $\partial_\mu \rightarrow D_\mu$ which varies with the basic unit of matter, also brings in the various gauge fields (i.e., the force fields). The complex scalar fields enter for the massive gauge fields. As the beginning, there is a quark group $SU_Q(3)$ for which the basic units of matter for quarks are triplets while the basic units of matter for leptons are singlets. The basic units of matter are described by the Dirac equations of the anti-commuting and closed 16-elements Dirac algebra. If we count everything carefully, all physics, including the hadron CP-violating phase, neutrino oscillations, etc., are taken into account.

6 The Origin of Mass

As pointed out in an early version of the paper [8], we may imagine that, in the U-gauge, the Standard-Model Higgs $\Phi(1, 2)$ looks like $(0, (v + \eta(x))/\sqrt{2})$ (*column*) and $\Phi^\dagger(\bar{3}, 2)\Phi(1, 2)$ would pick out the neutral sector naturally. In fact, the quartic term $(\Phi^\dagger(\bar{3}, 2)\Phi(1, 2))(\Phi^\dagger(1, 2)\Phi(3, 2))$ with a suitable sign, would modify the mass term for $\Phi(3, 2)$ field such that the neutral sector has SSB while the charged sector remains massive. This "projected-out" Higgs mechanism is what we need.

We may [8] write down the terms for potentials among the three Higgs fields, subject to (1) that they are renormalizable, and (2) that symmetries are only broken spontaneously (via the Higgs or induced Higgs mechanism). Thus, we write, from the earliest (17 April 2013) version of [8],

$$V = V_{SM} + V_1 + V_2 + V_3, \quad (8)$$

$$V_{SM} = \mu^2 \Phi^\dagger(1, 2)\Phi(1, 2) + \lambda(\Phi^\dagger(1, 2)\Phi(1, 2))^2, \quad (9)$$

$$\begin{aligned} V_1 = & M^2 \Phi^\dagger(\bar{3}, 2)\Phi(3, 2) + \lambda_1(\Phi^\dagger(\bar{3}, 2)\Phi(3, 2))^2 \\ & + \epsilon_1(\Phi^\dagger(\bar{3}, 2)\Phi(3, 2))(\Phi^\dagger(1, 2)\Phi(1, 2)) + \eta_1(\Phi^\dagger(\bar{3}, 2)\Phi(1, 2))(\Phi^\dagger(1, 2)\Phi(3, 2)) \\ & + \epsilon_2(\Phi^\dagger(\bar{3}, 2)\Phi(3, 2))(\Phi^\dagger(\bar{3}, 1)\Phi(3, 1)) + \eta_2(\Phi^\dagger(\bar{3}, 2)\Phi(3, 1))(\Phi^\dagger(\bar{3}, 1)\Phi(3, 2)) \\ & + (\delta_1 i \Phi^\dagger(3, 2) \times \Phi(3, 2) \cdot \Phi^\dagger(3, 1) + h.c.), \end{aligned} \quad (10)$$

$$\begin{aligned} V_2 = & \mu_2^2 \Phi^\dagger(\bar{3}, 1)\Phi(3, 1) + \lambda_2(\Phi^\dagger(\bar{3}, 1)\Phi(3, 1))^2 + (\delta_2 i \Phi^\dagger(3, 1) \cdot \Phi(3, 1) \times \Phi(3, 1) + h.c.) \\ & + \lambda'_2 \Phi^\dagger(\bar{3}, 1)\Phi(3, 1)\Phi^\dagger(1, 2)\Phi(1, 2), \end{aligned} \quad (11)$$

$$\begin{aligned} V_3 = & (\delta_3 i \Phi^\dagger(3, 2) \cdot \Phi(3, 2) \times (\Phi^\dagger(1, 2)\Phi(3, 2)) + h.c.) \\ & + (\delta_4 i (\Phi^\dagger(3, 2)\Phi(1, 2)) \cdot \Phi^\dagger(3, 1) \times \Phi(3, 1) + h.c.) \\ & + \eta_3(\Phi^\dagger(\bar{3}, 2)\Phi(1, 2)\Phi(3, 1) + c.c.). \end{aligned} \quad (12)$$

In obtaining the last expression, we maintain that every terms are naively renormalizable since the terms are of power four or less (in scalar fields). Note that the terms in $\delta_j i$ involve the so-called "SU(3) operations", as before.

This is *the first step* to decide what the Standard Model is going to be.

As shown earlier [9, 10], two triplets of complex scalar fields would make the eight family gauge bosons and four residual family Higgs particles all massive. In the $SU_f(3)$ family gauge theory treated alone, there are many ways of accomplishing such goal. In the present complicated case, the equivalence between two triplets is lost, but we discover that this modification would be a nice way to patch up that three neutrinos couple to independent vacuum expectation values (and the family Higgs), explaining neutrino oscillations among three generations.

The "principle" of renormalizability turns out to be very powerful. To the least, we may use it to define the calculability of the theory. It implies the existence of some branch of calculable mathematics (in quantum field theory).

On the other hand, there are too many terms - for example, there are three "ignition" terms; if there is only one, we must make the smart choice; etc. If we look into the lepton world or into the quark world, it is striking to realize that all the couplings are dimensionless in the 4-dimensional Minkowski space-time. In the Higgs and gauge sector,

we could take only one "ignition" term and take all the rest couplings to be dimensionless - then, the Standard Model is virtually a complete dimensionless theory (here "virtually" means "apart from the 'ignition' term").

This is *the second (last) step* to decide the Standard Model is going to be - to figure out what the dimensionless theory looks like.

Thus, "renormalizability" plus "dimensionless-ness in the 4-dimensional Minkowski space-time" leads to the final theory - is that elegant?

Before 2014, our understanding of mass generation for the building blocks of matter was still lacking, not mentioning the origin of mass; so, the same story for the mysterious Higgs mechanisms. The story has changed completely if we accept the recent proposal [11] on the origin of mass.

The Higgs lagrangian (i.e., Eqs. (8)-(12) above) is based on renormalizability. Physics-wise, we should think of its redundancy - for example, if there is a spontaneous symmetry breaking (SSB), there are three different channels for switch-on, depending on which complex scalar field gets turn on. On the other hand, when the temperature is high enough, those mass terms, except the switch-on of ignition, would become negligible; or, before the generation of masses.

Thus, the real world should be like this: There is at most one ignition channel for spontaneous symmetry breaking (SSB). When the temperature is high enough, all the mass terms should be zero (if necessary, by comparison).

In the 4-dimensional Minkowski space-time, the complex scalar field $\phi(x)$ have one unique feature - that the renormalizable term $\lambda(\phi^\dagger\phi)^2$ is allowed (only in the 4-dimensional space-time) [1]. This self-interaction is repulsive, explaining why we cannot see it unless under exceptions. Further, this term is dimensionless, so the coupling λ is a pure number that is determined by the 4-dimensional Minkowski space-time, *not* by the complex scalar field itself [11, 1].

Thus, one important beginning point is the 4-dimensional Minkowski space-time. The other point is whether the force-fields gauge symmetry $SU_c(3) \times SU_L(2) \times U(1) \times SU_f(3)$ should also be a part of this beginning point. As all the future entities should have these gauge group assignments (similar to quantum numbers), our suggestion is "yes" - that gives the assertion that "we live in the 4-dimensional Minkowski space-time with the force-fields gauge group $SU_c(3) \times SU_L(2) \times U(1) \times SU_f(3)$ built-in from the outset" [1].

Let us re-iterate the origin of mass [11]:

Suppose that, before the spontaneous symmetry breaking (SSB), the Standard Model does not contain any parameter that is pertaining to "mass", but, after the SSB, all particles in the Standard Model acquire the mass terms as it should - we call it "the origin of mass". In [11], we show that this is indeed the case - explaining the origin of mass. In this way, we sort of tie "the origin of mass" to the effects of the SSB, or the generalized Higgs mechanism.

In our opinion, this way of explaining the origin of mass is of fundamental importance. After all, we could not give some real meaning to the term "mass"; it makes sense to interpret masses as our proposal of the origin of mass [11]. According to [11], the masses would disappear altogether at the temperature high enough, if for example we are thinking of the early Universe. On other hand, when the matter system becomes so dense and so hot in the process of forming a "black hole", the mass parameters would lose its meaning altogether - making the interior inside the horizon of the Schwarzschild metric something

which we can't understand so far.

Starting from the Higgs sector of the Standard Model [8], we have the Standard-Model Higgs $\Phi(1,2)$, the purely family Higgs $\Phi(3,1)$, and the mixed family Higgs $\Phi(3,2)$, with the first label for $SU_f(3)$ and the second for $SU_L(2)$. We need another triplet $\Phi(3,1)$ since all eight family gauge bosons have to be massive [9].

We could say that it is the most interesting to put the different but related complex scalar fields (Higgs) in the same pot, letting them interact together. Two fields ϕ_a and ϕ_b can interact "directly" via $(\phi_a^\dagger \phi_a)(\phi_b^\dagger \phi_b)$ but, if related, via $((\phi_a^\dagger \phi_b)(\phi_b^\dagger \phi_a))$. Here we always talk about those things which are renormalizable. Under the renormalizable game, these are closely linked with the quartic self coupling such as $\lambda(\phi_a^\dagger \phi_a)^2$.

Besides, in the 4-dimensional Minkowski space-time, these terms have dimensionless couplings, or pure numbers. So, λ depends on the 4-dimensional Minkowski space-time, *not* on the field $\phi(x)$ itself. This is a very interesting aspect. *Thus, when we put three complex scalar fields $\Phi(1,2)$, $\Phi(3,2)$, and $\Phi(3,1)$ in this pot of the 4-dimensional Minkowski space-time all the quartic terms, including those crossed, are playing pure-number games among themselves. Consequently, the masses for the various Higgs emerge as the result. This discovery is closely linked to the origin of fields (point-like particles) [1].*

In fact we are "playing" with "energies", in terms of the complex scalar fields. For two related complex scalar fields, the cross term helps to lower the total energy. For two unrelated complex scalar fields, the cross term does not exist.

In the U-gauge, we choose to have

$$\Phi(1,2) = (0, \frac{1}{\sqrt{2}}(v+\eta)), \quad \Phi^0(3,2) = \frac{1}{\sqrt{2}}(u_1+\eta'_1, u_2+\eta'_2, u_3+\eta'_3), \quad \Phi(3,1) = \frac{1}{\sqrt{2}}(w+\eta', 0, 0), \quad (13)$$

all in columns. The five components of the complex triplet $\Phi(3,1)$ get absorbed by the $SU_f(3)$ family gauge bosons and the neutral part of $\Phi(3,2)$ has three real parts left - together making all eight family gauge bosons massive.

Before the mixing, the masses of the various Higgs are given by, using the general Higgs lagrangian (cf. Eqs. (8)-(12)),

$$\begin{aligned} \eta : & \quad (\mu^2/\lambda) + \frac{1}{4}(\epsilon_1 + \eta_1)u_i u_i + \frac{\lambda'_2}{4}w^2, \\ \eta' : & \quad (\mu_2^2/\lambda_2) + \frac{\epsilon_2}{4}u_i u_i + \frac{\eta_2}{4}u_1^2 + \frac{\lambda'_2}{4}v^2, \\ \eta'_1 : & \quad M^2 + \frac{1}{4}(\epsilon_1 + \eta_1)v^2 + \frac{\epsilon_2}{4}w^2 + (\lambda_1 - term), \\ \eta'_{2,3} : & \quad M^2 + \frac{1}{4}(\epsilon_1 + \eta_1)v^2 + \frac{\epsilon_2}{4}w^2 + (\lambda_1 - term), \\ \phi_1 : & \quad M^2 + \frac{1}{2}\epsilon_1 v^2 + \frac{1}{2}\epsilon_2 w^2 + \frac{1}{2}\eta_2 w^2 + \frac{\lambda_1}{2}u_i u_i, \\ \phi_{2,3} : & \quad M^2 + \frac{1}{2}\epsilon_1 v^2 + \frac{1}{2}\epsilon_2 w^2 + \frac{\lambda_1}{2}u_i u_i. \end{aligned} \quad (14)$$

The mixing term looks like, apart from some common factor:

$$2(\epsilon_1 + \eta_1)u_i \eta'_i v \eta + 2\epsilon_2 u_i \eta'_i w \eta' + 2\eta_2 u_1 \eta'_1 w \eta' + 2\lambda'_2 w \eta' v \eta. \quad (15)$$

There are the other mixings (such the mixing inside $\eta'_{1,2,3}$), to be discussed later in this paper. To work out on "the origin of mass", we would drop out all "mass" terms to begin with.

Let us try to fix the notations further. See Ch. 13, Ref. [?]. For η' going through SSB, we have the following terms, neglecting the small λ'_2 term,

$$\frac{\mu_2^2}{2}(\eta' + w)^2 + \left(\frac{\epsilon_2}{4}u_i u_i + \frac{\eta}{4}u_1^2\right)(\eta' + w)^2 + \frac{\lambda_2}{4}(\eta' + w)^4. \quad (16)$$

SSB means that all the linear terms add up to zero, resulting the change in sign of the mass term, $\frac{1}{2}(2\lambda_2 w^2(\eta')^2)$. Note that, for real fields, a factor of $\frac{1}{2}$ should be factored out.

The same applies to other SSB fields, even though the original minus signs are generated by other fields. This applies for the SM Higgs η .

In other word, the three "related" scalar (Higgs) fields $\Phi(1, 2)$, $\Phi(3, 2)$, and $\Phi(3, 1)$ should be "equivalent" among themselves. The spontaneous symmetry breaking (SSB) is happening for all of them, actively or passively. Thus, λ , λ_1 , and λ_2 should be the same. λ'_2 should be zero or vanishingly small.

All the mass terms, including those SSB-driving terms, should be absent, apart from one SSB-driving term. The scalar field (ϕ_a) could affect a different scalar field (ϕ_b), if they are triplets under $SU_f(3)$, etc. It is easy to see that only one SSB-driving term is enough for all the three Higgs fields – there may be several SSB's for the neutral fields - in our case, it works for all of them.

We are working with the lagrangian, i.e., the energy that should be bounded from below ("positive definiteness"). The combination $(\phi_a^\dagger \phi_a + \phi_b^\dagger \phi_b)^2 - 4(\phi_a^\dagger \phi_b)(\phi_b^\dagger \phi_a)$ would work very well. This in fact helps to fix $\epsilon_{1,2}$ and $\eta_{1,2}$. As we argued before, the three λ 's are the same.

As for the SSB-driving term, we decide to keep the purely family term, $\mu_2^2 \Phi^\dagger(3, 1)\Phi(3, 1)$. As we shall see that the symmetry breaking would occur much earlier (in the history of the early Universe), or at a much higher temperature.

Thus, we have

$$\begin{aligned} V_{Higgs} = & \mu_2^2 \Phi^\dagger(3, 1)\Phi(3, 1) + \lambda(\Phi^\dagger(1, 2)\Phi(1, 2) + \cos\theta_P \Phi^\dagger(3, 2)\Phi(3, 2))^2 \\ & + \lambda(-4\cos\theta_P)(\Phi^\dagger(\bar{3}, 2)\Phi(1, 2))(\Phi^\dagger(1, 2)\Phi(3, 2)) \\ & + \lambda(\Phi^\dagger(3, 1)\Phi(3, 1) + \sin\theta_P \Phi^\dagger(3, 2)\Phi(3, 2))^2 \\ & + \lambda(-4\sin\theta_P)(\Phi^\dagger(\bar{3}, 2)\Phi(3, 1))(\Phi^\dagger(3, 1)\Phi(3, 2)). \end{aligned} \quad (17)$$

These are two perfect squares minus the other extremes, to guarantee the positive definiteness, when the minus μ_2^2 was left out. (θ_P may be referred to as "Pauchy's angle".) ϵ_1 , η_1 , and ϵ_2 , η_2 are expressed in λ , a great simplification. Note that we only include the interference terms between those involving the same group, $SU_f(3)$ or $SU_L(2)$; thus $\lambda'_2 = 0$.

From the expressions of $u_i u_i$ and v^2 , we obtain

$$v^2(3\cos^2\theta_P - 1) = \sin\theta_P \cos\theta_P w^2. \quad (18)$$

And the SSB-driven η' yields

$$w^2(1 - 2\sin^2\theta_P) = -\frac{\mu_2^2}{\lambda} + (\sin 2\theta_P - \tan\theta_P)v^2. \quad (19)$$

These two equations show that it is necessary to have the driving term, since $\mu_2^2 = 0$ implies that everything is zero. Also, $\theta = 45^\circ$ is the (lower) limit.

The mass squared of the SM Higgs η is $2\lambda\cos\theta_P u_i u_i$, as known to be $(125 \text{ GeV})^2$. The famous v^2 is the number divided by 2λ , or $(125 \text{ GeV})^2/(2\lambda)$. Using PDG's for e , $\sin^2\theta_W$, and the W -mass [12], we find $v^2 = 255 \text{ GeV}$. So, $\lambda = \frac{1}{8}$, a simple model indeed.

The ratio of the VEV to its Higgs mass is determined by 2λ , whether the channel is not ignited or not. We might choose the channel of η' (the purely family Higgs) or that of η (the SM Higgs) as the ignition channel, but three Higgs channels have different labels. The three Lorentz-invariant scalar fields have different internal structures - an amusing question for further investigation.

The mass squared of η' is $-2(\mu_2^2 - \sin\theta_P u_1^2 + \sin\theta_P(u_2^2 + u_3^2))$. The other condensates are $u_1^2 = \cos\theta_P v^2 + \sin\theta_P w^2$ and $u_{2,3}^2 = \cos\theta_P v^2 - \sin\theta_P w^2$ while the mass squared of η'_1 is $2\lambda u_1^2$, those of $\eta'_{2,3}$ be $2\lambda u_{2,3}^2$. The mixings among η'_i themselves are neglected in this paper.

There is no SSB for the charged Higgs $\Phi^+(3, 2)$. The mass squared of ϕ_1 is $\lambda(\cos\theta_P v^2 - \sin\theta_P w^2) + \frac{\lambda}{2}u_i u_i$ while $\phi_{2,3}$ be $\lambda(\cos\theta_P v^2 + \sin\theta_P w^2) + \frac{\lambda}{2}u_i u_i$. (Note that a factor of $\frac{1}{2}$ appears in the kinetic and mass terms when we simplify from the complex case to that of the real field; see Ch. 13 of [6].)

A further look of these equations tells that $3\cos^2\theta_P - 1 > 0$ and $2\sin^2\theta_P - 1 > 0$. A narrow range of θ_P is allowed (greater than 45° while less than 57.4° , which is determined by the group structure). For illustration, let us choose $\cos\theta_0 = 0.6$ and work out the numbers as follows: (Note that $\lambda = \frac{1}{8}$ is used.)

$$\begin{aligned} 6w^2 &= v^2, & -\mu_2^2/\lambda &= 0.32v^2; \\ \eta : & 2\lambda\cos\theta_0 u_i u_i = (125 \text{ GeV})^2, & v^2 &= (250 \text{ GeV})^2; \\ \eta' : & \text{mass}^2 = (51.03 \text{ GeV})^2, & w^2 &= v^2/6; \\ \eta'_1 : & \text{mass}^2 = (107 \text{ GeV})^2, & u_1^2 &= 0.7333v^2; \\ \eta'_{2,3} : & \text{mass}^2 = (85.4 \text{ GeV})^2, & u_{2,3} &= 0.4667v^2; \\ \phi_1 : & \text{mass} = 100.8 \text{ GeV}; & \phi_{2,3} : & \text{mass} = 110.6 \text{ GeV}. \end{aligned} \quad (20)$$

All numbers appear to be reasonable. In the above, $\cos\theta_0$ is the only free parameter until one of the family Higgs particles $\eta'_{1,2,3}$ and η' is found experimentally. Since the new objects need to be accessed in the lepton world, it would be a challenge for our experimental colleagues.

As a footnote, our Standard Model predicts that the mass of the SM Higgs η is a half of the vacuum expectation value v - a prediction in the origin of mass [11].

As for the range of validity, $\frac{1}{3} \leq \cos^2\theta_P \leq \frac{1}{2}$. The first limit refers to $w^2 = 0$ while the second for $\mu_2^2 = 0$.

We may fix up the various couplings, using our common senses. The cross-dot products would be similar to κ , the basic coupling of the family gauge bosons. The electroweak coupling g is 0.6300 while the strong QCD coupling $g_s = 3.545$; my first guess for κ would be about 0.1. The masses of the family gauge bosons would be estimated by using $\frac{1}{2}\kappa \cdot w$, so slightly less than 10 GeV . (In the numerical example with $\cos\theta_P = 0.6$, we have $6w^2 = v^2$ or $w = 102 \text{ GeV}$. This gives $m = 5 \text{ GeV}$ as the estimate.) So, the range of the family forces, existing in the lepton world, would be 0.02 fermi .

Furthermore, the λ'_2 term (i.e., the interference between the spaces $(3, 1)$ and $(1, 2)$) could be neglected safely in these discussions. But similar discussions to justify our Standard Model could be presented. There is one interesting aspect that the scalar fields, when having same $SU_f(3)$ or $SU_L(2)$ group structure, should be dealt with together, like in this paper.

The term that ignites the SSB is chosen to be with η' , the purely family Higgs. This in turn ignites EW SSB and others. It explains the origin of all the masses, in terms of the spontaneous symmetry breaking (SSB). SSB in $\Phi(3, 2)$ is driven by $\Phi(3, 1)$, while SSB in $\Phi(1, 2)$ from the driven SSB by $\Phi(3, 2)$, as well. The different, but related, scalar fields can accomplish so much, to our surprise.

There is more reason that we could be optimistic. Besides the $SU_c(3)$ protection over the quark world, we now have another $SU_f(3)$ to protect the lepton world - the remaining part of the world in the Standard Model. The (conceptual) troubling QED Landau ghost is no longer there since QED alone is only part of the story; it is all asymptotically free, via $SU(3)$, and free of the ghosts. Maybe everything can be put together elegantly as a final complete and consistent theory.

On the experimental side, the family Higgs η'_1 , $\eta'_{2,3}$, and $\phi_{1,2,3}^+$ couple to only cross-generation leptons - making them difficult to observe. On the other hand, the purely family Higgs η' is very elusive - maybe hopeless to observe. It is an ideal candidate particle for the so-called dark matter. These all happen in the lepton world, if the Standard Model (of this paper) is valid.

In the (theoretical) processes of figuring out how to write the Standard Model, or what is the best way to write it down, which would be the first to spell out (i.e. the "background"), etc., it came out with "a very precise definition of the Standard Model" [1] and then the word "built-in from the outset" [1]. It is important to figure out these steps in detail - it should eventually let us understand the Nature.

Half a year before T.-M. Yan and I wrote the paper [5] to put six objects in the multiplet $\Psi_L(3, 2)$, I was rather reluctant to do so even though I did have some important breakthroughs [13] that eventually drove me till the end. One should remember that the processes of searching for the Standard Model is non-orthodox, varying from a physicist to another. To be honest, we are not sure that a specific Standard Model would survive, or would survive for how long. It is just amazing that the entire body of particle physics, or the Nature, could be summarized by the Standard Model - the description of the *smallest* units of matter, such as electrons, neutrinos, and quarks, on the basis of Einstein's relativity principle and the quantum principle.

7 The Statement of the Standard Model

To describe the motion of matter, we have to add the knowledge of the velocities on top of the space-time coordinates, according to Newton's doctrine. The gauge principle says $\partial_\mu \rightarrow D_\mu$ with D_μ containing the gauge fields. Thus, the information on the gauge fields should be given, *a priori*, the knowledge of the space-time coordinates.

To be precise, thus, we could state the Standard Model as follows:

"We should declare that we are living in the quantum 4-dimensional Minkowski space-time with, via the gauge principle, the force-fields gauge-group structure $SU_c(3) \times SU_L(2) \times U(1)$ (for the quark world) or $SU_L(2) \times U(1) \times SU_f(3)$ (for the lepton world) built-in from the very beginning. The input entries are members of the quark group $SU_Q(3)$, the so-called "the triple quark world" and each of them enters in a way both dimensionless and massless, of nuclear sizes and of the $SU_c(3) \times SU_L(2) \times U(1)$ gauge symmetry. Meanwhile, the lepton

world is dimensionless and massless, of atomic sizes and of the $SU_L(2) \times U(1) \times SU_f(3)$ gauge symmetry.”

The (123) gauge symmetry, the symmetry under $SU_c(3) \times SU_L(2) \times U(1)$, means that there exist $SU_Q(3)$ quark triplets, $\Psi_L(3, 2, 1)$, $\Psi_R^u(3, 1, 1)$, and $\Psi_R^d(3, 1, 1)$, as the Dirac entry point. The other (123) gauge symmetry, the symmetry under $SU_L(2) \times U(1) \times SU_f(3)$, means that there exist $SU_Q(3)$ lepton singlets, $\Psi_L(1, 2, 3)$, $\Psi_R(1, 1, 3)$, and $\Psi_R^C(1, 1, 3)$, as another Dirac entry point. Here the three numbers refer to the assignments under the gauge groups: the color group, the $SU_L(2)$, and the $SU_f(3)$. Usually, we do not mention the assignment under the color group, for the sake of simplicity.

Mathematically, the Standard Model [1] is a group theory - a group of elements, triplets and singlets, of the quark group $SU_Q(3)$; they interact among themselves through the symmetry of the gauge group, either $SU_c(3) \times SU_L(2) \times U(1)$ or $SU_L(2) \times U(1) \times SU_f(3)$; they play out on the background of the quantum 4-dimensional Minkowski space-time.

We remark that the meaning of the so-called ”generations” disappears altogether in this (new) Standard Model. The input of the quark group $SU_Q(3)$ and the introduction of the gauge symmetry $SU_f(3)$ kill the old notion of the ”generations”.

Even though both the triple quark world and the lepton world are dimensionless, the length scales, $10^{-13}cm$ versus $10^{-8}cm$, still have the vastly difference - one mystery of the quantum 4-dimensional Minkowski space-time.

First, we should remark on the gauge principle, of which the meaning could be rather obscure previously but, now, may be clarified. The substitution, $\partial_\mu \rightarrow D_\mu$, for each smallest unit of matter is required when the space-time coordinates x_μ move to a neighboring point. Thus, it has to be built-in from the very beginning. *The gauge principle is a fundamental mathematical principle.*

Second, renormalizability should be promoted to ”the principle of renormalizability”. Not just we should include $\lambda(\phi^\dagger\phi)^2$ for the complex scalar field $\phi(x)$, but also *all* other renormalizable terms, including the interactions among different fields, should be included. Again, the principle of renormalizability should be meaningful in the sense of mathematics.

Thirdly, the structure of the Standard Model [1] should be noted: Apart from the ”ignition” term, all the couplings are dimensionless in the 4-dimensional Minkowski space-time. *It is a dimensionless theory in the quantum 4-dimensional Minkowski space-time, if we could ignore the ”ignition” term.* It is a beautiful theory.

Lastly, the previous treatments of ultraviolet divergences all were based on the massive theories, resulting in divergences in $\ln(\frac{M^2}{m^2})$ or in $(\frac{M^2}{m^2})^n$, etc. [See, for example, the references [1, 6] for further discussions.] But the Standard Model is a massless theory apart from the *SSB* ”ignition” term. Maybe this opens the door to solve this century-old puzzle.

We should challenge our young colleagues in carrying out systematic calculations of these ultraviolet divergences, as we don’t feel easy toward why a beautiful theory (the Standard Model) would contain these diseases - maybe they are not there, or they are there for good reasons. The physicists of the 20th Century got beaten by these ultraviolet divergences, but those of the 21st Century may write a different chapter of the history.

8 Family Interactions for Leptons

The weak interactions involving leptons are determined by $-\bar{\Psi}(3, 2)\gamma_\mu D_\mu \Psi(3, 2)$ [with D_μ given by Eq. (5)] plus two curl terms [Eqs. (6) and (7)].

For the basic processes such as the muon decay, $\mu^- \rightarrow e^- + \bar{\nu}_e + \nu_\mu$, we may write, symbolically, the transition amplitude [6]:

$$iT = \frac{G_F}{\sqrt{2}} \bar{u}_e(p') \gamma_\lambda (1 + \gamma_5) v_e(k') \cdot \bar{u}_{\nu_\mu}(k) \gamma_\lambda (1 + \gamma_5) u_\mu(p) + \text{others}, \quad (21)$$

but this might be incorrect, since $u(p)$, $v(k)$, etc. are, by definition, on the mass shells. Neutrino oscillations tell us $u_{\nu_\mu}(k) \equiv U_{\mu i} u_i(k)$, with the left-hand side defined by " \equiv "; similarly for the antineutrino $v_e(k)$.

In fact, our language here is only for the mass-shell Dirac spinors, not for something which oscillates. So, we should write $\sum_i U_{ei} u^i(k)$ for the electron-like neutrino, etc., since $u^i(k)$'s are the mass-eigenstates - that is how we set up the Dirac equations for.

Similarly, for the muon or the electron, they should be on mass shells in our language - that is the way which we represent the muon or the electron. So, this seems against the off-diagonal mass terms - in fact, this requires the overall consistency check at least.

Thus, we should write, for the muon decay,

$$iT = \frac{G_F}{\sqrt{2}} \bar{u}_e(p') \gamma_\lambda (1 + \gamma_5) U_{ei} v_i(k') \cdot \bar{u}_j(k) U_{\mu j}^\dagger \gamma_\lambda (1 + \gamma_5) u_\mu(p) + \frac{G'}{\sqrt{2}} \bar{u}_e(p') (1 - \gamma_5) \times U_{\mu j} v_j(k') \cdot \bar{u}_i(k) U_{ei}^\dagger (1 - \gamma_5) \times u_\mu(p). \quad (22)$$

Here the second term is due to the charged family-Higgs exchange ϕ_1 . (Our ϕ_1 or η'_1 refers to the τ channel, by our convention.) So, G' [$\propto (h^C)^2/m(\phi^\dagger)^2$] is much smaller than the Fermi coupling G_F . The G' term arises from the product of the two cross-dot products; the Fierz reordering could be used in this context.

The important point is that all the Dirac spinors are on the mass shells - in order that the expression can be further calculated.

The differential cross sections, or the decay rates, can easily be calculated: The inference term between the dominant muon-decay amplitude and the family-Higgs exchange term is relatively tiny since the neutrino mass enters the numerator. The family-Higgs exchange term, when squared, may be detectable if h (or κ , essentially) is not too small - here the mass of η'_1 is 107 GeV in the above numerical example. In other words, the interference term is suppressed by the neutrino-mass effect while the square of the family-Higgs exchange term could have some observable effects. On a second thought, we should classify the experimental efforts to search for effects caused by the family Higgs in ordinary muon decays as "the precision experiments".

The other important example is the $\mu \rightarrow e$ conversions, say, $\mu^-(p) + p(q) \rightarrow e^-(p') + p(q')$. The amplitude is given by

$$iT = \frac{1}{(2\pi^4)} \int d^4 k \bar{u}_e(p') \cdot i \frac{1}{2\sqrt{2}} \frac{e}{\sin\theta_W} \cdot i \gamma_\lambda (1 + \gamma_5) \cdot \frac{1}{i} \sum_j U_{ej}^\dagger \frac{m_j - i\gamma \cdot k}{m_j^2 + k^2 - i\epsilon} \cdot i \cdot \frac{i}{2} h \cdot (-) \cdot \frac{1}{2} (1 - \gamma_5) \cdot u_i \frac{1}{\sqrt{2}} \cdot \frac{1}{i} \sum_l \frac{m_l - i\gamma \cdot k}{m_l^2 + k^2 - i\epsilon} U_{\mu l} \cdot i \frac{1}{2\sqrt{2}} \frac{e}{\sin\theta_W} \cdot 1 \gamma_\eta (1 + \gamma_5) \cdot u_\mu(p)$$

$$\begin{aligned} & \cdot \bar{u}(q') \cdot i \frac{1}{2\sqrt{2}} \frac{e}{\sin\theta_W} \cdot i\gamma_\eta(1 + \gamma_5) \cdot \frac{1}{i} \frac{m - i\gamma \cdot (q+k-p')}{m^2 + (q+k-p')^2 - i\epsilon} \\ & \cdot i \frac{1}{2\sqrt{2}} \frac{e}{\sin\theta_W} \cdot i\gamma_\lambda(1 + \gamma_5) \cdot u(q) \cdot \frac{1}{i} \frac{1}{m_W^2 + (k-p')^2 - i\epsilon} \cdot \frac{1}{i} \frac{1}{m_W^2 + (k+p)^2 - i\epsilon}, \end{aligned} \quad (23)$$

in the 't Hooft-Feynman gauge. This means that the amplitude is finite but not complete, but, as seen below, all are rather small.

Simplifying it further, we obtain

$$\begin{aligned} iT = & \left(\frac{1}{2\sqrt{2}} \frac{e}{\sin\theta_W} \right)^4 \cdot i(-) \frac{hu_1}{2\sqrt{2}} \cdot (-i \cdot 2)^2 \\ & \cdot \frac{1}{(2\pi)^4} \int d^4k \bar{u}_e(p') \gamma_\lambda \sum_{j,l} U_{ej}^\dagger \gamma \cdot km_l U_{\mu l} \gamma_\eta (1 + \gamma_5) u_\mu(p) \\ & \cdot \bar{u}(q') \gamma_\eta \gamma \cdot (q+k-p') \gamma_\lambda (1 + \gamma_5) u(q) \\ & \cdot \frac{1}{m_j^2 + k^2 - i\epsilon} \cdot \frac{1}{m_l^2 + k^2 - i\epsilon} \cdot \frac{1}{m^2 + (q+k-p')^2 - i\epsilon} \cdot \frac{1}{m_W^2 + (k-p')^2 - i\epsilon} \cdot \frac{1}{m_W^2 + (k+p)^2 - i\epsilon}. \end{aligned} \quad (24)$$

Comparing this amplitude to the dominant muon-decay amplitude, it is down by $u_1 m_l$ with the tiny neutrino mass m_l (and $u_1 = 107 \text{ GeV}$ in the above table) - about 10^{-9} GeV^2 . The remaining reduction is from the 2nd-order weak interaction - a factor of 10^{-5} . So, normally, we expect 10^{-28} smaller than the dominant modes.

We have discussed the $\mu \rightarrow e + \gamma$ before [5]. The decay rate is suppressed by the neutrino mass effect $u_1 m_l$ and further by gauge invariance (for a real photon). The mode turns out to be smaller than the current search limits by more than 20th order away (completely negligible).

To sum up, the modifications in the ordinary muon decays may be observed - it defines a new category of the high-precision experiments. The $\mu \rightarrow e$ conversions such as $\mu^- + p \rightarrow e^- + p$ and $\mu \rightarrow e + \gamma$ turn to be hopelessly small.

Nevertheless, in certain decays such as ${}^3H \rightarrow {}^3He + e^- + \bar{\nu}_e$, we have to rewrite the neutrino state as a sum of neutrino-mass eigen-states - as required by neutrino oscillations. These are another category of high-precision experiments.

In a short summary, we do expect to see some corrections in the ordinary muon decay $\mu^- \rightarrow e^- + \bar{\nu}_e + \nu_\mu$ but the branching ratios induced by the $\nu_\mu \rightarrow \nu_e$ conversion or vice versa, such as $p + \mu^- \rightarrow p + e^-$ or $\mu \rightarrow e + \gamma$, turn out to be rather small. In the muon decay or nuclear beta decays, such as ${}^3H \rightarrow {}^3He + e^- + \bar{\nu}_e$, the replacement, as required by neutrino oscillations, of flavor states by the mass eigen-states does produce some tiny observable effects which might be detected in the precision experiments of the next generation.

Besides the three Higgs, the primary prediction of our Standard Model is the existence of the force of a new kind - i.e., the family force mediated by the family gauge bosons. As said above, we could use $\frac{1}{2}\kappa w$ as an estimate of the mass(es) of the family gauge bosons. Our first guess is for some feeble force - $\kappa = 0.1$. The above numerical example corresponds to $w = 102 \text{ GeV}$, so as to the family gauge boson mass of 5 GeV .

The family gauge bosons would then be in the vicinity of 5 GeV or nearby, or the range of 0.04 fermi . Or, $0.04 \times 10^{-13} \text{ cm}$ in the effective range, between leptons (such as two electrons or an electron-positron pair) is too short to be detected for the entire atomic physics or the entire chemistry.

The precision experiments such as $g - 2$ would eventually detect the residual family effects, since the existing $g - 2$ calculations [14] is so far the QED calculation and should

be completed by inclusion of other effects with the emphasis on family gauge bosons. We are looking forward to prospects in this direction.

Of course, we need to examine the precision part of atomic physics when the story becomes clear; even though the effects are tiny, the evolutions usually come from the tiny effects to begin with.

9 Experimental Questions yet to be Answered

We are talking about the Standard Model [1] in that there is a (new) force, conducted by the family gauge bosons, that there are (new) family Higgs particles, and that the existing smallest units of matter, including electrons, neutrinos, and quarks, all of them are forming the triplets and the singlets of the quark group $SU_Q(3)$ to begin with. Mathematically, it is a group theory - a group theory of the Nature.

Surprisingly, the Standard Model [8, 1] is just a group theory, mathematically. The Standard Model is used to describe our Universe. So, our Universe can be described by, mathematically, a group theory. This recognition of our Universe being described by a group theory, mathematically, gives a lot of comforts to minds of philosophers, or of mathematicians, or of theoretical physicists.

This hypothetical fact should set the stage for testing the complicated theory from the experimental ends.

There are no longer the three "generations" of leptons, since they need something to make up entities of $SU_f(3)$. There are no more the three "generations" of quarks, since the basic entry entities are the triplets of the quark group $SU_Q(3)$. The CKM matrix [7] is naturally borne there with the triplets of $SU_Q(3)$. The misnomer of "three generations" comes from lack of the understandings in our previous knowledge.

The beauty of the Standard Model [1] should prompt the experimental physicists of the next generations, of many centuries onward - trying to test whether this is indeed the "final" truth. As a theorist, it is impossible to embrace, with the great joy, the "final" truth without the deep appreciation of its perfectness.

10 Concluding Remarks

We are suggesting [1] that we live in the quantum 4-dimensional Minkowski space-time with, *via the gauge principle*, the force-fields gauge-group structure $SU_c(3) \times SU_L(2) \times U(1)$ (at the fermi scale for quarks), or $SU_L(2) \times U(1) \times SU_f(3)$ (at the atomic scale for leptons), built-in from the very beginning. The quark world, with the entries from the triplets of the quark group $SU_Q(3)$, possesses the (123) gauge symmetry and can be seen, at the fermi scale, by this overall background. Meanwhile, the lepton world, with the entries from the singlets of the quark group $SU_Q(3)$, possesses another (123) gauge symmetry and can also be seen, at the atomic scale, by this overall background.

We are using Einstein's relativity principle and the quantum principle, established in the 20th Century, to describe the *smallest* units of matter, such as electrons, neutrinos, and quarks. It looks like an ultimate theory at its birth, the so-called "Standard Model" [1],

which we believe will eventually replace the thinkings during the Newton's classic era (of the past four hundred years).

Previously we underestimated the basic importance of the gauge principle and the fundamental importance of the matter group, or the quark group $SU_Q(3)$, in which the quark world are triplets and the lepton world are singlets. Once the gauge principle and the matter group, both mathematical stuffs, are taken into account, the theory, or the Standard Model, is a consistent and perhaps complete theory. The concept of "generations" should not be there and thus it is simply a misnomer.

In the quantum 4-dimensional Minkowski space-time, the complex scalar field $\phi(x)$ is described renormalizably by

$$\mathcal{L} = -(\partial_\mu \phi)^\dagger (\partial_\mu \phi) - M^2 \phi^\dagger \phi - \lambda (\phi^\dagger \phi)^2. \quad (25)$$

If $\lambda < 0$, the system collapses (i.e. unbounded from below). If $\lambda > 0$, it is repulsive so that the system cannot build up by itself. The interesting question is that λ is dimensionless - a pure number that characterizes the 4-dimensional Minkowski space-time (maybe $\lambda = \frac{1}{8}$, but we need the proof), *not* by the complex scalar fields.

The force fields are described by the gauge fields in either the group $SU_c(3) \times SU_L(2) \times U(1)$ (for quarks), or the group $SU_L(2) \times U(1) \times SU_f(3)$ (for leptons), *via the gauge principle* - a mathematical principle (for the completion of the description of motion). They need complex scalar fields to generate the masses of the gauge bosons via generalized Higgs mechanisms. The longitudinal components are missing in the purely gauge-fields description such that complex scalar fields are needed for the Higgs mechanisms. Thus, in the quantum 4-dimensional Minkowski space-time with, *via the gauge principle*, the force-fields gauge group $SU_c(3) \times SU_L(2) \times U(1)$ (for quarks), or in the gauge group $SU_L(2) \times U(1) \times SU_f(3)$ (for leptons), the world is already set up in terms of the force fields from the gauge fields and the complex scalar (Higgs) fields via the specific Higgs mechanisms, and this background world is called the "overall background".

It yields, and only yields, three Higgs fields $\Phi(1,2)$, $\Phi(3,2)$, and $\Phi(3,1)$. The "related" Higgs fields, such as $\Phi(1,2)$ and $\Phi(3,2)$ with $SU_L(2)$ doublets in common (or, $\Phi(3,2)$ and $\Phi(3,1)$ with $SU_f(3)$ triplets in common), can overcome the "self-repulsive" nature and become useful and be able to live and to sustain in this world.

On the lepton world, we know that they couple to the $SU_L(2) \times U(1)$ gauge sector (which makes them visible). To interpret the ordering via three "generations", we proposed the force-fields nature of the $SU_f(3)$ gauge symmetry. Neutrino oscillations provide a direct proof that "generations" can switch among themselves. By introducing the $SU_f(3)$ gauge symmetry to the original $SU_L(2) \times U(1)$, the lepton world is free from the Landau ghost and is asymptotically free. So, the lepton world is perfectly well-behaved.

On the quark world, it is already a perfect world since it couples to the $SU_c(3) \times SU_L(2) \times U(1)$ (i.e. the standard (123)) gauge symmetry. It exhibits the "size" effect, i.e., that the quark world exists only within a given volume of fermi sizes; or, it also exhibits the temperature effect, i.e., that it undergoes the phase transition (into something else). The entry inputs are triplet members of the "quark" group $SU_Q(3)$ - the entity with three "generations" altogether serves one entry input.

Thus, in this Standard Model, the old notion of "generations" disappears altogether. The family $SU_f(3)$ gauge symmetry and the presence of the $SU_Q(3)$ symmetry both kills the

notion of "three generations".

To sum up, we live in the quantum 4-dimensional Minkowski space-time with, *via the gauge principle*, the force-fields gauge group structure, $SU_c(3) \times SU_L(2) \times U(1)$ (for quarks) or $SU_L(2) \times U(1) \times SU_f(3)$ (for leptons), built-in from the very beginning [1]. Apart from the *SSB* "ignition" term, the overall theory is both dimensionless and massless [11]. Neutrino oscillations are there.

Appendix: Sideline Remarks

This world is, indeed, very special. It is based on the quantum 4-dimensional Minkowski space-time with, *via the gauge principle*, the force-fields gauge-group structure $SU_c(3) \times SU_L(2) \times U(1)$, or $SU_L(2) \times U(1) \times SU_f(3)$, built-in from the very beginning. We realize that the complex scalar field is self-repulsive (i.e. does not exist) if alone and that two "related" complex scalar fields could interact attractively (and so exist) and they become the very-much-wanted longitudinal components of the gauge fields. The quark world, operating under the quark group $SU_Q(3)$ (not a gauge symmetry), would be accepted because of the (123) gauge symmetry. Meanwhile, the lepton world could be accepted in view of another (123) symmetry.

In this language, everything carries the group characteristic. For example, $\nu_{e,L}$ belongs to the same multiplet as e_L , so the same characteristic - thus, the case would be ruled out that the electron is a Dirac particle while the neutrino is a Majorana particle.

A few years ago [15], it was proposed that we could work with two working rules: "Dirac similarity principle", based on ninety years of experience, and "minimum Higgs hypothesis", from the last fifty years of experience. Using these two working rules, the basic model [8] became rather unique in our choice - so, it is so much easier to check it against the experiments. To move forward in building up our knowledge, there are moments that we have to play conservatively - including the moments of using these two working rules.

At this point, the two working rules seem to be rather trivial.

We have to say that the phenomenon of three generations used to be one leading puzzle in particle physics; it is safe to add that neutrino oscillations used to be another related puzzle. When we write everything together, the need for introducing the Higgs $\Phi(3,2)$ becomes rather clear [5]. As this world does not have another massless gauge bosons (unless confined like gluons), it is also clear that there is another pure family Higgs $\Phi(3,1)$ to complete the story [9]. Is there any other possibility? It seems that it is unique.

We have to admit that the natural consequence is the $SU_f(3)$ family gauge theory - to put everything together, it leads to the (extended) Standard Model [8], with a very natural generalized Higgs mechanism for the three Higgs. It is in accord with the "minimum Higgs hypothesis". These complex scalar fields are meant together and interact together, and nothing more.

It is also of importance to point out that there is no experimental evidence to the assertion that neutrinos are point-like Dirac particles - "Dirac similarity principle" should be in experimental checks. Of course, they are in the same multiplet, like in [5], so that they both are Dirac particles of the same nature. So, the two working principles are justified by

our experiences for ninety or forty years. Surprising enough, it seems to work this way, at least so far.

Thus, we may declare that our world is the quantum 4-dimensional Minkowski space-time with, *via the gauge principle*, the force-fields gauge-group structure $SU_c(3) \times SU_L(2) \times U(1)$, or $SU_L(2) \times U(1) \times SU_f(3)$, built-in at the very beginning; in our world, the various Higgs "exist" such that all gauge bosons, except the photon, are either confined or massive - this provides the overall background to support the "triple" quark world as well as to support the lepton world. This is the origin of "point-like particles", or of "smallest particles", or of "fields".

To close up our discussions, there remain a few philosophical questions in our minds: First, why are the complex scalar fields so special in the 4-dimensional Minkowski space-time? The 4-dimensional Minkowski space-time (not the complex scalar fields themselves) would be the reason to determine the value of λ ($= \frac{1}{8}$?). In the other dimensions (different from four), the existence of the self-repulsive interaction $\lambda(\phi^\dagger\phi)^2$ would not have the same status. Second, why do we need the ignition channel, which turns out to be in the purely family channel ($\mu_2^2 < 0$)? Third, the dimensional regularization, or others, might not give whole story regarding the (leading) ultraviolet divergences. In the beginning of this 21th century, we might have this ghost (infinite or divergence) story to resurface again [16, 1]. But this might be destined to be so, as our knowledge accumulates in the process.

We may add that, under two working hypotheses or under our Standard Model (in the quantum 4-dimensional Minkowski space-time with the $SU_c(3) \times SU_L(2) \times U(1)$, or $SU_L(2) \times U(1) \times SU_f(3)$, gauge-group structure built in from the very beginning), we should be able to close the Universe; that is, all the dark-matter particles and all the ordinary-matter particles are accounted for. Our Standard Model provides a description of the entire matter world - i.e., the 25% dark-matter world and the 5% ordinary-matter world.

In our language, the *a priori* world (i.e., the overall background) is the quantum 4-dimensional Minkowski space-time with some force-fields gauge-group structure built-in from the very beginning. The triple quark world and the lepton world are something added on at a later stage (at the different scales), and in fact they may be the added-on at a different stage and at the fermi or atomic scales - that makes the study of leptons a subject by itself [17].

We would be curious about how the dark-matter world looks like, though it is difficult to verify experimentally. The first question would be: The dark-matter world, 25 % of the current Universe (in comparison, only 5 % in the ordinary matter), would clusterize to form the dark-matter galaxies? The dark-matter galaxies would then play the hosts of (visible) ordinary-matter galaxies, like our own galaxy, the Milky Way. Note that a dark-matter galaxy is by our definition a galaxy that does not possess any ordinary strong and electromagnetic interactions (with our visible ordinary-matter world). These fundamental question deserves some in-depth thoughts, for the evolution of our Universe.

The situation may be relatively simple, if we look closely at our Standard Model [1]. All the dark-matter candidates, except neutrinos and antineutrinos, will decay away with a lifetime shorter than a fraction of a sub-second [18]. Since neutrinos have masses, heaviest of 0.058 eV , the cosmic background neutrinos (CB ν 's) would cluster around the visual ordinary-matter centers. We believe that the CB ν 's account for all of the 25% dark matter.

Of course, we should remind ourselves that, in our ordinary-matter world, those quarks

can aggregate in no time, to hadrons, including nuclei, and the electrons serve to neutralize the charges also in no time. Then atoms, molecules, complex molecules, and so on. These serve as the seeds for the clusters, and then stars, and then galaxies, maybe in a time span of 1 Gyr (i.e., the age of our young Universe). The aggregation caused by strong and electromagnetic forces is fast enough to help giving rise to galaxies in a time span of 1 Gyr . *We believe that all these took place in the invisible environments of neutrino halos.*

In other words, the neutrino halos, the 25% dark matter, play the invisible roles of our Universe. They are there, very light but definitely massive, could not have the very heavy invisible centers, and use the visible heavy centers (such as planets, stars, etc.) as their centers. This would modify the Newton's gravitational law, a macroscopic law, with the invisible parts. The invisible stories may sound scary, but after certain clarifications everything goes its own way except that there is some invisible partner(s).

References

- [1] W-Y. Pauchy Hwang, The Universe, **4-4**, 7 (2016). "A Precise Definition of the Standard Model".
- [2] T. Yanadida, Phys. Rev. **D20**, 2986 (1979).
- [3] Yue-Liang Wu, Phys. Lett. **B 714**, 286 (2012).
- [4] W-Y. Pauchy Hwang, The Universe, **3-1**, 3 (2015).
- [5] W-Y. Pauchy Hwang and Tung-Mow Yan, The Universe, **1-1**, 5 (2013).
- [6] Ta-You Wu and W-Y. Pauchy Hwang, "Relativistic Quantum Mechanics and Quantum Fields" (World Scientific, 1991); W-Y. Pauchy Hwang and Ta-You Wu, *ibid.*, the 21st-Century Edition (World Scientific, 2017).
- [7] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. Japan, **49**, 652 (1973).
- [8] W-Y. Pauchy Hwang, arXiv:1304.4705v1 [hep-ph] 17 April 2013. This is the early version of this paper.
- [9] W-Y. Pauchy Hwang, Nucl. Phys. **A844**, 40c (2010); *ibid.*, International J. Mod. Phys. **A24**, 3366 (2009); *ibid.*, Intern. J. Mod. Phys. Conf. Series **1**, 5 (2011); *ibid.*, American Institute of Physics 978-0-7354-0687-2/09, pp. 25-30 (2009).
- [10] W-Y. P. Hwang, Phys. Rev. **D32**, 824 (1985).
- [11] W-Y. Pauchy Hwang, The Universe, **2-2**, 47 (2014). "The Origin of Mass".
- [12] Particle Data Group, "Review of Particle Physics", J. Phys. G: Nucl. Part. Phys. **37**, 1 (2010); and its biennial publications.
- [13] W-Y. Pauchy Hwang, arXiv:1207.6443v1 [hep-ph] 27 Jul 2012; *ibid.*, Hypefine Interactions **215**, 105 (2013); *ibid.*, arXiv:1209.5488v1 [hep-ph] 25 Sep 2012.

- [14] T. Kinoshita and W.B. Lindquist, Phys. Rev. Lett. **47**, 1573 (1981).
- [15] W-Y. P. Hwang, arXiv:11070156v1 [hep-ph] 1 Jul 2011; Plenary talk given at the 10th International Conference on Low Energy Antiproton Physics (Vancouver, Canada, April 27 - May 1, 2011).
- [16] W-Y. Pauchy Hwang, The Universe, **2-1**, 41 (2014).
- [17] W-Y. Pauchy Hwang, arXiv:1409.6296v1 [hep-ph] 22 Sep 2014. "The Family Collider".
- [18] W-Y. Pauchy Hwang, arXiv:1012.1082v6 [hep-ph] 13 Jan 2016.