

Two radiative inverse seesaw models, dark matter, and baryogenesis.

Iason Baldes,^a Nicole F. Bell,^a Kalliopi Petraki,^{a,b} and Raymond R. Volkas.^a

^aARC Centre of Excellence for Particle Physics at the Terascale,
School of Physics, The University of Melbourne, Victoria 3010, Australia

^bNikhef, Science Park 105, 1098 XG Amsterdam, The Netherlands

E-mail: i.baldes@student.unimelb.edu.au, n.bell@unimelb.edu.au,
kpetraki@nikhef.nl, raymondv@unimelb.edu.au

Abstract. The inverse seesaw mechanism allows the neutrino masses to be generated by new physics at an experimentally accessible scale, even with $\mathcal{O}(1)$ Yukawa couplings. In the inverse seesaw scenario, the smallness of neutrino masses is linked to the smallness of a lepton number violating parameter. This parameter may arise radiatively. In this paper, we study the cosmological implications of two contrasting radiative inverse seesaw models, one due to Ma and the other to Law and McDonald. The former features spontaneous, the latter explicit lepton number violation. First, we examine the effect of the lepton-number violating interactions introduced in these models on the baryon asymmetry of the universe. We investigate under what conditions a pre-existing baryon asymmetry does not get washed out. While both models allow a baryon asymmetry to survive only once the temperature has dropped below the mass of their heaviest fields, the Ma model can create the baryon asymmetry through resonant leptogenesis. Then we investigate the viability of the dark matter candidates arising within these models, and explore the prospects for direct detection. We find that the Law/McDonald model allows a simple dark matter scenario similar to the Higgs portal, while in the Ma model the simplest cold dark matter scenario would tend to overclose the universe.

Keywords: dark matter theory, baryon asymmetry, leptogenesis, neutrino properties

Contents

1	Introduction	1
2	The inverse seesaw mechanism	2
2.1	The generic mass matrix	2
2.2	ISS and baryogenesis	3
3	Law/McDonald radiative inverse seesaw	5
3.1	Review of the model	5
3.2	Constraints from BAU washout	7
3.3	Dark Matter	9
3.3.1	Double Higgs portal DM	10
3.3.2	DM relic abundance	12
3.3.3	Direct detection	14
3.3.4	Higgs signals at colliders	15
3.3.5	Fermionic DM	16
3.4	Concluding remarks on the Law/McDonald model	16
4	Ma radiative inverse seesaw model	17
4.1	Review of the model	17
4.2	Constraints from BAU washout	19
4.3	Resonant leptogenesis	22
4.4	The ρ parameter	22
4.5	Dark matter	23
4.5.1	DM candidate mass	23
4.5.2	Cold DM scenario	24
4.6	Concluding remarks on the Ma model	25
5	Conclusion	25

1 Introduction

The experimental evidence for physics beyond the standard model (SM) includes the baryon asymmetry of the universe (BAU), the existence of dark matter (DM), and the non-zero neutrino masses. It is possible that the mechanisms responsible for these observations are intrinsically linked. For example, the observed neutrino masses may arise from new interactions which at low energies generate the Weinberg operator [1], $\mathcal{L}_W = (1/\Lambda) \Phi \Phi_L l_L$, where Φ is the SM Higgs, l_L are the SM lepton doublets and Λ is the scale of new physics. $\mathcal{L}_W$ is the lowest-order (non-renormalisable) operator containing only SM fields which can give rise to neutrino masses. Moreover, it violates the lepton-number (L) symmetry of the SM, thus suggesting that the interactions which give rise to it may have also generated the BAU via leptogenesis at early times [2]. It is also plausible that the relic abundance of some of the degrees of freedom mediating $\mathcal{L}_W$ constitutes the DM of the universe today [3–9].

Another possibility is that the BAU was generated by a mechanism unrelated to the physics of neutrino masses. Well-motivated scenarios include the Affleck-Dine mechanism

in supersymmetric models [10, 11], and baryogenesis via a first-order phase transition [12]. Clearly, whatever mechanism of baryogenesis operated in the early universe, the existence of a baryon asymmetry today has to be consistent with the interactions which give rise to the neutrino masses. If the latter violate L , then they could pose a challenge for the BAU: L -violating interactions operating efficiently in the early universe, together with the rapid electroweak (EW) sphaleron processes [13, 14], which violate the baryon-plus-lepton number of the SM, can erase any baryon-number (B) asymmetry created before the EW phase transition. It is then important to understand what conditions the survival of a baryon asymmetry imposes on neutrino mass generation mechanisms.

The three minimal, tree level, UV completions of the Weinberg operator, the type-I [15–18], type-II [19–24], and type-III [25] seesaw mechanisms, have been studied extensively for their phenomenological consequences. In these scenarios, the scale of new physics ranges from ~ 100 GeV for Yukawa couplings of $\mathcal{O}(10^{-6})$, all the way up to $\sim 10^{14}$ GeV for Yukawa couplings of $\mathcal{O}(1)$. The small Yukawa couplings in the former case, and the very high energy scale in the latter case make these models difficult to test experimentally (although for the type-II and III seesaws gauge interactions allow the lower mass range to be tested at the LHC).

On the other hand, the inverse seesaw (ISS) mechanism [26, 27] can generate the observed neutrino masses via new physics at an experimentally accessible scale, and with $\mathcal{O}(1)$ Yukawa couplings. In the ISS, the smallness of neutrino masses is traced to the smallness of an L -violating parameter, which we shall denote by μ . This small lepton-number violation naturally suppresses the washout of a pre-existing baryon asymmetry [28]. To explain why μ is small, it is natural to suppose that it is generated radiatively [29–33]. (Other possibilities of explaining why μ is a *small* parameter are extra dimensions [34, 35], or a type-II seesaw-like mechanism [36, 37].)

Radiative ISS models introduce new fields and lepton-number violating interactions, which could potentially wash out the BAU. In this paper, we investigate the constraints on the parameters of the ISS implied by the BAU, or conversely, the conditions on baryogenesis assuming that neutrino masses are generated via the ISS. We consider two radiative ISS models: one in which lepton number is explicitly broken [32], and one in which it is spontaneously broken [29]. These models also feature accidental discrete symmetries which stabilise exotic particles. The latter can play the role of DM. We investigate the parameter space, derive conditions which ensure that the relic abundance of the new stable particles does not overclose the universe, and identify more specifically the regions which give rise to the observed DM abundance. Similar issues in a different implementation of the ISS have been studied in ref. [38].

The order of the paper is as follows. In section 2 we introduce the ISS, and review the natural suppression of washout in the ISS. In section 3 we introduce the Law/McDonald radiative ISS model [32], analyse washout with the additional fields present, and examine its DM candidates. In section 4 we repeat the analysis for the Ma radiative ISS model [29].

2 The inverse seesaw mechanism

2.1 The generic mass matrix

The ISS [26, 27] is generically realised with the additional terms in the Lagrangian

$$\mathcal{L} \supset \lambda_\nu \epsilon^{ab} \overline{l_{La}} \Phi_b^\dagger N_R + M_R \overline{N_R} S_L + \frac{1}{2} \mu \overline{(S_L)^c} (S_L) + \frac{1}{2} \tilde{\mu} \overline{(N_R)^c} (N_R) + H.c., \quad (2.1)$$

where $\Phi \sim (1, 2, 1/2)$ is the usual SM Higgs doublet, $l_L \sim (1, 2, -1/2)$ is the SM lepton doublet, and N_R and S_L are singlets under the SM gauge group. We leave the details of additional symmetries and symmetry breaking which results in the above terms [39] unspecified for now (a field redefinition ensures the absence of the $\bar{l}_L \Phi^\dagger (S_L)^c$ term). After Φ gains a vacuum expectation value (VEV) $\langle \Phi \rangle = (0, v_w/\sqrt{2})^T$ and breaks the EW symmetry $SU(2)_L \otimes U(1)_Y \rightarrow U(1)_{\text{EM}}$, the neutral-fermion mass matrix is (assuming one generation)

$$\mathcal{L} = \frac{1}{2} (\bar{\nu}_L \quad \bar{N}_R^c \quad \bar{S}_L) \begin{pmatrix} 0 & m_D & 0 \\ m_D & \tilde{\mu} & M_R \\ 0 & M_R & \mu \end{pmatrix} \begin{pmatrix} (\nu_L)^c \\ N_R \\ (S_L)^c \end{pmatrix}, \quad (2.2)$$

where $m_D \equiv \lambda_\nu v_w/\sqrt{2}$. Diagonalization of the mass matrix for $\mu, \tilde{\mu} \ll m_D, M_R$ leads to a light neutrino mass

$$m_\nu \approx \frac{m_D^2}{M_R^2} \mu. \quad (2.3)$$

The heavy Majorana neutrinos N_1 and N_2 , superpositions of mostly $(S_L)^c$ and N_R , have mass eigenvalues

$$M_N \approx M_R \pm \frac{\mu}{2}. \quad (2.4)$$

Due to the small mass splitting, they form a pseudo-Dirac pair. The mixing between the light and heavy states is approximately $\theta \simeq m_D/M_R$. In the limit $\mu \rightarrow 0$ the light neutrinos become massless at tree level, although loop corrections give finite contributions to their masses unless both $\mu, \tilde{\mu} \rightarrow 0$ [40]. In this limit one also recovers a global lepton number symmetry in the Lagrangian and hence small choices of μ , and $\tilde{\mu}$ are technically natural. This is the small parameter generated radiatively in the ISS models we return to below.

This discussion can be generalised to a realistic three generation case. The entries in the mass matrix (2.2), are promoted to 3×3 matrices, and the light neutrino mass matrix is then to first order,

$$M_\nu = M_D M_R^{-1} \mu (M_R^T)^{-1} M_D^T. \quad (2.5)$$

Various studies on the collider phenomenology [41, 42], flavour violating processes [43–46], and neutrinoless double beta decay [47, 48] of the ISS have appeared in the literature.

2.2 ISS and baryogenesis

Before the EW phase transition, the non-perturbative sphaleron processes rapidly wash out any existing $B+L$ asymmetry, while they preserve $B-L$. However, if there exist other rapid L violating processes, then both B and L will be driven to zero. In the ISS, the SM lepton number is violated by the parameters μ and $\tilde{\mu}$. Provided that these parameters are small enough, the B asymmetry is not washed out.

Let us examine the possible washout of a B asymmetry created at a temperature above the heavy neutrino mass scale, $T_{\text{BG}} > M_N$, where T_{BG} is the temperature of baryogenesis. In this case, the inverse decays to on-shell heavy Majorana neutrinos $l_L + \Phi \rightarrow N_i$ are not Boltzmann suppressed. Such inverse decays followed by decays $N_i \rightarrow \bar{l}_L + \Phi^*$ can change the lepton number by two units. This washout process is shown in figure 1.

It was found in ref. [28] that in the ISS such washout processes can be naturally suppressed, and we review the findings here. For simplicity we take from now on $\tilde{\mu} = 0$, as this parameter does not enter the neutrino mass eigenvalues to first order, and focus on μ




Figure 1. $\Delta L = 2$ process which can wash out the baryon asymmetry at high temperatures. Both heavy mass eigenstates N_1 and N_2 contribute, but destructive interference for small mass splittings $\mu \ll M_N$ crucially limits the washout rate.

instead. The more in-depth discussion of [28] allows both parameters to be non-zero with no major differences to the discussion here.

For typical mass choices the heavy neutrinos can decay into a component of the Higgs doublet and a lepton $N \rightarrow \phi + l_L$. For heavy neutrino masses well above the Higgs mass, $M_N \gg M_h$, the decay rate is

$$\Gamma_D = \frac{\lambda_\nu^2}{8\pi} M_N, \quad (2.6)$$

and for temperatures $T > M_N$ one includes a Lorentz factor (M_N/T). In the type I seesaw, the effectiveness of washout is well captured by the washout parameter [49]

$$K \equiv \frac{\Gamma_D}{H} \Big|_{T=M_N}, \quad (2.7)$$

where $H \sim g^{1/2} T^2 / M_{Pl}$ is the expansion rate of the universe, and g is the number of relativistic degrees of freedom. Typically washout processes are ineffective for $K \lesssim 1$, while any net B is erased for $K \gg 1$. In the intermediate regime $K \sim 1 - 100$ a more careful analysis is required. For our purposes however, we adopt the approximation that $K \lesssim 1$ is safe from washout.

In the ISS, however, the small mass splitting between the heavy pseudo-Dirac neutrinos leads to destructive interference in the scattering process depicted in figure 1. This leads to a suppressed washout rate [28]

$$K_{\text{eff}} = K \delta^2 \equiv \frac{\mu^2}{\Gamma_D H} \Big|_{T=M_N}, \quad (2.8)$$

where $\delta \equiv \mu / \Gamma_D$. As expected, in the limit of lepton number conservation, $\mu \rightarrow 0$, the heavy neutrinos form a Dirac pair, there is complete destructive interference in the $\Delta L = 2$ scattering process, and washout of the baryon asymmetry does not occur. Requiring $K_{\text{eff}} \lesssim 1$ to avoid washout translates into a bound:

$$\mu < \lambda_\nu \left(\frac{M_N}{1 \text{ TeV}} \right)^{3/2} \times 6 \text{ keV}. \quad (2.9)$$

For $\lambda_\nu = 0.1$, $M_N = 1 \text{ TeV}$, light neutrino masses $m_\nu = 0.1 \text{ eV}$ are explained by $\mu = 330 \text{ eV}$ (cf. eq. (2.3)). So the light neutrino masses can be explained while avoiding washout of the

BAU. Alternatively, one can express the above bound in terms of the mixing angle:

$$\theta \gtrsim 1.4 \times 10^{-2} \left(\frac{m_\nu}{0.1 \text{ eV}} \right)^{1/3} \left(\frac{1 \text{ TeV}}{M_N} \right)^{5/6} \left(\frac{100}{g} \right)^{1/6}. \quad (2.10)$$

To satisfy current experimental constraints (e.g. from universality of the weak interaction and rare leptonic decays) requires roughly, for M_N above the EW scale, a mixing angle $\theta \lesssim \mathcal{O}(10^{-2})$ [44, 50]. (The usually stringent constraint on the active-sterile mixing angle from neutrinoless double beta decay [51] is greatly weakened due to the small lepton number violation in the ISS [47, 48].) Future experimental tests at the $\theta \sim 10^{-3}$ level can therefore help test the parameter space corresponding to washout avoidance [52].

For baryogenesis temperatures below the heavy neutrino scale, $T_{\text{BG}} < M_N$, the number density of heavy states is Boltzmann suppressed. This means the thermally averaged rate of the net washout process $l_L + \Phi \rightarrow N_i \rightarrow \bar{l}_L + \Phi^*$ is suppressed by both a Boltzmann factor and the small μ . In this regime one is even safer from washout.

What about baryogenesis itself in the ISS? The obvious candidate is resonant leptogenesis [53]. The resonance itself is due to an almost degenerate pair of heavy neutrinos and exactly such pairs appear in the ISS. Detailed studies have shown resonant leptogenesis is indeed possible in the ISS framework [54], and it is even possible to link this with the DM abundance [55].

To explain the smallness of μ we now turn to models where it is generated radiatively. Because of the smallness of μ , washout of the BAU is suppressed in ISS models. However, to generate a finite value of μ radiatively requires the addition of new fields and L -violating interactions. We will see the effects of these new fields and interactions on washout of the BAU, and study the DM candidates in the radiative ISS models. We study the Law/McDonald [32] and Ma radiative ISS models in turn [29].

3 Law/McDonald radiative inverse seesaw

3.1 Review of the model

The Law/McDonald radiative ISS model [32] employs explicit violation of $(B - L)$, so that processes other than scattering through the heavy neutrinos crucially contribute to washout of the BAU. The model introduces an additional abelian gauge symmetry to the standard model gauge group

$$G = G_{\text{SM}} \otimes U(1)_d. \quad (3.1)$$

All the SM particles are neutral under this new charge. The following exotic fermions are introduced (with convention $Q_{\text{EM}} = I_3 + Y$):

$$E_{R,L} \sim (1, 3, -1)(0) \quad N_{R,L} \sim (1, 1, 0)(1). \quad (3.2)$$

Along with the SM Higgs $\Phi \sim (1, 2, 1/2)(0)$, the following scalars are introduced

$$\xi \sim (1, 3, -1)(-1), \quad \eta \sim (1, 1, 0)(1), \quad \chi \sim (1, 1, 0)(2). \quad (3.3)$$

The following Yukawa interactions and bare Dirac mass terms are allowed

$$\begin{aligned} -\mathcal{L} \supset & +y_E \bar{l}_L \Phi E_R + h_\xi \bar{E}_L \xi N_R + h'_\xi \bar{E}_R \xi N_L + h_\chi \bar{N}_R N_R^c + h'_\chi \bar{N}_L N_L^c \\ & + M_E \bar{E}_L E_R + M_N \bar{N}_L N_R + H.c. \end{aligned} \quad (3.4)$$

The VEV pattern is chosen to be

$$\langle \phi^0 \rangle = v_w / \sqrt{2} = 174 \text{ GeV}, \quad \langle \xi^0 \rangle, \langle \eta \rangle = 0, \quad \langle \chi \rangle = v_\chi / \sqrt{2} \neq 0. \quad (3.5)$$

The tree level mass matrix for the neutral fermions is then

$$\mathcal{L} = \frac{1}{2} \begin{pmatrix} \bar{\nu}_L & \overline{(E_R^0)^c} & \overline{E_L^0} \end{pmatrix} \begin{pmatrix} 0 & y_E v_w / \sqrt{2} & 0 \\ y_E v_w / \sqrt{2} & 0 & M_E \\ 0 & M_E & 0 \end{pmatrix} \begin{pmatrix} (\nu_L)^c \\ E_R^0 \\ (E_L^0)^c \end{pmatrix}. \quad (3.6)$$

Note $N_{L,R}$ are heavy Majorana fermions decoupled from the above matrix, and we have not yet taken into account the radiative corrections which will generate the terms $\tilde{\mu} \overline{(E_R^0)^c} E_R^0$ and $\mu \overline{(E_L^0)^c} E_L^0$ after symmetry breaking. The scalar potential is given by,

$$\begin{aligned} V_S = & (\Phi^\dagger \Phi) \left[\mu_\Phi^2 + \lambda_\Phi (\Phi^\dagger \Phi) + \lambda_{\phi\xi} (\xi^\dagger \xi) + \lambda_{\eta\phi} (\eta^\dagger \eta) + \lambda_{\phi\chi} (\chi^\dagger \chi) \right] \\ & + (\xi^\dagger \xi) \left[\mu_\xi^2 + \lambda_\xi (\xi^\dagger \xi) + \lambda_{\xi\eta} (\eta^\dagger \eta) + \lambda_{\xi\chi} (\chi^\dagger \chi) \right] \\ & + (\eta^\dagger \eta) \left[\mu_\eta^2 + \lambda_\eta (\eta^\dagger \eta) + \lambda_{\eta\chi} (\chi^\dagger \chi) \right] \\ & + (\chi^\dagger \chi) \left[\mu_\chi^2 + \lambda_\chi (\chi^\dagger \chi) \right] \\ & + \left[\lambda_{\xi\phi\eta} \xi \Phi \eta + \frac{1}{2} \mu_{\eta\chi} \eta \eta \chi^\dagger + \text{H.c.} \right]. \end{aligned} \quad (3.7)$$




Figure 2. Radiative mass generation in the Law/McDonald model.

The SM lepton number is violated explicitly by the combination of eq. (3.7) and eq. (3.4), but a Majorana mass term for the E_L and E_R fields is forbidden due to hypercharge. After χ and Φ gain VEVs, a radiative mass term $\mu \overline{(E_L^0)^c} E_L^0$ is generated, as shown in figure 2. A mass term $\tilde{\mu} \overline{(E_R^0)^c} E_R^0$ is also generated radiatively. The radiative mass is approximately

$$\mu \approx \frac{1}{16\pi^2} \frac{h_\xi^2 h_\chi \lambda_{\xi\phi\eta}^2 v_w^4 v_\chi^2}{120 \Lambda_{\text{ISS}}^6} \mu_{\eta\chi}, \quad (3.8)$$

where $\Lambda_{\text{ISS}} \sim M_N, M_\eta, M_\xi$ is the scale of the new physics, usually taken to be $\sim \mathcal{O}(\text{TeV})$, and one may choose $\mu_{\eta\chi} \sim \mathcal{O}(\text{TeV})$. This radiatively generated μ is exactly the small parameter required in the mass matrix for the ISS. The light neutrinos masses are then

$$m_\nu \approx \left(\frac{y_E^2 h_\xi^2 h_\chi \lambda_{\xi\phi\eta}^2}{10^{-5}} \right) \left(\frac{v_\chi}{M_E} \right)^2 \left(\frac{1 \text{ TeV}}{\Lambda_{\text{ISS}}} \right)^6 \left(\frac{\mu_{\eta\chi}}{1 \text{ TeV}} \right) \times 0.1 \text{ eV}. \quad (3.9)$$

3.2 Constraints from BAU washout

Previously we have seen that in the ISS framework, the washout of the BAU is naturally suppressed due to the smallness of the lepton-number violating parameter μ . In the Law/McDonald radiative ISS model, μ is in fact generated only after EW symmetry breaking and cannot itself cause washout. However, the SM lepton number is explicitly violated by the couplings which eventually generate μ and give rise to the radiative ISS. The interactions of eqs. (3.7) and (3.4) can then lead to washout. Here, we investigate under what conditions the BAU is not washed out.

The $(B-L)$ of the SM remains a good symmetry if any of the following sets of parameters vanishes

$$\{y_E\}, \{\mu_{\eta\chi}\}, \{\lambda_{\xi\phi\eta}\}, \{h_\xi, h'_\xi\}, \{h_\chi, h'_\chi\}. \quad (3.10)$$

(Note that M_E has to be large since $E_{R,L}$ are charged under the SM gauge group.) The BAU can change only if interactions which involve all of the above couplings are efficient. We show such a sequence of interactions in figure 3. In order to preserve the BAU, we require that the interactions mediated by at least one of these sets of couplings are out of equilibrium in the period between baryogenesis and the EW phase transition. We discern two cases, depending on the scale of baryogenesis: $T_{\text{BG}} > \Lambda_{\text{ISS}}$ and $T_{\text{EW}} < T_{\text{BG}} < \Lambda_{\text{ISS}}$.




Figure 3. An example of a sequence of interactions present in the early universe which shifts the lepton number.

- $T_{\text{BG}} > \Lambda_{\text{ISS}}$

- The Yukawa couplings of eq. (3.4) induce 2-body decays of the heaviest particle participating in each operator, e.g. $N_R \rightarrow E_L + \xi^*$. Assuming these decays are kinematically allowed, they proceed with thermally averaged decay rate

$$\Gamma_D \approx \frac{h_j^2 M_j}{16\pi} \left(\frac{M_j}{T} \right), \quad (3.11)$$

where $M_j \sim \Lambda_{\text{ISS}}$ is the mass of the decaying particle, h_j is the relevant Yukawa coupling, and from now on Lorentz factors, such as (M_j/T) , appearing in decay

rates are understood to be present only for temperatures above the mass of the decaying particle. To prevent washout one requires $(n_j/n_l)\Gamma_D \lesssim H \sim g^{1/2}T^2/M_{Pl}$, for $T_{EW} < T < T_{BG}$, where n_j (n_l) is the number density of species j (leptons). The strongest constraint arises from the latest time the density of the exotic particles is still significant: at $T \approx \Lambda_{ISS}$. This implies,

$$h_j \lesssim \left(16\pi g^{1/2} \frac{\Lambda_{ISS}^3}{M_{Pl}}\right)^{1/2} \approx 10^{-7} \left(\frac{g}{100}\right)^{1/4} \left(\frac{\Lambda_{ISS}}{\text{TeV}}\right)^{1/2}, \quad (3.12)$$

where g are the relativistic degrees of freedom at $T \sim M_j \sim \Lambda_{ISS}$.

If this condition holds for y_E , or h_ξ and h'_ξ , or h_χ and h'_χ , the lepton-number violating decays and inverse decays induced by these couplings are rare, and the BAU remains frozen.

- (ii) The thermally averaged rate of the decay mode $\chi \rightarrow \eta\eta$ is,

$$\Gamma_{D \chi \rightarrow \eta\eta} \approx \frac{\mu_{\eta\chi}^2}{32\pi\mu_\chi} \left(\frac{\mu_\chi}{T}\right). \quad (3.13)$$

Hence the lepton number of the universe is approximately conserved if:

$$\mu_{\eta\chi} \lesssim \left(32\pi g^{1/2} \frac{\Lambda_{ISS}^3}{M_{Pl}}\right)^{1/2} \approx \left(\frac{g}{100}\right)^{1/4} \left(\frac{\Lambda_{ISS}}{1 \text{ TeV}}\right)^{3/2} 10^{-4} \text{ GeV}. \quad (3.14)$$

- (iii) The cross-section for the scattering process $\eta^* + \xi^* \rightarrow \Phi + \Phi$ is

$$\sigma \sim \frac{\lambda_{\xi\phi\eta}^2}{16\pi} \frac{1}{(T^2 + \Lambda_{ISS}^2)}, \quad (3.15)$$

and the corresponding scattering rate is $\Gamma \sim n(T)\sigma$. In the relativistic regime, $\Gamma \propto T$, and hence increases with temperature slower the Hubble parameter. In the $T < \Lambda_{ISS}$ regime, Γ becomes Boltzmann suppressed. Thus, for sufficiently small coupling

$$\lambda_{\xi\phi\eta} \lesssim 10^{-7} \left(\frac{g}{100}\right)^{1/4} \left(\frac{\Lambda_{ISS}}{1 \text{ TeV}}\right)^{1/2}, \quad (3.16)$$

this process is never in equilibrium.

- (iv) Lepton-number violating processes may also occur via some of the ISS degrees of freedom propagating off-shell. For example the scattering process $N_R + \chi^* \rightarrow \overline{N}_R \rightarrow \overline{E}_L + \xi$ has an approximate cross section

$$\sigma \sim \frac{(h_\xi h_\chi)^2}{8\pi} \frac{T^2}{(T^2 + M_N^2)^2}, \quad (3.17)$$

which leads to a bound

$$(h_\xi h_\chi) \lesssim 10^{-7} \left(\frac{g}{100}\right)^{1/4} \left(\frac{\Lambda_{ISS}}{1 \text{ TeV}}\right)^{1/2}. \quad (3.18)$$

Satisfying *at least one* of the bounds of eqs. (3.12), (3.14), (3.16) and (3.18) ensures no washout of the BAU, if baryogenesis has occurred at temperatures $T_{\text{BG}} > \Lambda_{\text{ISS}}$. However, as can be seen from eq. (3.9), these bounds would imply that the contribution to the active neutrino masses generated via this mechanism is insignificant.

- $T_{\text{EW}} < T_{\text{BG}} < \Lambda_{\text{ISS}}$

The ISS degrees of freedom remain in thermal and chemical equilibrium even at temperatures much below the ISS scale, due to their gauge, Yukawa and scalar interactions. However, at $T_{\text{BG}} < \Lambda_{\text{ISS}}$, their density is Boltzmann suppressed. This implies that the exotic leptons carry only a small fraction of the net lepton number of the universe.

Consider for example again the 2-body decay rate of the ISS degrees of freedom via the Yukawa couplings of eq. (3.4). The decay rates at $T_{\text{BG}} < \Lambda_{\text{ISS}}$ are $\Gamma_j \approx h_j^2 M_j / 16\pi$, where as before M_j is the mass of the decaying particle and h_j is the Yukawa coupling that causes the decay. The lepton asymmetry will be approximately conserved if $\Gamma_j n_j \ll H n_l$, where $n_l \sim T^3$ is the lepton-number density, carried mostly by the light relativistic SM leptons, and $n_j \approx (M_j T / 2\pi)^{3/2} \exp(-M_j/T)$ is the number density of the non-relativistic exotic leptons.

For Yukawa couplings $h_j \sim \mathcal{O}(1)$, this yields the constraint

$$\frac{\Lambda_{\text{ISS}}}{T_{\text{BG}}} \gtrsim 42, \quad (3.19)$$

(where we set $M_j \sim \Lambda_{\text{ISS}}$). The interaction considered in the case (ii) gives a similar constraint, while the scattering rates in cases (iii) and (iv) are sufficiently suppressed for:

$$\frac{\Lambda_{\text{ISS}}}{T_{\text{BG}}} \gtrsim 25. \quad (3.20)$$

We therefore uncover an important feature of this model: baryogenesis has to occur below the ISS scale. If it occurs before the EW phase transition, the ISS scale has to be sufficiently high, as seen by eq. (3.19). If it occurs after the EW phase transition, no limit on Λ_{ISS} applies from considering the survival of the BAU today. Note that in this model, μ can only be generated after EW symmetry breaking (see eq. (3.8)), because the exotic fermions which gain a radiative Majorana mass carry hypercharge. This precludes resonant leptogenesis within this model, as leptogenesis must occur before the sphalerons switch off. It may, however, be possible that another of the ISS fields generates the asymmetry, with washout being suppressed by having the masses of at least one of the other exotic fields at a sufficiently high scale above T_{BG} . The construction of such a scenario, ensuring the satisfaction of the Sakharov conditions [56] and generation of the observed BAU, is beyond the scope of this work.

3.3 Dark Matter

In the Law/McDonald model the scalars, ξ and η , and the fermions, N_R and N_L , are odd under an accidental Z_2 symmetry. The lightest of these is a DM candidate. Assume for now that the lightest state is in the scalar sector (we will briefly discuss the fermionic case below). The neutral components of ξ and η mix and the masses of the real and imaginary components

split. The general mass squared matrix in the basis $(\text{Re}(\eta) \text{Re}(\xi^0) \text{Im}(\eta) \text{Im}(\xi^0))$ is,

$$\begin{pmatrix} M_\eta^2 + \mu_{\eta\chi}v_\chi/\sqrt{2} & \lambda_{\xi\phi\eta}v_w^2/4 & 0 & 0 \\ \lambda_{\xi\phi\eta}v_w^2/4 & M_\xi^2 & 0 & 0 \\ 0 & 0 & M_\eta^2 - \mu_{\eta\chi}v_\chi/\sqrt{2} & -\lambda_{\xi\phi\eta}v_w^2/4 \\ 0 & 0 & -\lambda_{\xi\phi\eta}v_w^2/4 & M_\xi^2 \end{pmatrix}, \quad (3.21)$$

where we have taken $\lambda_{\xi\phi\eta}$ and $\mu_{\eta\chi}$ to be real and positive without loss of generality as one can absorb the phase by redefining the fields, and

$$M_\eta^2 \equiv \mu_\eta^2 + \lambda_{\eta\phi}v_w^2/2 + \lambda_{\eta\chi}v_\chi^2/2, \quad (3.22)$$

$$M_\xi^2 \equiv \mu_\xi^2 + \lambda_{\xi\phi}v_w^2/2 + \lambda_{\xi\chi}v_\chi^2/2. \quad (3.23)$$

The lightest of the eigenstates forms the dark matter. If the admixture of ξ^0 in the DM field is small (which is the case for $M_\xi \gg M_\eta$) interactions with the Z boson will be suppressed. Scattering of DM with SM particles via the Z or Z' is also suppressed due to the mass splitting between the imaginary and real components of the DM field. (Gauge bosons interact off diagonally with the real and imaginary components of scalar fields.) Annihilation and scattering are dominated by the interactions with the Higgs fields, and for simplicity we will work in this regime. We then have a scalar Higgs portal DM candidate which has been studied extensively in the literature [57–60]. A possible difference from the standard Higgs portal arises when the heavier Higgs boson is light enough to influence the DM annihilation cross section [61–65]. We will explore this possibility in greater detail below in the context of the Law/McDonald model, though the discussion is largely applicable to generic Higgs portal models with an EW singlet Higgs mixing with the SM Higgs.

3.3.1 Double Higgs portal DM

We denote our (real scalar) DM candidate η , from now, and take its mass to be:

$$M_{DM}^2 = \mu_\eta^2 + \lambda_{\eta\phi}v_w^2/2 + \lambda_{\eta\chi}v_\chi^2/2 - \mu_{\eta\chi}v_\chi/\sqrt{2}, \quad (3.24)$$

where $\mu_{\eta\chi} > 0$. One can choose parameters for the scalar potential (3.7) so Φ , and χ gain the following VEVs: $\langle\Phi\rangle \equiv 1/\sqrt{2}(0, v_w)^T$, and $\langle\chi\rangle \equiv v_\chi/\sqrt{2}$. The VEVs satisfy the following equations:

$$\mu_\chi^2 + \frac{1}{2}\lambda_{\phi\chi}v_w^2 + \lambda_\chi v_\chi^2 = 0, \quad (3.25)$$

$$\mu_\phi^2 + \frac{1}{2}\lambda_{\phi\chi}v_\chi^2 + \lambda_\phi v_w^2 = 0. \quad (3.26)$$

To ensure that the potential is bounded from below and that $(\Phi, \chi, \eta) = (v_w/\sqrt{2}, v_\chi/\sqrt{2}, 0)$ is a minimum requires:

$$\lambda_\phi, \lambda_\chi, \lambda_\eta > 0, \quad (3.27)$$

$$\lambda_{\phi\chi}^2 < 4\lambda_\phi\lambda_\chi, \quad (3.28)$$

$$M_{DM}^2 > 0, \quad (3.29)$$

$$\lambda_{\eta\chi} > -2\sqrt{\lambda_\eta\lambda_\chi}, \quad (3.30)$$

$$\lambda_{\eta\phi} > -2\sqrt{\lambda_\eta\lambda_\phi}. \quad (3.31)$$

An additional condition for symmetry breaking is $\mu_\chi^2 < 0$, or $\mu_\phi^2 < 0$, and to ensure there is no deeper minimum (at least at zero temperature) one can take $\mu_\eta^2 > 0$. The above conditions are necessary but not sufficient to ensure the potential is bounded from below. An additional sufficient condition is:

$$\lambda_\eta > \frac{1}{4\lambda_\phi} \left\{ \frac{(4\lambda_\phi\lambda_{\eta\chi} - 2\lambda_{\phi\chi}\lambda_{\eta\phi})^2}{4(4\lambda_\phi\lambda_\chi - \lambda_{\phi\chi}^2)} + \lambda_{\eta\phi}^2 \right\}. \quad (3.32)$$

The squared mass matrix for the physical Higgs bosons ϕ and χ is given by:

$$\mathcal{L} = \frac{1}{2} \begin{pmatrix} \phi & \chi \end{pmatrix} \begin{pmatrix} 2\lambda_\phi v_w^2 & \lambda_{\phi\chi} v_w v_\chi \\ \lambda_{\phi\chi} v_w v_\chi & 2\lambda_\chi v_\chi^2 \end{pmatrix} \begin{pmatrix} \phi \\ \chi \end{pmatrix}. \quad (3.33)$$

One then finds the mass eigenstates ($m_H \geq m_h$),

$$m_h^2 = \lambda_\chi v_\chi^2 + \lambda_\phi v_w^2 - \sqrt{(\lambda_\chi v_\chi^2 - \lambda_\phi v_w^2)^2 + (\lambda_{\phi\chi} v_w v_\chi)^2}, \quad (3.34)$$

$$m_H^2 = \lambda_\chi v_\chi^2 + \lambda_\phi v_w^2 + \sqrt{(\lambda_\chi v_\chi^2 - \lambda_\phi v_w^2)^2 + (\lambda_{\phi\chi} v_w v_\chi)^2}. \quad (3.35)$$

The mass eigenstates are superpositions of ϕ and χ ,

$$\begin{pmatrix} h \\ H \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \phi \\ \chi \end{pmatrix}, \quad (3.36)$$

where the mixing angle θ is given by the following,

$$\tan \theta = \frac{\lambda_{\phi\chi} v_w v_\chi}{\lambda_\chi v_\chi^2 - \lambda_\phi v_w^2 + \sqrt{(\lambda_\chi v_\chi^2 - \lambda_\phi v_w^2)^2 + (\lambda_{\phi\chi} v_w v_\chi)^2}}. \quad (3.37)$$

Note h (H) couples to SM particles the same way as the SM Higgs but with an additional factor of $\cos \theta$ ($-\sin \theta$) at the vertex. We identify h with the Higgs-like particle discovered at the LHC.

In the limit where the H is very heavy the important DM interactions proceed through the SM-like Higgs, and one in effect returns to the standard Higgs portal scenario. If H is lighter, however, one now has annihilation to SM particles and interactions with quarks proceeding through both h and H propagators. We term this a double Higgs portal and we shall compare the two scenarios below. Similar ideas have been explored previously in the literature: refs. [61–63] consider two Higgs doublet models, while [64] considers fermionic DM with an exotic SM singlet Higgs. We consider scalar DM with an exotic singlet Higgs, a scenario which has been analysed previously in [65]. One important difference is that we focus more on the phenomenology of high DM masses, while the focus previously has been on the lower mass range $M_{DM} \lesssim 150$ GeV.

In terms of the Higgs mass eigenstates one finds the following interactions with the DM:

$$\begin{aligned} -\mathcal{L} \supset & \frac{1}{2} \eta^2 (\lambda_{\eta\phi} v_w \cos \theta + \lambda_{\eta\chi} v_\chi \sin \theta - \mu_{\eta\chi} \sin \theta) h \\ & + \frac{1}{2} \eta^2 (-\lambda_{\eta\phi} v_w \sin \theta + \lambda_{\eta\chi} v_\chi \cos \theta - \mu_{\eta\chi} \cos \theta) H \\ & + \frac{1}{2} \eta^2 (\lambda_{\eta\chi} \cos \theta \sin \theta - \lambda_{\eta\phi} \cos \theta \sin \theta) h H \\ & + \frac{1}{4} \eta^2 (\lambda_{\eta\phi} \sin^2 \theta + \lambda_{\eta\chi} \cos^2 \theta) H^2 \\ & + \frac{1}{4} \eta^2 (\lambda_{\eta\phi} \cos^2 \theta + \lambda_{\eta\chi} \sin^2 \theta) h^2. \end{aligned} \quad (3.38)$$

To reduce the number of parameters, and allow one to gain an appreciation of the phenomenology of the double Higgs portal, we study a special case where $\lambda_{\eta\chi} = \lambda_{\eta\phi} \equiv \lambda$ and $\mu_{\eta\chi} = |\lambda|\mu'_{\eta\chi}$, where we will vary λ but keep $\mu'_{\eta\chi} > 0$ and fixed. The interaction of the Higgs bosons with the DM then simplifies to:

$$\begin{aligned}
-\mathcal{L} \supset & \frac{1}{2} \left(v_w \cos \theta + v_\chi \sin \theta - \text{Sign}[\lambda] \cdot \mu'_{\eta\chi} \sin \theta \right) \lambda \eta^2 h \\
& + \frac{1}{2} \left(v_\chi \cos \theta - v_w \sin \theta - \text{Sign}[\lambda] \cdot \mu'_{\eta\chi} \cos \theta \right) \lambda \eta^2 H \\
& + \frac{1}{4} \lambda \eta^2 H^2 + \frac{1}{4} \lambda \eta^2 h^2.
\end{aligned} \tag{3.39}$$

To see the effect of the second Higgs boson, we present two cases. Case A is where the heavy Higgs boson is taken to be heavy enough and the mixing negligible, so it is effectively decoupled. In this limit one obtains the standard Higgs portal. For case B, the double Higgs portal, we choose some parameters, namely $M_H = 500$ GeV, $v_\chi = 1$ TeV, $\mu'_{\eta\chi} = 2$ TeV, and examine the change in phenomenology for different choices of the Higgs mixing parameter $\sin \theta$.

Furthermore, given the above choice of parameters, we have calculated λ_χ , λ_ϕ , and $\lambda_{\phi\chi}$ as functions of $\sin \theta$. One finds that condition (3.28) is satisfied for the entire range of the mixing angle ($-\pi/2 \leq \theta \leq \pi/2$), and that $\mu_\chi^2 < 0$ so that symmetry breaking can proceed.

As the heavy Higgs boson enters loop corrections to the W and Z boson propagators, a constraint on $\sin \theta$ comes from the oblique parameters: T , S , and U . By following the analysis of [65–67] one finds that for $M_H = 500$ GeV the mixing angle is constrained to be

$$|\sin \theta| \leq 0.43 \tag{3.40}$$

by EW precision observations at 95% C.L. (assuming the other exotic particles give only negligible contributions.)

3.3.2 DM relic abundance

One can then calculate the annihilation cross section of DM into standard model particles. The relevant Feynman diagrams are shown in figure 4. To obtain the velocity averaged cross section, one should integrate over the velocity distribution. However a good approximation is obtained with,

$$\langle \sigma v_{rel} \rangle \simeq (\sigma v_{rel})|_{s=4M_{DM}^2}. \tag{3.41}$$

One can then simply compare to the required annihilation cross section, for the observed relic abundance [68], which for DM masses greater than 10 GeV is approximately [69],

$$\langle \sigma v_{rel} \rangle_{relic} \approx 2.2 \times 10^{-26} \text{ cm}^3/\text{s}. \tag{3.42}$$

To obtain the desired M_{DM}^2 with a choice $\mu_\eta^2 > 0$ one wants to choose the required values of λ to be negative (particularly for small DM masses), as can be seen from eq. (3.24). Some degree of fine tuning is of course required for small DM masses given our choice $v_\chi = 1$ TeV.

We have calculated the required coupling $\lambda \equiv \lambda_{\eta\phi} = \lambda_{\eta\chi}$, with the constraint $\lambda < 0$. (The relevant expressions for the cross sections can be found in the appendix of ref. [60] for case A, and simple modifications for case B can be found by taking into account the Lagrangian in eq. (3.39).) The results for different choices of the mixing parameter are




Figure 4. Possible DM annihilation channels. The diagrams on the lower line give only a small correction and are not included in the calculations.

shown in figure 5. The observed relic abundance can be achieved with values of $\lambda \sim -0.1$ for most of the DM mass range. Note the falls in the required coupling at the points where $2M_{DM} = M_{h/H}$, as the intermediate Higgs boson is on-shell.

We have also checked that the constraints of eqs. (3.30-3.32) can be satisfied with perturbative choices of the quartic coupling λ_η (except for small regions of parameter space around $\sin \theta \approx 0.05$, where for low DM masses $|\lambda| > 1$).




Figure 5. Required coupling for the correct DM abundance for different values of $\sin \theta$ and Case A (single Higgs portal) for comparison. Parameters have been chosen so that $\lambda \equiv \lambda_{\eta\phi} = \lambda_{\eta\chi}$, $\mu'_{\eta\chi} = 2$ TeV, $M_H = 500$ GeV, and $v_\chi = 1$ TeV.

3.3.3 Direct detection




Figure 6. DM Nucleon scattering through t-channel Higgs exchange. The effective coupling of the Higgs with the Nucleon is indicated.

DM particles will scatter off nuclei through t-channel Higgs boson exchange (see figure 6). The spin independent DM-nucleon scattering cross section for our DM candidate η is given by,

$$\sigma_{DM-Nuc}^{SI} = \frac{\lambda^2 M_{Nuc}^4 f_N^2}{4\pi(M_{DM} + M_{Nuc})^2} \left(\frac{A}{m_H^2} + \frac{B}{m_h^2} \right)^2, \quad (3.43)$$

where $M_{Nuc} \simeq 0.94$ GeV is the nucleon mass, f_N parametrizes the effective Higgs-nucleon coupling, and the vertex factors are:

$$A = -\sin\theta \left\{ \frac{v_\chi}{v_w} \cos\theta - \sin\theta - \text{Sign}[\lambda] \cdot 2 \frac{\mu'_{\eta\chi}}{v_w} \cos\theta \right\}, \quad (3.44)$$

$$B = \cos\theta \left\{ \cos\theta + \frac{v_\chi}{v_w} \sin\theta - \text{Sign}[\lambda] \cdot 2 \frac{\mu'_{\eta\chi}}{v_w} \sin\theta \right\}. \quad (3.45)$$

Note that uncertainties in f_N result in roughly a factor of ~ 5 uncertainty in σ_{DM-Nuc}^{SI} [70]. We use a value of $f_N = 0.3$ [71–74].

Having calculated λ , we then find σ_{DM-Nuc}^{SI} for different values of $\sin\theta$. The results are presented in figure 7. XENON100 exclusion limits from 255 live days of data taking [75], labeled XENON100, and projected XENON1T exclusions for 2017 in the case of no WIMP-nucleon scattering [76], labeled XENON1T, are plotted for comparison. We see that $M_{DM} \gtrsim 80$ GeV is still largely viable, but that much of the remaining parameter space can be ruled out by XENON1T.

Note that due to different interferences the scattering rate can be above or below the zero mixing case. A minimum is reached for $\sin\theta = -0.052$, which is exactly where,

$$\left(\frac{A}{m_H^2} + \frac{B}{m_h^2} \right) = 0, \quad (3.46)$$

so that the DM-nucleon cross section is zero. Similar interference was noted in the case of a two Higgs doublet extension of the Higgs portal in ref. [62].




Figure 7. DM-Nucleon scattering cross section for case A and case B with different values of $\sin(\theta)$, the upper limit from XENON100, and the projected sensitivity for XENON1T in 2017.

3.3.4 Higgs signals at colliders

The double Higgs portal can be tested in a number of ways at colliders [62, 63, 77, 78], some of them common with the usual single Higgs portal [67, 70, 79, 80]. Mixing with the exotic Higgs will reduce the production of the SM-like Higgs,

$$\sigma_{pp \rightarrow h} = \cos^2 \theta \cdot \sigma_{pp \rightarrow h}^{SM}. \quad (3.47)$$

The total width will also be reduced by a factor of $\cos^2 \theta$, however the SM-width for $m_h = 125$ GeV, roughly 4 MeV, is already far below current experimental resolution. The discovery of a SM-like Higgs boson at the LHC [81, 82] allows us to constrain this scenario. The observed signal strength over the SM expectation and 1σ uncertainty is [83, 84]:

$$\sigma/\sigma_{SM} = 0.88 \pm 0.21 \quad (\text{CMS}), \quad (3.48)$$

$$\sigma/\sigma_{SM} = 1.3 \pm 0.13 \text{ (stat.)} \pm 0.14 \text{ (syst.)} \quad (\text{ATLAS}). \quad (3.49)$$

Combinations of the results put a limit on the universal suppression of all channels so that: $\sigma/\sigma_{SM} \gtrsim 0.8$ at the 2σ level [85–87]. This translates into a limit,

$$|\sin \theta| \leq 0.45, \quad (3.50)$$

if one assumes no invisible decays of the 125 GeV resonance. (An alternative, more conservative, treatment of the theoretical uncertainty in σ_{SM} modifies the above bound to be $|\sin \theta| \lesssim 0.7$ instead [88].) Equation (3.50) is comparable to the constraint from EW precision observables, eq. (3.40), however the limit on the mixing angle from measurement of the signal strength is generally applicable and not limited to our choice $M_H = 500$ GeV.

If the DM is light enough, the Higgs can decay invisibly to two DM particles. The invisible partial decay width of the Higgs ($h \rightarrow \eta\eta$) is given by,

$$\Gamma = \frac{\lambda^2}{32\pi m_h} \left(v_w \cos \theta + v_\chi \sin \theta - \text{Sign}[\lambda] \cdot 2\mu'_{\eta\chi} \sin \theta \right)^2 \left(1 - \frac{4m_\eta^2}{m_h^2} \right)^{1/2}. \quad (3.51)$$

This leads to a large invisible branching fraction $Br(h \rightarrow \eta\eta) \gtrsim 0.99$ for light DM masses $M_{DM} \lesssim 10$ GeV, except for a very narrow region around a mixing angle $\sin \theta = -0.049$ where the effective coupling of the $h\eta\eta$ term is zero. This helps to rule out such light DM candidates as the direct detection experiments lose sensitivity for low mass DM candidates.

If the mixing angle is large enough, the heavy state could also be detected. For a Higgs mass of 500 GeV, CMS data currently sets a 95% C.L. [89]:

$$\sigma/\sigma_{SM} \simeq 0.21. \quad (3.52)$$

The most stringent limit on the mixing angle will come from assuming $2M_{DM} > M_H$, so there is no invisible width. There is also an additional complication of $H \rightarrow hh$ decays [90], however, for the parameters we have chosen the relevant branching ratio turns out to be negligible. The limit on the mixing angle from high mass Higgs searches is then:

$$|\sin \theta| \leq 0.46. \quad (3.53)$$

Thus the most stringent limit on the mixing angle in case B with $M_H = 500$ GeV comes from considering EW precision observables, eq. (3.40), while the most widely applicable constraint comes from measurements of the signal strength of the 125 GeV resonance, eq. (3.50), as no assumptions about the mass of the heavier Higgs or VEV have to be taken as an input.

3.3.5 Fermionic DM

Alternatively one may consider fermionic Higgs portal DM. The important terms are then:

$$-\mathcal{L} \supset h_\chi \chi \overline{N_R} (N_R)^c + h'_\chi \chi \overline{N_L} (N_L)^c + M_N \overline{N_L} N_R + H.c., \quad (3.54)$$

where the lightest mass eigenstate of the N_L , $(N_R)^c$ admixture is the DM candidate (protected by the same accidental Z_2 as in the scalar DM case). In such a scenario DM is constrained to be very heavy from direct detection experiments, $M_{DM} \gtrsim 2$ TeV, unless the DM mass is at one of the two Higgs resonances, there is parity violation, or a Sommerfeld enhancement [64]. So while we have studied the scalar DM case above, either scalar or fermionic DM is viable in the Law/McDonald model. Further generalizations taking into account interactions with the Z' in the fermionic DM scenario are also possible, but we will not pursue the details here.

3.4 Concluding remarks on the Law/McDonald model

We have seen that in the Law/McDonald model, baryogenesis is severely constrained. The explicit lepton number violation will tend to wash out any baryon asymmetry created at a high scale. However at temperatures roughly a factor of 25-40 below the ISS mass scale, Boltzmann suppression of the thermally averaged reaction rates means that the ISS fields will not contribute to washout (though not necessarily *all* of the ISS fields must have such high masses). Unfortunately the small L violating parameter is only generated after EW symmetry

breaking, meaning that resonant leptogenesis is not possible in the Law/McDonald radiative ISS.

We found that the Law/McDonald model can easily accommodate the observed DM abundance through the Higgs portal. Also, in a simple extension, an additional Higgs state can also contribute to the DM annihilation rate and DM nucleon scattering rate. Apart from narrow regions of parameter space, either on resonance points for the annihilation rate, or near complete destructive interference for the DM nucleon scattering rates, these models can for the most part be tested by the XENON1T experiment.

4 Ma radiative inverse seesaw model

4.1 Review of the model

We now consider the Ma radiative ISS model [29]. An important difference in this model is that $(B - L)$ is a protected symmetry down to the ISS scale. First let us review the main features of this model. It employs an additional $U(1)$ which originates from a grand unified theory (GUT) with the following breaking pattern,

$$\begin{aligned} SO(10) &\rightarrow SU(5) \otimes U(1)_\chi \\ &\rightarrow SU(3)_C \otimes SU(2)_L \otimes U(1)_Y \otimes U(1)_\chi. \end{aligned} \quad (4.1)$$

The exotic charge is given by the following,

$$Q_\chi = 5(B - L) - 4Y = 5(B - L) + 4I_{3L} - 4Q_{EM}. \quad (4.2)$$

There are two scalars, η_1 and η_2 , that transform under $U(1)_\chi$ as per

$$Q_\chi(\eta_1) = 1, \quad Q_\chi(\eta_2) = 2, \quad (4.3)$$

and are neutral with respect to the SM gauge group. These will be used for the radiative ISS and to break the new $U(1)_\chi$. To allow the usual Yukawa interactions,

$$\mathcal{L} = \lambda_d \overline{Q_L} \cdot \Phi d_R + \lambda_u \epsilon^{ab} \overline{Q_{La}} \Phi_b^\dagger u_R + \lambda_e \overline{l_L} \cdot \Phi e_R + \lambda_\nu \epsilon^{ab} \overline{l_{La}} \Phi_b^\dagger N_R + H.c., \quad (4.4)$$

the SM Higgs doublet must carry a $U(1)_\chi$ charge,

$$\Phi = (\phi^+, \phi^0) \sim (1, 2, 1/2, -2). \quad (4.5)$$

The fermion multiplets of $SO(10)$ contain a right handed neutrino, $N_R \sim (1, 1, 0, -5)$, per generation of SM fermions. To these are added more SM neutral fermions with Q_χ given by,

$$1 \times S_{3L} \sim -3, \quad 4 \times S_{2L} \sim 2, \quad 5 \times S_{1L} \sim -1, \quad (4.6)$$

where the required number of copies has been indicated to form an anomaly-free set. The following Yukawa terms involving the exotic scalars are permitted,

$$\mathcal{L} \supset f_{R3} \overline{N_R} S_{3L} \eta_2^\dagger + f_{23} \overline{(S_{3L})^c} S_{2L} \eta_1 + f_{12} \overline{(S_{2L})^c} S_{1L} \eta_1^\dagger + f_{11} \overline{(S_{1L})^c} S_{1L} \eta_2 + H.c. \quad (4.7)$$

The scalar potential is given by,

$$\begin{aligned} V = & (A \eta_1^2 \eta_2^\dagger + H.c.) + \lambda_{11} (\eta_1^\dagger \eta_1)^2 + \lambda_{22} (\eta_2^\dagger \eta_2)^2 + \lambda_{12} (\eta_1^\dagger \eta_1) (\eta_2^\dagger \eta_2) \\ & + \mu_\Phi^2 \Phi^\dagger \Phi + \lambda_\Phi (\Phi^\dagger \Phi)^2 + \lambda_{1\Phi} (\Phi^\dagger \Phi) (\eta_1^\dagger \eta_1) + \lambda_{2\Phi} (\Phi^\dagger \Phi) (\eta_2^\dagger \eta_2) \\ & + \mu_1^2 (\eta_1^\dagger \eta_1) + \mu_2^2 (\eta_2^\dagger \eta_2), \end{aligned} \quad (4.8)$$

where A is a coupling constant with dimension of mass. The $U(1)_\chi$ symmetry is spontaneously broken by,

$$\langle \eta_2 \rangle \equiv v_\chi / \sqrt{2} \neq 0, \quad \langle \eta_1 \rangle = 0. \quad (4.9)$$

Before radiative corrections are taken into account the neutral fermion mass matrix is given by,

$$\mathcal{L} = \frac{1}{2} (\overline{\nu_L} \quad \overline{(N_R)^c} \quad \overline{S_{3L}} \quad \overline{S_{1L}}) \begin{pmatrix} 0 & m_D & 0 & 0 \\ m_D & 0 & M_N & 0 \\ 0 & M_N & 0 & 0 \\ 0 & 0 & 0 & M_1 \end{pmatrix} \begin{pmatrix} (\nu_L)^c \\ N_R \\ (S_{3L})^c \\ (S_{1L})^c \end{pmatrix}, \quad (4.10)$$

where $m_D = \lambda_\nu v_w / \sqrt{2}$, $M_N = f_{R3} v_\chi / \sqrt{2}$, and $M_1 = \sqrt{2} f_{11} v_\chi$. Note the S_{2L} are massless at tree level, and that the scalar interaction term,

$$A \eta_1 \eta_1 \eta_2^\dagger + A^* \eta_1^\dagger \eta_1^\dagger \eta_2 \rightarrow A \eta_1 \eta_1 v_\chi / \sqrt{2} + A^* \eta_1^\dagger \eta_1^\dagger v_\chi / \sqrt{2}, \quad (4.11)$$

induces a mass splitting between the mass of the real (m_R) and mass of the imaginary (m_I) components of η_1 ,

$$m_R^2 - m_I^2 = 2\sqrt{2} A v_\chi, \quad (4.12)$$

where we set A to be real and positive without loss of generality as any complex phase can be absorbed into the fields.




Figure 8. Radiative mass generation for S_2 and S_3 in the Ma model.

The exchange of $Re(\eta_1)$ and $Im(\eta_1)$ then give one-loop radiative Majorana masses to the S_{2L} as can be seen in figure 8:

$$M_2 = \frac{(f_{12})^2 M_1}{16\pi^2} \left[\frac{m_R^2}{m_R^2 - M_1^2} \ln \left(\frac{m_R^2}{M_1^2} \right) - \frac{m_I^2}{m_I^2 - M_1^2} \ln \left(\frac{m_I^2}{M_1^2} \right) \right]. \quad (4.13)$$

Similarly one obtains two-loop radiative Majorana masses for the S_{3L} ,

$$\mu \equiv M_3 = \frac{(f_{23})^2 M_2}{16\pi^2} \left[\frac{m_R^2}{m_R^2 - M_2^2} \ln \left(\frac{m_R^2}{M_2^2} \right) - \frac{m_I^2}{m_I^2 - M_2^2} \ln \left(\frac{m_I^2}{M_2^2} \right) \right], \quad (4.14)$$

which are exactly the small parameters required in the mass matrix of eq. (4.10) for the ISS.

Note η_1 and S_{2L} are odd under an accidental Z_2 symmetry, while all the other particles are even. We will see below, that for typical choices of the parameters, S_{2L} is lighter than η_1

as its mass arises radiatively and therefore S_{2L} is the DM candidate. One can find various approximate forms for the radiative masses depending on the sizes of various parameters. If we take $m_R^2 \gg m_R^2 - m_I^2 = 2\sqrt{2}Av_\chi, M_1$, eq. (4.13) can be approximated as,

$$M_2 \approx \frac{\sqrt{2}(f_{12})^2 Av_\chi M_1}{8\pi^2 m_I^2}, \quad (4.15)$$

while for natural choices of the parameters $m_R^2, m_I^2 \gg M_2^2$, one obtains from eq. (4.14):

$$\mu \equiv M_3 \approx \frac{(f_{12})^2 (f_{23})^2 A^2 v_\chi^2 M_1}{32\pi^4 m_I^4}. \quad (4.16)$$

To give a numerical example, for $v_\chi \sim M_1 \sim A \sim \mathcal{O}(10 \text{ TeV})$, and $m_I = 50 \text{ TeV}$, one finds $M_2 \approx (f_{12})^2 \times 7 \text{ GeV}$ and $M_3 \approx (f_{23})^2 (f_{12})^2 \times 5 \text{ MeV}$ (in very close agreement to what the exact formulas give).

4.2 Constraints from BAU washout

Until $U(1)_\chi$ is broken, $(B - L)$ is an exact symmetry of the Ma model, and generation or washout of the BAU will not take place.¹ The VEV v_χ breaks $Q_\chi = 5(B - L) - 4Y$, and hence the $(B - L)$ as defined above. Baryogenesis can then take place, either through resonant leptogenesis, or some other suitable baryogenesis scenario with $T_{\text{BG}} \leq T_\chi \sim v_\chi$. However, as shown in figure 9, the ISS fields can now also wash out the asymmetry.




Figure 9. Washout process involving the exotic fermions and scalars. Also shown are the Yukawa interactions which link this washout process to the SM sector. Exchange of the real and imaginary components of η_1 leads to destructive interference. Not shown are the interactions involving f_{12} and f_{11} which ensure S_{2L} cannot be assigned non-zero SM lepton number.

A new conserved global $(B - L)$ can be defined, even with $\langle \eta_2 \rangle \neq 0$, provided any of the following parameters vanishes:

$$\lambda_\nu, f_{R3}, f_{23}, f_{12}, f_{11}, A. \quad (4.17)$$

¹This constraint could be circumvented through sequestration of positive and negative $B - L$ charge into the visible sector and a hidden sector. This is a popular mechanism in asymmetric DM models, which aim to dynamically relate the visible and dark matter relic abundances [91–103]. In the context of the model studied here, it would require introducing additional fields and interactions, such that the $B - L$ symmetry encompasses two low-energy global $U(1)$ symmetries – the $(B - L)_v$ carried by the SM fields and a dark baryon number B_d . The construction of such a model in an $SO(10)$ GUT is beyond the scope of this paper.

If the interactions mediated by one of the above parameters are out of equilibrium the BAU is safe. We first examine what values the parameters must take so as to not wash out the BAU. In section (4.3) we examine the possibility for resonant leptogenesis, where the lightest pseudo-Dirac pair generates the BAU.

- $T_{BG} \approx \Lambda_{ISS} \approx v_\chi$

- (i) The decays and inverse decays $N_R \leftrightarrow l_L + \Phi$ are controlled by λ_ν . The decay rate is approximately,

$$\Gamma \approx \frac{\lambda_\nu^2 M_N}{8\pi} \left(\frac{M_N}{T} \right), \quad (4.18)$$

requiring $n_N \Gamma \lesssim n_l H$ where n_N (n_l) is the N_R (lepton) number density leads to a washout avoidance condition:

$$\lambda_\nu \lesssim 10^{-7} \left(\frac{g}{100} \right)^{1/4} \left(\frac{M_N}{10 \text{ TeV}} \right)^{1/2}. \quad (4.19)$$

- (ii) The mixing between the N_R and S_{3L} states is controlled by $f_{R3} = \sqrt{2}M_N/v_\chi$. Scattering $\Phi + l_L \rightarrow N_R/S_{3L} \rightarrow \eta_1^* + \overline{S_{2L}}$ has an approximate cross section,

$$\sigma \sim \frac{(\lambda_\nu f_{23})^2}{8\pi} \frac{M_N^2}{(T^2 + M_N^2)^2}. \quad (4.20)$$

The scattering rate $\Gamma \sim \sigma T^3$ is inefficient compared to H provided:

$$f_{R3} \lesssim 10^{-6} \left(\frac{1}{\lambda_\nu f_{23}} \right) \left(\frac{g}{100} \right)^{1/4} \left(\frac{\Lambda_{ISS}}{10 \text{ TeV}} \right)^{1/2}, \quad (4.21)$$

and for such values washout is avoided.

- (iii) The Yukawa couplings f_{23} and f_{12} induce two body decays of the heaviest particle involved in the relevant interaction (such decays are kinematically allowed given S_{2L} , which has a radiative and hence negligible mass compared to the other particles, can appear in the final state). We estimate the decay rate as,

$$\Gamma \approx \frac{(f_i)^2 M_j}{16\pi} \left(\frac{M_j}{T} \right), \quad (4.22)$$

where f_i denotes the relevant Yukawa coupling and M_j the mass of the heaviest particle. This leads to a washout avoidance condition of:

$$f_i \lesssim 10^{-6} \left(\frac{g}{100} \right)^{1/4} \left(\frac{\Lambda_{ISS}}{10 \text{ TeV}} \right)^{1/2}. \quad (4.23)$$

- (iv) Scattering processes $S_{2L} + \eta_1^* \rightarrow S_1/S_1^c \rightarrow \overline{S_{2L}} + \eta_1$ can be suppressed with small $f_{11} = M_1/\sqrt{2}v_\chi$. We estimate the cross section to be:

$$\sigma \sim \frac{(f_{12})^4}{8\pi} \frac{M_1^2}{(T^2 + M_1^2)^2}. \quad (4.24)$$

This gives a washout avoidance condition:

$$f_{11} \lesssim 10^{-7} \left(\frac{1}{f_{12}} \right)^2 \left(\frac{g}{100} \right)^{1/2} \left(\frac{\Lambda_{ISS}}{10 \text{ TeV}} \right)^{1/2}. \quad (4.25)$$

- (v) Washout may proceed through decays and inverse decays η_1 provided they are kinematically allowed. However, for a small enough A , and hence small enough mass splitting between $Re(\eta_1)$ and $Im(\eta_1)$, destructive interference leads to washout suppression. We estimate the decay rate as,

$$\Gamma \approx \frac{(|f_{23}|^2 + |f_{12}|^2) M_I}{16\pi} \frac{M_I}{T}, \quad (4.26)$$

and the effective washout parameter to be,

$$K_{\text{eff}} = \left(\frac{\Gamma}{H} \Big|_{T=M_I} \right) \cdot \delta^2, \quad (4.27)$$

where $\delta = (m_R - m_I)/\Gamma$. One finds this translates into a washout avoidance condition:

$$A \lesssim 10^{-4} \text{ GeV} \sqrt{|f_{23}|^2 + |f_{12}|^2} \left(\frac{g}{100} \right)^{1/4} \left(\frac{\Lambda_{\text{ISS}}}{10 \text{ TeV}} \right)^{3/2}. \quad (4.28)$$

If such decays are not kinematically allowed washout can instead proceed through a scattering process mediated by off-shell η_1 . We estimate the cross section as:

$$\sigma \sim \frac{(|f_{23}| + |f_{12}|)^4}{128\pi} \left(\frac{1}{T^2 + m_R^2} - \frac{1}{T^2 + m_I^2} \right)^2 T^2 \quad (4.29)$$

$$= \frac{(|f_{23}| + |f_{12}|)^4}{128\pi} \left(\frac{2\sqrt{2}Av_\chi}{(T^2 + m_R^2)(T^2 + m_I^2)} \right)^2 T^2. \quad (4.30)$$

This translates into a washout avoidance condition:

$$A \lesssim 10^{-2} \text{ GeV} \left(\frac{1}{|f_{23}| + |f_{12}|} \right)^2 \left(\frac{g}{100} \right)^{1/4} \left(\frac{\Lambda_{\text{ISS}}}{10 \text{ TeV}} \right)^{3/2}. \quad (4.31)$$

Satisfying any of the above washout bounds means the BAU is safe. However, similarly to the Law/McDonald model, such small choices for the couplings ruins the ISS mechanism.

- $T_{\text{BG}} < \Lambda_{\text{ISS}}$

So as to ensure the preservation of the BAU, while allowing for the ISS mechanism, we may suppress washout by taking one or more of the exotic particle masses to be higher than T_{BG} . This ensures that at T_{BG} the number density of that particle will be sufficiently Boltzmann suppressed for it to not play a role in washout. For decay processes the requirement on the mass scale is:

$$\frac{\Lambda_{\text{ISS}}}{T_{\text{BG}}} \gtrsim 42, \quad (4.32)$$

while for scattering processes the constraint is relaxed to roughly,

$$\frac{\Lambda_{\text{ISS}}}{T_{\text{BG}}} \gtrsim 25. \quad (4.33)$$

This is again similar to the washout in the Law/McDonald model. For $T_{\text{BG}} < T_{\text{EW}}$ the bounds of eqs. (4.32) and (4.33) do not apply as the sphalerons are no longer active.

4.3 Resonant leptogenesis

For typical parameter choices, the lightest pseudo-Dirac pair of neutrinos, N_α will generate the BAU in resonant leptogenesis. There is an interplay between the CP violation and washout, which both depend on μ , the mass splitting of the N_α pair. So as to show that this model can generate the right order of magnitude for the parameters, let us take as a guide the example of [28], which found the observed BAU can be generated with $\delta = \mu/\Gamma_\alpha \approx 10^{-5}$ (Γ_α is the width of the N_α and μ the mass splitting of the pair), heavy neutrino mass $M_{N_\alpha} \sim \mathcal{O}(1)$ TeV, and Yukawa coupling $\lambda_\nu \approx 5 \times 10^{-2}$.

The scalar mass m_I can be taken to a sufficiently high scale so that the other interactions of the ISS do not interfere with the generation of the BAU. Let us take $m_I \sim 50$ TeV and $v_\chi \sim A \sim M_1 \sim 10$ TeV, then $M_N \sim \mathcal{O}(1)$ TeV can be achieved with $f_{R3} \sim 0.1$. Note that solutions of the Boltzmann equations taking into account decay and scattering processes show the BAU is generated at $T_{BG} \approx M_{N_\alpha}/10$, which allows the out-of-equilibrium condition for N_α to be met even with a relatively low Z' mass: $M_{Z'} \gtrsim 2M_{N_\alpha}$ [54].

Choices of $\delta = 10^{-5}$, $\lambda_\nu = 0.05$, and $M_{N_\alpha} = 2$ TeV imply: $\mu = 0.2 \times 10^{-5}$ GeV. This is compatible with a radiative origin in the Ma model as, for these parameters, eq. (4.16) gives $\mu \approx (f_{12})^2(f_{23})^2 \times 5 \times 10^{-3}$ GeV. The light neutrino mass is $m_\nu \approx 10^{-2}$ eV.

In the $\delta \ll 1$ regime, the CP asymmetry of N_α decays is [28],

$$\epsilon \simeq -\frac{\lambda_\nu^2}{16\pi} \frac{\mu \tilde{\mu}}{(\mu^2 + \tilde{\mu}^2 + 2\mu\tilde{\mu}\cos\alpha)} \sin\alpha, \quad (4.34)$$

where α is a physical phase of the ISS mass matrix and $\tilde{\mu}$ is the Majorana mass $\tilde{\mu}\overline{N_R}(N_R)^c$. In this model one typically expects $\tilde{\mu} \sim 10^{-3}\mu$, due to suppression by additional Yukawa couplings in the radiative generation of $\tilde{\mu}$, which allows for $\epsilon \sim 10^{-8}$. The washout factor is $K_{\text{eff}} \approx 6$. Taking into account a sphaleron reprocessing and dilution factor, the final baryon number density to photon number density is, $\eta_B \sim 10^{-2}|\epsilon|$, and can easily be made to match the observed value $\eta_B \approx 6 \times 10^{-10}$ [104] in this scenario.

A more detailed study finding the full form of the 3×3 matrices m_D and μ , and mass spectrum for the pseudo-Dirac pairs compatible with the observed oscillation parameters, BAU, and $SO(10)$ origin of the model is beyond the scope of this paper.

4.4 The ρ parameter

In the Ma model, the Higgs doublet responsible for EW symmetry breaking also carries $U(1)_\chi$ charge. This leads to a tree level correction to the Z boson mass, and consequently to the ρ parameter,

$$\rho \equiv \frac{M_W^2}{M_Z^2 \cos^2 \theta_W}, \quad (4.35)$$

where the Weinberg angle is defined in terms of the weak isospin, g , and weak hypercharge, g_Y , coupling constants as $\tan \theta_W \equiv g_Y/g$. Experimental observations give a value $\rho - 1 = 4_{-4}^{+3} \times 10^{-4}$ [105]. The masses of the neutral gauge bosons are given by,

$$m_\pm^2 = \frac{v_W^2}{8} \left\{ g^2 + g_Y^2 + 16g_\chi^2 \left[1 + \frac{v_\chi^2}{v_W^2} \right] \pm \sqrt{\left(g^2 + g_Y^2 - 16g_\chi^2 \left[1 + \frac{v_\chi^2}{v_W^2} \right] \right)^2 + 4[4g_\chi g_Y \sin \theta_W + 4g_\chi g \cos \theta_W]^2} \right\}, \quad (4.36)$$

so to remain within the 2σ limit of the ρ parameter, one requires $v_\chi \gtrsim 7.7$ TeV (given $g_\chi \sim g_Y$).² The $Z - Z'$ mixing angle is given by,

$$\tan 2\phi_Z = \frac{2(g_\chi g_Y \sin \theta_W + g g_\chi \cos \theta_W)}{g^2 + g_Y^2 - 4g_\chi^2(1 + v_\chi^2/v_w^2)}, \quad (4.37)$$

so one finds $|\phi_Z| \simeq (v_w/v_\chi)^2 \lesssim 10^{-3}$ given $v_\chi \gtrsim 7.7$ TeV. We will see the consequences of this for our DM candidate below.

So far we have dealt with mixing coming from the mass matrix. As the theory contains particles charged under both $U(1)_Y$ and $U(1)_\chi$, kinetic mixing between the respective field strength tensors will be generated radiatively. The full mass matrix for the Z and Z' bosons, taking into account both kinetic and mass mixing, can then be found [109]. By following this procedure we have checked that including such a term makes no qualitative difference to our discussion above. Furthermore one typically expects radiatively generated kinetic mixing to be small [110] and therefore negligible for mixing in this model.

There is also a competing limit from the LHC. For example, using the ATLAS limit on dilepton resonances [111] and a choice of $g_\chi = 0.1$, one finds a lower bound of $M_{Z'} \gtrsim 2.7$ TeV. Given that $M_{Z'} \simeq 2g_\chi v_\chi$, this corresponds to $|\phi_Z| \lesssim 6 \times 10^{-4}$. Larger g_χ means stronger collider limits on $M_{Z'}$. However, ATLAS and CMS currently only provide limits for Z' resonances up to 3.5 TeV [111, 112], and as the limit on the possible signal strength is rising for high mass regions, we do not pursue the details here. Suffice to say the bound on the mixing angle is approximately $|\phi_Z| \lesssim 10^{-3}$.

4.5 Dark matter

4.5.1 DM candidate mass

In the Ma radiative ISS, there exists an accidental Z_2 symmetry under which only η_1 and S_{2L} transform. The lightest of these species is stable and forms a DM candidate. By examining the scalar potential in eq. (4.8) we see that the masses of the real and imaginary components of η_1 after symmetry breaking are given by,

$$m_R^2 = (\mu_{\eta_1}^2 + \lambda_{1\phi} v_w^2/2 + \lambda_{12} v_\chi^2/2)^2 + \sqrt{2} A v_\chi \quad (4.38)$$

$$m_I^2 = (\mu_{\eta_1}^2 + \lambda_{1\phi} v_w^2/2 + \lambda_{12} v_\chi^2/2)^2 - \sqrt{2} A v_\chi, \quad (4.39)$$

where $A > 0$ and for reasons of naturalness one typically expects $\mu_{\eta_1} \sim v_\chi \sim A$. The fermion S_{2L} gains a radiative Majorana mass, M_2 , at 1-loop order, and one typically expects,

$$m_I \gg M_2, \quad (4.40)$$

if fine-tuning is absent. To provide a sense of the typical values for M_2 , we have plotted M_2 as a function of $\sqrt{A v_\chi}$, with $M_1 = \sqrt{A v_\chi}$ for different choices of m_I in figure 10. One can see that as one moves away from the fine tuned limit $m_I = 0$, the radiative mass M_2 decreases to be in the range,

$$M_2 \approx (10^{-2} - 10^{-3})(f_{12})^2 v_\chi. \quad (4.41)$$

²The constraint can also be satisfied with $g_\chi \lesssim g_Y/95$ and $v_\chi \lesssim 5$ TeV. Given the GUT origin of the model, we consider such small choices for g_χ unattractive. Furthermore the interactions of the resultant light Z' are constrained to be so weak by observations of Caesium transitions [106–108], that they would also lead to overclosure of the universe.




Figure 10. $M_2/(f_{12})^2$ as a function of $\sqrt{Av_\chi}$, with $M_1 = \sqrt{Av_\chi}$, for different choices of m_I .

4.5.2 Cold DM scenario

Given the discussion of the ρ parameter one requires $v_\chi \gg v_w$. We will discuss the DM relic abundance in light of this constraint here. Let us assume the S_{2L} species have a thermal abundance in the early universe, as will be the case unless $g_\chi \lesssim 10^{-8} \ll g_Y$. There is furthermore an absence of any interactions of the form $S_{2L}S_{2L}(\sigma)$ where σ represents any of the scalar fields in the theory. As S_{2L} is charged only under $U(1)_\chi$, it can annihilate only through the Z' , and due to mixing of the gauge bosons, through the Z boson [113] (see figure 11).




Figure 11. Annihilation process of S_{2L} which determines its relic density.

So as not to overclose the universe one requires the annihilation cross section to be larger than $\langle\sigma|v|\rangle_{DM} \approx 1.9 \times 10^{-9} \text{ GeV}^{-2}$ [69]. However annihilation through the Z boson is suppressed by the small $Z - Z'$ mixing angle. For example, consider the $2M_2 \lesssim M_Z$ regime. It suffices for our purposes to use a simple estimate for the annihilation cross section:

$$\langle\sigma|v|\rangle \approx \frac{\sin^2 \phi_Z G_F^2 M_2^2}{2\pi}, \quad (4.42)$$

where ϕ_Z is the $Z - Z'$ mixing angle and G_F is the Fermi constant. Requiring $\langle \sigma|v| \rangle \gtrsim \langle \sigma|v| \rangle_{DM}$, and substituting for the mixing angle $\phi_Z \simeq (v_w/v_\chi)^2$, one finds:

$$M_2 \gtrsim \frac{\sqrt{2\pi\langle \sigma|v| \rangle_{DM}}}{G_F} \left(\frac{v_\chi}{v_w} \right)^2. \quad (4.43)$$

Given the constraint from the ρ parameter, $v_\chi \gtrsim 7$ TeV, this demands $M_2 \gtrsim 7.5$ TeV, which contradicts our original assumption of $2M_2 \lesssim M_Z$. One also finds a too small cross section in the $2M_2 \gtrsim M_Z$ regime.

Annihilations occurring through the Z' are suppressed by the large mass $M_{Z'}$. So annihilations through the Z' are also too weak.

Finally we note that the cross section will be much higher at a resonance point, where $2M_2 \simeq M_Z$ or $2M_2 \simeq M_{Z'}$ [113], and this could avoid overclosure. However, this corresponds to a fine tuning of the DM mass.

From this discussion we see that for a simple cold DM scenario the S_{2L} DM candidate overcloses the universe if we demand no fine tuning. One would have to go to more convoluted DM scenarios such as warm dark matter to perhaps find viable regions of parameter space. The simple cold DM case is ruled out.

4.6 Concluding remarks on the Ma model

The Ma radiative ISS model has several positive features for baryogenesis and washout avoidance. Resonant leptogenesis may be accommodated, as the mass splitting of the heavy neutrinos is generated before the EW phase transition, allowing the interplay between heavy neutrino decays and sphalerons crucial for leptogenesis. Also, washout due to the other ΔL interactions required for the radiative ISS can be suppressed with suitable choices of mass parameters.

On the other hand we have also seen a drawback of the Ma model with respect to cold DM. We found that the constraints from measurements of the ρ parameter translated into DM annihilations through the Z and Z' gauge bosons which are too weak to avoid overclosure. These simple cold DM scenarios are therefore ruled out for the Ma radiative ISS (apart for possibly small areas of parameter space corresponding to resonant enhancements). One would therefore have to go to more convoluted scenarios such as warm dark matter to find a suitable DM scenario.

5 Conclusion

Motivated originally by how new physics explanations for the baryon asymmetry, dark matter, and neutrino oscillations can overlap with each other, we began by looking at the properties of the inverse seesaw. We saw how the inverse seesaw mechanism offers various advantages, namely a scale low enough for direct experimental tests, and the possibility of resonant leptogenesis, or at least a suppressed washout rate if some other mechanism is responsible for the baryon asymmetry.

The inverse seesaw mechanism does come with a cost: the small lepton number violating parameter that must enter the mass matrix. Despite its technical naturalness, there may be a more appealing mechanism to generate it. One of the options is a radiative origin. We therefore continued our investigation by focusing on two radiative inverse seesaw models.

It was found in the case of the Law/McDonald radiative inverse seesaw model, the DM abundance is easily explained with the Higgs portal mechanism, or the double Higgs portal.

The advantages of the inverse seesaw with respect to baryogenesis, however, are lost due to the extra fields present which break lepton number explicitly with their interactions. In case of the Ma radiative inverse seesaw model, the situation is reversed: it nicely accommodates resonant leptogenesis, but does not feature a simple cold dark matter candidate.

Acknowledgements

We thank S. S. C. Law and K. L. McDonald for their useful comments. IB was supported by the Commonwealth of Australia. NFB, KP, and RRV were supported by the Australian Research Council. KP was also supported by the Netherlands Foundation for Fundamental Research of Matter (FOM) and the Netherlands Organisation for Scientific Research (NWO).

References

- [1] S. Weinberg, *Baryon and Lepton Nonconserving Processes*, *Phys. Rev. Lett.* **43** (1979) 1566–1570.
- [2] M. Fukugita and T. Yanagida, *Baryogenesis without grand unification*, *Phys. Lett.* **B174** (1986), no. 1 45 – 47.
- [3] S. Dodelson and L. M. Widrow, *Sterile-neutrinos as dark matter*, *Phys. Rev. Lett.* **72** (1994) 17–20, [[hep-ph/9303287](#)].
- [4] X.-D. Shi and G. M. Fuller, *A New dark matter candidate: Nonthermal sterile neutrinos*, *Phys. Rev. Lett.* **82** (1999) 2832–2835, [[astro-ph/9810076](#)].
- [5] A. Kusenko, *Sterile neutrinos, dark matter, and the pulsar velocities in models with a Higgs singlet*, *Phys. Rev. Lett.* **97** (2006) 241301, [[hep-ph/0609081](#)].
- [6] K. Petraki and A. Kusenko, *Dark-matter sterile neutrinos in models with a gauge singlet in the Higgs sector*, *Phys. Rev.* **D77** (2008) 065014, [[arXiv:0711.4646](#)].
- [7] A. Kusenko, F. Takahashi, and T. T. Yanagida, *Dark Matter from Split Seesaw*, *Phys. Lett.* **B693** (2010) 144–148, [[arXiv:1006.1731](#)].
- [8] L. Canetti, M. Drewes, and M. Shaposhnikov, *Sterile Neutrinos as the Origin of Dark and Baryonic Matter*, [arXiv:1204.3902](#).
- [9] L. Canetti, M. Drewes, T. Frossard, and M. Shaposhnikov, *Dark Matter, Baryogenesis and Neutrino Oscillations from Right Handed Neutrinos*, [arXiv:1208.4607](#).
- [10] I. Affleck and M. Dine, *A new mechanism for baryogenesis*, *Nucl. Phys.* **B249** (1985), no. 2 361 – 380.
- [11] M. Dine, L. Randall, and S. D. Thomas, *Baryogenesis from flat directions of the supersymmetric standard model*, *Nucl. Phys.* **B458** (1996) 291–326, [[hep-ph/9507453](#)].
- [12] A. Riotto and M. Trodden, *Recent progress in baryogenesis*, *Ann. Rev. Nucl. Part. Sci.* **49** (1999) 35–75, [[hep-ph/9901362](#)].
- [13] F. R. Klinkhamer and N. S. Manton, *A saddle-point solution in the weinberg-salam theory*, *Phys. Rev.* **D30** (Nov, 1984) 2212–2220.
- [14] V. Kuzmin, V. Rubakov, and M. Shaposhnikov, *On anomalous electroweak baryon-number non-conservation in the early universe*, *Phys. Lett.* **B155** (1985), no. 12 36 – 42.
- [15] P. Minkowski, *$\mu \rightarrow e\gamma$ at a Rate of One Out of 10^9 Muon Decays?*, *Phys. Lett.* **B67** (1977) 421.
- [16] T. Yanagida, *Horizontal symmetry and masses of neutrinos*, *Conf. Proc.* **C7902131** (1979) 95.

- [17] M. Gell-Mann, P. Ramond, and R. Slansky, *Complex spinors and unified theories*, *Conf. Proc.* **C790927** (1979) 315–321.
- [18] R. N. Mohapatra and G. Senjanovic, *Neutrino Mass and Spontaneous Parity Violation*, *Phys. Rev. Lett.* **44** (1980) 912.
- [19] M. Magg and C. Wetterich, *Neutrino mass problem and gauge hierarchy*, *Phys. Lett.* **B94** (1980) 61.
- [20] J. Schechter and J. Valle, *Neutrino Masses in $SU(2) \otimes U(1)$ Theories*, *Phys. Rev.* **D22** (1980) 2227.
- [21] C. Wetterich, *Neutrino Masses and the Scale of B-L Violation*, *Nucl. Phys.* **B187** (1981) 343.
- [22] G. Lazarides, Q. Shafi, and C. Wetterich, *Proton Lifetime and Fermion Masses in an $SO(10)$ Model*, *Nucl. Phys.* **B181** (1981) 287.
- [23] R. N. Mohapatra and G. Senjanovic, *Neutrino Masses and Mixings in Gauge Models with Spontaneous Parity Violation*, *Phys. Rev.* **D23** (1981) 165.
- [24] T. Cheng and L.-F. Li, *Neutrino Masses, Mixings and Oscillations in $SU(2) \times U(1)$ Models of Electroweak Interactions*, *Phys. Rev.* **D22** (1980) 2860.
- [25] R. Foot, H. Lew, X. He, and G. C. Joshi, *Seesaw neutrino masses induced by a triplet of leptons*, *Z. Phys.* **C44** (1989) 441.
- [26] R. N. Mohapatra and J. W. F. Valle, *Neutrino mass and baryon-number nonconservation in superstring models*, *Phys. Rev.* **D34** (Sep, 1986) 1642–1645.
- [27] J. Bernabu, A. Santamaria, J. Vidal, A. Mendez, and J. Valle, *Lepton flavour non-conservation at high energies in a superstring inspired standard model*, *Phys. Lett.* **B187** (1987), no. 34 303 – 308.
- [28] S. Blanchet, T. Hambye, and F.-X. Josse-Michaux, *Reconciling leptogenesis with observable $\mu \rightarrow e\gamma$ rates*, *JHEP* **1004** (2010) 023, [[arXiv:0912.3153](#)].
- [29] E. Ma, *Radiative inverse seesaw mechanism for nonzero neutrino mass*, *Phys. Rev.* **D80** (2009) 013013, [[arXiv:0904.4450](#)].
- [30] S. Khalil, *TeV-scale gauged B-L symmetry with inverse seesaw mechanism*, *Phys. Rev.* **D82** (2010) 077702, [[arXiv:1004.0013](#)].
- [31] F. Bazzocchi, *Minimal Dynamical Inverse See Saw*, *Phys. Rev.* **D83** (2011) 093009, [[arXiv:1011.6299](#)].
- [32] S. S. C. Law and K. L. McDonald, *Inverse seesaw and dark matter in models with exotic lepton triplets*, *Phys. Lett.* **B713** (2012), no. 45 490 – 494, [[arXiv:1204.2529](#)].
- [33] G. Guo, X.-G. He, and G.-N. Li, *Radiative Two Loop Inverse Seesaw and Dark Matter*, *JHEP* **1210** (2012) 044, [[arXiv:1207.6308](#)].
- [34] S. C. Park, K. Wang, and T. T. Yanagida, *Neutrino mass from a hidden world and its phenomenological implications*, *Phys. Lett.* **B685** (2010) 309–312, [[arXiv:0909.2937](#)].
- [35] C. S. Fong, R. N. Mohapatra, and I. Sung, *Majorana Neutrinos from Inverse Seesaw in Warped Extra Dimension*, *Phys. Lett.* **B704** (2011) 171–178, [[arXiv:1107.4086](#)].
- [36] A. Dias, C. de S. Pires, and P. R. da Silva, *How the Inverse See-Saw Mechanism Can Reveal Itself Natural, Canonical and Independent of the Right-Handed Neutrino Mass*, *Phys. Rev.* **D84** (2011) 053011, [[arXiv:1107.0739](#)].
- [37] A. Dias, C. de S. Pires, P. Rodrigues da Silva, and A. Sampieri, *A Simple Realization of the Inverse Seesaw Mechanism*, *Phys. Rev.* **D86** (2012) 035007, [[arXiv:1206.2590](#)].
- [38] F.-X. Josse-Michaux and E. Molinaro, *A Common Framework for Dark Matter, Leptogenesis and Neutrino Masses*, *Phys. Rev.* **D84** (2011) 125021, [[arXiv:1108.0482](#)].

- [39] S. S. C. Law and K. L. McDonald, *Generalized Inverse Seesaws*, [arXiv:1303.4887](#).
- [40] P. B. Dev and A. Pilaftsis, *Minimal Radiative Neutrino Mass Mechanism for Inverse Seesaw Models*, [arXiv:1209.4051](#).
- [41] P. B. Dev, R. Franceschini, and R. Mohapatra, *Bounds on TeV Seesaw Models from LHC Higgs Data*, [arXiv:1207.2756](#).
- [42] A. Das and N. Okada, *Inverse Seesaw Neutrino Signatures at LHC and ILC*, [arXiv:1207.3734](#).
- [43] A. Abada, C. Biggio, F. Bonnet, M. Gavela, and T. Hambye, *Low energy effects of neutrino masses*, *JHEP* **0712** (2007) 061, [[arXiv:0707.4058](#)].
- [44] M. Malinsky, T. Ohlsson, Z.-z. Xing, and H. Zhang, *Non-unitary neutrino mixing and CP violation in the minimal inverse seesaw model*, *Phys. Lett.* **B679** (2009) 242–248, [[arXiv:0905.2889](#)].
- [45] F. del Aguila, J. de Blas, and M. Perez-Victoria, *Effects of new leptons in Electroweak Precision Data*, *Phys. Rev.* **D78** (2008) 013010, [[arXiv:0803.4008](#)].
- [46] H. Zhang and S. Zhou, *The minimal seesaw model at the tev scale*, *Phys. Lett.* **B685** (2010), no. 45 297 – 301.
- [47] J. Lopez-Pavon, S. Pascoli, and C.-f. Wong, *Can heavy neutrinos dominate neutrinoless double beta decay?*, [arXiv:1209.5342](#).
- [48] A. Ibarra, E. Molinaro, and S. Petcov, *TeV Scale See-Saw Mechanisms of Neutrino Mass Generation, the Majorana Nature of the Heavy Singlet Neutrinos and $(\beta\beta)_{0\nu}$ -Decay*, *JHEP* **1009** (2010) 108, [[arXiv:1007.2378](#)].
- [49] E. W. Kolb and M. S. Turner, *The Early universe*, *Front. Phys.* **69** (1990) 1–547.
- [50] S. Antusch, J. P. Baumann, and E. Fernandez-Martinez, *Non-Standard Neutrino Interactions with Matter from Physics Beyond the Standard Model*, *Nucl. Phys.* **B810** (2009) 369–388, [[arXiv:0807.1003](#)].
- [51] A. Y. Smirnov and R. Zukanovich Funchal, *Sterile neutrinos: Direct mixing effects versus induced mass matrix of active neutrinos*, *Phys. Rev.* **D74** (2006) 013001, [[hep-ph/0603009](#)].
- [52] S. Antusch, M. Blennow, E. Fernandez-Martinez, and J. Lopez-Pavon, *Probing non-unitary mixing and CP-violation at a Neutrino Factory*, *Phys. Rev.* **D80** (2009) 033002, [[arXiv:0903.3986](#)].
- [53] M. Garny, A. Kartavtsev, and A. Hohenegger, *Leptogenesis from first principles in the resonant regime*, [arXiv:1112.6428](#).
- [54] S. Blanchet, P. B. Dev, and R. Mohapatra, *Leptogenesis with TeV Scale Inverse Seesaw in $SO(10)$* , *Phys. Rev.* **D82** (2010) 115025, [[arXiv:1010.1471](#)].
- [55] S. Davidson and M. Elmer, *Similar Dark Matter and Baryon abundances with TeV-scale Leptogenesis*, *JHEP* **1210** (2012) 148, [[arXiv:1208.0551](#)].
- [56] A. Sakharov, *Violation of CP Invariance, C Asymmetry, and Baryon Asymmetry of the Universe*, *Pisma Zh. Eksp. Teor. Fiz.* **5** (1967) 32–35.
- [57] V. Silveira and A. Zee, *SCALAR PHANTOMS*, *Phys. Lett.* **B161** (1985) 136.
- [58] J. McDonald, *Gauge singlet scalars as cold dark matter*, *Phys. Rev.* **D50** (1994) 3637–3649, [[hep-ph/0702143](#)].
- [59] C. Burgess, M. Pospelov, and T. ter Veldhuis, *The Minimal model of nonbaryonic dark matter: A Singlet scalar*, *Nucl. Phys.* **B619** (2001) 709–728, [[hep-ph/0011335](#)].
- [60] K. Cheung, Y.-L. S. Tsai, P.-Y. Tseng, T.-C. Yuan, and A. Zee, *Global Study of the Simplest Scalar Phantom Dark Matter Model*, *JCAP* **1210** (2012) 042, [[arXiv:1207.4930](#)].

- [61] M. Aoki, S. Kanemura, and O. Seto, *Multi-Higgs portal dark matter under the CDMS II results*, *Phys. Lett.* **B685** (2010) 313–317, [[arXiv:0912.5536](#)].
- [62] Y. Cai, X.-G. He, and B. Ren, *Low Mass Dark Matter and Invisible Higgs Width In Darkon Models*, *Phys. Rev.* **D83** (2011) 083524, [[arXiv:1102.1522](#)].
- [63] X.-G. He, B. Ren, and J. Tandean, *Hints of Standard Model Higgs Boson at the LHC and Light Dark Matter Searches*, *Phys. Rev.* **D85** (2012) 093019, [[arXiv:1112.6364](#)].
- [64] L. Lopez-Honorez, T. Schwetz, and J. Zupan, *Higgs portal, fermionic dark matter, and a Standard Model like Higgs at 125 GeV*, *Phys. Lett.* **B716** (2012) 179–185, [[arXiv:1203.2064](#)].
- [65] M. Gonderinger, H. Lim, and M. J. Ramsey-Musolf, *Complex scalar singlet dark matter: Vacuum stability and phenomenology*, *Phys. Rev.* **D86** (Aug, 2012) 043511.
- [66] S. Profumo, M. J. Ramsey-Musolf, and G. Shaughnessy, *Singlet Higgs phenomenology and the electroweak phase transition*, *JHEP* **0708** (2007) 010, [[arXiv:0705.2425](#)].
- [67] V. Barger, P. Langacker, M. McCaskey, M. J. Ramsey-Musolf, and G. Shaughnessy, *Cern lhc phenomenology of an extended standard model with a real scalar singlet*, *Phys. Rev.* **D77** (Feb, 2008) 035005.
- [68] **WMAP Collaboration**, E. Komatsu et al., *Seven-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation*, *Astrophys. J. Suppl.* **192** (2011) 18, [[arXiv:1001.4538](#)].
- [69] G. Steigman, B. Dasgupta, and J. F. Beacom, *Precise Relic WIMP Abundance and its Impact on Searches for Dark Matter Annihilation*, *Phys. Rev.* **D86** (2012) 023506, [[arXiv:1204.3622](#)].
- [70] Y. Mambrini, *Higgs searches and singlet scalar dark matter: Combined constraints from xenon 100 and the lhc*, *Phys. Rev.* **D84** (Dec, 2011) 115017.
- [71] **MILC Collaboration**, D. Toussaint and W. Freeman, *The Strange quark condensate in the nucleon in 2+1 flavor QCD*, *Phys. Rev. Lett.* **103** (2009) 122002, [[arXiv:0905.2432](#)].
- [72] R. D. Young and A. W. Thomas, *Recent results on nucleon sigma terms in lattice QCD*, *Nucl. Phys.* **A844** (2010) 266C–271C, [[arXiv:0911.1757](#)].
- [73] J. M. Alarcon, J. Martin Camalich and J. A. Oller, *The chiral representation of the πN scattering amplitude and the pion-nucleon sigma term*, *Phys. Rev.* **D 85** (2012) 051503, [[arXiv:1110.3797](#)].
- [74] J. M. Alarcon, L. S. Geng, J. M. Camalich and J. A. Oller, *On the strangeness content of the nucleon*, [[arXiv:1209.2870](#)].
- [75] **XENON100 Collaboration**, E. Aprile et al., *Dark Matter Results from 225 Live Days of XENON100 Data*, [[arXiv:1207.5988](#)].
- [76] **XENON1T Collaboration**, E. Aprile, *The XENON1T Dark Matter Search Experiment*, [[arXiv:1206.6288](#)].
- [77] P. Bandyopadhyay, E. J. Chun, H. Okada, and J.-C. Park, *Higgs Signatures in Inverse Seesaw Model at the LHC*, [[arXiv:1209.4803](#)].
- [78] C. Englert, T. Plehn, M. Rauch, D. Zerwas, and P. M. Zerwas, *LHC: Standard Higgs and Hidden Higgs*, *Phys. Lett.* **B707** (2012) 512–516, [[arXiv:1112.3007](#)].
- [79] A. Djouadi, O. Lebedev, Y. Mambrini, and J. Quevillon, *Implications of LHC searches for Higgs–portal dark matter*, *Phys. Lett.* **B709** (2012) 65–69, [[arXiv:1112.3299](#)].
- [80] A. Djouadi, A. Falkowski, Y. Mambrini, and J. Quevillon, *Direct Detection of Higgs-Portal Dark Matter at the LHC*, [[arXiv:1205.3169](#)].

- [81] **CMS Collaboration**, S. Chatrchyan et al., *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett.* **B716** (2012) 30–61, [[arXiv:1207.7235](#)].
- [82] **ATLAS Collaboration**, G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett.* **B716** (2012) 1–29, [[arXiv:1207.7214](#)].
- [83] **CMS Collaboration**, *Combination of standard model Higgs boson searches and measurements of the properties of the new boson with a mass near 125 GeV*, *CMS-PAS-HIG-12-045*, .
- [84] **ATLAS Collaboration**, *Combined coupling measurements of the higgs-like boson with the atlas detector using up to 25 fb⁻¹ of proton-proton collision data*, Tech. Rep. ATLAS-CONF-2013-034, CERN, Geneva, Mar, 2013.
- [85] J. Ellis and T. You, *Updated Global Analysis of Higgs Couplings*, [arXiv:1303.3879](#).
- [86] A. Falkowski, F. Riva, and A. Urbano, *Higgs At Last*, [arXiv:1303.1812](#).
- [87] P. P. Giardino, K. Kannike, I. Masina, M. Raidal, and A. Strumia, *The universal Higgs fit*, [arXiv:1303.3570](#).
- [88] A. Djouadi and G. Moreau, *The couplings of the Higgs boson and its CP properties from fits of the signal strengths and their ratios at the 7+8 TeV LHC*, [arXiv:1303.6591](#).
- [89] **CMS Collaboration**, S. Chatrchyan et al., *Search for a standard-model-like Higgs boson with a mass of up to 1 TeV at the LHC*, [arXiv:1304.0213](#).
- [90] M. Bowen, Y. Cui, and J. D. Wells, *Narrow trans-TeV Higgs bosons and $H \rightarrow hh$ decays: Two LHC search paths for a hidden sector Higgs boson*, *JHEP* **0703** (2007) 036, [[hep-ph/0701035](#)].
- [91] S. Nussinov, *Technoc cosmology: Could a technibaryon excess provide a ‘natural’ missing mass candidate?*, *Phys. Lett.* **B165** (1985) 55.
- [92] H. Davoudiasl, D. E. Morrissey, K. Sigurdson, and S. Tulin, *Hylogenesis: A unified origin for baryonic visible matter and antibaryonic dark matter*, *Phys. Rev. Lett.* **105** (2010) 211304, [[arXiv:1008.2399](#)].
- [93] B. von Harling, K. Petraki, and R. R. Volkas, *Affleck-Dine dynamics and the dark sector of pangenesis*, *JCAP* **1205** (2012) 021, [[arXiv:1201.2200](#)].
- [94] N. F. Bell, K. Petraki, I. M. Shoemaker, and R. R. Volkas, *Pangenesis in a Baryon-Symmetric Universe: Dark and Visible Matter via the Affleck-Dine Mechanism*, *Phys. Rev.* **D84** (2011) 123505, [[arXiv:1105.3730](#)].
- [95] C. Cheung and K. M. Zurek, *Affleck-Dine Cogenesis*, *Phys. Rev.* **D84** (2011) 035007, [[arXiv:1105.4612](#)].
- [96] S. Dodelson and L. M. Widrow, *Baryon-symmetric baryogenesis*, *Phys. Rev. Lett.* **64** (Jan, 1990) 340–343.
- [97] V. A. Kuzmin, *A Simultaneous solution to baryogenesis and dark matter problems*, *Phys. Part. Nucl.* **29** (1998) 257–265, [[hep-ph/9701269](#)].
- [98] R. Kitano and I. Low, *Dark matter from baryon asymmetry*, *Phys. Rev.* **D71** (2005) 023510, [[hep-ph/0411133](#)].
- [99] P.-H. Gu, M. Lindner, U. Sarkar, and X. Zhang, *WIMP Dark Matter and Baryogenesis*, *Phys. Rev.* **D83** (2011) 055008, [[arXiv:1009.2690](#)].
- [100] P.-H. Gu, *Origin of matter in the universe*, *Phys. Lett.* **B657** (2007) 103–106, [[arXiv:0706.1946](#)].

- [101] J. J. Heckman and S.-J. Rey, *Baryon and Dark Matter Genesis from Strongly Coupled Strings*, *JHEP* **1106** (2011) 120, [[arXiv:1102.5346](#)].
- [102] G. R. Farrar and G. Zaharijas, *Dark matter and the baryon asymmetry*, *Phys. Rev. Lett.* **96** (2006) 041302, [[hep-ph/0510079](#)].
- [103] K. Petraki, M. Trodden, and R. R. Volkas, *Visible and dark matter from a first-order phase transition in a baryon-symmetric universe*, *JCAP* **1202** (2012) 044, [[arXiv:1111.4786](#)].
- [104] **Planck Collaboration**, P. Ade et al., *Planck 2013 results. XVI. Cosmological parameters*, [arXiv:1303.5076](#).
- [105] **Particle Data Group**, J. Beringer et al., *Review of particle physics*, *Phys. Rev.* **D86** (Jul, 2012) 010001.
- [106] C. Wood, S. Bennett, D. Cho, B. Masterson, J. Roberts, et al., *Measurement of parity nonconservation and an anapole moment in cesium*, *Science* **275** (1997) 1759–1763.
- [107] J. Ginges and V. Flambaum, *Violations of fundamental symmetries in atoms and tests of unification theories of elementary particles*, *Physics Reports* **397** (2004), no. 2 63 – 154.
- [108] C. Bouchiat and P. Fayet, *Constraints on the parity-violating couplings of a new gauge boson*, *Phys. Lett.* **B608** (2005) 87–94, [[hep-ph/0410260](#)].
- [109] R. Foot and X.-G. He, *Comment on Z-Z' mixing in extended gauge theories*, *Phys. Lett.* **B267** (1991), no. 4 509 – 512.
- [110] M. Baumgart, C. Cheung, J. T. Ruderman, L.-T. Wang, and I. Yavin, *Non-Abelian Dark Sectors and Their Collider Signatures*, *JHEP* **0904** (2009) 014, [[arXiv:0901.0283](#)].
- [111] *Search for high-mass dilepton resonances in 20 fb⁻¹ of pp collisions at $\sqrt{s} = 8$ tev with the atlas experiment*, Tech. Rep. ATLAS-CONF-2013-017, CERN, Geneva, Mar, 2013.
- [112] *Search for resonances in the dilepton mass distribution in pp collisions at $\sqrt{s} = 8$ tev*, Tech. Rep. CMS-PAS-EXO-12-061, CERN, Geneva, 2013.
- [113] D. E. Brahm and L. J. Hall, *U(1)' dark matter*, *Phys. Rev.* **D41** (Feb, 1990) 1067–1076.