

# When the Higgs meets the Top: Search for $t \rightarrow ch^0$ at the LHC

Kai-Feng Chen<sup>a</sup>, Wei-Shu Hou<sup>a</sup>, Chung Kao<sup>a,b</sup>, and Masaya Kohda<sup>a</sup>

<sup>a</sup>*Department of Physics, National Taiwan University, Taipei, Taiwan 10617*

<sup>b</sup>*Homer L. Dodge Department of Physics and Astronomy,  
University of Oklahoma, Norman, OK 73019, USA*

The newly discovered “Higgs” boson  $h^0$ , being lighter than the top quark  $t$ , opens up new probes for flavor and mass generation. In the general two Higgs doublet model, new  $ct$ ,  $cc$  and  $tt$  Yukawa couplings could modify  $h^0$  properties. If  $t \rightarrow ch^0$  occurs at the percent level, the observed  $ZZ^*$  and  $\gamma\gamma$  signal events may have accompanying  $cbW$  activity coming from  $t\bar{t}$  feeddown. We suggest that  $t \rightarrow ch^0$  can be searched for via  $h^0 \rightarrow ZZ^*$ ,  $\gamma\gamma$ ,  $WW^*$  and  $b\bar{b}$ , perhaps even  $\tau^+\tau^-$  modes in  $t\bar{t}$  events. Existing data might be able to reveal some clues for  $t \rightarrow ch^0$  signature, or push the branching ratio  $\mathcal{B}(t \rightarrow ch^0)$  down to below the percent level.

**PACS numbers:** 12.15.Mm, 12.60.Fr, 14.65.Ha, 14.80.Ec

## I. INTRODUCTION AND MOTIVATION

With the landmark discovery [1, 2] of a 126 GeV boson in 2012 by the ATLAS and CMS experiments at the Large Hadron Collider (LHC), efforts have shifted towards Higgs property studies, to either confirm that this is indeed the Higgs boson of the Standard Model (SM), or to find deviations indicating that it may not be *the* Higgs boson. With the Higgs boson as the “mass giver”, it is natural to ask whether it reveals any special effects associated with the heaviness of the top quark. A related question is, analogous to the generation repetition seen with fermions, if we have seen one Higgs boson, could there be others.

We wish to explore a possible  $tch^0$  coupling, where  $h^0$  stands for the 126 GeV boson. Absent within SM, this coupling has not been the focus of attention. At the one loop level, the SM branching ratio (for  $m_{h^0} = 115$  GeV)  $\mathcal{B}(t \rightarrow ch^0) \simeq 3 \times 10^{-15}$  [3] is extremely suppressed. But with  $m_{h^0} < m_t$ ,  $t \rightarrow ch^0$  decay [4–7] can readily be searched for in  $t\bar{t}$  production. In the post-discovery era, we must search for  $t \rightarrow ch^0$  at the LHC as part of the Higgs, and top, property programs.

The  $B \rightarrow D^{(*)}\tau\nu$  “anomaly” uncovered by the BaBar experiment [8] suggests the need of a multi-Higgs boson sector. The  $B \rightarrow D^{(*)}\tau\nu$  rates are found to deviate from SM expectations, with combined significance over  $3\sigma$ . More intriguing, it cannot [8] be explained by the usual two Higgs doublet model (2HDM), type II, the popular Higgs extension realized in minimal supersymmetry (SUSY). One explanation is to adopt [9, 10] a general 2HDM, called 2HDM-III, allowing for flavor changing neutral Higgs (FCNH) couplings that modify the charged Higgs couplings, which could resolve the BaBar anomaly. The Natural Flavor Conservation (NFC) condition was proposed over 35 years ago [11] to forbid such couplings, but the FCNH couplings of Ref. [10] go even beyond the Cheng-Sher ansatz [12], an approach designed to tame FCNH couplings involving lighter quarks. One basically lets data decide the strength of possible FCNH couplings, which cannot be argued against. Note that it was within the Cheng-Sher ansatz that  $t \rightarrow ch^0$  (or  $h^0 \rightarrow t\bar{c}$ ) decay

was first proposed [4] as the leading effect.

To account for the BaBar anomaly, the FCNH coupling  $\rho_{ct}$  of the exotic heavy Higgs doublet need to be of order 1 in strength [10]. For our purpose, keeping notation of the usual 2HDM-II [13], the observed boson  $h^0$  may contain a small admixture  $\cos(\beta - \alpha)$  of the exotic neutral Higgs, hence the  $tch^0$  coupling [14]

$$\rho_{ct} \cos(\beta - \alpha) \bar{c}th^0 + \text{h.c.}, \quad (1)$$

which can induce  $t \rightarrow ch^0$  decay. In Fig. 1 we illustrate the branching ratio  $\mathcal{B}(t \rightarrow ch^0)$  vs  $\rho_{ct} \cos(\beta - \alpha)$ . The question is: Considering all available data, what is the allowed  $\mathcal{B}(t \rightarrow ch^0)$ , or equivalently,  $\rho_{ct} \cos(\beta - \alpha)$  value? What are the signatures to pursue?

We note that, if we take  $\rho_{ct} \sim 1$ , which is not quite explored because it is suppressed by  $\cos(\beta - \alpha)$  for couplings involving  $h^0$  (but not suppressed for couplings involving the heavy exotic Higgs bosons  $H^0$ ,  $A^0$  and  $H^\pm$ ), then the analogous parameters  $\rho_{tt}$ ,  $\rho_{cc}$ ,  $\rho_{bb}$  and  $\rho_{\tau\tau}$  will enter the Higgs property study program, as we shall elucidate. An existing study of multi-lepton final states finds a bound [7] of  $\mathcal{B}(t \rightarrow ch^0) < 2.7\%$ . But this should be taken with some caution, as it assumed SM branching ratios for  $h^0 \rightarrow WW^*$ ,  $ZZ^*$  and  $\tau\tau$  final states. The study took an effective field theory approach to isolate the  $tch^0$  coupling, hence is not a full theory.




FIG. 1.  $\mathcal{B}(t \rightarrow ch^0)$  vs  $\rho_{ct} \cos(\beta - \alpha)$ , with 2% indicated.




FIG. 2. Allowed  $\rho_{\tau\tau}$ - $\rho_{ct}$  region for 2HDM-III to solve the  $B \rightarrow D^{(*)}\tau\nu$  anomaly, taking  $m_{H^+} = 700$  GeV (and  $\rho_{bb} = 0$ ). The shaded-green area is the combined result from  $\mathcal{R}(D)$  (solid-blue lines) and  $\mathcal{R}(D^*)$  (dashed-red lines), while the dotted-purple lines illustrate the  $h^0 \rightarrow \tau\tau$  bound by taking  $c_{\beta-\alpha} = 0.2$  in Eq. (3).

## II. BABAR ANOMALY, $h^0 \rightarrow \tau\tau$ , AND $b \rightarrow s\gamma$

The BaBar experiment measured the ratios  $\mathcal{R}(D^{(*)}) = \Gamma(\bar{B} \rightarrow D^{(*)}\tau\nu)/\Gamma(\bar{B} \rightarrow D^{(*)}\ell\nu)$ , finding them both larger than SM expectations, with a combined significance of  $3.4\sigma$ . In 2HDM-II, this implied [8] that  $\tan\beta/m_{H^+} = 0.44 \pm 0.02 \text{ GeV}^{-1}$  and  $0.75 \pm 0.04 \text{ GeV}^{-1}$  from  $\mathcal{R}(D)$  and  $\mathcal{R}(D^*)$ , respectively. The impressive precision is because many uncertainties, both from measurement and from theory, cancel. The two numbers are incompatible with each other, hence “excludes the type II 2HDM charged Higgs boson with a 99.8% confidence level for any value of  $\tan\beta/m_{H^+}$ ” [8]. Either  $\tan\beta/m_{H^+}$  value, however, would over-enhance [15]  $B \rightarrow \tau\nu$ , which is found in agreement (in order of magnitude) with SM expectation, spelling further trouble.

Employing FCNH parameters in 2HDM-III to account [9, 10] for the BaBar “anomaly”, a new  $c$ - $t$  coupling (and  $u$ - $t$  for  $B \rightarrow \tau\nu$ ), heretofore forbidden by the NFC condition [11], needs to be of order 1. For the lepton sector, Ref. [10] assumed the usual 2HDM-II coupling, i.e.  $\rho_{\tau\tau} = -\tan\beta\sqrt{2}m_\tau/v$ , where  $v$  is the weak scale.

We shall use the following notation for the Yukawa couplings of the 2HDM-III Higgs sector [16],

$$\begin{aligned}
 & -\frac{1}{\sqrt{2}} \sum_{f=u,d,\ell} \bar{f} [(\kappa^f s_{\beta-\alpha} + \rho^f c_{\beta-\alpha}) h^0 \\
 & + (\kappa^f c_{\beta-\alpha} - \rho^f s_{\beta-\alpha}) H^0 - i \text{sgn}(Q_f) \rho^f \gamma_5 A^0] f \\
 & - [\bar{u} (V \rho^d R - \rho^u V L) d H^+ + \bar{\nu} \rho^\ell R \ell H^+ + \text{h.c.}], \quad (2)
 \end{aligned}$$

where  $s_{\beta-\alpha}$  ( $c_{\beta-\alpha}$ ) stands for  $\sin(\beta-\alpha)$  ( $\cos(\beta-\alpha)$ ) in the 2HDM-II notation [13] for sake of comparison (even though  $\beta$  is no longer physical). The diagonal  $\kappa$  terms relate to mass generation, while the hermitian [14]  $\rho$  terms come from the second Higgs doublet, and can have off-diagonal terms allowed by data. It is the combined effect of  $\rho_{ct}$  and  $\rho_{\tau\tau}$ , both entering through the  $H^+$  couplings, that can account [10] for  $B \rightarrow D^{(*)}\tau\nu$ . Stringent constraints from down quark sector imply that only  $\rho_{bb}$  needs




FIG. 3. Constraint on  $\rho_{bb}$  from  $b \rightarrow s\gamma$ , assuming  $\rho_{ct} = 1$ ,  $\rho_{cc} = 0.2$  and  $m_{H^+} = 700$  GeV.

to be considered; note that the superscript for  $\rho$  can be dropped for flavor-specific elements. We shall only keep  $\rho_{\tau\tau}$  from lepton sector, but it is  $\rho_{ct}$ ,  $\rho_{tt}$  and  $\rho_{cc}$  from up quark sector that are of interest.

In the so-called “decoupling limit” [13, 17],  $\cos(\beta-\alpha) \rightarrow 0$  and  $\sin(\beta-\alpha) \rightarrow 1$ ,  $h^0$  becomes just the Higgs boson of SM, while  $H^0$ ,  $A^0$  and  $H^\pm$  form an exotic heavy scalar doublet with FCNH couplings. This limit was tacitly assumed in Ref. [10], which advocated  $H^0$ ,  $A^0 \rightarrow t\bar{c}$  search. But we entertain finite  $c_{\beta-\alpha}$  values for sake of  $t \rightarrow ch^0$  decay, hence would need to consider  $h^0 \rightarrow \tau\tau$  data. The latter from vector boson fusion (VBF) production is within a factor of 2 [18] from SM expectations,

$$|s_{\beta-\alpha} + (\rho_{\tau\tau} v / \sqrt{2} m_\tau) c_{\beta-\alpha}| \lesssim \sqrt{2}. \quad (3)$$

With  $\rho_{\tau\tau} \simeq -0.5$  [10], even a small  $c_{\beta-\alpha}$  could upset this bound. To illustrate further, we plot in Fig. 2 the range for  $\rho_{\tau\tau}$ - $\rho_{ct}$  allowed by BaBar anomaly for the typical value of  $m_{H^+} = 700$  GeV. The point  $\rho_{\tau\tau} \simeq -0.5$ ,  $\rho_{ct} \sim 1$  of Ref. [10], far outside the plot, would require  $c_{\beta-\alpha}$  to be rather small. If we take  $c_{\beta-\alpha} = 0.2$  (i.e.  $s_{\beta-\alpha} \simeq 0.98$ ) in Eq. (3), then  $-0.12 < \rho_{\tau\tau} < 0.02$  would push  $\rho_{ct}$  to become very large, as seen from Fig. 2.

Thus,  $h^0 \rightarrow \tau\tau$  data imply either one goes to the decoupling limit of  $c_{\beta-\alpha} \rightarrow 0$ , where  $t \rightarrow ch^0$  vanishes, or one has to entertain nonperturbative values for  $\rho_{ct}$  [19]. As further analysis [20] of  $q^2$  ( $\tau\nu$  pair mass) dependence of  $B \rightarrow D\tau\nu$  favors New Physics from spin-1 particles, we will not strongly advocate the link to the BaBar anomaly, but use it to illustrate that  $\rho_{ct}$  can be of order 1, and focus on probing  $tch^0$  coupling directly at the LHC.

Before turning to the LHC, however, we explore one other piece of B physics:  $b \rightarrow s\gamma$  (Fig. 3). We have already seen how  $h^0 \rightarrow \tau\tau$  data pushes down  $\rho_{\tau\tau}$ . We now show that, if one takes  $\rho_{ct} \sim 1$ , the well-known  $b \rightarrow s\gamma$  process constrains  $\rho_{bb}$  to be rather tiny (noted recently in Ref. [21] in a different way).

In the notation of Ref. [22], the  $H^+$  loop gives

$$\begin{aligned}
 \delta C_{7,8} \simeq & \frac{1}{3} \left( \rho_{tt} + \frac{V_{cs}^*}{V_{ts}^*} \rho_{ct} \right) \left( \rho_{tt}^* + \frac{V_{cb}}{V_{tb}} \rho_{ct}^* \right) \frac{F_{7,8}^{(1)}(y)}{2m_t^2/v^2} \\
 & - \left( \rho_{tt} + \frac{V_{cs}^*}{V_{ts}^*} \rho_{ct} \right) \rho_{bb} \frac{F_{7,8}^{(2)}(y)}{2m_t m_b / v^2}, \quad (4)
 \end{aligned}$$

evaluated at matching scale  $\mu_W \sim M_W$ , where  $y = m_t^2/M_{H^\pm}^2$  and  $F_{7,8}^{(1,2)}(y)$  are given in Ref. [22]. The effect through  $\rho_{bb}$  is enhanced by  $m_t/m_b$  as well as quark mixing elements, such that even a tiny  $\rho_{bb}$  could affect  $b \rightarrow s\gamma$ . We illustrate  $\rho_{bb}$  vs  $\rho_{tt}$  in Fig. 3, where we take  $\rho_{ct} = +1$ ,  $m_{H^\pm} = 700$  GeV, and constrain  $\mathcal{B}(B \rightarrow X_s\gamma)$  to be within 50% of SM expectation. The “wrong-sign”  $C_7^{\text{eff}}$  case has been included for comparison. Assuming  $C_7^{\text{eff}}$  does not change sign,  $|\rho_{bb}|$  is constrained to be considerably less than 0.01.

### III. PROBING $tch^0$ COUPLING AT LHC: GENERAL 2HDM-III

With  $\rho_{\tau\tau}$  and  $\rho_{bb}$  separately constrained to be small, we are still left with the up-sector exotic couplings, the FCNH  $\rho_{ct}$ , as well as the diagonal  $\rho_{cc}$  and  $\rho_{tt}$ , which we turn to constrain with present data. How large can  $\mathcal{B}(t \rightarrow ch^0)$  be when  $\rho_{ct} \sim 1$ ? I.e., what constraint do we have on  $\cos(\beta - \alpha)$ ? We now take a direct search approach, namely, from knowledge of top quark physics.

It is clear that  $\mathcal{B}_{ch} \equiv \mathcal{B}(t \rightarrow ch^0)$  cannot be too large, for otherwise we should have seen deviations in  $t\bar{t}$  measurements during the past two decades. The best measured  $t\bar{t}$  production cross section is  $\sigma_{t\bar{t}}^{\ell\ell} \simeq 162$  pb by CMS [23] at  $\sqrt{s} = 7$  TeV via dileptons, with an experimental error of 5%. On the theoretical side, studies are approaching NNLO (next-to-next-to-leading order) QCD corrections. Comparing the two calculations of Refs. [24] and [25], theoretical errors up to 10% appear allowed. However, a more recent full NNLO result [26] has reached much better theoretical control. In any case, given that  $\sigma_{t\bar{t}}^{\ell\ell}$  would be diluted by  $(1 - \mathcal{B}_{ch})^2$ ,  $\mathcal{B}_{ch}$  of order several percent seems still allowed.

A multi-lepton analysis of Ref. [7], based on 7 TeV data from CMS [27], gives a slightly more stringent bound of 2.7%, or  $[\sigma \cdot \mathcal{B}](pp \rightarrow t\bar{t} \rightarrow ch^0 bW) \lesssim 9$  pb (at 7 TeV). We have commented that this study assumed SM branching ratios for  $h^0$  decay, which should be taken with caution. However, it illustrates possible feeddown effects to observable Higgs boson decay modes, *given the large  $t\bar{t}$  production cross section at the LHC*. Our chief suggestion is to inspect the clean  $ZZ^* \rightarrow 4\ell$  samples of Higgs search data, which we now elaborate.

What has been observed so far at the LHC is

$$\sigma_{gg \rightarrow h^0} \cdot \frac{\Gamma_{h^0 \rightarrow ZZ^*}}{\Gamma_{h^0}^{\text{SM}}} \cdot \frac{\Gamma_{h^0}^{\text{SM}}}{\Gamma_{h^0}} \simeq [\sigma \cdot \mathcal{B}]_{ZZ^*}^{\text{SM}}, \quad (5)$$

where we assume  $h^0$  is produced dominantly through gluon-gluon fusion. We have separated respective pieces where  $h^0$ -properties may deviate from SM. Both experiments find consistency with the expected 15–20  $\ell\ell'\ell'$  signal events expected from full 2011–2012 data set. However,  $\sigma_{t\bar{t}}$  is of order 220 pb at 8 TeV [25]. If one takes  $\mathcal{B}_{ch} \simeq 2.7\%$ , this amounts to  $\sim 12$  pb into  $t\bar{t} \rightarrow ch^0 bW$ , which should be compared with [28]  $\sim 20$  pb for  $gg$ -fusion production of a 126 GeV SM Higgs boson! An excess

TABLE I. Light Higgs  $h^0$  properties in 2HDM-III with  $\rho_{ct} \sim 1$ . Widths are in MeV units, with  $\Gamma_{h^0}^{\text{SM}} \simeq 4.55$  MeV [29].

	$\mathcal{B}^{\text{SM}}$	$\Gamma^{\text{SM}}$	$\Gamma$	Comment
$WW^*$	21.5%	0.98	hard to change	$\sin(\beta - \alpha) \simeq 1$
$ZZ^*$	2.7%	0.12	hard to change	$\sin(\beta - \alpha) \simeq 1$
$\gamma\gamma$	0.24%	0.011	hard to change	$W$ -loop dom.
$b\bar{b}$	59.4%	2.70	hard to change	$b \rightarrow s\gamma$
$\tau\tau$	5.7%	0.26	within fac. 2	direct
$c\bar{c}$	2.6%	0.12	up to $\sim \Gamma_{b\bar{b}}$	not measured ( $\rho_{cc} \lesssim 0.2$ )
$g\bar{g}$	7.7%	0.35	up to fac. 2	$\rho_{tt} \sim 1$

could have appeared already in  $ZZ^*$  mode, except that each of the three product factors in Eq. (5) could deviate from SM. For example,  $\sigma_{gg \rightarrow h^0}$  may be smaller, or  $\Gamma_{h^0} > \Gamma_{h^0}^{\text{SM}}$  might dilute direct production.

Compared with Ref. [7], Eq. (2) allows us a more complete treatment of  $h^0$ -properties with  $\rho_{ct} \sim 1$ , hence understand what SM-like observation of  $ZZ^*$  may imply. Our study also illustrates how 2HDM-III with FCNH could alter several Higgs properties, driving in the importance of their measurement.

With  $h^0$  dominantly the SM Higgs boson, its  $WW^*$  and  $ZZ^*$  decay rates, proportional to  $\sin^2(\beta - \alpha)$ , are hardly changed. Likewise, the  $h^0 \rightarrow \gamma\gamma$  rate, dominated by  $W$ -loop, is also SM-like. For fermions, the mass generating  $\kappa$  terms are close to SM, while a small  $\cos(\beta - \alpha)$  (we are close to decoupling limit) dilutes the effect of  $\rho$ -type couplings. The consistency of  $h^0 \rightarrow \tau\tau$  with SM constrains  $\rho_{\tau\tau}$  to be small, while  $\rho_{bb}$  is constrained by  $b \rightarrow s\gamma$  to be tiny if  $\rho_{ct} \sim 1$ . Further diluted by  $\cos(\beta - \alpha)$ , the  $b\bar{b}$  rate arises from  $\kappa_{bb}$  and is SM-like.

We are left with potential  $\rho_{cc}$  and  $\rho_{tt}$  effects. The  $c\bar{c}$  mode is extremely hard to search for, hence there are no limits so far. With  $\cos(\beta - \alpha) \sim 0.2$ ,  $\rho_{cc} \sim 0.2$  [30] would bring  $\Gamma_{c\bar{c}} \sim \Gamma_{b\bar{b}} \simeq \Gamma_{b\bar{b}}^{\text{SM}}$ , and the enhanced  $\Gamma_{h^0}$  would dilute the Higgs signal. This can be partially compensated for by  $\rho_{tt}$ , as this parameter should naturally be of order 1 if  $\rho_{ct} \sim 1$ , since  $\kappa_{tt} \simeq 1$  also. With some suppression by  $\cos(\beta - \alpha)$ , nevertheless it could bring  $\sigma_{gg \rightarrow h^0}$  up or down by a factor of  $\sim 2$ .

We summarize in Table I possible effects of our constrained 2HDM-III (with  $\rho_{ct} \sim 1$ ). While  $\Gamma_{h^0 \rightarrow ZZ^*}/\Gamma_{h^0}^{\text{SM}}$  is similar to  $\mathcal{B}_{ZZ^*}^{\text{SM}}$ ,  $\sigma_{gg \rightarrow h^0}$  could change by a factor of 2 and  $\Gamma_{h^0}$  could be enhanced. We comment:

- If  $\rho_{cc}$  is small and  $h^0$  branching ratios are SM-like (except for  $gg$  mode), then the bound of  $\mathcal{B}(t \rightarrow ch^0) < 2.7\%$  from Ref. [7] would apply;
- For enhanced  $\sigma_{gg \rightarrow h^0}$ , then dilution of  $\mathcal{B}_{ZZ^*}$  would be necessary, implying enhanced  $h^0 \rightarrow c\bar{c}$ ;
- If  $\sigma_{gg \rightarrow h^0}$  is suppressed, or  $\mathcal{B}_{ZZ^*}$  is diluted, then more  $ZZ^*$  events may come from  $t\bar{t}$  feeddown!

Given the clean signature of the  $ZZ^*$  or  $\ell\ell'\ell'$  mode, the searches at CMS and ATLAS have been carried out

in an inclusive way, i.e. simply reconstruct four charged leptons without looking into any associated byproducts. The experimental results can thus be used to constrain any other Higgs production process by looking into the extra activities in the events, which already have a clean peak around 126 GeV. There may well be some fraction of  $\ell\ell'\ell' + cbW$  events, with up to 4 jets.

The CMS preliminary result with full 7 and 8 TeV data [31] shows 13, 8, and 4 events with 0, 1, and 2 jets, respectively, after selecting events with  $m_{4\ell} \in (121.5, 130.5)$  GeV. There is no indication for higher associated jet activity. To extract a bound on  $\mathcal{B}(t \rightarrow ch^0)$ , we assume  $\sigma_{gg \rightarrow h^0} \cdot \mathcal{B}(h^0 \rightarrow ZZ^*)$  takes SM value. By inserting the CMS data points, together with the background histograms provided in the same plot [31], and jet multiplicity distribution from top events, an upper limit on the top-Higgs contribution is estimated based on the standard  $\text{CL}_s$  method [32] used at the LHC. The resulting 95% confidence level limit on the relative signal strength between  $t \rightarrow ch^0$  and inclusive Higgs production is around 31%, which can be converted to a limit of 6.5 pb on the effective cross section of  $t \rightarrow ch^0$  at 8 TeV, or a branching ratio limit around 1.5%. This result is based on simple jet counting, with no simulation done.

Interestingly, there is in fact one  $\ell\ell'\ell' + 4j$  event observed [33] by ATLAS for full 7 and 8 TeV data, although no jet-multiplicity plot is given. This event passed the VBF selection, but all 4 jets (in addition to the 4 leptons) are basically in the central rapidity region. As an exercise, we simply add 1 more event to the  $N_{\text{jet}} = 4$  bin, and without changing anything else, we obtain an upper limit of 2.2%, instead of 1.5%. If we add 2 events to the  $N_{\text{jet}} = 4$  bin, the upper limit becomes 2.8%.

It is clear that a genuine analysis is best left to the experiments, as data is already at hand. The ATLAS event reminds one to carefully check whether there is any bias towards lower number of jets, as VBF production is a measurement target. We remark that, except for our simplifying assumption of  $\sigma_{gg \rightarrow h^0} \cdot \mathcal{B}(h^0 \rightarrow ZZ^*)$  being SM-like, this is in fact a model-independent search for  $t \rightarrow ch^0$  ( $h^0 \rightarrow ZZ^*$ ) in  $t\bar{t}$  events.

Our argument can be applied to the other mode,  $\gamma\gamma$ , that drives the Higgs boson discovery. But this mode is not so clean, and clearly carries a bias for VBF event selection of extra jets. However, for  $\gamma\gamma + 4j$  events from  $t\bar{t}$  feeddown, with  $m_{\gamma\gamma}$  in the  $m_{h^0}$  window, the background should be completely different from the case when jet number is no more than 2, and should be rather promising. For the  $h^0 \rightarrow WW^*$  final state, the multi-lepton analysis of Ref. [7] should be redone, while a specific  $\tau\tau + 4j$  analysis can also be pushed. There is one final “steadfast” analysis that one could do, which is searching for  $h^0 \rightarrow b\bar{b}$  mode in  $t\bar{t} \rightarrow ch^0bW \rightarrow cb\bar{b}b + \ell\nu$ . It has

been shown [6] that, through heavy use of  $b$ -tagging and mass reconstruction, one should be able to push down to 1% sensitivity with 2011-2012 data. Here,  $\mathcal{B}(h^0 \rightarrow b\bar{b})$  might get diluted by  $h^0 \rightarrow c\bar{c}$ , which was not considered in Ref. [6], but perhaps the actual experimental analysis could do better than the theoretical study.

It is assuring that, if  $h^0$  behaves SM-like except for inducing  $t \rightarrow ch^0$  decay, we have multiple methods to probe  $\mathcal{B}(t \rightarrow ch^0)$  down to the 1% level. The combined result of the above multi-channel analysis should reach the sub-percent level, which becomes comparable with  $t \rightarrow cZ$  search [34]. If the ATLAS  $4\ell + 4j$  event is any guide, we could even make a discovery.

#### IV. CONCLUSION

It is of great interest to search for the link between the top quark  $t$  and the Higgs boson  $h^0$ . As we have illustrated with  $t\bar{t} \rightarrow ch^0bW \rightarrow 4l + nj$ , it is quite impressive that the intense efforts of Higgs search in the past two years could already push the limit on  $t \rightarrow ch^0$  down to the percent level. Actual experimental studies of  $h^0$  production from  $t\bar{t}$  feeddown, incorporating  $h^0 \rightarrow ZZ^*$ ,  $\gamma\gamma$ ,  $WW^*$ ,  $b\bar{b}$  and  $\tau^+\tau^-$  modes, should be able to push the limit to below the percent level. A discovery of the  $t \rightarrow ch^0$  process with present data would not only imply the existence of an extended Higgs sector, but one beyond the usual 2HDM-II of minimal SUSY.

**Acknowledgement.** KFC is supported by the National Science Council grant NSC 101-2112-M-002-009; WSH is supported by the Academic Summit grant NSC 101-2745-M-002-001-ASP, as well as by grant NTU-EPR-102R8915; CK is supported in part by the U.S. Department of Energy under Grant No. DE-FG02-04ER41305; MK is supported under NTU-ERP-102R7701 and the Laurel program. CK thanks the High Energy Physics group of National Taiwan University for excellent hospitality and support during a sabbatical visit. We are grateful to Yang Bai and Bill Murray, as well as others, for useful discussions.

**Note Added** After submission of this letter, Ref. [35] appeared in arXiv which also addresses the  $tch^0$  coupling at the LHC. This study finds a far better sensitivity reach for the  $tch^0$  coupling compared to our results, but we do not understand how it is achieved. In addition, after this letter was accepted, we learned about the ATLAS search for  $t \rightarrow ch^0$ , with  $h^0 \rightarrow \gamma\gamma$ , in  $t\bar{t}$  events, finding the limit [36] of  $\mathcal{B}(t \rightarrow ch^0) < 0.83\%$  at 95% C.L.

[1] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012).

[2] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012).

- [3] G. Eilam, J.L. Hewett and A. Soni, Phys. Rev. D **44**, 1473 (1991) [Erratum-ibid. D **59**, 039901 (1999)]; B. Mele, S. Petrarca and A. Soddu, Phys. Lett. B **435**, 401 (1998); J.A. Aguilar-Saavedra, Acta Phys. Polon. B **35**, 2695 (2004).
- [4] W.-S. Hou, Phys. Lett. B **296**, 179 (1992).
- [5] J.A. Aguilar-Saavedra and G.C. Branco, Phys. Lett. B **495**, 347 (2000).
- [6] C. Kao, H.-Y. Cheng, W.-S. Hou and J. Sayre, Phys. Lett. B **716**, 225 (2012).
- [7] N. Craig *et al.*, Phys. Rev. D **86**, 075002 (2012). We note that  $t \rightarrow qh$  has recently be discussed by R. Harnik, J. Kopp and J. Zupan, JHEP **1303**, 026 (2013), with Craig *et al.* as the main reference.
- [8] J.P. Lees *et al.* [BaBar Collaboration], Phys. Rev. Lett. **109**, 101802 (2012).
- [9] S. Fajfer, J.F. Kamenik, I. Nisandzic and J. Zupan, Phys. Rev. Lett. **109**, 161801 (2012).
- [10] A. Crivellin, C. Greub and A. Kokulu, Phys. Rev. D **86**, 054014 (2012).
- [11] S.L. Glashow and S. Weinberg, Phys. Rev. D **15**, 1958 (1977).
- [12] T.P. Cheng and M. Sher, Phys. Rev. D **35**, 3484 (1987).
- [13] J.F. Gunion, H.E. Haber, G.L. Kane and S. Dawson, Front. Phys. **80**, 1 (2000); *The Higgs Hunter's Guide* (Addison-Wesley, Redwood City, CA, 1990).
- [14] For simplicity and to reduce the number of parameters, we assume FCNH couplings to be hermitian.
- [15] G.W.-S. Hou, Theory Summary talk of “Flavor Physics and CP Violation” conference, Hefei, China, May 2012, arXiv:1207.7275 [hep-ph].
- [16] F. Mahmoudi and O. Stål, Phys. Rev. D **81**, 035016 (2010).
- [17] J.F. Gunion and H.E. Haber, Phys. Rev. D **67**, 075019 (2003).
- [18] See, e.g. ATLAS-CONF-2012-160 [ATLAS Collaboration].
- [19] If we adopt smaller  $m_{H^+}$ , e.g.  $\sim 300$  GeV, the required  $\rho_{ct}$  can be pulled down to perturbative regime. With small  $m_{H^+}$  and  $\rho_{ct} \sim 1$ ,  $b \rightarrow s\gamma$  data forces  $\rho_{bb}$  and  $\rho_{tt}$  to be significantly smaller than  $\kappa_{bb} = \sqrt{2}m_b/v$  and  $\kappa_{tt} = \sqrt{2}m_t/v$ , barring cancellations. We do not pursue such special possibilities in this paper.
- [20] J.P. Lees *et al.* [BaBar Collaboration], arXiv:1303.0571 [hep-ex].
- [21] A. Crivellin, C. Greub and A. Kokulu, Phys. Rev. D **87**, 094031 (2013).
- [22] M. Ciuchini, G. Degrassi, P. Gambino and G.F. Giudice, Nucl. Phys. B **527**, 21 (1998).
- [23] S. Chatrchyan *et al.* [CMS Collaboration], JHEP **1211**, 067 (2012).
- [24] S. Moch, P. Uwer and A. Vogt, Phys. Lett. B **714**, 48 (2012).
- [25] M. Czakon and A. Mitov, JHEP **1301**, 080 (2013).
- [26] M. Czakon, P. Fiedler and A. Mitov, arXiv:1303.6254 [hep-ph].
- [27] S. Chatrchyan *et al.* [CMS Collaboration], JHEP **1206**, 169 (2012).
- [28] See <https://twiki.cern.ch/twiki/bin/view/LHCPhysics/CERNYellowReportPageAt8TeV>.
- [29] Estimates were made using HDECAY, A. Djouadi, J. Kalinowski and M. Spira, Comput. Phys. Commun. **108**, 56 (1998), which can be contrasted with Ref. [28].
- [30] This is not inconsistent with  $|\rho_{cc}| \lesssim 0.2$  found by Ref. [21] using  $D_s \rightarrow \tau\nu$ ,  $D_{(s)} \rightarrow \mu\nu$  and  $d_n$  (neutron EDM) constraints, which suffer from hadronic uncertainties.
- [31] See <https://twiki.cern.ch/twiki/pub/CMSPublic/Hig13002TWiki/jets.pdf> [CMS Collaboration].
- [32] A.L. Read, In \*Geneva 2000, Confidence limits\* 81-101. A brief description, especially suitable for theorists, can be found in Sec. II.C of J. Alwall, T. Enkhbat, W.-S. Hou and H. Yokoya, Phys. Rev. D **86**, 074029 (2012).
- [33] See Fig. 38 of ATLAS-CONF-2013-013 [ATLAS Collaboration]. We thank Bill Murray for bringing this to our attention.
- [34] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **718**, 1252 (2013).
- [35] D. Atwood, S. K. Gupta and A. Soni, arXiv:1305.2427 [hep-ph].
- [36] ATLAS-CONF-2013-081 [ATLAS Collaboration].