On the equal time two-point distribution of the one-dimensional KPZ equation by replica

T. Imamura ^{*}, T. Sasamoto ^{†‡}, H. Spohn [‡]

February 27, 2024

Abstract. In a recent contribution, Dotsenko establishes a Fredholm determinant formula for the two-point distribution of the KPZ equation in the long time limit and starting from narrow wedge initial conditions. We establish that his expression is identical to the Fredholm determinant resulting from the Airy₂ process.

1 Introduction

The Kardar-Parisi-Zhang (KPZ) equation is a stochastic evolution equation for a height function h(x, t), which models a surface growing through random deposition [1]. It was noted early on that the N-th exponential moment of the height function can be expressed through the propagator of N quantum particles interacting with a short range attractive potential [2]. In one spatial dimension, this would be the attractive Lieb-Liniger Hamiltonian. For its propagator we have now rather concise expressions, which allows one to carry out sophisticated replica computations so to obtain exact distribution functions for the height [3–11]. The various cases are distinguished by the initial conditions (narrow wedge, flat, two-sided Brownian motion in space variable x), by the location of the space-time reference points, and by possibly taking a scaling limit (e.g. $t \to \infty$).

In the recent contribution [12], Dotsenko develops a novel summation formula, which he uses to study narrow wedge initial conditions. In [13] he considers jointly h(0, t), h(0, 2t), while in [12] he considers jointly $h(x_1, t), h(x_2, t)$, both in the scaling limit $t \to \infty$. It is this latter case which is of interest to us. Dotsenko obtains a Fredholm determinant for the joint distribution. Based on universality, Dotsenko's expression should agree with the corresponding Fredholm determinant of the Airy₂ process, but merely a direct inspection does not suffice to confirm the conjecture.

^{*}Research Center for Advanced Science and Technology, The University of Tokyo, Japan, email: imamura@jamology.rcast.u-tokyo.ac.jp

[†]Department of mathematics and informatics, Chiba University, Japan, email: sasamoto@math.s.chiba-u.ac.jp

[‡]Zentrum Mathematik, Technische Universität Munchen, Germany, email: sasamoto@ma.tum.de, spohn@ma.tum.de

In our note we close the gap and prove the desired identity. As a by-product, the factorization assumption in [10] can be understood *a posteriori* as omitting the "cross terms" of Dotsenko's summation formula.

2 One-point distribution

The notations in [10,12] differ from each other. To make our contribution self-contained and to fix the notation, we briefly recall the replica computation for the one-point distribution.

In dimensionless form the KPZ equation reads

$$\frac{\partial}{\partial t}h = \frac{1}{2}\left(\frac{\partial h}{\partial x}\right)^2 + \frac{1}{2}\frac{\partial^2 h}{\partial x^2} + \eta, \qquad (2.1)$$

where h = h(x, t) is the height function at location $x \in \mathbb{R}$ and time $t \ge 0$ and η is normalized Gaussian white noise. Through the Cole-Hopf transformation, $Z = e^{h}$, (2.1) turns into the stochastic heat equation,

$$\frac{\partial}{\partial t}Z = \frac{1}{2}\frac{\partial^2 Z}{\partial x^2} + \eta Z \,. \tag{2.2}$$

We will consider only the narrow wedge initial condition $Z(x, 0) = \delta(x)$.

Let us set

$$h(x,t) = -\frac{1}{24}t + \gamma_t \tilde{h}(x,t) , \qquad (2.3)$$

 $\gamma_t = (t/2)^{1/3}$, and introduce the generating function

$$G_{t,x}(s) = \left\langle \exp\left(-e^{\gamma_t \tilde{h}(x,t)-s}\right) \right\rangle = \left\langle \exp\left(-e^{(t/24)-s}Z(x,t)\right) \right\rangle.$$
(2.4)

Since $\lim_{t\to\infty} \exp(-e^{-\gamma_t x}) = \theta(x)$, θ the step function, $\theta(x) = 1$ for x > 0, $\theta(x) = 0$ for x < 0, it holds

$$\lim_{t \to \infty} G_{t,2\gamma_t^2 x}(\gamma_t(s-x^2)) = \lim_{t \to \infty} \left\langle \theta(s-\tilde{h}(2\gamma_t^2 x,t)-x^2) \right\rangle = \lim_{t \to \infty} \operatorname{Prob}\left[\tilde{h}(2\gamma_t^2 x,t)+x^2 \le s\right].$$
(2.5)

 $G_{t,x}(s)$ is a moment generating function for Z(x,t),

$$G_{t,x}(s) = \sum_{N=0}^{\infty} \frac{(-\mathrm{e}^{-s})^N}{N!} \langle Z(x,t)^N \rangle \mathrm{e}^{tN/24} \,.$$
(2.6)

The N-th moment, $\langle Z(x,t)^N \rangle$, can be written using the eigenvalues and eigenfunctions of the δ -Bose gas with N particles,

$$\langle Z(x,t)^N \rangle = \langle x | e^{-H_N t} | 0 \rangle = \sum_z \langle x | \Psi_z \rangle \langle \Psi_z | 0 \rangle e^{-E_z t}, \qquad (2.7)$$

where $|x\rangle = |x, \ldots, x\rangle$, N-times, H_N is the Hamiltonian of the δ -Bose gas of N particles, and z labels the eigenvalues E_z and eigenstates $|\Psi_z\rangle$. Using the usual parametrization of the eigenstates [3,7], for fixed N one writes

$$E_z = \frac{1}{2} \sum_{j=1}^N z_j^2 = \frac{1}{2} \sum_{\alpha=1}^M n_\alpha q_\alpha^2 - \frac{1}{24} \sum_{\alpha=1}^M (n_\alpha^3 - n_\alpha), \qquad (2.8)$$

$$\langle x|\Psi_z\rangle\langle\Psi_z|0\rangle = |\langle 0|\Psi_z\rangle|^2 \mathrm{e}^{\mathrm{i}\sum_{\alpha=1}^M n_\alpha q_\alpha x}, \qquad (2.9)$$

$$|\langle 0|\Psi_z\rangle|^2 = N! \det\left(\frac{1}{\frac{1}{2}(n_\alpha + n_\beta) + i(q_\alpha - q_\beta)}\right)_{\alpha,\beta=1}^M, \qquad (2.10)$$

$$\sum_{z} = \sum_{M=1}^{N} \frac{1}{M!} \prod_{\alpha=1}^{M} \sum_{n_{\alpha}=1}^{\infty} \int_{-\infty}^{\infty} \frac{dq_{\alpha}}{2\pi} \delta_{\sum_{\alpha=1}^{M} n_{\alpha}, N}.$$
 (2.11)

Hence

$$G_{t,x}(s) = \sum_{M=0}^{\infty} \frac{(-1)^M}{M!} \prod_{\alpha=1}^M \int_{-\infty}^{\infty} \frac{dq_{\alpha}}{2\pi} \sum_{n_{\alpha}=1}^{\infty} (-1)^{n_{\alpha}-1} \mathrm{e}^{n_{\alpha}^3 t/24 + n_{\alpha}(\mathrm{i}q_{\alpha}x - \frac{1}{2}q_{\alpha}^2 t - s)} \times \det\left(\frac{1}{\frac{1}{2}(n_{\alpha} + n_{\beta}) + \mathrm{i}(q_{\alpha} - q_{\beta})}\right)_{\alpha,\beta=1}^M$$
(2.12)

with the understanding that the term with M = 0 equals 1.

Using

$$\frac{1}{\frac{1}{2}(n_{\alpha}+n_{\beta})+\mathrm{i}(q_{\alpha}-q_{\beta})} = \int_{0}^{\infty} d\omega_{\alpha} \mathrm{e}^{-(\frac{1}{2}(n_{\alpha}+n_{\beta})+\mathrm{i}(q_{\alpha}-q_{\beta}))\omega_{\alpha}}$$
(2.13)

and a simple identity for determinants, one has

$$G_{t,x}(s-x^2/(2t)) = \sum_{M=0}^{\infty} \frac{(-1)^M}{M!} \prod_{\alpha=1}^M \int_0^\infty d\omega_\alpha \int_{-\infty}^\infty \frac{dq_\alpha}{2\pi} \sum_{n_\alpha=1}^\infty (-1)^{n_\alpha-1} \\ \times \mathrm{e}^{n_\alpha^3 t/24 + n_\alpha (\mathrm{i}q_\alpha x - \frac{1}{2}q_\alpha^2 t - s + x^2/2t)} \det \left(\mathrm{e}^{-\frac{1}{2}n_\alpha (\omega_\alpha + \omega_\beta) - \mathrm{i}q_\alpha (\omega_\alpha - \omega_\beta)}\right)_{\alpha,\beta=1}^M.$$
(2.14)

We rescale $s \to \gamma_t s, x \to 2\gamma_t^2 x, \omega_\alpha \to \gamma_t \omega_\alpha, q_\alpha \to q_\alpha/\gamma_t$. Shifting $q_\alpha \to q_\alpha + ix$ and noting that $e^{x(\omega_\alpha - \omega_\beta)}$ inserted in the determinant does not change its value, we arrive at

$$G_{t,2\gamma_t^2 x}(\gamma_t(s-x^2)) = \det(1-K_{t,s}), \qquad (2.15)$$

where the kernel of $K_{t,s}$ is given by

$$K_{t,s}(\omega,\omega') = \int_{-\infty}^{\infty} \frac{dq}{2\pi} e^{iq(\omega'-\omega)} \sum_{n=1}^{\infty} (-1)^{n-1} e^{n^3t/24 - \gamma_t n \left(q^2 + \frac{1}{2}(\omega+\omega') - s\right)}.$$
 (2.16)

Using

$$e^{\lambda^3 n^3/3} = \int_{-\infty}^{\infty} dy \operatorname{Ai}(y) e^{\lambda n y}$$
(2.17)

with $\lambda = t^{1/3}/2 = \gamma_t/2^{2/3}$ and shifting $y \to y + q^2 + \frac{1}{2}(\omega + \omega')$, one obtains

$$K_{t,s}(\omega,\omega') = 2^{2/3} \int_{-\infty}^{\infty} \frac{dq}{2\pi} e^{iq(\omega'-\omega)} \int dy \operatorname{Ai} \left(2^{2/3} (y+q^2+\frac{1}{2}(\omega+\omega')) \right) \\ \times \sum_{n=1}^{\infty} (-1)^{n-1} e^{-\gamma_t n(s-y)} .$$
(2.18)

Summing over n and using the identity

$$2^{2/3} \int_{-\infty}^{\infty} \frac{dq}{2\pi} e^{2iqy} \operatorname{Ai}\left(2^{2/3}(q^2+x)\right) = \operatorname{Ai}(x+y) \operatorname{Ai}(x-y), \qquad (2.19)$$

one obtains

$$K_{t,s}(\omega,\omega') = \int_{-\infty}^{\infty} dy \operatorname{Ai}(\omega+y) \operatorname{Ai}(\omega'+y) \frac{\mathrm{e}^{\gamma_t(y-s)}}{1+\mathrm{e}^{\gamma_t(y-s)}}.$$
 (2.20)

In the $t \to \infty$ limit,

$$\frac{\mathrm{e}^{\gamma_t(y-s)}}{1+\mathrm{e}^{\gamma_t(y-s)}} \to \theta(y-s) \tag{2.21}$$

and the kernel of $K_{t,s}$ turns into the kernel for the GUE Tracy-Widom distribution.

3 Two-point distribution and factorization assumption

Next we study the joint distribution of $\tilde{h}(x_1,t), \tilde{h}(x_2,t)$ for $x_2 - x_1 = \mathcal{O}(t^{2/3})$ in the limit $t \to \infty$. Following the strategy of Section 2, we consider the generating function

$$G_{t,x_1,x_2}(s_1,s_2) = \left\langle \exp\left(-e^{\gamma_t \tilde{h}(x_1,t)-s_1} - e^{\gamma_t \tilde{h}(x_2,t)-s_2}\right) \right\rangle = \left\langle \exp\left(-e^{(t/24)-s_1} Z(x_1,t) - e^{(t/24)-s_2} Z(x_2,t)\right) \right\rangle.$$
(3.1)

It holds

$$\lim_{t \to \infty} G_{t,2\gamma_t^2 x_1,2\gamma_t^2 x_2} (\gamma_t(s_1 - x_1^2), \gamma_t(s_2 - x_2^2)) = \lim_{t \to \infty} \operatorname{Prob} \left[\tilde{h}(x_1, t) + x_1^2 \le s_1, \tilde{h}(x_2, t) + x_2^2 \le s_2 \right].$$
(3.2)

By expanding in e^{-s_1} , e^{-s_2} ,

$$G_{t,x_1,x_2}(s_1,s_2) = \sum_{L=0}^{\infty} \sum_{R=0}^{\infty} \frac{(-\mathrm{e}^{-s_1})^L}{L!} \frac{(-\mathrm{e}^{-s_2})^R}{R!} \langle Z(x_1,t)^L Z(x_2,t)^R \rangle \mathrm{e}^{(L+R)t/24} , \qquad (3.3)$$

where the term with L = 0 = R equals 1. Expanding in the eigenstates of the δ -Bose gas one arrives at

$$\left\langle Z(x_1,t)^L Z(x_2,t)^R \right\rangle = \sum_z |\langle 0|\Psi_z \rangle|^2 \mathrm{e}^{-E_z t} \frac{\langle x_1, x_2; L, R|\Psi_z \rangle}{\langle 0|\Psi_z \rangle}, \qquad (3.4)$$

where $|x_1, x_2; L, R\rangle = |x_1, \ldots, x_1, x_2, \ldots, x_2\rangle$ with L times the argument x_1 and R times the argument x_2 .

By Dotsenko's summation formula, the last factor in the summation can be written as

$$\frac{\langle x_1, x_2; L, R | \Psi_z \rangle}{\langle 0 | \Psi_z \rangle} \delta_{\sum_{\alpha=1}^M n_\alpha, L+R} = {\binom{L+R}{L}}^{-1} \sum_{m_\alpha+l_\alpha>0} \prod_{\alpha=1}^M {\binom{n_\alpha}{m_\alpha}} \times e^{ix_1m_\alpha q_\alpha + ix_2l_\alpha q_\alpha - m_\alpha l_\alpha x/2} \delta_{m_\alpha+l_\alpha, n_\alpha} \delta_{\sum_{\alpha=1}^M m_\alpha, L} \delta_{\sum_{\alpha=1}^M l_\alpha, R} \times CT , \qquad (3.5)$$

where $x = x_2 - x_1$ and the summation over m_{α}, l_{α} is for all nonnegative values except for $m_{\alpha} = 0 = l_{\alpha}$. The factor *CT* denotes the cross term which involves a product over all pairs α, β , explicitly given by (*cf.* (26) in [12])

$$CT = (3.6)$$

$$\prod_{\alpha \neq \beta} \frac{\Gamma[1 + \frac{1}{2}(m_{\alpha} + m_{\beta} + l_{\alpha} + l_{\beta}) + i(q_{\alpha} - q_{\beta})]\Gamma[1 + \frac{1}{2}(-m_{\alpha} + m_{\beta} + l_{\alpha} - l_{\beta}) + i(q_{\alpha} - q_{\beta})]}{\Gamma[1 + \frac{1}{2}(-m_{\alpha} + m_{\beta} + l_{\alpha} + l_{\beta}) + i(q_{\alpha} - q_{\beta})]\Gamma[1 + \frac{1}{2}(m_{\alpha} + m_{\beta} + l_{\alpha} - l_{\beta}) + i(q_{\alpha} - q_{\beta})]}.$$

Substituting in (3.3), we obtain

$$G_{t,x_{1},x_{2}}(s_{1},s_{2}) = \sum_{M=0}^{\infty} \frac{(-1)^{M}}{M!} \prod_{\alpha=1}^{M} \int_{-\infty}^{\infty} \frac{dq_{\alpha}}{2\pi} \sum_{m_{\alpha}+l_{\alpha}>0} (-1)^{m_{\alpha}+l_{\alpha}-1} \binom{m_{\alpha}+l_{\alpha}}{m_{\alpha}} \times \det\left(\frac{1}{\frac{1}{2}(m_{\alpha}+l_{\alpha}) + \frac{1}{2}(m_{\beta}+l_{\beta}) + i(q_{\alpha}-q_{\beta})}\right)_{\alpha,\beta=1}^{M} \times e^{(m_{\alpha}+l_{\alpha})^{3}t/24 - \frac{1}{2}(m_{\alpha}+l_{\alpha})q_{\alpha}^{2}t - m_{\alpha}s_{1} - l_{\alpha}s_{2} + ix_{1}m_{\alpha}q_{\alpha} + ix_{2}l_{\alpha}q_{\alpha} - m_{\alpha}l_{\alpha}x/2} \times CT. \quad (3.7)$$

On the other hand, the expression (4.8) in [10] after factorization reads, upon replacing $x \to x/2$ there to match with our notation¹,

$$G_{t,x_1,x_2}^{\#}(s_1,s_2) = \sum_{M=0}^{\infty} \frac{(-1)^M}{M!} \prod_{\alpha=1}^M \int_{-\infty}^{\infty} \frac{dq_\alpha}{2\pi} \sum_{n_\alpha=1}^{\infty} (-1)^{n_\alpha-1} \det\left(\frac{1}{\frac{1}{2}n_\alpha + \frac{1}{2}n_\beta + i(q_\alpha - q_\beta)}\right)_{\alpha,\beta=1}^M \times \prod_{\alpha=1}^M e^{n_\alpha^3 t/24 - \frac{1}{2}n_\alpha q_\alpha^2 t} e^{-\frac{1}{2}x\partial_1\partial_2} \left(e^{ix_1q_\alpha - s_1} + e^{ix_2q_\alpha - s_2}\right)^{n_\alpha},$$
(3.8)

¹Note that the definitions of x in [12] and [10] differ by factor of 2. Also 1/2 in (3.21), (3.23), (3.27) of [10] should rather read 1/4.

where $\partial_i = \partial_{s_i}$, i = 1, 2. Binomially expanding the last factor on the RHS, we note that

$$e^{-\frac{1}{2}x\partial_{1}\partial_{2}}\left(e^{ix_{1}q_{\alpha}-s_{1}}+e^{ix_{2}q_{\alpha}-s_{2}}\right)^{n_{\alpha}}$$
$$=\sum_{m_{\alpha}=0}^{n_{\alpha}}e^{-\frac{1}{2}xm_{\alpha}(n_{\alpha}-m_{\alpha})}\binom{n_{\alpha}}{m_{\alpha}}e^{m_{\alpha}(ix_{1}q_{\alpha}-s_{1})}e^{(n_{\alpha}-m_{\alpha})(ix_{2}q_{\alpha}-s_{2})}.$$
(3.9)

Rewriting the summation over n_{α} in (3.8) by setting $l_{\alpha} = n_{\alpha} - m_{\alpha}$, this expression is the same as (3.7) except for the last factor CT. Hence the factorization assumption in [10] amounts to set the cross term CT = 1.

4 Dotsenko's formula yields the Airy₂ process

In [10] it is established that under factorization the limit $t \to \infty$ of the expression in the last section yields indeed the two-point distribution of the Airy₂ process. On the other hand in [12] Dotsenko argues that, after a suitable analytic continuation, $CT \to 1$ as $t \to \infty$ and then writes a Fredholm determinant expression for the limiting two-point distribution. One expects that Dotsenko's formula should coincide with the two-point function of the Airy₂ process. In this section we establish rather straightforwardly the equivalence between the expression (51) in [12] and the $t \to \infty$ limit of (4.19) in [10].

Substituting $s_i \to \gamma_t s_i, x_i \to 2\gamma_t^2 x_i, i = 1, 2, cf.$ [10] Eq. (2.31), (4.19) in [10] turns into

$$G_{t,2\gamma_t^2 x_1,2\gamma_t^2 x_2}^{\#}(\gamma_t(s_1 - x_1^2), \gamma_t(s_2 - x_2^2)) = \det(1 - N), \qquad (4.1)$$

where the kernel of N is given by

$$N(z_{1}, z_{2}) = \mathbb{1}_{\{z_{1}, z_{2} > 0\}} \int_{-\infty}^{\infty} du e^{-2x\partial_{1}\partial_{2}} e^{-(x_{1}\partial_{1} + x_{2}\partial_{2})(\partial_{z_{1}} - \partial_{z_{2}})} \Phi(\gamma_{t}(u - s_{1}), \gamma_{t}(u - s_{2}))$$

$$\times \operatorname{Ai}(z_{1} + u) \operatorname{Ai}(z_{2} + u).$$
(4.2)

Here

$$\Phi(u,v) = \frac{e^u + e^v}{e^u + e^v + 1} = \sum_{l+m>0} (-1)^{l+m-1} \binom{l+m}{m} e^{mu+lv}.$$
(4.3)

Therefore

$$\lim_{t \to \infty} \Phi(\gamma_t(u - s_1), \gamma_t(u - s_2))$$

= $\theta(u - s_1) + \theta(u - s_2) - \theta(u - s_1)\theta(u - s_2) =: \Phi_{\infty}(s_1, s_2, u)$ (4.4)

and hence

$$\lim_{t \to \infty} \operatorname{Prob}\left[\tilde{h}(2\gamma_t^2 x_1, t) + x_1^2 \le s_1, \tilde{h}(2\gamma_t^2 x_2, t) + x_2^2 \le s_2\right] = \det(1 - N_\infty), \qquad (4.5)$$

where

$$N_{\infty}(z_1, z_2) = \mathbb{1}_{\{z_1, z_2 > 0\}} \int_{-\infty}^{\infty} du e^{-x\partial_1 \partial_2} e^{-(x_1\partial_1 + x_2\partial_2)(\partial_{z_1} - \partial_{z_2})} \Phi_{\infty}(s_1, s_2, u) \operatorname{Ai}(z_1 + u) \operatorname{Ai}(z_2 + u) .$$
(4.6)

The derivative with respect to s_1 , s_2 may seem rather formal, since it is acting on a step function. However, in later sections of [10] there are equivalent expressions for N_{∞} , for which such difficulties do not arise, compare with Section 5.

Turning to Eq. (51) in [12], with the understanding that one should replace $x \to 2^{4/3}x$, $s \to 2^{2/3}s$ to conform with the notations up to now, the probability in (4.5) is written as a Fredholm determinant with operator kernel

$$\begin{aligned} A(\omega_{1},\omega_{2}) \\ &= \int_{0}^{\infty} dy \int_{-\infty}^{\infty} \frac{dq}{2\pi} \operatorname{Ai} \left(y + q^{2} + s_{1} + \omega_{1} + \omega_{2} - \operatorname{i} q x_{1} \right) e^{\operatorname{i} q (\omega_{2} - \omega_{1})} \\ &+ \int_{0}^{\infty} dy \int_{-\infty}^{\infty} \frac{dq}{2\pi} \operatorname{Ai} \left(y + q^{2} + s_{2} + \omega_{1} + \omega_{2} + \operatorname{i} q x_{2} \right) e^{\operatorname{i} q (\omega_{2} - \omega_{1})} \\ &- \int_{-\infty}^{\infty} dy \int_{-\infty}^{\infty} \frac{dq}{2\pi} (2\pi)^{-3/2} \int d\xi_{1} d\xi_{2} d\xi_{3} \operatorname{Ai} \left(y + q^{2} + \omega_{1} + \omega_{2} - \operatorname{i} \xi_{3} \sqrt{x} \right) e^{\operatorname{i} q (\omega_{2} - \omega_{1})} \\ &\times e^{-\frac{1}{2} (\xi_{1} - \operatorname{i} q x_{1} / \sqrt{x})^{2} - \frac{1}{2} (\xi_{2} - \operatorname{i} q x_{2} / \sqrt{x})^{2} - \frac{1}{2} \xi_{3}^{2}} \theta \left(y - s_{1} + \xi_{1} \sqrt{x} \right) \theta \left(y - s_{2} + \xi_{2} \sqrt{x} \right) \end{aligned}$$

$$= \int_{-\infty}^{\infty} \frac{dq}{2\pi} \int dy \operatorname{Ai} \left(y + q^{2} + \omega_{1} + \omega_{2} \right) e^{\operatorname{i} q (\omega_{2} - \omega_{1})} \left\{ \theta \left(y - s_{1} + \operatorname{i} q x_{1} \right) + \theta \left(y - s_{2} + \operatorname{i} q x_{2} \right) \right\}$$

$$- (2\pi)^{-3/2} \int d\xi_{1} d\xi_{2} d\xi_{3} e^{-\frac{1}{2} (\xi_{1}^{2} + \xi_{2}^{2} + \xi_{3}^{2})} e^{-(\xi_{1} + \operatorname{i} \xi_{3}) \sqrt{x} \partial_{1}} e^{-(\xi_{2} + \operatorname{i} \xi_{3}) \sqrt{x} \partial_{2}}$$

$$\times \theta \left(y - s_{1} + \operatorname{i} q x_{1} \right) \theta \left(y - s_{2} + \operatorname{i} q x_{2} \right) \bigg\} . \tag{4.7}$$

Here the step functions with complex arguments are to be understood as resulting from a limit in analogy to (2.21). The last term can be rewritten as

$$(2\pi)^{-3/2} \int d\xi_1 d\xi_2 d\xi_3 e^{-\frac{1}{2}(\xi_1^2 + \xi_2^2 + \xi_3^2)} e^{-(\xi_1 + i\xi_3)\sqrt{x}\partial_1} e^{-(\xi_2 + i\xi_3)\sqrt{x}\partial_2} \theta (y - s_1 + iqx_1) \times \theta (y - s_2 + iqx_2) = e^{-x\partial_1\partial_2} \theta (y - s_1 + iqx_1) \theta (y - s_2 + iqx_2).$$
(4.8)

Proof. It is easy to see that

$$e^{-\frac{1}{2}x\partial_{1}\partial_{2}}e^{-ms_{1}-ls_{2}} = e^{-ms_{1}-ls_{2}-\frac{1}{2}xml}$$

$$= (2\pi)^{-3/2} \int d\xi_{1}d\xi_{2}d\xi_{3}e^{-\frac{1}{2}(\xi_{1}^{2}+\xi_{2}^{2}+\xi_{3}^{2})+m\sqrt{x}\xi_{1}+l\sqrt{x}\xi_{2}+i(m+l)\sqrt{x}\xi_{3}}e^{-ms_{1}-ls_{2}}$$

$$= (2\pi)^{-3/2} \int d\xi_{1}d\xi_{2}d\xi_{3}e^{-\frac{1}{2}(\xi_{1}^{2}+\xi_{2}^{2}+\xi_{3}^{2})}e^{-(\xi_{1}+i\xi_{3})\sqrt{x}\partial_{1}}e^{-(\xi_{2}+i\xi_{3})\sqrt{x}\partial_{2}}e^{-ms_{1}-ls_{2}}.$$
(4.9)

Multiplying both sides by $(-1)^{l+m-1} e^{mu+lv+i(mx_1+lx_2)q} {m+l \choose m}$, substituting $s_i \to \gamma_t s_i, x_i \to \gamma_t s_i$

 $2\gamma_t^2 x_i$, and summing over l, m, we obtain

$$e^{-x\partial_{1}\partial_{2}}\Phi\left(\gamma_{t}(u-s_{1}+ix_{1}q),\gamma_{t}(u-s_{2}+ix_{2}q)\right)$$

= $(2\pi)^{-3/2}\int d\xi_{1}d\xi_{2}d\xi_{3}e^{-\frac{1}{2}(\xi_{1}^{2}+\xi_{2}^{2}+\xi_{3}^{2})}$
 $\times e^{-(\xi_{1}+i\xi_{3})\sqrt{x}\partial_{1}}e^{-(\xi_{2}+i\xi_{3})\sqrt{x}\partial_{2}}\Phi\left(\gamma_{t}(u-s_{1}+ix_{1}q),\gamma_{t}(u-s_{2}+ix_{2}q)\right).$ (4.10)

Taking the limit $t \to \infty$, we arrive at the desired equality.

Using Eq. (4.8) one writes

$$A(\omega_{1},\omega_{2}) = \int_{-\infty}^{\infty} \frac{dq}{2\pi} \int dy \operatorname{Ai}(y+q^{2}+\omega_{1}+\omega_{2}) e^{\mathrm{i}q(\omega_{2}-\omega_{1})} \\ \times e^{-x\partial_{1}\partial_{2}} \Phi_{\infty}(s_{1}-\mathrm{i}x_{1}q,s_{2}-\mathrm{i}x_{2}q,y) \\ = 2 \int_{-\infty}^{\infty} \frac{dq}{2\pi} e^{\mathrm{i}q2^{1/3}\left((\omega_{2}-\omega_{1})-(x_{1}\partial_{1}+x_{2}\partial_{2})\right)} \int dy \operatorname{Ai}(2^{2/3}(y+q^{2}+2^{-2/3}(\omega_{1}+\omega_{2})) \\ \times e^{-x\partial_{1}\partial_{2}} \Phi_{\infty}(s_{1},s_{2},2^{2/3}y).$$

$$(4.11)$$

We switch to x, s as in (4.6) and use (2.19) to obtain

$$A(\omega_1, \omega_2) = 2^{1/3} \int dy \operatorname{Ai}(y + 2^{1/3}\omega_1 - (x_1\partial_1 + x_2\partial_2)) \operatorname{Ai}(y + 2^{1/3}\omega_2 + (x_1\partial_1 + x_2\partial_2)) \times e^{-x\partial_1\partial_2} \Phi_{\infty}(s_1, s_2, y).$$
(4.12)

Since Φ_{∞} can be written as a limit of a sum of exponentials, see (4.3),(4.4), the derivatives ∂_i , i = 1, 2, inside the Airy functions can be regarded as numbers. Finally one can rewrite (4.12) as

$$A(\omega_{1}, \omega_{2}) = 2^{1/3} \int dy e^{-x\partial_{1}\partial_{2}} e^{-2^{-1/3}(x_{1}\partial_{1}+x_{2}\partial_{2})(\partial_{\omega_{1}}-\partial_{\omega_{2}})} \\ \times \operatorname{Ai}(y+2^{1/3}\omega_{1})\operatorname{Ai}(y+2^{1/3}\omega_{2})\Phi_{\infty}(s_{1},s_{2},y), \qquad (4.13)$$

which is the same as the kernel N_{∞} of (4.6) with the identification $z_i = 2^{1/3} \omega_i$, i = 1, 2.

We draw one conclusion from our computation. Both in [12] and [10] one has to linearize the factor e^{-xml} , see (4.9). [12] uses a threefold Gaussian integration, whereas [10] uses the differential operator of (4.9). If already in [12] Eq. (34) one would use the latter method, then N_{∞} would appear without further efforts.

5 A simplified proof for the Fredholm determinant of N to determine the Airy₂ process

In [10] it is proved that the Fredholm determinant of N equals the one of the two-point distribution of the Airy₂ process as $t \to \infty$. The arguments of [10] are somewhat involved. Here we provide a more straightforward proof.

We start from an expression $G_{t,x_1,x_2}^{\#}(s_1 - x_1^2/2t, s_2 - x_2^2/2t) = \det(1 - \tilde{N})$, which follows from multiplying out the last factor of (3.8), with the kernel of \tilde{N} given by

$$\tilde{N}(z,z') = \mathbb{1}_{\{z,z'>0\}} \sum_{n=0}^{\infty} \sum_{k=0}^{n} (-1)^n \binom{n}{k} e^{-(s_1 - x_1^2/2t)k - (s_2 - x_2^2/2t)(n-k)} \\ \times e^{-\frac{1}{2}xk(n-k)} \int du \operatorname{Ai}(u+z+y) \operatorname{Ai}(u+z'-y) e^{\gamma_t u n}$$
(5.1)

with $y = (x_1k + x_2(n-k))/2\gamma_t$. Let us rewrite the integral over u as

$$\int du dv \delta(u-v) e^{\gamma_t u n} \operatorname{Ai}(u+z+y) \operatorname{Ai}(v+z'-y)$$

=
$$\int du dv \delta(u-v) e^{\gamma_t n w} e^{\gamma_t u (n-k)} \operatorname{Ai}(u+z+(n-k)x/2\gamma_t) e^{\gamma_t v k} \operatorname{Ai}(v+z'+kx/2\gamma_t), (5.2)$$

where the integrals over u, v have both been shifted by $w = (-x_1k + x_2(n-k))/2\gamma_t$. We notice the identity

$$e^{au}\operatorname{Ai}(u+\lambda) = e^{-\tau(\lambda+a^2)}e^{a\tau^2}e^{-\tau H}e^{au}\operatorname{Ai}(v+\lambda-2\tau a)$$
(5.3)

with the Airy operator $H = -(\partial_u)^2 + u$, see Appendix B of [10].

Proof. Since

$$(H + \lambda)\operatorname{Ai}(u + \lambda) = 0, \qquad (5.4)$$

one arrives at

$$e^{au} \operatorname{Ai}(u+\lambda) = e^{-\tau(H+\lambda)} e^{\tau(H+\lambda)} e^{au} e^{-\tau(H+\lambda)} \operatorname{Ai}(u+\lambda).$$
(5.5)

Using

$$u(\tau) = e^{\tau H} u e^{-\tau H} = u + \tau^2 - 2\tau \partial_u,$$

$$e^{au\tau} = e^{a\tau^2} e^{au} e^{-2a\tau \partial_u} e^{-\frac{1}{2}[au, -2a\tau \partial_u]} = e^{a\tau^2 - a^2\tau} e^{au} e^{-2a\tau \partial_u},$$
(5.6)

(5.3) follows from

$$e^{au}\operatorname{Ai}(u+\lambda) = e^{-\tau(\lambda+a^2)}e^{a\tau^2}e^{-\tau H}e^{au}e^{-2\tau a\partial_u}\operatorname{Ai}(u+\lambda).$$
(5.7)

With the help of identity (5.3) the kernel of \tilde{N} can be rewritten as

$$\tilde{N}(z,z') = \mathbb{1}_{\{z,z'>0\}} \sum_{n=0}^{\infty} \sum_{k=0}^{n} (-1)^{n-1} \binom{n}{k} e^{-s_1 k - s_2 (n-k)} e^{(x_1 z' - x_2 z)/2\gamma_t^2} \\ \times \int du dv \langle u | e^{-xH/2\gamma_t^2} | v \rangle e^{\gamma_t (n-k)u + \gamma_t k v} \operatorname{Ai}(u+z) \operatorname{Ai}(v+z') .$$
(5.8)

Using the cyclicity of the determinant, $\det(1 - AB) = \det(1 - BA)$, it holds that $\det(1 - \tilde{N}) = \det(1 - \tilde{L})$ with the kernel of \tilde{L} given by

$$\tilde{L}(u,v) = \sum_{n=0}^{\infty} \sum_{k=0}^{n} (-1)^{n-1} \binom{n}{k} e^{-s_1 k - s_2 (n-k)} \langle u | e^{-xH/2\gamma_t^2} | v \rangle$$
$$\times e^{\gamma_t (n-k)u + \gamma_t k v} \int_0^\infty \operatorname{Ai}(u+z) \operatorname{Ai}(v+z) e^{-xz/2\gamma_t^2} dz \,.$$
(5.9)

From this expression it is easy to see that the Fredholm determinant of \hat{L} yields the one of the Airy₂ process for $t \to \infty$, see the arguments in Section 2.4 of [10].

Acknowledgement. We are grateful to Victor Dotsenko for useful comments and for communicating his preprint [12].

References

- M. Kardar, G. Parisi and Y. C. Zhang. Dynamic scaling of growing interfaces. *Phys. Rev. Lett.*, 56:889–892, 1986.
- [2] M. Kardar. Replica Bethe ansatz studies of two-dimensional interfaces with quenched random impurities. Nucl. Phys. B., 290:582–602, 1987.
- [3] V. Dotsenko. Bethe ansatz derivation of the Tracy-Widom distribution for onedimensional directed polymers. *Euro. Phys. Lett.*, 90:200003, 2010.
- [4] P. Calabrese, P. Le Doussal, A. Rosso. Free-energy distribution of the directed polymer at high temperature. *Euro. Phys. Lett.*, 90:200002, 2010.
- [5] P. Calabrese, P. Le Doussal. An exact solution for the KPZ equation with flat initial conditions. *Phys. Rev. Lett.*, 106:250603, 2011.
- [6] P. Calabrese, P. Le Doussal. The KPZ equation with flat initial condition and the directed polymer with one free end. J. Stat. Mech., page P06001, 2012.
- [7] T. Imamura and T. Sasamoto. Replica approach to the KPZ equation with half Brownian motion initial condition. J.Phys. A, 44:385001, 2011.
- [8] T. Imamura and T. Sasamoto. Exact solution for the stationary KPZ equation. *Phys. Rev. Lett.*, 108:190603, 2012.
- [9] T. Imamura and T. Sasamoto. Stationary correlations for the 1D KPZ equation. J. Stat. Phys., 150:908–939, 2013.
- [10] S. Prolhac and H. Spohn. Two-point generating function of the free energy for a directed polymer in a random medium. J. Stat. Mech., page P01031, 2011.

- [11] S. Prolhac and H. Spohn. The one-dimensional KPZ equation and the Airy process. J. Stat. Mech., page P03020, 2011.
- [12] V. Dotsenko. Two-point free energy distribution function in (1+1) directed polymers, arXiv:1304.6571.
- [13] V. Dotsenko. Two-time free energy distribution function in (1+1) directed polymers, arXiv:1304.0626.